In vitro processing of glutamyl endopeptidase proenzymes from *Enterococcus faecalis* and importance of N-terminal residue in enzyme catalysis

Shakh M. A. Rouf^{1,2,*}, Y. Ohara-Nemoto¹, T. Ono¹, Y. Shimoyama³, S. Kimura³, T. K. Nemoto¹

¹Department of Oral Molecular Biology, Courses of Medical and Dental Science, Nagasaki University Graduate School of Biomedical Sciences, Nagasaki 852-8588, Japan

²Department of Applied Nutrition & Food Technology, Islamic University, Kushtia-7003, Bangladesh

³Division of Molecular Microbiology, Iwate Medical University, Yahaba-cho 028-3694, Japan

Email address:

abdurrouf_7@yahoo.com (S. M. A. Rouf)

To cite this article:

Shakh M. A. Rouf, Y. Ohara-Nemoto, T. Ono, Y. Shimoyama, S. Kimura, T. K. Nemoto. In Vitro Processing of Glutamyl Endopeptidase Proenzymes from *Enterococcus faecalis* and Importance of N-terminal Residue in Enzyme Catalysis. *Advances in Biochemistry*. Vol. 1, No. 5, 2013, pp. 73-80. doi: 10.11648/j.ab.20130105.11

Abstract: Glutamyl endopeptidase from *Enterococcus faecalis*, designated SprE, is one of the important virulence factors secreted as zymogen. In the present study we expressed recombinant SprE proenzyme (pro-SprE) in *Escherichia coli* and investigated the *in vitro* processing to mature SprE. It was found that trypsin could efficiently produce the active form of SprE with the N-terminus Ser¹ through cleavage between Arg⁻¹ and Ser¹ bond, which was subsequently auto-degraded into inactive species through the cleavage at the Glu⁶-Asp⁷ and Glu¹¹-Val¹² bonds. Although thermolysin could produce SprE with the N-terminus Leu², but possessed no proteolytic activity. In contrast to the absolute requirement of the N-terminal Val¹ in *staphylococcal* glutamyl endopeptidases, the N-terminal Ser¹ of mature SprE could be substituted by other amino acids despite that Ser showed the maximal activity. Substitution of penultimate Leu² of SprE to Val² also reduced the activity to 40% of the wild type. Taken together, we conclude that pro-SprE was converted to mature form with the N-terminus Ser¹ by a protease with specificity of trypsin and the length of the N-terminal region rather than specific residue is absolutely required for enzyme activity.

Keywords: Glutamyl Endopeptidase, Proenzyme Processing, SprE, *Enterococcus faecalis*, GluV8

1. Introduction

Enterococcus faecalis is a gram-positive cocci and common inhabitant of human gastrointestinal and genitourinary tract [1]. It causes a wide variety of diseases in human, infecting the urinary tract, endocardium, abdomen, biliary tract, burn wounds and indwelling foreign devices [2]. In dentistry, *E. faecalis* has been associated with caries lesions, chronic periodontitis, recurrent root canal infection and persistent apical periodontitis [3, 4]. Candidate virulence factors of *E. faecalis* include: cytolysin and proteolytic enzyme (gelatinase and serine protease), adhesions (aggregation substance, enterococci surface protein or Esp, collagen adhesion protein or Ace, antigen A or EfaA) and capsular and cellular wall polysaccharides [2, 5]. Serine protease, gelatinase and collagen-binding protein (Ace) facilitate *E. faecalis* bind to dentin [6]. Several

studies have been undertaken to investigate the possible role of gelatinase and serine proteases in the disease processes caused by *E. faecalis* and they were found to contribute to the pathogenesis through biofilm formation, facilitating bacterial invasions, degradation of immune peptides required for host response [7, 8, 9,10]. Gelatinase (GelE) and serine protease (SprE) expressed in *E. faecalis* were reported to be important for enterococcal virulence in mouse peritonitis model [11]. Previous studies also indicated that the presence of extracellular proteases, GelE and SprE, in the *E. faecalis* raised mortality in animal models [12, 13].

Glutamyl endopeptidase (EC 3.4.21.19) from *Staphylococcus aureus* V8 strain (GluV8) is a serine protease with unique substrate specificity to Glu-X and Asp-X with the higher preference to the former [14,15]. This family protease from *E. faecalis* SprE, which has been shown to contribute to pathogenesis in animal models [16, 11, 17]. Kawalec et al. [18] purified several isoforms of

mature SprE, starting with N-terminal Ser¹ and Leu² from wild-type *E. faecalis* OG1RF (TX4002) and a gelatinase-null mutant TX5264, among which SprE starting with Ser¹ was super active. Moreover, they suspected the instability of active SprE (Ser¹-SprE form) apparently due to auto-degradation.

SprE shows sequence homology with GluV8 (27% identity, 49% similarity) [19] and that from *S. epidermidis* (GluSE) (26% identity, 49% similarity) [20]. It has been reported that the proenzyme of GluV8 is processed by a thermolysin-family metalloprotease, aureolysin *in vivo* [21,22]. The prosequence of *Staphylococcal* GluV8-family protease was reported to sequentially remove to shorter proenzymes in an autocatalytic manner by cleaving at Glu-X and Gln-X in the prosegment and finally matured by aureolysin at N-terminal Asn¹-Val¹ [23, 24]. Consistently recombinant GluV8 expressed in *E. coli* was efficiently processed to mature GluV8 by thermolysin *in vitro* [25]. Val¹ is conserved at the N-termini of mature glutamyl endopeptidases from *Staphylococcus epidermidis* [26, 27], *Staphylococcal warneri* [28], *Staphylococcal cohnii* and *Staphylococcal caprae* [29].

The N-terminus amino acid residue of mature GluV8-family endopeptidases was found to be critical for maturation by aureolysin and for the stabilization of mature structure resistant to further processing at the Val¹-Ile² and Ile²-Leu³ bonds [25]. Furthermore, Val¹ is directly involved in the protease activity itself, because Val¹ cannot be substituted by other amino acids, even if the processing was correctly mediated with Arg¹-X¹-substituted forms by trypsin (X=Ala, Phe, Gly and Ser) [25]. Exceptionally, the Val¹ substitution to Leu¹ could partially (30%) retained proteolytic activity of the wild type, further indicating the significance of Val¹ for the protease activity [29]. Sequence alignment indicates that Val¹ of GluV8 corresponds to Leu¹ of SprE, whereas it has been reported that SprE with N-terminal Leu² (Leu²-SprE) had a significantly negligible activity compared to that with N-terminal Ser¹ [18]. Moreover, the processing mechanism of pro-SprE still remains unknown. Therefore, we here investigated the *in vitro* processing of pro-SprE expressed in *E. coli* and addressed the roles of N-terminal amino acids in the enzyme catalysis.

2. Materials and Methods

2.1. Materials

The materials used and their sources were as follow: expression vector pQE60, from Qiagen Inc. (Chatsworth, CA, USA); low-molecular-weight markers, from GE Healthcare (Buckinghamshire, England); restriction enzymes and DNA-modifying enzymes, from Nippon Gene (Tokyo, Japan); KOD Plus DNA polymerase, from Toyobo (Tokyo, Japan); Talon metal-affinity resin, from Clontech Laboratories Inc. (Palo Alto, CA, USA); Z-Leu-Leu-Glu-MCA (LLE-MCA),

Ac-Thr-Val-Ala-Asp-MCA (TVAD-MCA), Z-Ala-Ala-Asn-MCA (AAN-MCA) and Boc-Gln-Ala-Arg-MCA (QAR-MCA) from the Peptide Institute Inc. (Osaka, Japan); Leu-Asp-MCA (LD-MCA) and Z-Leu-Leu-Gln-MCA (LLQ-MCA), synthesized by Thermo Fisher Scientific (Ulm, Germany); thermolysin from *Bacillus thermoproteolyticus* rokko, trypsin porcine pancreas, bovine serum albumin (BSA), were from Sigma-Aldrich (St. Louis, MO, USA); and gelatin from the Nacalai Tesque Inc. (Osaka, Japan).

2.2. Expression Vector for the Expression of SprE

The SprE gene was amplified by PCR using KOD Plus DNA polymerase and chromosomal DNA (0.1 µg) of *E. faecalis* NCTC 775 as template. The synthetic oligonucleotides primers (5'-ATGGATCCAAAAAGTTCTCCATACGAAAAATTA G-3') and (5'-GTGGATCCCGCTGCAGGCACAGCGGATAAACG-3') containing *Bam*HI sites (underlined) were designed on the basis of the DNA sequence of SprE (GeneBank accession no. Z12296), used for SprE gene amplification. PCR-amplified 0.8-kb fragment (without stop codon) was cleaved with *Bam*HI, and then inserted into the *Bam*HI site of pQE60 to yield pQE60-SprE expression vector.

2.3. Expression and Purification of Recombinant Proteases

In order to minimize the modification in the N-terminal preprosequence of SprE, the expression vector pQE60 that encoded an affinity tag, [Gly-Ser-Arg-Ser-(His)₆] at the C-terminus was used (Fig. 1A). In addition Gly-Gly-Ser derived from the vector was present between 1st-Met and 2nd-Lys of the N-terminal prepropeptide. *Escherichia coli* XL1-blue was transformed with the pQE60-SprE expression plasmids. C-terminal His₆-tagged recombinant proteins were expressed and purified as described previously [25]. Protein concentrations were determined by the bicinchoninic acid method using BSA as the standard (Pierce, Rockford, IL, USA). Purified proteins were stored at -80 °C until used.

2.4. Amino Acid Numbering and In Vitro Mutagenesis

The N-terminal Ser of the active SprE was numbered as the first amino acid residue (Ser¹). *In vitro* mutagenesis was performed as reported previously [30] by PCR with mutated primer(s) to substitute 3 amino acids in the prosequence (Glu⁻¹⁵Ser, Glu⁻¹⁴Lys, Glu⁻⁸Ile, designated as SprE-mut), 4 amino acids in the mature region (Glu¹¹Gln, Glu⁶Gln, Ser¹Thr/Ala/Val and Leu²Val), and an essential Ser¹⁸⁰ to Ala. All mutations were confirmed by DNA sequencing.

2.5. SDS-PAGE and Zymography

Recombinant proteins (1 µg) were separated by SDS-PAGE at a polyacrylamide concentration of 12.5% (w/v), and then stained with Coomassie Brilliant Blue

(CBB). Band intensities were quantified with *Image J* software [31]. Zymography was performed by using a polyacrylamide gel containing 1mg/ml of gelatin (instead of azocasein) according to the method previously described [32]. For zymography, thermolysin- or trypsin-treated samples (0.25µg) were separated on a 12.5% polyacrylamide gel containing 1% (w/v) of gelatin by SDS-PAGE, the gel was incubated twice with 100 ml of 2.5% (v/v) of Triton X100 at 25 °C for 20 min, twice for 10 min each time with 100 ml of 50 mMTris-HCl (pH 7.8) containing 30 mMNaCl, and then incubated overnight at 37 °C in 100 ml of the new batch of the same buffer containing 0.001% NaN₃. Finally, non-hydrolyzed gelatin in the gel was visualized by CBB staining.

2.6. In Vitro Processing of Recombinant Proteases and the Measurement of the Protease Activity

Recombinant proteins (10 µg) were incubated for 2 h in 0.1 ml of buffer A [10 mM sodium borate (pH 8.0) containing 0.005% (v/v) Triton X100, and 2 mM CaSO₄] without or with 0.3 µg of thermolysin (molar ratio of 33:1) and 0.003µg of trypsin (molar ratio 3300:1) at 37 °C, unless otherwise stated. Thereafter, proteins (0.25 µg as proform) were incubated at 37 °C for 1 h with 20µM MCA peptides in 0.2 ml of 50 mMTris-HCl (pH 8.0) containing 5 mM EDTA. The fluorescence was measured with excitation at 380 nm and emission at 460 nm with a Fluorescence Photometer F-4000 (Hitachi, Tokyo, Japan).

2.7. N-terminal Amino Acid Sequencing

N-terminal amino acid sequences of recombinant SprE and its derivatives were determined after separation by SDS-PAGE (2µg) and transference to a polyvinylidenedifluoride membrane (Sequi-Blot PVDF Membrane, Bio-Rad, Hercules, CA, USA). After having been stained with CBB, the bands were excised; and directly sequenced with a Procise 49XcLC protein sequencer (ABI, Foster City, CA, USA).

3. Results

3.1. Expression and In Vitro Processing of SprE

When the full-length pro-SprE was expressed in *E. coli*, a mixture of 34- and 35-kDa proteins was recovered by affinity purification (Fig. 1a, upper panel, lane 1). Amino acid sequencing revealed that the 35-kDa band started at Glu⁻¹⁴ and 34-kDa one at Ser⁻⁷, indicating the autocatalytic prosequence cleavage between the Glu⁻¹⁵-Glu⁻¹⁴ and Glu⁻⁸-Ser⁻⁷ bonds, respectively (Table 1). The 34-35-kDa SprE showed negligible glutamic acid-specific peptidase activity, as recombinant SprE still carried part of the propeptide (Fig. 1b,c upper; d). *Staphylococcal* GluV8-family proteases were reported to be processed into mature form by thermolysin cleaving at Asn⁻¹-Val¹ in vitro [29]. To investigate heterocatalytic maturation of pro-SprE, we incubated 34-35-kDa recombinant pro-SprE with trypsin

or thermolysin, expected that may induce processing at Arg⁻¹-Ser¹ and at Ser¹-Leu² sites respectively. Consequently, trypsin treatment accompanying a slight decrease in the apparent molecular mass induced an acquisition of the gelatinase activity (Fig. 1c, lanes 3-10). In contrast, thermolysin treatment did not accompany an apparent increased gelatinase activity, although the apparent molecular mass was decreased. Very faint gelatinase activities were found for thermolysin treated or untreated samples (Fig. 1b lower). In consistent to these findings, the Glu-specific peptidase activity was measured for the trypsin treated SprE in a dose-dependent manner (Fig. 1d).

Fig 1. A schematic presentation of pro-SprE expressed in *E. coli*. (a) Amino acid sequences of Pre-pro and N-terminal regions of mature SprE. The open (V) and closed (▼) arrow-head showing the trypsin (Tryp) and thermolysin (Th) processing site respectively, arrow(↓) indicates the site of autodegradation in the pro- and mature region. (b,c) In vitro processing of the SprE proenzymes and Z-LLE-MCA cleavage activity of the mature SprE. Pro-SprE (10µg) were incubated for 2 h at 0°C (lane1) or at 37°C (lane2) without Thermolysin (Th)/Trypsin (Tryp) and with Th-0.001µg/Tryp- 0.0001µg (lane3), Th- 0.003/ Tryp- 0.0003µg (lane4), Th-0.01µg/Tryp- 0.001 (lane5), Th- 0.03µg/Tryp- 0.003µg (lane6), Th- 0.1µg/ Tryp-0.01µg (lane7), Th- 0.3µg/ Tryp- 0.03µg (lane8), Th- 1.0µg/ Tryp-0.1µg (lane9), and Th- 3.0µg/ Tryp- 0.3µg (lane10). Aliquots (1µg or 0.25µg) of each Thermolysin/ Trypsin treated samples were subjected to SDS-PAGE and then stained with CBB (b,c upper) or subjected to collagen-Zymography (b,c lower) respectively. (d) The proteolytic activities (means ± SD; n=3) towards Z-LLE-MCA of the thermolysin (Th) and trypsin (Tryp) treated samples were determined as described in the materials and methods.

N-terminal sequencing of thermolysin-processed 31-kDa species revealed a cleavage at the Ser¹-Leu² bond, whereas the N-terminus of trypsin-processed species was a mixture of SprE with N-terminus Val¹² (major) and Ser¹ (minor) (Table 1). Since the Glu¹¹-Val¹² bond should not be processed by trypsin, but by glutamyl endopeptidase itself, it was reasonably suggested that mature SprE with N-terminal

Ser¹ (Ser¹-SprE) produced by trypsin readily auto-degraded to the Val¹²-SprE. However, the possibility that Val¹²-SprE represents a genuine mature species with the peptidase activity was not completely eliminated, if considered that N-terminal Val¹ was commonly observed in *Staphylococcal* mature glutamyl endopeptidases [29]. To address these possibilities, we tried to increase recovery of Ser¹-SprE quantitatively after trypsin processing by introducing amino acid substitutions in the N-terminal region of the mature SprE as in the following section.

3.2. Suppression of Auto-Degradation of Mature SprE

The autodegradation of *Staphylococcal* glutamyl endopeptidases occurring within the prosequence region was efficiently suppressed by the substitution of Glu and Asp in the prosequences, to Gln, Asn or other amino acids [25, 30]. Here, this strategy was introduced at the N-terminal region of mature SprE.

When Glu¹¹ of SprE was substituted to Gln (SprE-Glu11Gln), the 31-kDa mature SprE band was more intensively recovered after trypsin treatment of the purified proenzyme (Fig-2a upper lane6), which accompanied by an increased peptidase activity in the fraction (Fig-2a lower).

Fig 2. Prevention of auto-degradation and recovery enhancement of mature SprE. Proenzymes (10 µg) of SprE wild-type (wt), single and double amino acid substitution mutants Glu6Gln and Glu6Gln/Glu11Gln respectively were incubated for 2 h at 37°C without thermolysin (Th)/ trypsin (Tryp) or with Th-0.3 µg (Th)/ Tryp- 0.003 µg (Tryp) as described in the materials and methods section. (a) Proteins (1 µg) were separated by SDS-PAGE, stained with CBB and proteolytic activities (mean ± SD; n=3) of the samples were determined with LLE-MCA as per materials and methods section. (b) The band intensities of the trypsin treated mature SprE wild-type (wt) (□), Glu6Gln (▼) and Glu6Gln/Glu11Gln (●) (1 µg) were determined by using image J software [31]. Band intensities of pro-SprE set as 100%.

Because there was Glu⁶ at position nearer to the N-terminus, this residue was also substituted by Gln (SprE-Glu6Gln/Glu11Gln). As a result, the recovery was further increased after trypsin treatment (Fig-2a upper lane9). It is noticeable that the specific activity of the mutant form was not varied (Fig-2b), whereas the recovery after the processing was increased by suppression of the auto-degradation. Even when thermolysin treatment was conducted on these mutants, no increase in recovery (Fig-2a upper lane 5 & 8) as well as in the activity (Fig-2a lower panel) was observed again. The N-terminus of trypsin-processed SprE-Glu11Gln was a mixture of Ser¹ and Asp⁷ and that of SprE-Glu6Gln/Glu11Gln was Ser¹. Taken together with the activity measurement, it was concluded that Ser¹-SprE was the active and mature form and that Val¹²-SprE as well as Asp⁷-SprE was an inactive auto-degraded product (Table 1).

3.3. Substrate Specificity of Pro- and Mature SprE

Glutamyl endopeptidase cleaved peptide bond between Glu-X with highest efficiency and much less efficiently between Asp-X [15]. We investigated the substrate specificity of pro- and mature SprE with several peptidyl-MCA substrates to know whether any possibility of propeptide processing in an autocatalytic manner. Mature Ser¹-SprE specifically hydrolyzed LLE-MCA, and did not hydrolyze TVAD-, LD-, AAN-MCA as well as GAR-MCA, whereas it possessed very faint activity for LLQ-MCA after 1h incubation (Fig-3).

Fig 3. Substrate specificity of SprE. Substrate specificity of pro- and mature SprE was determined with different MCA-peptides as shown in the figure below. Proteolytic activity towards Z-LLE-MCA of trypsin processed mature SprEwt set as positive control and considered 100%. Pro-SprE (10 µg) was incubated for 2h at 37°C without or with 0.003 µg trypsin (molar ratio 3300:1), there after 0.25 µg of each sample was used for different MCA-peptides proteolytic assay as discussed in the materials and methods.

Table 1. N-terminal amino acid sequences of recombinant SprE and its derivatives

SprE derivatives	Thermolysin (Th)/Trypsin (Tryp)	Cleavage site (X-/X)
pro-SprE	- ^a	E ⁻¹⁵ /E ⁻¹⁴ YIVPAE ⁻⁸ /S ⁻⁷ RQKRSLLDPED
pro-SprEmut	-	SHSQ ⁻³ /K ⁻² RSLLDPEDRRQ
	Th	SHSQKRS ¹ /L ² LDPEDRRQ
SprEwt	Tryp ^b	SHSQKR ⁻¹ /S ¹ LLDPEDRRQ
		RRQE ¹¹ /V ¹² ADTTEA
SprE-Glu11Gln	Th	KRS ¹ /L ² LDPEDRRQ
	Tryp ^c	SHSQKR ⁻¹ /S ¹ LLDPEDRRQ
		KRSLLDPE ⁶ /D ⁷ RRQQVADT
SprE-Glu6Gln/Glu11Gln	Tryp	SHSQKR ⁻¹ /S ¹ LLDPQDRRQQ
SprE-G6Gln/Glu11Gln/Ser1Thr	Tryp	SHSQKR ⁻¹ /T ¹ LLDPQDRRQQ
SprE-G6Gln/Glu11Gln/Ser1Ala	Tryp	SHSQKR ⁻¹ /A ¹ LLDPQDRRQQ
SprE-G6Gln/Glu11Gln/Ser1Val	Tryp	SHSQKR ⁻¹ /V ¹ LLDPQDRRQQ
SprE-Leu2Val	Th	SHSQKRS ¹ /V ² LDPEDRRQ
	Tryp ^b	SHSQKR ⁻¹ /S ¹ VLDPEDRRQ
		RRQE ¹¹ /V ¹² ADTTEA

^{abc} A mixture of two polypeptides; Detected N-terminal amino acids of the SprE species were underlined.

The substrate specificity of SprE strongly suggested that pro-SprE could not be processed to active SprE as either pro- or mature SprE unable to cleave GAR-MCA. More interestingly, pro-SprE possessed a faint activity for LLE- and LLQ-MCA. These results strongly suggested that, the propeptide shortening via cleavages at the Glu⁻¹⁵-Glu⁻¹⁴ or Glu⁻⁸-Ser⁻⁷ bond observed in the purified pro-SprE (Fig-1 a,b lane 1,2) was mediated by pro-SprE, but not by mature SprE and finally maturation achieved by cleaving between Arg⁻¹-Ser¹ in a hetero catalytic manner.

3.4. Significance of Ser¹ and Leu² in the Catalytic Processes

As Val¹ is highly conserved in the mature form of all *Staphylococcal* glutamyl endopeptidases and is indispensable for enzyme catalysis [26, 29], we examined the requirement of the N-terminal Ser¹ on the activity of SprE. Because the wild-type SprE subsequently underwent auto-degradation after trypsin treatment, we used SprE-Glu6Gln/Glu11Gln instead of wild-type SprE as control and for mutagenesis. When Ser¹ of SprE-Glu6Gln/Glu11Gln was mutated to Thr, Ala and Val, the activity of SprE after trypsin treatment was decreased to 55%, 36%, and 31% respectively (Fig. 4a). Thus, the amino acid Ser¹ was required at N-terminal position for the maximal activity and amino acids similar to Ser are more preferable, whereas the extent is not so absolute compared to the Val¹ requirements of GluV8 [26,29]. The substitution of Leu² to Val also reduced the activity to 44% of that of wild-type (Fig.4b).

Altogether, these results indicated the N-terminal amino acids involved with the substrate interaction in the enzyme catalysis.

Fig 4. Comparison of proteolytic activities of SprE substitution mutants. (a) Proenzymes of SprE-Glu6Gln/Glu11Gln and its Ser¹ substitution derivatives (Ser1Thr, Ser1Ala and Ser1Val) (10μg) were incubated at 37°C without or with 0.003μg trypsin (Tryp), as described in the materials and methods section. The proteolytic activities (mean±SD; n=3) of the samples were determined with LLE-MCA. (b) Proenzymes of SprEwt and SprE-Leu2Val (10μg) were incubated at 37°C without or with 0.3μg thermolysin (Th) and 0.003μg trypsin (Tryp) for 2h and their proteolytic activities (mean±SD; n=3) were measured as discussed in the materials and methods.

4. Discussion

We purified mixture of two SprE proenzymes starting Glu¹⁴ and Ser⁷ when expressed in *E. coli* and these proenzymes showed trace Glu-specific activity towards LLE- and LLQ-MCA. When all of the three susceptible Glu in the prosequence [Glu¹⁵, Glu¹⁴ and Glu⁸] to stop auto degradation in the prosequence, then only one proenzymes starting with Lys² (Lys² pro-SprE) was purified (data not shown). However, no such cleavage was found when catalytically inert SprE (active site Ser¹⁸⁰ was mutated to Ala) was purified from *E. coli* lysate (unpublished data). Taken together, we proposed that full-length pro-SprE truncated its prosequence to shorter zymogens (Glu¹⁴ pro-SprE, Ser⁷ pro-SprE or Lys² pro-SprE) by autocatalysis to facilitate final heterocatalytic maturation. Furthermore, absence of any R-X or S-X cleavage activity by the SprE (both pro- and mature) reiterated the inability of final maturation by autocatalysis.

We have been reported for the first time the in vitro processing of glutamyl endopeptidase proenzymes from *Enterococcus faecalis*, (pro-SprE) expressed in *E. coli*. In this study we showed that trypsin (molar ratio 3300:1) could efficiently process pro-SprE to mature SprE in dose-dependent manner, whereas thermolysin (molar ratio 33:1) could also process leaving nearly inactive SprE (Fig-1d). Sequencing data confirmed that thermolysin processed between Ser¹-Leu². Whereas taken together the sequencing data of trypsin processed SprEwt, SprE-Glu11Gln and SprE-Glu6Gln/Glu11Gln, it was confirmed that trypsin processed between Arg¹-Ser¹ (Table 1) leaving fully active SprE, however the wild-type SprE subsequently underwent auto degradation preferentially between Glu¹¹-Val¹² and Glu⁶-Asp⁷ after maturation. *E. faecalis* strain V583 and TX0411 were reported to carry the *htrA* gene that produces the trypsin-like serine protease [34,35]. Thus we hypothesize, *E. faecalis* strains expressing trypsin might efficiently process secreted pro-SprE zymogens to fully active mature SprE in vivo and might be more pathogenic than strains which do not express trypsin.

It was reported that N-terminal Val¹ truncated GluV8 (GluV8 starting with Ile²) loose enzyme activity [29]. Moreover, it is known that docking of an N-terminal residue of a chymotrypsin family serine protease affected substrate-binding sites of the enzyme [36]. Thus the length of N-terminal region was indispensable for the glutamyl endopeptidase activity and in absence of trypsin-like protease; other thermolysin-like protease(s) might process SprE at Ser¹-Leu² leaving Leu² as N-terminal amino-acids that still retained very faint residual gelatinase activity.

As the wild-type active SprE rapidly underwent autodegradation at Glu¹¹-Val¹² and Glu⁶-Asp⁷ (Table-1), we mutated the susceptible Glu- residues (Glu¹¹ and Glu⁶) to Gln- and found that the recovery of mature SprE following trypsin treatment was greatly enhanced (approximately two-fold increase when Glu¹¹ was mutated and about 2.4-fold when both Glu¹¹ and Glu⁶ were mutated) (Fig-2). However we did not find any auto-degradation between

Glu¹⁷-Ala¹⁸ that was reported to be the primary auto-degradation site [18]. We supposed that auto degradation first attacked at Glu¹¹-Val¹² and then the enzymatically inactive degraded products underwent further sequential degradation at many points including Glu¹⁷-Ala¹⁸.

Staphylococcal GluV8 family proteases were reported to conserve Val¹ as N-terminal amino acid and indispensable in enzyme catalysis [14,25,29]. To investigate such importance of Ser¹ of mature SprE, we mutated Ser¹ to Thr, Ala and Val. In our surprise we found such substitutions were not significantly affect the enzyme activity as in the case of Val¹ of GluV8. Moreover Leu² substitution to Val also reduced Glu-specific activity (SprE-Leu2Val showed 44% activity of that of the SprEwt) after maturation by trypsin. However, thermolysin again could not raise protease activity even after processing at Val² position. The possible effects of substitution of N-terminal Ser¹ or Leu² on SprE activity might be due to the difference in structure as well as in hydrophobicity with respect to Ser¹ and Leu². Increase hydrophobicity might bury the N-terminal residue more compactly making less available for substrate interaction to S1 site. Thus we hypothesized the N-terminal amino acid residue of the glutamyl endopeptidase is evolutionally conserved for facilitating heterocatalytic maturation and to favour substrate interaction to the active site.

In conclusion, we reported in vitro processing of SprE by trypsin, and investigated the significance of N-terminal residue that may enlighten further study on the in vivo processing and pathogenic importance of SprE.

Acknowledgements

We greatly acknowledge T. Kobayakawa (Nagasaki University) for the technical assistance. This work was supported by grants-in-aid for scientific research from the Ministry of Education, Science, Sports, and Culture of Japan (to Y. O.-N., S. K. and T. K. N.).

Abbreviations

GluV8 & SprE, glutamyl endopeptidase from *Staphylococcus aureus* and *Enterococcus faecalis* respectively; ac-, acetyl; boc, t-butyloxycarbonyl-[(2S)-2-amino-3-(benzyloxycarbonyl)propionyl]; Z-, benzyloxycarbonyl-; MCA, 4-methylcoumaryl-7-amide, Th, thermolysin; Tryp, trypsin

References

- [1] Murry BE, "The life and times of the *enterococcus*" Clin Microbiol Rev, 1990, Vol-3, pp 46-65
- [2] Jett BD, Huycke MM, Gilmore MS, "Virulence of *Enterococci*" Clin Microbiol Rev, 1994, Vol 7, pp 472-478
- [3] Sedgley CM, Molander A, Flannagan SE, et al, "Phenotype, genotype and virulence of endodontic enterococci" Oral Microbiol Immunol 2005, Vol 20, pp 10- 19

- [4] Stuart CH, Schwartz SA, Beeson TJ, Owatz CB, "Enterococcus faecalis: its role in root canal treatment failure and current concepts in retreatment" J Endod 2006, Vol 32, pp 93-98
- [5] Rich RL, Kreikemeyer B, Owens RT, et al, "Ace is a collagen-binding MSCRAMM from *Enterococcus faecalis*" J Biol Chem, 1999, Vol 274, pp 26939-26945.
- [6] Hubble TS, Hatton JF, Nallapareddy SR, Murray BE, Gillespie MJ, "Influence of *Enterococcus faecalis* proteases and the collagen binding protein, Ace, on adhesion to dentin" Oral Microbiol Immunol, 2003, Vol 18, pp 121-126.
- [7] Wang L, Dong M, Zheng J, Song Q, Yin W, Li J and Niu W, "Relationship of biofilm formation and *gelE* gene expression in *Enterococcus faecalis* recovered from root canals in patients requiring endodontic retreatment" JOE, 2011, Vol 37, pp631-636
- [8] Park SY, Kim KM, Lee JH, Seo SJ and Lee IH, "Extra cellular gillatinase of *Enterococcus faecalis* destroys a defense system in insect hemolymph and human serum" Infect Immun, 2007 Vol 75, pp 1861-1869.
- [9] Potempa J, and Pike RN, "Corruption of innate immunity by bacterial proteases" J. Innate Immun, 2009 Vol 1, pp 70-87
- [10] Schmidtchen A, Frick IM, Anderson E, Tapper H and Bjorck L, "Proteinases of common pathogenic bacteria degrade and inactivate the antibacterial peptide LL-37" Mol. Microbiol, 2002, Vol 46, pp 157-168
- [11] Qin X, Shing K V, Weinstock GM and Murray BE. "Effects of *Enterococcus faecalis* *fsr* genes on production of gillatinase and a serine protease and virulence" Infect. Immun, 2000, Vol 68, pp 2579-2586.
- [12] Gutschik E, Moller S and Christensen N, "Experimental endocarditis in rabbits.3. Significance of the proteolytic capacity of the infecting strain of *Enterococcus faecalis*". Acta Pathol. Microbiol. Scand, 1979, Vol B87, pp 353-362.
- [13] Shing KV, Nallapareddy SR, Nannini EC, and Murray BE, "Fsr-independent production of protease(s) may explain the lack of attenuation of an *Enterococcus faecalis* *fsr* mutant versus a *gelE-sprE* in induction of endocarditis" Infect. Immun, 2005, Vol 73, pp 4888-4894.
- [14] Prasad L, Leduce Y, Hayakawa K and Delbaere L.T. "The structure of a universally employed enzyme: V8 protease from *Staphylococcus aureus*" 2004, Acta Crystallogr. Sect. D, Vol 60, pp 256-259.
- [15] Nemoto T K, Ono T, Shimoyama Y, Kimura S and Ohara-Nemoto, Y, "Determination of three amino acids causing alteration of proteolytic activities of *Staphylococcal* glutamyl endopeptidase" J. Biol. Chem, 2009, Vol 390, pp 277-275.
- [16] Engelbert M, Mylonakis E, Ausubel FM, Calderwood SB and Gilmore MS. "Contribution of gillatinase, serine protease, and *fsr* to the pathogenesis of *Enterococcus faecalis* endophthalmitis" Infect. Immun, 2004, Vol 72, pp 3628-3633.
- [17] Sifri CD, Mylonakis E, Shing KV, Qin X, Garsin DA, Murry BE, Ausubel FM, and Calderwood, SB. "Virulence effects of *Enterococcus faecalis* protease genes and the quorum-sensing locus *fsr* in *Caenorhabditis elegans* and mice" Infect. Immun, 2002, Vol 70, pp5647-5650.
- [18] Kawalec M, Potempa J, Moon JL, Travis J and Murray BE, "Molecular diversity of a putative virulence factor: Purification and characterization of an extracellular serine glutamyl endopeptidase of *Enterococcus faecalis* with different enzymatic activities" J Bacteriol, 2005, Vol 187, pp266-275.
- [19] Yoshikawa K, Tsuzuki H, Fujiwara T, Nakamura E, Iwamoto H, Matsumoto K, Shin M, Yoshida N and Teraoka H, " Purification, Characterization and gene cloning of a novel glutamic acid-specific endopeptidase from *Staphylococcus aureus* ATCC12600" Biochim. Biophys. Acta, 1992, Vol 1121, pp 221-228.
- [20] Ohara-Nemoto Y, Ikeda Y, Kobayashi M, Sasaki M, Tajika S and Kimura S, "Characterization and molecular cloning of a glutamyl endopeptidase from *Staphylococcus epidermidis*" Microb. Pathog, 2002, Vol 33, pp33-41.
- [21] Drapeau GR, "Role of metalloprotease in activation of the precursor of *Staphylococcal* protease" J bacterial, 1978, Vol 136, pp607-613.
- [22] Shaw L, Golonka E, Potempa J and Foster S J, "The role and regulation of the extracellular proteases of *Staphylococcus aureus*" Microbiology, 2004, Vol 150, pp217-228.
- [23] Nickerson NN, Prasad L, Jacob L, Delbaere LT and McGavin MJ, "Activation of the SspA serine protease zymogen of *Staphylococcus aureus* proceeds through unique variations of a trypsinogen mechanism and is dependent on both auto-catalytic and metallo protease-specific processing" J. Biolo. Chem, 2007, Vol 47, pp34129-34138.
- [24] Rouf SMA, Ohara-Nemoto Y, Shimoyama Y, Kimura S, Ono T and Nemoto, TK, "Propeptide processing and proteolytic activity of proenzymes of the *Staphylococcal* and *Enterococcal* GluV8-family protease" Indian J Biochem Biophys, 2012, Vol 49(6), pp421-7
- [25] Nemoto TK, Ohara-Nemoto Y, Ono T, Kobayakawa T, Shimoyama Y, Kimura S, and Takagi T, "Characterization of the glutamyl endopeptidase from *Staphylococcus aureus* expressed in *Escherichia coli*" FEBS.J, 2008, Vol 275, pp573-587.
- [26] Ohara-Nemoto Y, Ono T, Shimoyama Y, Kimura S and Nemoto T K, "Homologous and heterologous expression and maturation processing of extracellular glutamyl endopeptidase of *Staphylococcus epidermidis*" J.Biol. Chem, 2008, Vol 389, pp 1209-1217.
- [27] Dubin G, Chmiel D, Mak P, Rakwalska M, Rzychon M and Dubin A, "Molecular cloning and biochemical characterization of proteases from *Staphylococcus epidermidis*" J.Biol. Chem, 2001, Vol 382, pp 1575-1582.
- [28] Yokoi K, Kakikawa M, Kimoto H, Watanabe K, Yasukawa H, Yamakawa A, Taketo A and Kodaira K, "Genetic and Biochemical characterization of glutamyl endopeptidase of *Staphylococcus warneri*" M Gene, 2001, Vol 281, pp115-122.
- [29] Ono T, Ohara-Nemoto Y, Shimoyama Y, Okawara H, Kobayakawa T, Baba T T, Kimura S and Nemoto T K. "Amino acid residues modulating the activities of *Staphylococcal* glutamyl endopeptidases" J. Biol. Chem, 2010, Vol 391, pp1221-1232.

- [30] Ono T, Nemoto TK Shimoyama Y, Kimura S and Ohara-Nemoto Y, “ An *Escherichia coli* expression system for glutamyl endopeptidases optimized by complete suppression of autodegradation” Analytical Biochemistry. 2008, Vol 381, pp74-80.
- [31] Rasband WS., ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA, <http://imagej.nih.gov/ij/>, 1997-2011.
- [32] Rice K, Peralta R, Bast D, de Azavedo J and McGavin MJ, “Description of staphylococcus serine protease (*ssp*) operon in *Staphylococcus aureus* and nonpolar inactivation of *sspA*-encoded serine protease” Infect. Immun, 2001, Vol 69, pp159-169.
- [33] Qin X, Shing KV, weinstock GM and Murray BE. “Characterization of *fsr*, a regulator controlling expression of gelatinase and serine protease in *Enterococcus faecalis* OG1RF” J Bacteriol, 2001, Vol 183, pp3372-3382.
- [34] Paulsen I, Banerjee L, Myers G et al, “Role of mobile DNA in the evaluation of venomycin-resistant *Enterococcus faecalis*” Science, 2003, Vol 299 (5615), pp2071-2074.
- [35] Weinstock G, Sodergren E et al. Direct Submission, 2010, Genome sequencing center, Washington university school of medicine, 4444 Forest park, Saint-Louis, MO 63108, USA.
- [36] Bode W, and Huber R, “Induction of the bovine trypsinogen-trypsin transition by peptides sequentially similar to the N-terminus of trypsin” FEBS Lett, 1976, Vol 68, pp231-236.