
Diversity and Abundance of Arbuscular Mycorrhizal Fungi Under Different Plant and Soil Properties in Sidama, Southern Ethiopia

Beyene Dobo^{1,*}, Fassil Asefa², Zebene Asfaw³

¹Department of Natural Resources Management and Environmental Sciences, Haramaya University, Haramaya, Ethiopia

²Department of Microbial, Cellular and Molecular Biology, Addis Ababa University, Addis Ababa, Ethiopia

³Wondo Genet College of Forestry and Natural Resources, Hawassa University, Hawassa, Ethiopia

Email address:

beyeneashl@yahoo.co.uk (D. Beyene), asefafasil2013@gmail.com (A. Fassil), zebeneasfaw@yahoo.co.uk (A. Zebene)

*Corresponding author

To cite this article:

Beyene Dobo, Fassil Asefa, Zebene Asfaw. Diversity and Abundance of Arbuscular Mycorrhizal Fungi Under Different Plant and Soil Properties in Sidama, Southern Ethiopia. *Advances in Bioscience and Bioengineering*. Vol. 4, No. 3, 2016, pp. 16-24.

doi: 10.11648/j.abb.20160403.11

Received: June 13, 2016; **Accepted:** July 1, 2016; **Published:** July 21, 2016

Abstract: In Sidama, agroforestry represents land-use systems with deliberate management of multipurpose trees and shrubs that grow in intimate association with annual and perennial agricultural crops and/or livestock. The interaction of microbiota with the trees, shrubs and crops make the system fertile, productive and sustainable. One of the beneficial microbiota which has symbiotic association with most of the plants in agroforestry is arbuscular mycorrhizal fungi (AMF). In November and December of 2012, root and rhizosphere soil samples of 21 plant species from nine peasant associations (PAs)(villages within districts where 300-500 families live) were collected from the agroforestry practices in Sidama of Southern Ethiopia for the determination of diversity and abundance of AMF under selected soil parameters and plant species density. Findings on the diversity of AMF based on soil properties showed that at moderate to low P and N concentrations the rate of AMF root colonization and spore density was high in comparison with the rhizosphere soils with the highest P and N concentration. The highest percentage of total AMF colonization was recorded for shade trees *Millettia ferruginea* (84%) and *Erythrina brucei* (80%) followed by intercropped perennial crops *Ensete ventricosum* (86%), *Catha edulis* (85%) and *Coffea arabica* (80%) and the lowest percentage AMF colonization was recorded for *Rhamnus prinoides* (53%) and *Colocasia esculenta* (52%). Though found in almost all homegarden agroforestry practices and with broad coverage in Sidama agroforestry, some crops and vegetables such *Brassica integrifolia* and *Cucurbita pepo*, grown intercropped were found to be non-mycorrhizal as none of the AMF structures were recorded. The highest number of AMF spore population was recorded in rhizosphere soils of *Croton macrostachyus* (1066±19.33) and *Catha edulis* (1054±53.12) and the lowest spore density was recorded for *Dioscorea alata* (100.00±2.89) spore per 100 g of dry soil. The percentage fungal colonization in any individual plant species and spore population in the rhizosphere soils of that species did not correlate to each other and percentage AM root colonization and spore density of all plants in the agroforestry of Sidama were found significantly different at P<0.05 level.

Keywords: AMF, Rhizosphere, Parameters, Colonization, Density

1. Introduction

Forests are cleared for agriculture and settlement in different parts of Ethiopia. Agroforestry is an age old traditional practice which gradually replaced forests. It is a

sustainable management system for land that combines agricultural crops, trees, forest plants and/or animals simultaneously or sequentially, and applies management practices that are compatible with the cultural patterns of the local population [1]. It is practiced on Homegarden [24] and [40], cropland [31], and forest lands [10]; [39]. Although the

role of agroforestry in conserving biological diversity is being appreciated in many parts, the sustainability of these practices is a major concern in Ethiopia.

Low soil fertility, acidity and deficiency in major nutrients such as phosphorus and nitrogen are some of the problems related to successful agroforestry systems. The conversion of forest to different agricultural systems causes loss of plant species that expose the soil to water and wind erosion. Besides, the removal of plant species from forests and agroforests may be accounted for the loss of biodiversity, not only of plants, but also the microbiota important for the soil health as a whole. Therefore, maintenance of floristic diversity in agroforestry is number one crucial condition for sustainable productivity of the agroforests. One of such situations is planting of mycorrhizal seedlings that can be a good measure to increase the yield of the plants for the sustainable management for land that will increase the overall production.

Arbuscular mycorrhizal fungi are widespread and symbiotic copious members of the soil biota, a generally acknowledged key factor in agricultural ecosystem functioning and sustainability [39]. These symbiotically associated fungi play an important role to the plants in agroforestry practices by enhancing establishment and growth, increasing nutrient and water uptake, maintaining diversity by accelerating the ability of host plants to compete for resources, contributing to efficient recycling of nutrients and thus to long term stability and stabilization of the soil [16, 35]. AMF are widespread in tropical soils and associate with a wide range of plant species, including most commercial crops and trees [6, 34]. They are keystone organisms and form an interface between soils and plant roots, and are sensitive to changes in soil physicochemical properties and plant conditions [28].

Under the insufficient concentration of nutrients in the root zone of plants, exploring further in the soil with its extended extra radical hyphae, AMF supply phosphorus, macro and micro soil nutrients to roots which has symbiosis with AMF. In these symbiotic associations both partners benefit from each other under certain conditions [4, 8, 19, 30]. Fungi take some organic matter and carbohydrates from the plant. In return, AMF supply nutrients such as phosphorus, nitrogen (N), calcium (Ca), copper (Cu), manganese (Mn), sulphur (S) and zinc (Zn) [26, 34]. But when the soil is rich in necessary

nutrients, plants may not waste more energy for the association with AMF and go for the cheapest source of energy, and individual plant species and plant communities in natural and farming systems affect the distribution and diversity of AMF species [7, 17].

Therefore, according to the previous investigators, AMF spore density, diversity and rates of root colonization are dependent on soil physicochemical properties [22] and plant species density and richness [1]. Though there are some research reports [40] on mycorrhizal association of agroforestry trees, information is scarce on spore density and diversity of AMF based on soil parameters and type of plant species density and richness in Sidama agroforestry. Therefore, the objective of the present investigation is to study diversity of AMF based on selected soil parameters and plant species density in selected sites of Sidama agroforestry practices.

2. Materials and Methods

2.1. Sampling Site

This study was conducted in Sidama Administrative region of Southern Ethiopia during dry season in November and December of 2012. It lies between 06°45'33" and 06°54'713" N and 038°27'432" and 038°31'788"E and 1740 to 2135 m above the sea level. The study area is characterized by a moist to sub-humid warm sub-tropical climate with annual rainfall of 1000–1800 mm and mean annual temperature of 15–20°C. The main agricultural system in the region is the Tree-enset-coffee based home-garden agroforestry systems commonly practiced in Southern Ethiopia. The study was undertaken in 36 home gardens located in 9 peasant association (PAs, smallest Ethiopian administrative unit, where 300-500 families live) distributed over two woredas or district. Each woreda (district) has 40–60 PAs where these agroforestry systems are practiced. In this study, we selected only 9 PAs (Table 1) because these PAs are located at closest proximity to the original forest relics from where the agroforestry practices were emerged. From each PAs, 4 homegardens were selected randomly and soil and root samples were collected from 23 commonly occurring plant species in selected agroforestry practice (Table 2).

Table 1. Geographical location and altitudinal ranges of the Peasant Associations (PAs).

No	Study site (PAs)	Latitude	Longitude	Altitude
1.	Ferro I	06°46'30"N	038°29'00"E	1795-1896
2.	Ferro II	06°49'602"N	038°29'135"E	1860-1940
3.	Halekena	06°45'333"N	038°27'432"E	1740-1825
4.	Taramesa	06°086'055"N	038°45'605"E	1745-1868
5.	Tellamo	06°083'629"N	038°48'489"E	1825-1906
6.	Gonowagoda	06°084'609"N	038°50'916"E	1855-1990
7.	AbelaLida	06°54'993"N	038°29'317"E	1945-1960
8.	Galakohireye	06°54'864"N	038°30'315"E	2006-2011
9.	Arossa	06°54'713"N	038°31'788"E	2111-2135

Table 2. Plant species present in Sidama agroforestry systems the time of sampling.

S. No	Latin names	Family	Amharic	Sidamigna
1.	<i>Ensete ventricosum</i> (Welw.) Cheesman	Musaceae	Enset	Wesse
2.	<i>Catha edulis</i> (vahl.) Forssk. exEndl	Celastraceae	Khat	Khat
3.	<i>Millettia ferruginea</i> Hochst	Fabaceae	Birbira	Hengedicho
4.	<i>Erythrina brucei</i> Schweinf	Fabaceae	Korch	Wellako
5.	<i>Cordia africana</i> Lam.	Boraginaceae	Wanza	Wadicho
6.	<i>Croton macrostachyus</i> (HochstExDel.)	Euphorbiaceae	Bisana	Mesincho
7.	<i>Prunus africana</i> (Hook. f.) Kalkm.	Rosaceae	Tikurlnchet	Garbicho
8.	<i>Vernonia amygdalina</i> Del.	Asteraceae	Grawa	Hecho
9.	<i>Persea americana</i> Mill.	Lauraceae	Avokado	Abukato
10.	<i>Mengifera indica</i> L.	Anacardiaceae	Mango	Mango
11.	<i>Saccharum officinarum</i> L.	Poaceae	Shenkorageda	Shenkora
12.	<i>Ricinus communis</i> L.	Euphorbiaceae	Castor	Qenboo
13.	<i>Coffea arabica</i> L.	Rubiaceae	Coffee	Buna
14.	<i>Zea mays</i> L.	Poaceae	Bokolo	Bedela
15.	<i>Phaseolus vulgaris</i> L.	Fabaceae	Common bean	Wahe
16.	<i>Ipomoea batatas</i> (L.) Lam	Convolvulaceae	Sikuardinich	Metatessa
17.	<i>Solanum tuberosum</i> L.	Solanaceae	Potato	Dinich
18.	<i>Cucurbita pepo</i> L.	Cucurbitaceae	Pumpkin	Baqula
19.	<i>Brassica integrifolia</i> (West) O. E. schulz	Brassicaceae	Kale	Shana
20.	<i>Cappisicum annuum</i> L.	Solanaceae	Chilly	Mitmita
21.	<i>Rhamnus prinoides</i> L' herit	Rhamnaceae	Gesho	Taddo
22.	<i>Dioscorea alata</i> L.	Dioscoraceae	Boyna	Bohe
23.	<i>Colocasia esculenta</i> (L.) Schoot.	Araceae	Godere	Qolchoma

2.2. Soil Sampling and Analysis

Rhizosphere soils from under the canopy of the 23 plant species which included trees, crops and vegetable species were collected during dry season (November-December) in year 2012 from each sampling site (PAs). Soil sample collection was carried out at four 10m x 10m quadrats from each PAs. The soil was sampled to a depth of 0–15cm in 828 sampling points using a soil auger. A composite soil sample of 500 g was collected from each plant species. In total (23 x 9 x 4) samples were collected for analysis from all the 9 PAs (sampling sites), i. e. 23 x 4 samples from each sampling site. The soil samples were air-dried at room temperature for two weeks and preserved at 4°C for analysis of soil physicochemical properties, AMF spore abundance and AMF diversity. About 0.5mg of fine root samples from each plant were also collected, washed with tap water, preserved in 50% ethanol, and stored at 4°C for percentage root colonization analysis.

Soil analysis were undertaken at the Southern Nations, Nationalities and Peoples Region (SNNPR) Soil Laboratory and Debrezeit Agricultural Research Center following standard procedures and methods: Soil textural fractions were analyzed following the hydrometric method after removing organic matter using H₂O₂ and dispersing the soils with sodium hexameta-phosphate [2]. Soil pH was determined by potentiometric methods using 1:2.5 soil: water ratio. Soil organic carbon (SOC) was determined by the Walkley-Black oxidation method [33]. Total nitrogen (TN) was determined using the Kjeldahl distillation method [5], and available phosphorous (AP) was determined using Olsen's extraction method (UV/visible Spectrometer, Lambda EZ 201) [25]. Available potassium (Av. K) was determined by Sodium Acetate method [18].

2.3. Root Colonization

The root pieces collected and preserved were stained according to [27] with some modifications. The root pieces were cleared in 10% KOH solution for 15-20 minutes at 90°C in a boiling water bath. Then, they were rinsed in water several times and acidified with 1% HCl solution. Pigmented roots were bleached in H₂O₂ for about 20 minutes and again washed with water. Finally the root segments were stained for 10 minutes in 0.05% trypan blue at 90°C for about 10 minutes and subsequently destained at room temperature in acidic glycerol.

The stained root segments were mounted in acidic glycerol on slides. The roots were then observed under the compound microscope (200x objective magnification). The presence of colonization in a root segment was recorded if hyphae (only), vesicle or arbuscule were found. About 100-150 intersections were examined for each sample. Total root colonization was calculated using the following formula: % Colonization = Total number of positive segments/Total number of segments studied × 100.

2.4. Spore Population

From each soil sample 100 g soil was mixed in a 2 liter capacity beaker containing 1.5 liter of water. The soil in water was agitated by stirring vigorously by hand and left to settle down for about five minutes. The suspension was then sequentially (480, 100, 50 & 38µm) sieved following the wet sieving and decanting method [13]. The last pellet (38µm) was suspended in 60% sucrose solution and thoroughly mixed and centrifuged at 2000rpm for 1 minute and the spores were rinsed carefully with tap water and transferred into plastic petri-dishes.

The AMF spores and sporocarps of each sample were

counted under 4x stereomicroscope. The spore densities were expressed as the numbers of spores and sporocarps per 100 g⁻¹ of dry soil. Representative Morphospecies were mounted on slides with polyvinyl-lactic acid-glycerol (PVLG) or PVLG mixed with Melzer's reagent (1:1 v/v). Taxonomic differentiation was made based on the descriptions of the International Culture Collection of Vesicular/Arbuscular Mycorrhizal Fungi (<http://invam.caf.wvu.edu>; 2005), and following descriptions by [32].

2.5. Statistical Analysis

Data on spore abundance and root colonization was log (x)

3. Results

3.1. Soil Physicochemical Properties Under Different Plant Species

Table 3. Rhizosphere soil chemical properties of plants.

S. No	Plant species	pH (H ₂ O)	OC%	TN%	AP (Olson) (mg/kg)	Kcmol (+) /kg
1.	<i>Ensete ventricosum</i> (Welw.) Cheesman	6.49±.12b	2.41±.23a	0.26 ±0.02c	10.20±1.3e	0.61±0.33a
2.	<i>Catha edulis</i> (vahl.) Forssk. exEndl	5.02±0.00a	2.04±0.02a	0.23±0.00 b	7.12±0.01b	0.88±0.01a
3.	<i>Milletia ferruginea</i> Hochst	6.530.03b	2.46±0.01a	0.36 ±0.01d	9.13±0.01d	1.01±0.00b
4.	<i>Erythrina bruce</i> Schweinf	6.11±0.01b	2.48±0.03a	0.31 ±0.00d	9.10±0.05d	1.16±0.01b
5.	<i>Cordia africana</i> Lam.	7.12±0.01c	3.45±0.03a	0.36 ±0.01d	15.33±0.02g	1.22±0.01b
6.	<i>Croton macrostachyus</i> (HochstExDel)	6.50±0.06b	2.50±0.06a	0.33 ±0.02d	8.15±0.01c	0.98±0.01ab
7.	<i>Prunus africana</i> (Hook. f.) Kalkm.	5.71±0.32a	2.64±0.01ab	0.29±0.00 d	16.12±0.01h	0.55±0.01a
8.	<i>Vernonia amygdalina</i> Del.	5.60±0.08a	3.15±0.00c	0.34±0.01d	17.07±0.06i	1.19±0.05b
9.	<i>Persea americana</i> Mill.	6.40±0.12b	2.63±0.00ab	0.29 ±0.01d	18.25±0.14j	0.88±0.01a
10.	<i>Mengifera indica</i> L.	5.20±0.23a	2.74±0.01ab	0.36±0.02 d	16.50±0.06h	0.75±0.02a
11.	<i>Saccharum officinarum</i> L.	5.35±0.02a	2.31±0.00a	0.29±0.01d	9.23±0.03c	1.35±0.0b
12.	<i>Ricinus communis</i> L.	5.38±0.03a	2.63±0.04ab	0.29 ±0.00d	8.21±0.01c	0.55±0.06a
13.	<i>Coffea arabica</i> L.	6.50±0.06b	3.21±0.01c	0.32 ±0.01d	7.68±0.01b	1.21±0.02b
14.	<i>Zea mays</i> L.	5.63±0.04a	2.51±0.29a	0.30 ±0.01d	8.93±0.04c	0.71±0.02a
15.	<i>Phaseolus vulgaris</i> L.	5.51±0.29a	3.14±0.02c	0.33 ±0.01d	7.68±0.28b	1.14±0.01b
16.	<i>Ipomoea batatas</i> (L.) Lam	5.14±0.09a	2.53±0.00a	0.30 ±0.01d	7.35±0.03b	2.18±0.01c
17.	<i>Solanum tuberosum</i> L.	5.33±0.09a	2.71±0.02ab	0.35 ±0.00d	6.75±0.03a	1.15±0.01b
18.	<i>Capsicum annum</i> L.	6.05±0.48b	2.47±0.02a	0.30 ±0.01d	14.52±0.25f	0.95±0.03ab
19.	<i>Rhamnus prinoides</i> L' herit	5.67±0.22a	2.65±0.00ab	0.34 ±0.01d	16.01±1.16h	1.20±0.02b
20.	<i>Dioscorea alata</i> L.	5.93±0.08ab	2.37±0.07a	0.19 ±0.01a	17.09±0.03i	1.34±0.01b
21.	<i>Colocasia esculenta</i> (L.) Schoot.	5.23±0.09a	2.78±0.01ab	0.27 ±0.01c	16.60±0.12h	2.22±0.01c
	Mean Total	5.87±0.00ab	2.73±0.05ab	0.30 ±0.01d	12.39±0.54ef	1.10±0.05a

Key: OC, organic carbon; TN, total nitrogen; AP, available phosphors, K, potassium. Similar letters in columns show not significant difference between plant species at p<0.05.

The physicochemical properties of soil samples of each plant are shown in Table 3. The soil in the study area was slightly acidic with mean pH of 5.87; mean organic carbon, total nitrogen, available phosphorus and potassium being 2.73, 0.30, 12.39 and 1.10 respectively. When individual plant species were considered, all plant species studied significantly affected the rhizosphere soil chemical properties (Table 3). They showed variation in soil pH (H₂O) from 5.0 (acidic) – 7.0 (neutral), whereas soil nitrogen content was in between 0.19 (medium)-0.36% (high). Similarly, phosphorus (Olson) content varied from the lowest 6.75 mg P/kg (*Solanum tuberosum*) to the highest 18.25 mg P/kg (*Persea americana*). Potassium was highest in the rhizosphere soil of *Ipomoea batatas* (2.18cmol (+) /kg) and the lowest (0.55

and arcsine (the inverse sine of the square root of the proportion) transformed using PAST3 (ver. 1.0.0.0) and SPSS software package (version 20.0) respectively, prior to analysis to meet assumptions of ANOVA such as normality and homogeneity of variance.

Significance of differences in AM fungal spore abundance and percentage of root colonization between the samples was tested using Duncan's Multiple Range Test at p <0.05 after one-way analysis of variance (ANOVA) with the SPSS software package (version 20.0) [12]. Pearson's correlation analysis was used to study the relationship between AMF and soil parameters.

cmol (+)/Kg) was recorded from *Prunus africana* and *Ricinus communis*. Organic carbon (OC) content of soil sample fall in between 2.04 (high) to 3.45 (very high).

3.2. AMF Root Colonization and Spore Density

The AMF root colonization on the different agroforestry plant species is presented in Table 4. The highest percentage root length colonization (RLC) was recorded in *Ensete ventricosum* (86%) and the lowest in *Rhamnus prinoides* (53%) and *Colosica esculenta* (52%) respectively (Table 4). All plants were infected with AMF arbuscules and vesicles except *Brassica integrifolia* and *Cucurbita pepo* species (data not shown) (Table 4).

In case of vesicle formation, the highest colonization was

found in *Ensete ventricosum* and *Coffea arabica* each 20.33% followed by *Zea mays* 18.67% and *Catha edulis* (17%) and the lowest in *Persea americana* (9.67%), *Mangifera indica* (9.33) and *Dioscorea alata* (9%). Considering arbuscule formation, the highest percentage was observed in *Ricinus communis* (18%), *Persea americana* (18%) followed by *Ensete ventricosum* (15%) and *Solanum tuberosum* (15.33%). The lowest arbuscular colonization was recorded for *Colocasia esculenta* (5%). In most of the

records arbuscular and vesicular colonization are not significantly different at $P < 0.05$ level.

The highest number of spore population/100 g dry soil was recorded in the rhizosphere soil of *Croton macrostachyus* (1066) followed by *Catha edulis* (1054) and *Coffea arabica* (995) and the lowest density was recorded in *Dioscorea alata* (100 spores/100 g soil) (Table 4). From the data (Table 4), it is also observed that the percentage root colonization and spore population do not correlate to each other.

Table 4. Root colonization and spore density of AMF in agroforestry plant species.

S. No	Plant species	AMF Structural Colonization (%) and spore density			
		AC	VC	RLC	SD/100 g ⁻¹ dry soil
1.	<i>Ensete ventricosum</i> (Welw.) Cheesman	15.00±2.08ab	20.33±0.33a	85.73±3.73a	630.00±2.90d
2.	<i>Catha edulis</i> (vahl.) Forssk. exEndl	16.67±2.33a	17.00±1.15b	84.74±1.24a	1054.00±53.12a
3.	<i>Milletia ferruginea</i> Hochst	11.67±0.88b	16.67±1.76b	83.80±2.22ab	686.00±2.31d
4.	<i>Erythrina brucei</i> Schweinf	13.67±1.67abc	14.67±1.33bc	80.23±1.04b	768.00±39.26cd
5.	<i>Cordia africana</i> Lam.	12.00±2.31c	14.00±3.00bc	72.33±1.57c	880.00±49.07c
6.	<i>Croton macrostachyus</i> (Hochst ExDel.)	10.00±0.58c	14.33±0.33c	64.11±3.20d	1066.00±19.63a
7.	<i>Prunus africana</i> (Hook. f.) Kalkm.	11.67±0.33b	9.67±0.88e	57.67±2.69d	675.00±14.43d
8.	<i>Vernonia amygdalina</i> Del.	7.00±1.53c	13.67±2.40cbc	77.78±1.91b	600.00±46.19d
9.	<i>Persea americana</i> Mill.	18.33±3.84a	15.33±3.18bc	80.32±2.47ab	550.00±21.94e
10.	<i>Mangifera indica</i> L.	10.33±1.20c	9.33±0.88de	79.97±0.77b	580.00±69.28d
11.	<i>Saccharum officinarum</i> L.	13.67±4.18abc	16.00±4.04ab	68.87±2.25b	625.00±14.43d
12.	<i>Ricinus communis</i> L.	18.00±5.69ab	14.00±4.1bcd	74.25±4.31bc	800.00±86.60c
13.	<i>Coffea arabica</i> L.	11.33±1.45c	20.33±2.19a	80.14±1.89b	995.00±2.89b
14.	<i>Zea mays</i> L.	12.67±0.33b	18.67±1.76ab	80.59±1.82b	700.00±46.19c
15.	<i>Phaseolus vulgaris</i> L.	11.33±1.67c	15.00±0.58b	73.27±1.10c	495.00±83.72d
16.	<i>Ipomoea batatas</i> (L.) Lam	12.00±0.58b	15.00±0.58b	70.62±1.96c	665.00±39.84d
17.	<i>Solanum tuberosum</i> L.	15.33±5.55ab	12.00±2.52c	73.04±1.63c	520.00±11.55e
18.	<i>Cappisicum annum</i> L.	12.33±1.45b	10.00±3.00dcd	72.67±1.76c	632.00±27.71d
19.	<i>Rhamnus prinoides</i> L'herit	9.67±1.20c	9.67±0.88de	53.33±5.02d	751.00±5.20c
20.	<i>Dioscorea alata</i> L.	11.33±1.45c	9.00±1.53de	60.52±4.67d	100.00±2.89f
21.	<i>Colocasia esculenta</i> (L.) Schoot	5.00±1.0d	14.33±7.17c	51.50±7.17d	660.00±92.38cd

Key: AC, Arbuscular Colonization; VC, Vesicular Colonization; RLC, total Root Length Colonization; SD, Spore Density. Similar letters in columns show not significant difference between groups at $p < 0.05$.

Current result in this study reveals that under highest P and N concentrations spore density decreased in soils from rhizospheres of some plant species, although the opposite is true for the other species. For soils from the rhizospheres of *Persea Americana* (P, 19.25; N 0.29), *Vernonia amygdalina* (P, 17.07; N 0.34), *Mangifera indica* (P, 16.50; N, 0.36), and *Dioscorea alata* (P, 17.09; N, 0.19) was recorded, 550 & 80, 600 & 78, 580 & 80 and 100 & 66 spores per 100 g soil and percentage root colonization respectively.

On the other hand for agroforestry plant species with low to medium P concentration such as *Catha edulis* (P, 7.12; N, 0.23) *Croton macrostachyus* (P, 8.15; N, 0.33, *Coffea arabica* (P, 7.68; N, 0.32) was recorded 1054 & 85, 1066 & 64 and 995 & 80 spores per 100 g soil and percentage root colonization respectively. This result indicates that different plant species respond to AMF differently under low, medium and high concentrations of nutrients in their rhizospheres.

Pearson's correlation coefficient showed that spore density was not significantly correlated with soil pH, OC, total nitrogen & potassium and significant negative correlation with available phosphorus ($r = -0.346$, $p = 0.005$) and electron conductivity ($r = -0.441$, $p = 0.00$) at $P < 0.01$ level (2-tailed). Spore density was also negatively correlated with total

nitrogen, but not significantly different.

Pearson's correlation coefficient between root colonization and soil parameters showed that it is negatively correlated to available phosphorus ($r = -0.324$, $p = 0.010$) and potassium ($r = -0.301$, $p = 0.017$) and significantly different at $P < 0.01$ and $P < 0.05$ level (2-tailed) respectively. It was positive with pH and total nitrogen but not significantly different and was negative but significantly different between root colonization and organic carbon, and negative but not significantly different in between EC and root colonization.

3.3. AMF Spore Diversity

In general 8 different AMF genera were identified (Table 5): *Acaulospora*, *Claroidioglomus*, *Funneliformis*, *Glomus*, *Gigaspora*, *Rhizophugus*, *Septoglosum*, and *Scutellospora* of which *Glomus* and *Acaulospora* were the dominant genera found in all species. The highest percentage of *Glomus* was recorded in *Coffea arabica* (15.27%) and *Ensete ventricosum* (12.68%) the lowest being recorded in *Cappisicum annum* and *Colocasia esculenta* (0.29%). The highest percentage of *Acaulospora* was recorded in three species; *Ensete ventricosum*, *Coffea arabica*, and *Zea mays* with 14.93%, 10.85% and 11.76% respectively. The highest percentage was

recorded for *Claroideoglossum* (24.18%) in *Croton macrostachyus*, *Funneliformis* (33.9%) in *Ensete ventricosum*, *Gigaspora* (22.94%) in *Catha edulis*, *Rhizophogus* (28.67%) in *Ensete ventricosum*, *Septoglossum* (20%) in *Saccharum officinarum*, and *Scutellospora* (16.67%) in *Ensete ventricosum*.

Table 5. Percentage AMF diversity in the 21 plant species in Sidama agroforestry.

S. No	Plant species	Different Genera of AMF (%)								
		<i>Ac.</i>	<i>Clar.</i>	<i>Fun.</i>	<i>Glo.</i>	<i>Giga.</i>	<i>Rhi.</i>	<i>Sep.</i>	<i>Scut.</i>	<i>Sp. No</i>
1.	<i>Ensete ventricosum</i> (Welw.) Cheesman)	14.93	14.38	33.9	12.68	17.65	28.67	13.33	16.67	8
2.	<i>Catha edulis</i> (vahl.) Forssk. exEndl	3.18	-	10.17	9.8	22.94	16	-	2.38	6
3.	<i>Milletia ferruginea</i> Hochst	4.07	14.38	11.86	11.82	2.35	12.67	6.67	5.95	8
4.	<i>Erythrina brucei</i> Schweinf	4.07	5.23	16.95	2.31	3.53	3.33	-	1.19	7
5.	<i>Cordia africana</i> Lam.	3.62	5.23	5.08	0.86	3.53	0.67	-	2.38	7
6.	<i>Croton macrostachyus</i> (HochstExDel)	9.95	24.18	6.78	9.51	1.76	7.33	13.33	8.33	8
7.	<i>Prunus africana</i> . (Hook. f.) Kalkm	3.17	1.31	-	1.73	1.18	6	-	4.76	6
8.	<i>Vernonia amygdalina</i> Del.	2.71	2.61	3.39	4.9	1.18	6	13.33	4.76	8
9.	<i>Persea americana</i> Mill.	4.07	-	-	1.73	1.18	2.67	-	3.57	5
10.	<i>Mengifera indica</i> L.	1.36	1.31	-	3.17	-	2	-	3.57	5
11.	<i>Saccharum officinarum</i> L.	3.62	2.61	-	3.46	0.58	3.33	20	7.14	7
12.	<i>Ricinus communis</i> L.	3.18	-	3.39	1.44	2.53	0.67	-	3.57	6
13.	<i>Coffea arabica</i> L.	10.85	17.64	-	15.27	17.65	2	-	13.1	6
14.	<i>Zea mays</i> L.	11.76	-	-	12.1	14.1	0.67	13.33	4.76	6
15.	<i>Phaseolus vulgaris</i> L.	4.07	1.31	3.39	1.44	2.53	3.33	-	4.76	7
16.	<i>Ipomoea batatas</i> (L.) Lam	2.26	3.92	-	2.59	2.94	2	-	3.57	6
17.	<i>Solanum tuberosum</i> L.	1.8	-	3.39	1.73	1.76	-	13.33	5.95	6
18.	<i>Cappisicum annum</i> L.	3.18	1.31	-	0.29	-	-	-	-	3
19.	<i>Rhanmu sprinoides</i> L'herit	2.26	1.96	1.69	0.58	-	-	-	-	4
20.	<i>Dioscorea alata</i> L.	1.36	1.31	-	0.86	1.18	-	6.67	1.19	6
21.	<i>Colocasia esculenta</i> (L.) School.	0.45	1.31	-	0.29	1.18	-	-	-	4
	<i>Mean</i>	9.91	7.29	2.814	16.29	8.05	6.95	0.71	3.90	

Key: *Ac.*, *Acaulospora*; *Claro.*, *Claroideoglossum*; *Fun.*, *Funneliformis*; *Glo.*, *Glomus*; *Giga.*, *Gigaspora*; *Rhi.*, *Rhizophogus*; *Sep.*, *Septoglossum*; *Scut.*, *Scutellospora*.

In this study, observed Plant species density (Figure 1) in the agroforestry does not induced consistent spore formation and percentage root colonization values in all plant species. *Colocassia esculenta* with its relative abundance value of 0.1% (982) showed percentage colonization of 52%, spore density of 660 and for *Dioscorea alata* with its relative abundance value of 0.05% (511) was recorded root colonization of 61%, more than *Colocasia esculenta* (Figure 1) and the least spore density of 100 spore/100 g soil.




Figure 1. Effect of plant species density on root colonization and spore density.

Key: PD, plant density; RLC, total root length colonization; SD, Spore density, En, *Ensete*; Ca, *Catha*; Mi, *Milletia*; Er, *Erythrina*; Co, *Cordia*; Cr, *Croton*; Pr, *prunus*; Ve, *Vernonia*; Pe, *Persea*; Me, *Mengifera*; Sa, *Saccharum*; Ri, *Ricinus*; Cof, *Coffee*; Z, *Zea*; Ph, *Phaseolus*; Ip, *Ipomoea*; So, *Solanum*; Ca, *Cappisicum*; Rh, *Rhamnus*; Di, *Discorea*; Col, *Colocasia*

On the other hand for species with the highest value of percentage abundance 0.22% (2107) in *Ensete ventricosum*, was recorded the highest percentage colonization (86%) and medium number of spore density (630 spores/100-g dry soil). For *Croton macrostachyus* with percentage abundance value of about 0.008% (75), was recorded 1066 spores/100 g soil and 64.11 percentage root colonization. The next highest spore density (1054 100-1 g soil) and root colonization (85%) was recorded for perennial crop *Catha edulis* with its percentage abundance value of about 0.02% (188), followed by *Coffea arabica*, 995 spores 100-1 g soil and 80% colonization with its abundance value of 0.12% (1147) respectively. This indicates that in some plant species (e.g. *Ensete ventricosum*) density decreased the spore production but in some other plant species such as *Coffea arabica* plant species density has increased spore production.

4. Discussion

The effects of plant species density and its associated soil chemical properties in Sidama agroforestry homegardens were studied to determine their influence on AMF diversity, spore density and root colonization. The arbuscular mycorrhizal fungi spore population and root colonization pattern varied in different plant species in Sidama homegarden agroforestry systems. The study on diversity of AMF in relation to soil parameters and plant species density did not show significant differences in AMF diversity indices (Shannon–Weaver diversity index,

Simpson's Index of Diversity (data not shown)) due to plant species density and soil physical and chemical properties. However, there were differences in AMF spore density and root colonization between plant species. Shifts in AMF species composition with increasing or decreasing plant species density may be explained in part by differential responses of individual AMF species to individual host plant species and edaphic factors in the rhizospheres of each plant species.

According to [20], three mechanisms have been proposed to account for plant density-dependent regulation of mycorrhizal colonization. According to their report, higher plant together with higher root density causes, a) an increase in overlap of phosphorus depletion zone, b) a competition for light results in less carbohydrate available for mycorrhizal fungi and c) an increase in cost-benefit ratio of mycorrhizal colonization. As plant density increase, competition for light increases and photosynthesis per plant declines and plants become limited more by carbon than by P reducing the root colonization. However, in some cases, increasing plant density has been associated with increasing AMF colonization [1, 11].

Most of the plants included in this study are multiple purpose trees and perennial crops with coarse root systems. The variation in the percentage of colonization in the roots and the AM population in the rhizosphere soils of different plants recorded in the present study, therefore might explain the generally held view that plants with coarse roots gain more AMF [22] compared to those with fine roots. These differences might be due to the presence of diverse type of AMF in the rhizosphere soils of individual plant species or might be a manifestation of greater host susceptibility to AMF [23].

In this study 8 different AMF genera were identified: *Acaulospora*, *Claroidioglomus*, *Funneliformis*, *Glomus*, *Gigaspora*, *Rhizophugus*, *Septoglomus*, and *Scutellospora* of which *Glomus* and *Acaulospora* were the dominant genera found in all species. This is not surprising since it was previously shown that *Glomus* and *Acaulospora* species are most abundant among the glomeromycotan genera in tropical areas [3, 14, 37, 41]. *Glomus* species is widely distributed regardless of the type and intensity of disturbance in the different ecosystems, whereas *Acaulospora* is dominant in least disturbed agro ecosystems [36].

In our study, the total spore density, in general, did not correlate with mycorrhizal colonization, possibly because of the presence of a diverse population of AM fungal species and it could also be due to the fact that there are AMF species that rely more on extensive formation of hyphal networks instead of survival through spore formation as primary infective propagules.

The relationship between spore numbers and mycorrhizal colonization has been found positive by several workers [15] but negative by others [21]. Some workers have also found no relationship between mycorrhizal colonization and spore density of AMF [9]. The factors like edaphic or climatic condition; host fungus compatibility, root properties and soil microorganisms might influence the abundance of spore population and mycorrhizal associations with a particular tree species.

It is noticeable in the present study that the wide spread presence of *Glomus* and *Acaulospora* in the rhizospheres of selected plants, makes them favorable fungi for mass multiplication as well as seedling inoculation for their better establishment. The present survey of AM fungi on Sidama agroforestry plant species reveals that AM fungi are common, occurring in 21 of the 23 plant species examined. However, when establishing new agroforestry practices the tree species must possess adequate amounts of mycorrhizal colonization at the planting stage in order to survive better and perform well in adverse agroforestry sites. More studies are needed to select the suitable indigenous AM fungal strains for the production of quality plant and expanding the awareness of the role of AMF in agroforestry systems to the small holder peasants for better organic and sustainable production should be initiated.

5. Conclusion

The investigation reveals that AM fungi are common, occurring in 21 out of 23 plant species examined. The study also demonstrated the influence of different plant densities on the diversity of arbuscular mycorrhizal fungi. As to the dependency of AMF on soil parameters, plants exhibited variable patterns; some with high concentration of P showed lower spore density and root colonization, while others shows the opposite phenomena. *Glomus* and *Acaulospora* were dominant in this study. They could be used for mass multiplication as well as seedling inoculation for better establishment of major trees and perennial crops which can also serve as inoculums sources. Mycorrhization of agroforestry plants has gained considerable attention over the last few years because of their role as bio-fertilizers for improving host growth. More studies are needed to select the suitable indigenous AM fungal strains for the production of quality plant and expand the awareness of the role of mycorrhiza in agroforestry systems to the small holder farmers.

Acknowledgment

We would like to acknowledge Hawassa College of Teacher Education for financial and logistic supports; Department of Microbiology, Cellular and Molecular Biology, Addis Ababa University and College of agriculture, Hawassa University for their support with chemicals and laboratory equipments.

References

- [1] Allsopp, N. and Stock W. F. (1992). Density dependent interactions between VA mycorrhizal fungi and even-aged seedlings of two perennial *Fabaceae* species. *Oecologia*, 91, 281-287. doi: 10.1007/BF00317797.
- [2] Black, C., Evans, A. D., White, J. L., Ensminger, L. E. and Clark, F. E. (1965). *Methods of Soil Analysis Part 1. Physical and Mineralogical Properties Including Statistics of Measurement and Sampling*, American Society of Agronomy, Madison, Wis, USA.

- [3] Blaszkowski, J. (1989). The occurrence of the Endogonaceae in Poland. *Agril. Ecosy. Environ.*, 29: 45-50.
- [4] Bolan, N. S., Robson, A. D. and Barrow, N. J. (1987). Effects of vesicular-arbuscular mycorrhiza on the availability of iron phosphates to plants. *Plant Soil* 99: 401-410.
- [5] Bremner, J. M. and Mulvaney, S. C. (1982). "Nitrogen-Total," in *Methods of Soil Analysis*, A. L. Page, Eds., vol. 2, pp. 595–624, American Society of Agronomy, Madison, Wis, USA.
- [6] da Silva Sousa, C., Cezar Menezes, R. S., de Sá Barreto Sampaio, E. V., de Sousa Lima, F. and Costa Maia, L. (2013). Arbuscular mycorrhizal fungi within agroforestry and traditional land use systems in semi-arid Northeast Brazil. *Acta Scient. Agron.*, 35: 307-314.
- [7] Dalpe, Y., Plenchette, C. and Gueye, M. (2000). Glomales species associated with surface and deep rhizosphere of *Faidherbia albida* in Senegal. *Mycorrhiza*, 10, 125-129.
- [8] Demir, S., Kaya, I., Şavur, O. B. and Özkan, O. U. (2008). Determination of Arbuscular-Mycorrhizal Fungus (AMF) from the Plants Belonging to *Gramineae* Family in Van Province, Yüzüncü Yıl University, Faculty of Agriculture, *J. Agric. Sci.*, 18 (2): 103-111.
- [9] Diaz, G. and Honrubia, M. (1994). A mycorrhizal survey of plants growing on mine wastes in Southern Spain. *Arid Soil Res. Rehabilitation*, 8: 59- 68.
- [10] Emiru, Birhanea, Thomas, W. K., Frank, J. S. and Frans, B. (2010). Arbuscular mycorrhizal associations in *Boswellia papyrifera* (frankincense-tree) dominated dry deciduous woodlands of Northern Ethiopia. *Forest Ecology and Management*, 260: 2160–2169.
- [11] Eissenstat, D. M. and Newman, E. I. (1990). Seedling establishment near large plants: effects of vesicular-arbuscular mycorrhizas on the intensity of plant competition. *Funct Ecol.*, 4, 95-99.
- [12] Fisher, R. A. and Yates, F. (1970). In *Statistical tables for biological, agriculture and medical research*. 6th ed. Davien: Hafner.
- [13] Gerdemann, J. W. and Nicolson, T. H. (1963). Spores of mycorrhizal *Endogone* species extracted from soil by wet sieving and decanting. *Transactions of the British Mycological Society*, 46, 235-244.
- [14] Gerdemann, J. W. and Trappe, M. J. (1974). The Endogonaceae in the Pacific Northwest. *Mycologia Memoir*, No. 5. The New York Botanical Garden, New York. Pp 76.
- [15] Giovannetti, M and Nicolson, H. T. (1983). Vesicular-arbuscular Mycorrhizas in Italian sand dunes. *Trans. Br. Mycol. Soc.*, 80: 552-555.
- [16] Jasper, D. A. (1992). Management of mycorrhizas in revegetation. In: Robson, A. D.; L. K. Abbott and N. Malajczuk (eds.), *Proceedings of International Symposium on Management of Mycorrhizas in Agriculture, Horticulture and Forestry*.
- [17] Jefwa, J. M., Sinclair, R. and Maghembe, J. A. (2006). Diversity of glomale mycorrhizal fungi in maize/sesbania intercrops and maize monocrop systems in Southern Malawi, Agroforestry systems, 67: 107-114.
- [18] Jones, B. J. J. (2001). *Laboratory Guide for Conducting Soil Tests and Plant Analysis*, CRC press, LLC.
- [19] Li, X. L., Marschner, H. and George, E. (1999). Extension of the phosphorus depletion zone in VA mycorrhizal white clover in a calcareous soil. *Plant and Soil*, Vol: 135, p: 41-48.
- [20] Koide, R. T. and Dickie, I. A. (2002). Effects of mycorrhizal fungi on plants populations. *Plant Soil* 244, 307-317. doi: 10.1023/A:1020204004844.
- [21] Louis, L. and Lim G. 1987. Spore density and root colonization of vesicular-arbuscular Mycorrhizas in tropical soil. *Trans. Br. Mycol. Soc.*, 88: 207-212.
- [22] Mahmud, R. M., Mridha, U. A., Osman, T. K., Xu, L. H. and Umemura, H. (1999). Relationship between edaphic factors and arbuscular mycorrhizal fungi in soils of rubber plantation. The 207th Annual Meeting of Japanese Society of Crop Science, April 2-3, 1999, Tokyo. *Jap. J. Crop Sci.*, 68 (Extra 1): 244-245.
- [23] Mehrotra, V. S. 1998. Arbuscular mycorrhizal association of plants colonizing coal mine spoils in India. *J. Agril. Sci., Cambridge*. 130: 123-133.
- [24] Millat-e-Mustafa, M. (1997). Tropical Homegardens: An overview. In: Alam, M. K., F. U. Ahmed and A. M. Ruhul Amin (eds.), *Agroforestry: Bangladesh Perspective*, BARC. Pp 18-33.
- [25] Olsen, S. R. and Dean L. A. (1965). "Phosphorous," in *Methods of Soil Analysis Part 2: Chemical and Microbiological Properties*, C. A. Black, Ed., vol. 9, pp. 1035–1049, American Society of Agronomy, Madison, Wis, USA.
- [26] Ortaş, İ. 2002. Do Plants depend on mycorrhizae in terms of nutrient requirement?, International conference on sustainable land use and management. Çanakkale.
- [27] Phillips, J. M. and Hayman D. S. (1970). Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. Br. Mycol. Soc.*, 55: 158–161.
- [28] Power M. E. and Mills L. S. (1995). The keystone cops meet in Hilo. *Tree* 10, 182-184.
- [29] Raintree, J. B. (1997). Agroforestry concepts. In: Alam, M. K.; F. U. Ahmed and A. M. Ruhul Amin (eds.), *Agroforestry: Bangladesh Perspective*, BARC. Pp 1-17.
- [30] Rhodes, L. H. (1980). The use of mycorrhizae in crop production systems. *Outlook on Agriculture*, 10 (6): 275 - 281.
- [31] Roy, I. (1997). Cropland Agroforestry: The experience of the village and farm forestry programme. In: Alam, M. K.; F. U. Ahmed and A. M. Ruhul Amin (eds.), *Agroforestry: Bangladesh Perspective*, BARC. pp. 98-11.
- [32] Schenck, N. C. and Pérez, Y. (1990). *Manual for the Identification of VA Mycorrhizal Fungi*. Synergistic-Publications, Gainesville, Florida.
- [33] Schnitzer, M. 1982. "Total carbon, organic matter and carbon," in *Methods of Soil Analysis-Part 2, Agronomy Monograph*, A. L. Page, R. H. Miller, and D. R. Keeney, Eds., vol. 9, pp. 539- 577, American Society of Agronomy, Madison, Wis, USA, 2nd edition.
- [34] Sieverding, E. 1991. Vesicular-arbuscular mycorrhiza management in tropical agro systems. German Technical cooperation (GTZ) Eschborn, pp 371. ISBN 3 88085 462 9.
- [35] Smith, S. E. and Read, D. J. (1997). *Mycorrhizal symbiosis*, 2nd edn. Academic Press, London, U. K.

- [36] Snoeck, D., Abolo, D. and Jagoret, P. (2010). Temporal changes in VAM fungi in the cocoa agroforestry systems of central Cameroon. *Agroforestry Syst.*, 78: 323–328.
- [37] Talukdar, N. C. and Germida, J. J. (1993). Occurrence and isolation of vesicular-arbuscular mycorrhizae in cropped field soils of Saskatchewan. *Can. J. Microbiol.*, 39: 576-586.
- [38] Verbruggen, E. and Toby Kiers, E. (2010). Evolutionary ecology of mycorrhizal functional diversity in agricultural systems. *Evolutionary Appl.*, 3: 547–560.
- [39] Wang, F. Y., Liu, R. J. and Lin, X. G. (2004). Arbuscular mycorrhizal status of wild plants in saline-alkaline soils of the yellow river delta. *Mycorrhiza*, 14, 133–137.
- [40] Zebene Asfaw, (2003). Tree species diversity, Top soil conditions and Arbuscular mycorrhizal Association in the Sidama Traditional agroforestry land use, southern Ethiopia, Doctorial Thesis Department of Forest management and products, SLU. *Acta Universitatis Sueciae. Silverstria*, p. 263.
- [41] Zerihun Belay, Fassil Asefa, and Mauritz V. (2013). Diversity and abundance of arbuscular mycorrhizal fungi associated with acacia trees from different land use systems in Ethiopia. *African Journal of Microbial Research*, 7 (48), pp. 550.