A variational method in the Sturm-Liouville problem with the Neumann and Dirichlet boundary values

Khapaeva Tatiana Mikhailovna^{1,2}

¹Department of Calculative Mathematic and Cybernatic, Moscow, Russia

²Moscow State Lomonosov's University, MSU, Moscow, Russia

Email address:

tmhapa@yahoo.com

To cite this article:

Khapaeva Tatiana Mikhailovna. A Variational Method in the Sturm-Liouville Problem with the Neumann and Dirichlet Boundary Values. *Applied and Computational Mathematics*. Vol. 3, No. 4, 2014, pp. 117-120. doi: 10.11648/j.acm.20140304.11

Abstract: A variational method for calculation of the eigenfunctions and eigenvalues in the Sturm-Liouville problem with the Neumann boundary values is offered. The method is based on a functional, which is introduced in this work. An appropriate numerical algorithm is developed. Calculations for the three potentials are produced: $\sin((x-\pi)^2/\pi)$, $\cos(4x)$ and the high not isosceles triangle. The method is applied to the Sturm-Liouville problem with the Dirichlet boundary values. Some suppositions about the inverse Sturm-Liouville problem are made.

Keywords: Sturm-Liouville Problem, Neumann Boundary Values, Dirichlet Boundary Values, Eigenfunctions, Eigenvalues, Variational Method, Functional, Inversed Sturm-Liouville problem, Algorithm

1. Introduction

The aim of this work is calculation of the eigenfunctions (EFs) $y_1, \dots, y_n$ and eigenvalues (EVs) $\lambda_1, \dots, \lambda_n$ in the Sturm-Liouville (SL) problem in the form of:

$$y'' - q(x)y = -\lambda y \quad (1)$$

with the Neumann boundary values:

$$y(0) = y'(\pi) = 0 \quad (2)$$

Several methods of calculation of the EFs and EVs in the Sturm-Liouville problem exist. The bibliography on this question you can find in [1,2]. In this work a variational method based on a functional is offered:

$$\int_0^\pi [y'' + (\lambda - q(x))y]^2 dx + \left| \sqrt{\int_0^\pi y^2 dx} - 1 \right| + \gamma \sum_{i=2}^{n-1} \left| \int_0^\pi y_i y dx \right| + \beta |y'(\pi)|$$

where γ and β are chosen sufficiently small.

2. The Main Point of the Variational Method for the Neumann Boundary Values

An algorithm is based on the well-known properties of the EFs and EVs:

1. orthogonality of the EFs:

$$\int_0^\pi y_i y_j dx = 0, \quad i \neq j \quad (3)$$

2. monotony and positivity of the EVs:

$$0 < \lambda_1 < \dots < \lambda_n < \lambda_{n+1} < \dots \quad (4)$$

3. normality condition for EFs:

$$\int_0^\pi y_n^2 dx = 1, \quad \forall n \quad (5)$$

Let's break the segment $[0, \pi]$ into k pieces. $h = \pi / (k-1)$, $y' = 0$ in view of the boundary values.

Let's suppose that the first $n-1$ EFs and EVs are known. Let's base the algorithm construction on the following statement:

Theorem. Let the functional be defined:

$$J[y^2, \dots, y^k, \lambda] = h \sum_{j=2}^{k-1} \left[\frac{y^{j-1} - 2y^j + y^{j+1}}{h^2} + (\lambda - q^j(x))y^j \right]^2 + \left| \sqrt{h \sum_{j=2}^k y^{j2} - 1} \right| + \gamma h \sum_{i=1}^{n-1} \left| \sum_{j=2}^k y_i^j y^j \right| + \beta \frac{|y^k - y^{k-1}|}{h} \quad (6)$$

where γ and β are sufficiently small.

If $\min_{y^2, \dots, y^k, \lambda} J[y^2, \dots, y^k, \lambda] = 0$ subject to $\lambda > \lambda_{n-1}$ and $k \rightarrow \infty$, then the function $\tilde{y} = [0, \tilde{y}^2, \dots, \tilde{y}^k]$ and the value $\tilde{\lambda}$, bringing this minimum, are the EF and the corresponding EV for the SL problem (1)–(2). At that, these EF and EV are different from the previous ones.

Proof. Let the functional (6) reach a zero minimum by $\tilde{y} = [0, \tilde{y}^2, \dots, \tilde{y}^k]$ and $\tilde{\lambda}$. I.e.,

$$h \sum_{j=2}^{k-1} \left[\frac{\tilde{y}^{j-1} - 2\tilde{y}^j + \tilde{y}^{j+1}}{h^2} + (\tilde{\lambda} - q^j(x))\tilde{y}^j \right]^2 + \left| \sqrt{h \sum_{j=2}^k \tilde{y}^{j2} - 1} \right| + \gamma h \sum_{i=1}^{n-1} \left| \sum_{j=2}^k \tilde{y}_i^j \tilde{y}^j \right| + \beta \frac{|\tilde{y}^k - \tilde{y}^{k-1}|}{h} = 0 \quad (7)$$

Let's direct k to ∞ . The sums over k in (7) by that transform into the integrals by the trapezium rule with a negligibly small remainder. The expression $\frac{\tilde{y}^k - \tilde{y}^{k-1}}{h}$ transforms into the difference statement of the first derivative, and then $\frac{\tilde{y}^{j-1} - 2\tilde{y}^j + \tilde{y}^{j+1}}{h^2}$ transforms into the difference statement of the second derivative. We derive:

$$\int_0^\pi [\tilde{y}'' + (\tilde{\lambda} - q(x))\tilde{y}]^2 dx + \left| \sqrt{\int_0^\pi \tilde{y}^2 dx - 1} \right| + \gamma \sum_{i=1}^{n-1} \left| \int_0^\pi y_i \tilde{y} dx \right| + \beta |\tilde{y}'(\pi)| = 0$$

The expression under the modulus is the normality condition for EFs, and the integrals under the sum are the scalar product of the EFs (3). Thereby, we derive: $\int_0^\pi [\tilde{y}'' + (\tilde{\lambda} - q(x))\tilde{y}]^2 dx = 0$ and $|\tilde{y}'(\pi)| = 0$, or $\tilde{y}'' + (\tilde{\lambda} - q(x))\tilde{y} = 0$ and $\tilde{y}'(\pi) = 0$.

I.e., $\tilde{y}$ meets the SL problem (1)–(2), and $\tilde{y}$ is a EF, and $\tilde{\lambda}$ is a corresponding EV.

Which required to be proved.

The distinction between the derived EF, EV and the

previous ones follows from monotony of the EVs (4) and the condition $\tilde{\lambda} > \lambda_{n-1}$.

Thereby, at the first step of our algorithm the following functional

$$J[y] = \int_0^\pi [y'' + (\lambda - q(x))y]^2 dx + \left| \sqrt{\int_0^\pi y^2 dx - 1} \right| + |y'(\pi)|$$

is being minimized in y and λ under the condition $\lambda > 0$. As a result we derive a EF y_l and a corresponding EV λ_l .

In order to derive the second pair of the EF and EV we'll minimize the functional

$$J[y] + \left| \int_0^\pi y_1 y dx \right|$$

in y and λ under the condition: $\lambda > \lambda_1$, etc.

The global minimum search was carried out by the Random search method (the procedure *Nminimize* of the famous software *Wolfram Mathematica*).

It will be observed, that not always one can at once find the zero minimum with a sufficient accuracy. In this case one has to move enough far away from the previous EV and then make sequentially a few steps until a given accuracy is achieved. At that, the step must be sufficiently small.

If the step isn't small enough, you can miss the proper minimum. The step was being selected experimentally. The distance from the previous EV was being selected experimentally too.

The described calculations were carried out for the potential $q(x) = \cos(4x)$ (Fig. 2) and asymmetrical functions $\sin((x-\pi)^2/\pi)$ (Fig. 1) and a not isosceles triangle, whose altitude is much longer then the base.

The results are tabulated in the end of the article.

In Fig. 3, 4, 5 you can see the first, second and the twentieth EFs for the not isosceles triangle.

In Fig. 6, 7, 8 you can see the first, second and the twentieth EFs for $\sin((x-\pi)^2/\pi)$.

In Fig. 9, 10, 11 you can see the first, second and the twentieth EFs for $\cos(4x)$.

The interval $[0, \pi]$ was partitioned in 100 pieces ($n=100$). By such conditions the minimum of the functional is of the order of 10^{-5} . Unfortunately, author's computational capability allows only first 20 EV and EF.

The following EVs one can easily calculate with the help of an asymptotic formula [2] (*Theorem 4.14*):

$$\lambda_n = \left(n + \frac{1}{2}\right)^2 + \frac{1}{\pi} \int_0^\pi q(t) dt - \frac{1}{\pi} \int_0^\pi q(t) \cos((2n+1)t) dt + O\left(\frac{1}{\pi}\right)$$

3. Briefly about the Variational Method for the Dirichlet boundary values

The variational method is useful for the Sturm-Liouville problem with the Dirichlet boundary values. In this case the base functional looks:

$$\int_0^\pi [y'' + (\lambda - q(x))y]^2 dx + \left| \sqrt{\int_0^\pi y^2 dx} - 1 \right| + \sum_{i=2}^{n-1} \left| \int_0^\pi y_i y dx \right|$$

The algorithm is constructed similarly. On the first step of our algorithm the following functional

$$J[y] = \int_0^\pi [y'' + (\lambda - q(x))y]^2 dx + \left| \sqrt{\int_0^\pi y^2 dx} - 1 \right|$$

is being minimized in y and λ under the condition $\lambda > 0$. The successive EF and EV are being calculated consequentially from the previous ones.

4. Briefly about the Variational Method for the Inverse Sturm-Liouville Problem

So, the variational method provides the ability to solve the direct Sturm-Liouville problem for the various boundary values.

Let's mention briefly the inversed Sturm-Liouville problem: to restore the potential $q(x)$ by the known EFs and Evs. According to *Theorem P* from [3], in order to determine a potential identically you must have two sets of Evs for the tasks with the different boundary values. For example, one task with the Dirichlet boundary values and another one with the Neumann boundary values.

Possibly, the variational method can help to solve the inversed Sturm-Liouville problem.

The problem is incorrect and can be solved by the regularization method, which was developed by Tikhonov A. N. [4].

5. Conclusions

It is significant, that in particular the asymmetrical potentials are of interest. Symmetrical potentials were considered earlier by different methods.

So, we have obtained not only theoretical computations, but the results of calculations for the concrete functions too.

This work demonstrates a use of a present-day software to obtain series of EFs and Evs.

The variational method can be useful in some problems of mathematical physics, based on the Sturm-Liouville problem. Maybe, not only for direct, but for inversed problems too.

6. Results

	$\cos(4x)$	$\sin((x-\pi)^2/\pi)$	triangle
λ_1	0.547	0.855	1.534
λ_2	2.102	3.139	10.936
λ_3	6.531	7.122	15.43
λ_4	11.391	13.163	19.947
λ_5	18.195	21.235	28.034
λ_6	26.95	31.282	38.101
λ_7	38.146	43.325	49.993
λ_8	51.714	57.349	63.714
λ_9	66.937	73.336	79.802
λ_{10}	84.155	91.268	97.824
λ_{11}	103.677	111.126	117.71
λ_{12}	125.503	132.896	138.932
λ_{13}	148.728	148.728	162.786
λ_{14}	174.439	182.068	188.007
λ_{15}	201.706	209.42	215.223
λ_{16}	230.988	238.578	244.455
λ_{17}	261.731	269.515	275.167
λ_{18}	294.457	302.199	307.895
λ_{19}	328.689	336.597	342.129
λ_{20}	364.896	372.673	377.849

Figure 1. $\sin((x-\pi)^2/\pi)$.

Figure 2. $\cos(4x)$.

Figure 3. The first EF for the not isosceles triangle.

Figure 4. The second EF for the not isosceles triangle.

Figure 5. The twentieth EF for the not isosceles triangle.

Figure 6. The first EF for $\sin((x-\pi)^2/\pi)$.

Figure 7. The second EF for $\sin((x-\pi)^2/\pi)$.

Figure 8. The twentieth EF for $\sin((x-\pi)^2/\pi)$.

Figure 9. The first EF for $\cos(4x)$.

Figure 10. The second EF for $\cos(4x)$.

Figure 11. The twentieth EF for $\cos(4x)$.

References

- [1] L. D. Akulenko and S.V. Nesterov "Jeffektivnyj metod issledovanija kolebanij sushhestvenno neodnorodnyh raspredelennyh system", Prikladnaja matematika i mehanika, vol. 61:3, 1997, pp. 466–478
- [2] A. Kirsch "An Introduction to the Mathematical Theory of Inverse Problems", 2nd ed, Applied Mathematical Sciences, vol. 120, Springer, 2011
- [3] Borg G. "Eine Umkehrung der Sturm-Liouvilleschen Eigenwertaufgabe. Bestimmung der Differentialgleichung durch die Eigenwerte", Acta Mathematica, vol. 78, 1946, pp. 1–96
- [4] A. N. Tikhonov and V. Ya. Arsenin, "Methods for Solving Ill-Posed Problems", Nauka, Moscow, 1986 [in Russian]