

General Distance Energies and General Distance Estrada Index of Random Graphs

Nan Gao

College of Science, Xi'an Shiyou University, Xi'an, China

Email address:

gaonankk@163.com

To cite this article:

Nan Gao. General Distance Energies and General Distance Estrada Index of Random Graphs. *Applied and Computational Mathematics*. Vol. 7, No. 3, 2018, pp. 173-179. doi: 10.11648/j.acm.20180703.24

Received: July 2, 2018; **Accepted:** July 31, 2018; **Published:** August 13, 2018

Abstract: In 2000s, Gutman and Güngör introduced the concept of distance energy and the distance Estrada index for a simple graph G respectively. Moreover, many researchers established a large number of upper and lower bounds for these two invariants. But there are only a few graphs attaining the equalities of those bounds. In this paper, however, the exact estimates to general distance energy are formulated for almost all graphs by probabilistic and algebraic approaches. The bounds to general distance Estrada index are also established for almost all graphs by probabilistic and algebraic approaches. The results of this paper generalize the results of the distance energy and distance Estrada of random graph.

Keywords: E-R Random Graph, General Distance Matrix, General Distance Energy, General Distance Estrada Index

1. Introduction

Let G be a simple connected graph with vertex set $\{v_1, v_2, \dots, v_n\}$. Denote by $A = A(G) \in \mathbb{R}^{n \times n}$ the adjacency matrix of G , and $\lambda_1, \lambda_2, \dots, \lambda_n$ are eigenvalues of $A(G)$. The distance between vertices v_i and v_j of G , denoted by d_{ij} , is defined to be the length (i.e., the number of edges) of the shortest path from v_i to v_j . The distance matrix of G , denoted by $D(G)$, is the $n \times n$ matrix whose (i, j) -entry is equal to d_{ij} ($i, j = 1, 2, \dots, n$), (see [1], [2]). Note that $d_{ii} = 0$, $i = 1, 2, \dots, n$. The eigenvalues of $D(G)$ are said to be the D -eigenvalues of G . Since $D(G)$ is a real symmetric matrix, the D -eigenvalues are real and can be ordered in non-increasing order, $\rho_1 \geq \rho_2 \geq \dots \geq \rho_n$. The Harary matrix $H(G)$ of G (see [3]), which can be regarded as a generalization of $D(G)$ and is initially called reciprocal distance matrix, is a $n \times n$ matrix whose (i, j) -entry is equal to $\frac{1}{d_{ij}}$ if $i \neq j$ and 0 otherwise. The eigenvalues of the Harary matrix $H(G)$ are denoted by $\beta_1, \beta_2, \dots, \beta_n$ and are said to be the H -eigenvalues of G . Since the Harary matrix is symmetric, its eigenvalues are real and can be ordered as

$$\beta_1 \geq \beta_2 \geq \dots \geq \beta_n.$$

For the Hückel molecular orbital approximation, the total π -electron energy in conjugated hydrocarbons is given by the sum of absolute values of the eigenvalues corresponding to the molecular graph G in which the maximum degree is not more than three in general. In 1970s, Gutman [4] extended the concept of energy to all simple graphs G , and defined the energy of G as

$$\mathcal{E}(G) = \sum_{i=1}^n |\lambda_i|.$$

In theoretical chemistry, the energy of a given molecular graph is related to the total π -electron energy [5] of the molecule represented by that graph. The graph energy has been studied extensively by many mathematicians and chemists, and many results have been obtained on this invariant of graphs (see [6]).

The distance energy, $D\mathcal{E}(G)$, of G is defined as

$$D\mathcal{E}(G) = \sum_{i=1}^n |\rho_i| \quad (1)$$

The concept of distance energy, Eq. (1), was recently

introduced [7]. This definition was motivated by the much older [8] and nowadays extensively studied [9, 10, 11, 12] graph distance energy.

In [13], Estrada introduced another graph-spectrum-based invariant of graphs, which was later called the Estrada index, defined as

$$EE(G) = \sum_{i=1}^n e^{\lambda_i}.$$

Since then, there were various applications of the Estrada index. Initially, it was used to quantify the degree of folding of long chain polymeric molecules, especially those of proteins [14, 15]. And later, a connection between $EE(G)$ and the concept of extended atomic branching was established [16], which was an attempt to apply $EE(G)$ in quantum chemistry. In addition, Estrada and Rodríguez-Velázquez showed that $EE(G)$ provides a measure of the centrality of complex networks [17, 18]. Recently, a information-theoretical application of $EE(G)$ was put forward. Carbó-Dorca endeavored to find connections between $EE(G)$ and the Shannon entropy [19].

The distance Estrada index of a connected graph G was introduced in [20] as

$$DEE(G) = \sum_{i=1}^n e^{\rho_i}.$$

Some upper and lower bounds for $DEE(G)$ were established in [20] and [21].

In 2010, Güngör introduced the Harary energy and Harary Estrada index of a graph [22]:

$$HE(G) = \sum_{i=1}^n |\beta_i|, \quad HEE(G) = \sum_{i=1}^n e^{\beta_i}.$$

The Erdős-Rényi random graph $\mathcal{G}_n(p)$, named after Erdős and Rényi [23], consists of all graphs on n vertices in which the edges are chosen independently with probability p , where p is a constant and $0 < p < 1$. Recently, there have been a lot of work on chemical indices of random graphs. In [24], Du et al. have considered the energy of Erdős-Rényi model $\mathcal{G}_n(p)$. They obtained that almost every graph $G_n(p)$ in $\mathcal{G}_n(p)$ satisfies

$$\mathcal{E}(G_n(p)) = \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \quad a.s. \quad (2)$$

In [25], Chen et al. have considered the Estrada index of Erdős-Rényi model $\mathcal{G}_n(p)$. They obtained that almost every graph $G_n(p)$ in $\mathcal{G}_n(p)$ satisfies

$$EE(G_n(p)) = e^{np} \left(e^{O(\sqrt{n})} + o(1) \right) \quad a.s. \quad (3)$$

Du et al. also presented exact estimates of Laplacian-energy like invariant, incidence energy, and distance energy, and a tight bound of signless Laplacian energy for almost all graphs (see [26]). They obtained that almost every graph $G_n(p)$ in $\mathcal{G}_n(p)$ satisfies

$$RDE(G_n(p)) = \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \quad a.s. \quad (4)$$

In [27], Shang established better lower and upper bounds to $DEE(G)$ for almost all graphs. He obtained that almost every graph $G_n(p)$ in $\mathcal{G}_n(p)$ satisfies

$$\begin{aligned} & \left[e^{np} (e^{n^2} + 1) + n - 2 \right] e^{-3O(\sqrt{n})} \\ & \leq DEE(G_n(p)) \\ & \leq \left[e^{n^2} (n - 1) + 1 \right] e^{-3O(\sqrt{n})} \quad a.s. \end{aligned} \quad (5)$$

Throughout this paper, following the term introduced in Bollobás's book [28]. Almost every (a.e. for short) graph $G_n(p)$ in $\mathcal{G}_n(p)$ has a certain property Q if the probability that $G_n(p)$ has Q converges to 1 as n becomes infinite. Sometimes, "almost all" can replace "almost every". An event in a probability space holds asymptotically almost surely (a.s. for short) if its probability goes to one as n tends to infinity. Evidently, almost every graph in $\mathcal{G}_n(p)$ has Q if the probability of random graphs satisfying Q converges almost surely.

Denote by $\Delta(G)$ the diameter of G . The diameter of a random graph $G_n(p)$ has the following properties.

Lemma 1 ([1]). Suppose that $p^2 n - 2 \log n \rightarrow \infty$ and $n^2(1-p) \rightarrow \infty$ as $n \rightarrow \infty$. Then

$$\Delta G_n(p) = 2 \quad a.s.$$

Since p is a constant with $0 < p < 1$ in this paper, it follows from Lemma 1 that a.e. graph $G_n(p)$ has diameter 2. The distance matrix of a random graph $G_n(p) \in \mathcal{G}_n(p)$ is denoted as

$$\mathbf{D} = \mathbf{D}(G_n(p)).$$

Recall that the diameter of a graph G is the greatest distance between two vertices of G . Let $\mathcal{G}'_n(p) \subseteq \mathcal{G}_n(p)$ be a subset containing all graphs with diameter 2 and $G_n(p) \in \mathcal{G}'_n(p)$. Evidently, the entries of $\mathbf{D}(G_n(p))$ satisfy the following [14]

$$\mathbf{D}(ij) = \begin{cases} 0, & \text{if } v_i = v_j; \\ 1, & \text{if } v_i \text{ and } v_j \text{ are adjacent;} \\ 2, & \text{if } v_i \text{ and } v_j \text{ are nonadjacent.} \end{cases}$$

In order to study the more general form of distance energy

of random graphs, we introduce the concept of general distance matrix of a random graph $RD_\alpha(G_n(p))$, where $G_n(p) \in \mathcal{G}'_n(p)$ and $\alpha \neq 0$ is a real number. The entries of $RD_\alpha(G_n(p))$ satisfy the following

$$RD_\alpha(ij) = \begin{cases} 0, & \text{if } v_i = v_j; \\ 1, & \text{if } v_i \text{ and } v_j \text{ are adjacent;} \\ 2^\alpha, & \text{if } v_i \text{ and } v_j \text{ are nonadjacent,} \end{cases} \quad (6)$$

Clearly, $RD_1(G_n(p)) = D(G_n(p))$. Similar to the definition of the Harary matrix $H(G)$ of G , we can give a definition of the Harary matrix $RD_{-1}(G_n(p))$ of a random graph $G_n(p) \in \mathcal{G}'_n(p)$.

The goal of this paper is to formulate the exact estimates to general distance energy and establish bounds to general distance Estrada index for almost all graphs constructed from Erdős-Rényi random graph model.

In Section 2, two exact estimates are formulated to $RD\mathcal{E}_\alpha(G_n(p))$ for almost all $G_n(p) \in \mathcal{G}_n(p)$, where $\mathcal{G}_n(p)$ is Erdős-Rényi model.

i If $\alpha > 0$, then

$$RD\mathcal{E}_\alpha(G_n(p)) = (2^\alpha - 1) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \text{ a.s.}$$

ii If $\alpha < 0$, then

$$RD\mathcal{E}_\alpha(G_n(p)) = (1 - 2^\alpha) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \text{ a.s.}$$

In Section 3, In Section 2, two asymptotically inequation are formulated to $RDEE_\alpha(G_n(p))$ for almost all $G_n(p) \in \mathcal{G}_n(p)$.

i If $\alpha > 0$, then

$$\begin{aligned} & \left[e^{np(2^\alpha-1)}(e^{n2^\alpha} + 1) + n - 2 \right] e^{-2^\alpha - (2^\alpha-1)O(\sqrt{n})} \\ & \leq RDEE_\alpha(G_n(p)) \\ & \leq \left[e^{n2^\alpha}(n-1) + 1 \right] e^{-2^\alpha - (2^\alpha-1)O(\sqrt{n})} \text{ a.s.} \end{aligned}$$

ii If $\alpha < 0$, then

$$\begin{aligned} & \left[e^{n2^\alpha} + n - 1 \right] e^{-2^\alpha + (1-2^\alpha)O(\sqrt{n})} \\ & \leq RDEE_\alpha(G_n(p)) \\ & \leq \left[e^{np(1-2^\alpha)}(e^{n2^\alpha} + 1) + (n-2)e^{n2^\alpha} \right] e^{-2^\alpha + (1-2^\alpha)O(\sqrt{n})} \text{ a.s.} \end{aligned}$$

2. General Distance Energies of Random Graphs

This section starts by the definition of the adjacency matrix

of the random graph and an important lemma which will play an important role in the proofs of our main results. General distance energies $RD\mathcal{E}_\alpha(G_n(p))$ for random graphs shall be estimated.

Let $A_n := A(G_n(p))$ denote the adjacency matrix of the random graph $G_n(p) \in \mathcal{G}_n(p)$. Apparently, $A(G_n(p))$ is a symmetric random matrix in which the diagonal entries are zeros while a_{ij} ($i < j$) is 1 or 0, with probability p or $1-p$ respectively. Evidently, the matrix (6) can be rewritten as

$$RD_\alpha(G_n(p)) = 2^\alpha(J_n - I_n) - (2^\alpha - 1)A_n, \quad (7)$$

where J_n is $n \times n$ matrix in which all entries equal 1 and I_n is the unit $n \times n$ matrix.

Lemma 2 (Fan Ky's inequality [29]). Let X, Y, Z be real symmetric matrices of order n such that $X + Y = Z$. Then

$$\mathcal{E}(X) + \mathcal{E}(Y) \geq \mathcal{E}(Z).$$

Theorem 1. Let $G_n(p) \in \mathcal{G}_n(p)$ and $\alpha \neq 0$ be a fix real number. Then general distance energies of $G_n(p) \in \mathcal{G}_n(p)$ can be given as follows.

i If $\alpha > 0$, then

$$RD\mathcal{E}_\alpha(G_n(p)) = (2^\alpha - 1) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \text{ a.s.}$$

ii If $\alpha < 0$, then

$$RD\mathcal{E}_\alpha(G_n(p)) = (1 - 2^\alpha) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \text{ a.s.}$$

Proof: Let $\mathcal{G}'_n(p) \subseteq \mathcal{G}_n(p)$ be a subset containing all graphs with diameter 2. In view of Lemma 1, it show to show our result for the graph $G_n(p) \in \mathcal{G}'_n(p)$. This problem can be explained according to the following proof. i If $\alpha > 0$, (7) is equivalent to

$$A_n = \frac{2^\alpha}{2^\alpha - 1}(J_n - I_n) - \frac{RD_\alpha}{2^\alpha - 1}. \quad (8)$$

The following results can be obtained from (7), (8) and Lemma 2.

$$\mathcal{E}(RD_\alpha) \leq 2^\alpha \mathcal{E}(J_n - I_n) + (2^\alpha - 1)\mathcal{E}(-A_n)$$

and

$$\mathcal{E}(-A_n) \leq \frac{2^\alpha}{2^\alpha - 1} \mathcal{E}(J_n - I_n) + \frac{1}{2^\alpha - 1} \mathcal{E}(-RD_\alpha).$$

Thus,

$$\begin{aligned} & (2^\alpha - 1)\mathcal{E}(A_n) - 2^\alpha \mathcal{E}(J_n - I_n) \leq \mathcal{E}(RD_\alpha) \\ & \leq (2^\alpha - 1)\mathcal{E}(-A_n) + 2^\alpha \mathcal{E}(J_n - I_n). \end{aligned}$$

It is easy to see that $\mathcal{E}(J_n - I_n) = 2(n-1)$. By (2), the following result can be obtained.

$$(2^\alpha - 1) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} - 2^{\alpha+1}(n-1) \leq \mathcal{E}(RD_\alpha) \leq (2^\alpha - 1) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} + 2^{\alpha+1}(n-1).$$

Consequently, a.e. graph $G_n(p)$ in $\mathcal{G}_n(p)$ satisfies

$$RD\mathcal{E}_\alpha(G_n(p)) = \mathcal{E}(RD_\alpha) = (2^\alpha - 1) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \text{ a.s.} \tag{9}$$

iii If $\alpha < 0$, (7) is equivalent to

$$RD_\alpha(G_n(p)) = 2^\alpha(J_n - I_n) + (1 - 2^\alpha)A_n \tag{10}$$

and

$$A_n = \frac{RD_\alpha}{1 - 2^\alpha} - \frac{2^\alpha}{1 - 2^\alpha}(J_n - I_n). \tag{11}$$

The following two results can be obtained from (9), (10) and Lemma 2.

$$\mathcal{E}(RD_\alpha) \leq 2^\alpha \mathcal{E}(J_n - I_n) + (1 - 2^\alpha)\mathcal{E}(A_n)$$

and

$$\mathcal{E}(A_n) \leq \frac{2^\alpha}{1 - 2^\alpha} \mathcal{E}(J_n - I_n) + \frac{1}{1 - 2^\alpha} \mathcal{E}(RD_\alpha).$$

Thus,

$$(1 - 2^\alpha)\mathcal{E}(A_n) - 2^\alpha \mathcal{E}(J_n - I_n) \leq \mathcal{E}(RD_\alpha) \leq (1 - 2^\alpha)\mathcal{E}(A_n) + 2^\alpha \mathcal{E}(J_n - I_n),$$

i.e.

$$(1 - 2^\alpha) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} - 2^{\alpha+1}(n-1) \leq \mathcal{E}(RD_\alpha) \leq (1 - 2^\alpha) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} + 2^{\alpha+1}(n-1).$$

Consequently, a.e. graph $G_n(p)$ in $\mathcal{G}_n(p)$ satisfies

$$RD\mathcal{E}_\alpha(G_n(p)) = \mathcal{E}(RD_\alpha) = (1 - 2^\alpha) \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \text{ a.s.}$$

This completes the proof.

It is easy to see that

$$RD\mathcal{E}_1(G_n(p)) = \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \text{ a.s.} = D\mathcal{E}(G_n(p)).$$

The results of Theorem 1 generalize the results of Du et al. [25] on the distance energy of random graph.

The concept of Harary matrix $H(G_n(p))$ of random graph can be introduced such as the Harary matrix $H(G)$ of G . The entries of $H(G_n(p))$ satisfy the following

$$H(ij) = RD_{-1}(ij) = \begin{cases} 0, & \text{if } v_i = v_j; \\ 1, & \text{if } v_i \text{ and } v_j \text{ are adjacent;} \\ 1/2, & \text{if } v_i \text{ and } v_j \text{ are nonadjacent,} \end{cases} \tag{12}$$

Clearly, $RD\mathcal{E}_{-1}(G_n(p))$ is Harary energy $H\mathcal{E}(G_n(p))$ for random graphs in $\mathcal{G}_n(p)$.

By Theorem 1, the estimator of Harary energy of a random graph is obtained for the first time.

Corollary 1. Let $G_n(p) \in \mathcal{G}_n(p)$. Then

$$H\mathcal{E}(G_n(p)) = \frac{1}{2} \left(\frac{8}{3\pi} \sqrt{p(1-p)} + o(1) \right) n^{3/2} \text{ a.s.}$$

3. General Distance Estrada Index of Random Graphs

In this section, general distance Estrada indices $RDEE_\alpha(G_n(p))$ for random graphs shall be estimated. Lemma 3 present the explicit information about the eigenvalues of $A_n(G)$.

Lemma 3 (Chen [25]). Let A_n is the adjacency matrix $A(G_n(p))$ of the random graph $G_n(p) \in \mathcal{G}_n(p)$. Then, the eigenvalues of $A_n(G)$ satisfies

$$\lambda_1(A_n) = np + O(\sqrt{n}) \text{ a.s.}$$

and for $i = 2, \dots, n$,

$$\lambda_i(A_n) = O(\sqrt{n}) \text{ a.s.}$$

The result of Lemma 4 will play an important role in the proofs of our main results.

Lemma 4 (Weyl's inequality [30]). Let X, Y and Z be all $n \times n$ real symmetric matrices such that $X = Y + Z$ Suppose that X, Y, Z have eigenvalues, respectively,

$$\lambda_1(X) \geq \lambda_2(X) \geq \dots \geq \lambda_n(X), \lambda_1(Y) \geq \dots \geq \lambda_n(Y),$$

$\lambda_1(Z) \geq \dots \geq \lambda_n(Z)$. Then for $i = 1, 2, \dots, n$ the following inequalities hold:

$$\lambda_i(Y) + \lambda_n(Z) \leq \lambda_i(X) \leq \lambda_i(Y) + \lambda_1(Z)$$

Theorem 2. Let $G_n(p) \in \mathcal{G}_n(p)$ and $\alpha \neq 0$ be a fix real number. Then general distance Estrada indices of

$G_n(p) \in \mathcal{G}_n(p)$ can be given as follows.

i If $\alpha > 0$, then

$$\begin{aligned} & \left[e^{np(2^\alpha-1)}(e^{n2^\alpha} + 1) + n - 2 \right] e^{-2^\alpha - (2^\alpha-1)O(\sqrt{n})} \\ & \leq RDEE_\alpha(G_n(p)) \\ & \leq \left[e^{n2^\alpha}(n-1) + 1 \right] e^{-2^\alpha - (2^\alpha-1)O(\sqrt{n})} \quad a.s. \end{aligned}$$

ii If $\alpha < 0$, then

$$\begin{aligned} & \left[e^{n2^\alpha} + n - 1 \right] e^{-2^\alpha + (1-2^\alpha)O(\sqrt{n})} \\ & \leq RDEE_\alpha(G_n(p)) \\ & \leq \left[e^{np(1-2^\alpha)}(e^{n2^\alpha} + 1) + (n-2)e^{n2^\alpha} \right] e^{-2^\alpha + (1-2^\alpha)O(\sqrt{n})} \quad a.s. \end{aligned}$$

Proof: The general distance Estrada index $RDEE_\alpha(G_n(p))$ can be evaluated once the eigenvalues of RD_n are known.

i If $\alpha > 0$, applying lemma 4 to (7)

$$\begin{aligned} & 2^\alpha \lambda_i(J_n - I_n) + (2^\alpha - 1)\lambda_n(-A_n) \\ & \leq \lambda_i(RD_\alpha) \\ & \leq 2^\alpha \lambda_i(J_n - I_n) + (2^\alpha - 1)\lambda_1(-A_n) \end{aligned}$$

can be obtained.

That is

$$\begin{aligned} & 2^\alpha \lambda_i(J_n - I_n) - (2^\alpha - 1)\lambda_1(A_n) \\ & \leq \lambda_i(RD_\alpha) \\ & \leq 2^\alpha \lambda_i(J_n - I_n) - (2^\alpha - 1)\lambda_n(-A_n). \\ & 2^\alpha(n-1) - (2^\alpha - 1)(np + O(\sqrt{n})) \\ & \leq \lambda_i(RD_\alpha) \\ & \leq 2^\alpha(n-1) - (2^\alpha - 1)O(\sqrt{n}) \end{aligned} \tag{13}$$

and

$$\begin{aligned} & -2^\alpha - (2^\alpha - 1)(np + O(\sqrt{n})) \\ & \leq \lambda_n(RD_\alpha) \\ & \leq -2^\alpha - (2^\alpha - 1)O(\sqrt{n}) \end{aligned} \tag{14}$$

can be deduced from Lemma 3 and $\lambda_1(J_n - I_n) = n - 1$,

$\lambda_i(J_n - I_n) = -1$ for $i = 2, \dots, n$.

Owing to Lemma 4 again,

$$\begin{aligned} & 2^\alpha \lambda_n(J_n - I_n) - (2^\alpha - 1)\lambda_{n+1-i}(A_n) \\ & \leq \lambda_i(RD_\alpha) \\ & \leq 2^\alpha \lambda_1(J_n - I_n) - (2^\alpha - 1)\lambda_{n+1-i}(A_n) \end{aligned}$$

is very easy to get.

Consequently,

$$\begin{aligned} & -2^\alpha - (2^\alpha - 1)O(\sqrt{n}) \\ & \leq \lambda_i(RD_\alpha) \\ & \leq 2^\alpha(n-1) - (2^\alpha - 1)O(\sqrt{n}) \end{aligned} \tag{15}$$

for $i = 2, \dots, n - 1$.

Combining (13), (14) and (15) the conclusion can be proved. That is

$$\begin{aligned} RDEE_\alpha(G_n(p)) &= \sum_{i=1}^n e^{\lambda_i(RD_\alpha)} \\ &\geq e^{2^\alpha(n-1) - (2^\alpha-1)(np+O(\sqrt{n}))} + \\ & \quad (n-2)e^{-2^\alpha - (2^\alpha-1)O(\sqrt{n})} + \\ & \quad e^{-2^\alpha - (2^\alpha-1)(np+O(\sqrt{n}))} \\ &= \left[e^{np(2^\alpha-1)}(e^{n2^\alpha} + 1) + n - 2 \right] e^{-2^\alpha - (2^\alpha-1)O(\sqrt{n})} \quad a.s. \end{aligned}$$

and

$$\begin{aligned} RDEE_\alpha(G_n(p)) &= \sum_{i=1}^n e^{\lambda_i(RD_\alpha)} \\ &\leq e^{2^\alpha(n-1) - (2^\alpha-1)O(\sqrt{n})} + (n-2)e^{2^\alpha(n-1) - (2^\alpha-1)O(\sqrt{n})} + \\ & \quad e^{-2^\alpha - (2^\alpha-1)O(\sqrt{n})} \\ &= \left[e^{n2^\alpha}(n-1) + 1 \right] e^{-2^\alpha - (2^\alpha-1)O(\sqrt{n})} \quad a.s. \end{aligned}$$

ii If $\alpha < 0$, applying lemma 4 to (7)

$$\begin{aligned} & 2^\alpha \lambda_i(J_n - I_n) + (1 - 2^\alpha)\lambda_n(A_n) \\ & \leq \lambda_i(RD_\alpha) \\ & \leq 2^\alpha \lambda_i(J_n - I_n) + (1 - 2^\alpha)\lambda_1(A_n) \end{aligned}$$

can be obtained.

Then,

$$\begin{aligned} & 2^\alpha(n-1) + (1 - 2^\alpha)O(\sqrt{n}) \\ & \leq \lambda_1(RD_\alpha) \\ & \leq 2^\alpha(n-1) + (1 - 2^\alpha)(np + O(\sqrt{n})) \end{aligned} \tag{16}$$

and

$$\begin{aligned} & -2^\alpha + (1 - 2^\alpha)O(\sqrt{n}) \\ & \leq \lambda_n(RD_\alpha) \\ & \leq -2^\alpha + (1 - 2^\alpha)(np + O(\sqrt{n})) \end{aligned} \tag{17}$$

Owing to Lemma 4 again,

$$\begin{aligned} & 2^\alpha \lambda_n(J_n - I_n) + (1 - 2^\alpha)\lambda_{n+1-i}(A_n) \\ & \leq \lambda_i(RD_\alpha) \\ & \leq 2^\alpha \lambda_1(J_n - I_n) + (1 - 2^\alpha)\lambda_{n+1-i}(A_n) \end{aligned}$$

is very easy to get.

Consequently,

$$\begin{aligned} & -2^\alpha + (1-2^\alpha)O(\sqrt{n}) \\ & \leq \lambda_i(RD_\alpha) \\ & \leq 2^\alpha(n-1) + (1-2^\alpha)O(\sqrt{n}) \end{aligned} \tag{18}$$

for $i = 2, \dots, n-1$.

Combining (16), (17) and (18) the conclusion can be proved. That is

$$\begin{aligned} RDEE_\alpha(G_n(p)) &= \sum_{i=1}^n e^{\lambda_i(RD_\alpha)} \\ &\geq e^{2^\alpha(n-1) + (1-2^\alpha)O(\sqrt{n})} + \\ & (n-2)e^{-2^\alpha + (1-2^\alpha)O(\sqrt{n})} + e^{-2^\alpha + (1-2^\alpha)O(\sqrt{n})} \\ &= \left[e^{n2^\alpha} + n-1 \right] e^{-2^\alpha + (1-2^\alpha)O(\sqrt{n})} \quad a.s. \end{aligned}$$

and

$$\begin{aligned} RDEE_\alpha(G_n(p)) &= \sum_{i=1}^n e^{\lambda_i(RD_\alpha)} \\ &\leq e^{2^\alpha(n-1) + (1-2^\alpha)(np+O(\sqrt{n}))} + (n-2)e^{2^\alpha(n-1) + (1-2^\alpha)O(\sqrt{n})} + \\ & e^{-2^\alpha + (1-2^\alpha)(np+O(\sqrt{n}))} \\ &= \left[e^{np(1-2^\alpha)}(e^{n2^\alpha} + 1) + (n-2)e^{n2^\alpha} \right] e^{-2^\alpha + (1-2^\alpha)O(\sqrt{n})} \quad a.s. \end{aligned}$$

This completes the proof.

It is easy to see that

$$\begin{aligned} & \left[e^{np}(e^{n2} + 1) + n-2 \right] e^{-3O(\sqrt{n})} \\ & \leq RDEE_1(G_n(p)) = DEE(G_n(p)) \\ & \leq \left[e^{n2}(n-1) + 1 \right] e^{-3O(\sqrt{n})} \quad a.s. \end{aligned}$$

The results of Theorem 2 generalize the results of Shang [27] on the distance Estrada index of random graph.

Clearly, $RDEE_{-1}(G_n(p))$ is Harary Estrada index

$HEE(G_n(p))$ for random graphs in $\mathcal{G}_n(p)$. By Theorem 2, the asymptotically inequation of Harary Estrada index of a random graph is obtained for the first time.

Corollary 2. Let $G_n(p) \in \mathcal{G}_n(p)$. Then

$$\begin{aligned} & \left[e^{\frac{1}{2}} + n-1 \right] e^{-\frac{1}{2} + \frac{1}{2}O(\sqrt{n})} \\ & \leq HEE(G_n(p)) \\ & \leq \left[e^{\frac{np}{2}}(e^{\frac{n}{2}} + 1) + (n-2)e^{\frac{n}{2}} \right] e^{-\frac{1}{2} + \frac{1}{2}O(\sqrt{n})} \quad a.s. \end{aligned}$$

4. Conclusion

In 1970s, Gutman introduced the concept of energy $\mathcal{E}(G)$ for a simple graph G , which is defined as the sum of the

absolute values of the eigenvalues of G and can be used to estimate the total π -electron energy in conjugated hydrocarbons. The concept attracted lots of attention and furthermore, some other similar notions were also considered such as Laplacian energy $LE(G)$, signless Laplacian energy, $LE^+(G)$, incidence energy $IE(G)$, and distance energy $DE(G)$. Moreover, many researchers established a large number of upper and lower bounds for those invariants. But there are only a few graphs attaining the equalities of those bounds. In the present paper, however, we present exact estimates of general distance energy for almost all graphs by probabilistic and algebraic approaches.

In spite of the fact that the Estrada index of G , $EE(G)$ has numerous practical applications, investigations of its basic properties started only short time ago. The concept attracted lots of attention and furthermore, some other similar notions were also considered such as Laplacian

Estrada index $LEE(G)$, signless Laplacian Estrada index $LEE^+(G)$ incidence Estrada index $IEE(G)$, and distance Estrada index $DEE(G)$. It is rather hard, as well-known, to compute the eigenvalues for a large matrix even for $A(G)$. So, in order to estimate these invariants, researchers established some lower and upper bounds by algebraic approaches in the last few years. However, there are, as examples given below, only a few classes of graphs attaining the equalities of those bounds. Consequently, one can hardly see the major behavior of these invariants for most graphs. In this paper, however, we establish bounds to general distance Estrada index for almost all graphs by probabilistic and algebraic approaches.

Acknowledgements

We wish to thank the referee for valuable suggestion and advice, which led to an improvement of this paper.

This paper supported by the Scientific Research Program Funded by Shaanxi Provincial Education Department (No.17JK0613).

References

- [1] F. Buckley, F. Harary, "Distance in Graphs", Addison-Wesley, Redwood, 1990.
- [2] D. M. Cvetkovi, M. Doob, H. Sachs, "Spectra of Graphs-Theory and Application", Academic Press, New York, 1980.
- [3] O. Ivanciuc, T. S. Balaban, A. T. Balaban, Design of topological indices. Part 4. Reciprocal distance matrix, related local vertex invariants and topological indices, J. Math. Chem. 12 (1993), pp.309-318.
- [4] I. Gutman, The energy of a graph, Ber. Math. Statist. Sect. Forschungsz. Graz, 103 (1978), 1-22.
- [5] K. Yates, Hückel Molecular Orbital Theory, Academic Press, New York, 1978.

- [6] X. Li, Y. Shi, I. Gutman, Graph Energy, Springer, New York, 2012.
- [7] G. Indulal, I. Gutman, A. Vijaykumar, On the distance energy of a graph, MATCH Commun. Math. Comput. Chem., 60 (2008), pp. 461–472.
- [8] Gutman I. The Energy of a Graph: Old and New Results [M]// Algebraic Combinatorics and Applications. Springer Berlin Heidelberg, 2001, pp.196-211.
- [9] Díaz R C, Rojo O. Sharp upper bounds on the distance energies of a graph. Linear Algebra and Its Applications, 545 (2018), pp: 55-75.
- [10] So W. A shorter proof of the distance energy of complete multipartite graphs. Special Matrices, 5 (2017), pp:61-63.
- [11] Andjelic M, Koledin T, Stanic Z. Distance spectrum and energy of graphs with small diameter. Applicable Analysis & Discrete Mathematics, 11 (2017), pp: 108-122.
- [12] G. Indulal, Sharp bounds on the distance spectral radius and the distance energy of graphs, Linear Algebra Appl., 430 (2009), pp. 106–113.
- [13] E. Estrada, Characterization of 3D molecular structure, Chem. Phys. Lett., 319 (2000), pp. 713–718.
- [14] E. Estrada, Characterization of the folding degree of proteins, Bioinformatics 18 (2002), pp: 697–704.
- [15] E. Estrada, Characterization of the amino acid contribution to the folding degree of proteins, Proteins 54 (2004), pp: 727–737.
- [16] E. Estrada, J. A. Rodríguez-Velázquez, M. Randić, Atomic branching in molecules, Int. J. Quantum Chem. 106 (2006), pp: 823–832.
- [17] E. Estrada, J. A. Rodríguez-Velázquez, Subgraph centrality in complex networks, Phys. Rev. E71 (2005) 056103-1-9.
- [18] E. Estrada, J. A. Rodríguez-Velázquez, Spectral measures of bipartivity in complex networks, Phys. Rev. E72 (2005) 046105-1-6.
- [19] R. Carbó-Dorca, Smooth function topological structure descriptors based on graph-spectra, J. Math. Chem. 44 (2008), pp: 373–378.
- [20] A. D. Güngör, Ş. B. Bozkurt, On the distance Estrada index of graphs, Hacettepe J. Math. Stat., 38 (2009), pp.277–283.
- [21] Ş. B. Bozkurt, D. Bozkurt, Bounds for the distance Estrada index of graphs. AIP Conf. Proc., 1648 (2015), pp. 1351–1359.
- [22] Güngör A D, Sinan A. On the Harary energy and Harary Estrada index of a graph. MATCH Commun. Math. Comput. Chem., 64 (2010), pp: 270-285.
- [23] P. Erdős, A. Rényi, On random graphs I, Publ. Math. Debrecen., 6 (1959), pp. 290–297.
- [24] W. Du, X. Li, Y. Li, The energy of random graphs, Linear Algebra Appl., 435 (2009), pp. 2334–2346.
- [25] Z. Chen, Y. Fan, W. Du, Estrada index of random graphs, MATCH Commun. Math. Comput. Chem., 68 (2012), pp. 825–834.
- [26] W. Du, X. Li, Y. Li, Various energies of random graphs, MATCH Commun. Math. Comput. Chem., 64 (2010), pp.251–260.
- [27] Y. Shang, Distance Estrada index of random graphs, Linear Multilinear Algebra, 63 (2015), pp. 466–471.
- [28] B. Bollobás, Random Graphs, Cambridge Univ. Press, Cambridge, 2001.
- [29] K. Fan, Maximum properties and inequalities for the eigenvalues of completely continuous operators, Proc. Natl. Acad. Sci. USA 37 (1951), pp. 760–766.
- [30] H. Weyl, Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen, Math. Ann., 71 (2010), pp. 441–479.