

GC/GC-MS Analysis, Isolation and Identification of Bark Essential Oil Components from *Cinnamomum culilawan*, Blume

Hanoch Julianus Sohilait¹, Healthy Kainama²

¹Departement of Chemistry, Faculty of Mathematic and Natural Science, Pattimura University, Ambon, Indonesia

²Departement of Chemistry Education, Faculty of Training and Education, Pattimura University, Ambon, Indonesia

Email address:

nokesohilait@yahoo.com (H. J. Sohilait)

To cite this article:

Hanoch Julianus Sohilait, Healthy Kainama. GC/GC-MS Analysis, Isolation and Identification of Bark Essential Oil Components from *Cinnamomum culilawan*, Blume. *American Journal of Applied Chemistry*. Vol. 4, No. 4, 2016, pp. 157-160.

doi: 10.11648/j.ajac.20160404.16

Received: July 1, 2016; **Accepted:** July 13, 2016; **Published:** August 2, 2016

Abstract: The essentials oils (EO) in *Cinnamomum culilawan* bark have been isolated by steam distillation and evaluated by GC/GC-MS, IR and NMR methods. The GC/GC-MS analysis identified 12 compounds. The major components are eugenol (66.477%), safrole (12.432%) and methyleugenol (6.972%). Repeated purification of the two main compounds yielded eugenol (51.66%), and safrole (6.71%).

Keywords: *Cinnamomum culilawan*, Bark Essential Oil, Chemical Composition, Eugenol, Safrole

1. Introduction

The lawang tree, *Cinnamomum culilawan* (Lauraceae), grows naturally in Moluccas Islands, and west Papua, East Indonesia [1, 2]. The local name of this tree in Moluccas Islands is *lawang*; the trunk of tree can grow as high as 12-20 meters, on altitudes of 300 m above sea level. The *culilawan* oil is produced through steam distillation of the tree bark and the oil of *culilawan* is usually used as a pain killer for aches and pains in Moluccas Islands. Also, the *lawang* bark extract is used for tradisional medicine such as tooth aches, muscular pains and stomach aches in Papua [3]. While many plants from some genera have been studied and some even commercially explored for essential oils, some of those are *Cinnamomum sintoc* [4, 5], *Cinnamomum carolinense* [6] and *Cinnamomum bodinieri* [7], the species *Cinnamomum culilawan* has been rarely investigated. Isolation of safrole from *culilawan* oil from other places in West Seram, Moluccas and Papua has been reported [8, 9], but the chemical constituent of this oil not presented in those papers yet.

This paper reports results of chemical composition of *culilawan* oil obtained from original trees grown in Ambon Island.

2. Materials and Methods

2.1. Materials

The barks of lawang three were collected from Allang Village, Ambon Island, Moluccas in January 2016 and were deposited in the Organic Chemistry Laboratory, Pattimura University, Ambon. The raw material was air-dried until constant weight and chopped into small pieces.

2.2. Chemicals and Equipment

The chemicals in the study are: sodium hydroxide p.a (E. Merck), petroleum ether 35-60°C p.a (J. T. Baker), anhydrous sodium sulfate p.a (E. Merck), sodium chloride p.a (E. Merck), hydrochloric acid 30% p.a (E. Merck). The tools used in this study are, a set of fractional distillation under reduced pressure, electric heaters, evaporators and tools Buchi laboratory glassware, Gas Chromatography GC-2010, Shimadzu, an Infra Red spectrophotometer (FTIR-8400S, Shimadzu), ¹H-NMR, Spectrophotometer (JEOL-MY 400, MHz), ¹³C-NMR, Spectrophotometer (JEOL-MY 100, MHz), MassSpectrophotometer (GC-MS QP-2010 Plus, Shimadzu).

2.3. Isolation of Essential Oil from Bark of *Cinnamomum culilawan*, Bl

The dry plant material (2.0 kg) from bark of *Cinnamomum culilawan* was steam-distilled using conventional steam distiller for 6 hours to obtain the essential oil. The essential oil were then dried by anhydrous sodium sulfate (Na_2SO_4) and cool-stored in brown bottle for further analysis.

2.4. Gas Chromatography (GC) Analysis

GC analysis of the *culilawan* oil was performed on a Shimadzu QP-2010, equipped with a FID and Rtx-5 using a fused silica capillary column (30 m x 0.25 mm ID, film thickness 1.0 μm). Oven temperature was from 60°C for 5 minutes and programmed heating from 60 to 180°C at a rate of 20°C for 5 minutes, and from 180°C for 4 minutes, injector temperature 270°C, detector temperature 280°C, pressure of carrier nitrogen gas at inlet 7 psi, split 20 and volume was 0.5 μL .

2.5. Gas Chromatography-Mass Spectrometry (GC-MS)

GC-MS analyses of bark *culilawan* oil were performed using Shimadzu QP-2010 Plus at 70 eV and 320°C with auto sampler, system equipped with Rtx-5 fused silica capillary column (30 m x 0.25 mm ID, film thickness 1.0 μm). Oven temperature was from 60°C for 5 minutes, then programmed heating from 60 to 220°C at a rate of 10°C for 5 minutes, and from 220°C for 4 min, injector temperature 270°C. The carrier gas was helium at flow rate of 1.40 mL/min. Spectra were scanned from 40 to 600 m/z , split ratio 20, ion source temperature 225°C. The spectrum of the unknown component was compared with that of known components

3.2. Chemical Composition of Bark Essential Oil of *Cinnamomun culilawan*

Table 1. Chemical composition of bark essential oils of *Cinnamomun culilawan*, Bl.

Peak No.	Retention time (min)	Compound Name	M. W	Formula	Conc(%) ^{a)}	Base peaks	Main Fragment ion (m/z)
1	5.919	α -thujene	136	$\text{C}_{10}\text{H}_{16}$	0.516	93	41, 65, 77, 93, 136
2	6.083	α -pinene	136	$\text{C}_{10}\text{H}_{16}$	1.999	93	41, 67, 93, 121, 136
3	6.942	β -pinene	136	$\text{C}_{10}\text{H}_{16}$	0.631	93	41, 69, 79, 93, 121, 136
4	7.733	o-cymene	134	$\text{C}_{10}\text{H}_{14}$	1.590	119	65, 77, 91, 119, 134
5	7.800	Limonena	136	$\text{C}_{10}\text{H}_{16}$	0.856	93	41, 53, 68, 79, 93, 107, 121, 136
6	7.850	1,8-cineol	154	$\text{C}_{10}\text{H}_{18}\text{O}$	2.431	43	41, 43, 69, 81, 84, 108, 139, 154
7	8.758	Linalool	154	$\text{C}_{10}\text{H}_{18}\text{O}$	4.852	71	41, 43, 69, 71, 93, 121, 136
8	9.700	4-terpineol	154	$\text{C}_{10}\text{H}_{18}\text{O}$	0.651	71	41, 43, 71, 93, 111, 154
9	10.775	Safrole ^{b)}	162	$\text{C}_{10}\text{H}_{10}\text{O}_2$	12.432	162	51, 77, 104, 131, 162
10	11.450	Eugenol ^{b)}	164	$\text{C}_{10}\text{H}_{12}\text{O}_2$	66.477	164	55, 77, 91, 103, 131, 149, 164
11	11.700	Methyleugenol	178	$\text{C}_{11}\text{H}_{14}\text{O}_2$	6.972	178	41, 65, 77, 91, 107, 135, 147, 163, 178
12	12.908	Myristicin	192	$\text{C}_{11}\text{H}_{12}\text{O}_3$	0.592	192	65, 77, 91, 119, 131, 147, 161, 192

a) Concentration of compounds base on GC-FID peaks (see Figure 1) and the isolated of compounds were highlighted in bold face.

b) Identified by FTIR, $^1\text{H-NMR}$, $^{13}\text{C-NMR}$ and MS

The essential oil, Bl was analyzed by GC and GC-MS. Twelve components were identified and quantified, as shown in Table 1, according to their elution order on the Rtx-5 column. The major components are safrole (12.432%), eugenol (66.477%) and methyleugenol (6.972%), (Figure 1, Table 1).

stored in the Wiley 7 library. Name, molecular weight and structure of the component of the test materials were ascertained.

2.6. Isolation of Two Main Components from *Cullilawan* Oil

In a 500 mL flask, NaOH (20.0 g), aquades (150 mL) and *culilawan* oil (100.0 g) were charged and the mixture were stirred until forming two layers. The upper layer (A) was separated from the bottom layer (B). The upper layer was extracted with 25 mL NaOH 20% and aqueous layer was combined to bottom layer (B). The organic layer (A) was washed with water until neutral and dried with Na_2SO_4 anhydrous and the residue was distilled under reduced pressure at 15 mmHg/120°C. The bottom layer (B) was acidified with HCl 20% until pH reached 3 and extracted twice with 100 mL petroleum ether. The organic layer was washed until neutral and dried over anhydrous Na_2SO_4 . After removing the petroleum ether, the residue was distilled under reduced pressure at 15 mmHg/140°C. The purity of compounds was tested by GC and the elucidation of the structures employed FTIR, $^1\text{H-NMR}$, $^{13}\text{C-NMR}$ and GC-MS methods.

3. Results and Discussion

3.1. Performance of Bark Essential Oil of *Cinnamomun culilawan*, Bl.

The bark essential oils of *C. culilawan* was distilled from dry plant material on equipment of the steam distillation. *Culilawan* oil obtained from steam-distillation revealed that oil from the bark have a yield of 4.12% w/w.

The minor constituents of the oil included the monoterpenes hydrocarbon (5.592%), oxygenated monoterpenes (7.934%). Between the minor constituents of the oil, the oxygenated monoterpenes linalool (4.852%) and 1,8-cineol (2.431%) were the most representatives, totaling other 7.283% of the oil. The phenylpropanoid compounds

(86.473%) with myristicin did at low concentration (0.592%).

Due to very limited amount of published data on *Cinnamomum culilawan*, BI essential oil, it is difficult to compare results of this study with essential oil composition of plant grown in other countries.

3.3. Isolation of Two Main Components from Culilawan Oil

The percent contents of the EO were determined on the basis of their FID responses upon GC (Figure 1). Eugenol (1) was the major component (66.477%) of the EO (Figure 1, Table 1) followed by Safrole (12.432%), and methyleugenol (6.972%).




Figure 1. GC-FID peaks for% oil and the structure for isolated compounds (1 and 2) from the bark essential oil of *Cinnamomum culilawan*.

The isolation of eugenol (1) from culilawan oil is effected by using sodium hydroxide solution to obtain sodium eugenolate followed by regeneration with 20% hydrochloric acid. The mixture was extracted with petroleum ether and the solution was washed with water until neutral and dried over anhydrous Na_2SO_4 . After removing the petroleum ether, the residue was distilled under reduced pressure at 15 mmHg/140°C yield (51.66%). Spectrum IR (cm^{-1}): 3475, 2976, 2842, 1638, 1613, 1432, 1149, 1034. Spectrum $^1\text{H-NMR}$ (400 MHz, CDCl_3 , ppm), δ : 3.41 (d $-\text{CH}_2-$, $J = 6.8$ Hz), 3.86 (s $-\text{OCH}_3$), 5.16 (d, $=\text{CH}_2$), 5.20 (s, $-\text{OH}$), 6.05 (m, $-\text{CH}=\text{C}$, $J = 6.8$ Hz), 6.76 (s, H, C3Ar), 6.78 (d, H, C6Ar), 6.98 (d, H5-Ar). $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ : 40.01 ($-\text{CH}_2-$), 55.95 ($-\text{OCH}_3$), 111.20 ($=\text{CH}_2$), 114, 37 (C2-Ar), 115.65 (C5-Ar), 121.26 (C6-Ar), 132.03 (C1-Ar), 137.95 ($-\text{CH}=\text{C}$), 143.95 (C4-Ar), 146.53 (C3-Ar). The spectral data matched that given in a previous report [10]. The Infra-red spectrum of eugenol (1) showed absorption band in the 3475 cm^{-1} ($-\text{OH}$), in the region 2976 cm^{-1} which is the absorption $\text{Csp}^3\text{-H}$, this was confirmed by the appearance of absorption at 1432 cm^{-1} for $-\text{CH}_2-$ (methylene). The range of $\text{C}=\text{C}$ aliphatic absorption appeared at 1638 cm^{-1} , absorption at 1613 cm^{-1} for $\text{C}=\text{C}$ aromatic and supported by absorption at $2976\text{-}2842 \text{ cm}^{-1}$ which is absorption band for $=\text{Csp}^2\text{-H}$ (aliphatic/aromatic). Absorption band at 1149 cm^{-1} and 1034 cm^{-1} region showed the range of C-O-C (ether). The $^1\text{H-NMR}$ spectrum, signal doublet at 3.41 ppm ($-\text{CH}_2-$, $J = 6.8$ Hz) and signal multiplet

at 6.06 ppm ($-\text{CH}=\text{C}$, $J = 6.8$ Hz), signal singlet at 3.86 ppm of one methoxy group. The $^{13}\text{C-NMR}$ spectrum showed 10 nonequivalent carbon resonances and the MS spectrum showed a molecular ion peak $[\text{M}^+]$ at m/z 164, $\text{C}_{10}\text{H}_{12}\text{O}_2$ (see table 1).

The upper layer containing safrole (2) was processed as follows; the solution was washed with water until neutral followed by drying over anhydrous Na_2SO_4 . After removing petroleum ether, the residue was distilled under reduced pressure 15 mmHg/120°C yield (6.71%). Spectrum IR (cm^{-1}): 2977, 2842, 1639, 1608, 1432, 1246, 1034. Spectrum $^1\text{H-NMR}$ (400 MHz, CDCl_3 , ppm), δ : 3.33 (d $-\text{CH}_2-$, $J = 6.4$ Hz), 5.10 (m, $=\text{CH}_2$), 5.95 (s, $-\text{OCH}_2\text{O}-$), 6.00 (m, $-\text{CH}=\text{C}$, $J = 6.4$ Hz), 6.67 (d, H-C5Ar), 6.74 (s, C3-Ar), 6.84 (d, H-C6-Ar). $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ : 40.06 ($-\text{CH}_2-$), 100.95 ($-\text{OCH}_2\text{O}-$), 108.30 (C3-Ar), 109.23 (C5-Ar), 115.82 ($=\text{CH}_2$), 121.44 (C6-Ar), 133.96 (C1-Ar), 137.80 ($-\text{CH}=\text{C}$), 145.99 (C4-Ar), 147.90 (C3-Ar). The spectral data matched that given in a previous report [11, 12].

The Infra red spectrum of safrole (2) showed absorption band in the region 2977 cm^{-1} which is the absorption $\text{Csp}^3\text{-H}$. This was confirmed by the appearance of absorption at 1432 cm^{-1} for $-\text{CH}_2-$ (methylene). Untake range of $\text{C}=\text{C}$ aliphatic absorption appeared at 1639 cm^{-1} , absorption at 1608 cm^{-1} for $\text{C}=\text{C}$ aromatic and supported by absorption at $2977\text{-}2842 \text{ cm}^{-1}$ which is absorption band for $=\text{Csp}^2\text{-H}$ (aliphatic/aromatic). Absorption band at 1246 cm^{-1} and 1034 cm^{-1} region showed

the range of C-O-C (ether). The ¹H-NMR spectrum, signal doublet at 3.33 ppm (-CH₂-, *J* = 6.4Hz) and signal multiplet at 6.00 ppm (-CH=, *J* = 6.4Hz), signal singlet at 5.95 ppm of one methylenedioxy group. The ¹³C-NMR spectrum showed 10 nonequivalent carbon resonances and the MS spectrum showed a molecular ion peak [M+.] at *m/z* 162, C₁₀H₁₀O₂ (see table 1). The high concentration of eugenol in bark essential oil of *Cinnamomun culilawan* makes it potentially useful for the preparation of herbal and modern medicines because they exhibit antibacterial, antifungal, anti-inflammatory, insecticidal and antioxidant activities [11-15].

4. Conclusions

The chemical composition of essential oil from *Cinnamomun culilawan*, Bl, was obtained from steam distillation method, and its chemical composition was determined by GC and GC-MS. The result indicated that the essential oil is mainly phenylpropanoid compound (86.473%), the minor compounds are oxygenated monoterpenes (7.934%) and monoterpenes hydrocarbon (5.592%). The purification of the two main compounds yielded eugenol (51.66%), and safrole (6.71%).

Acknowledgement

This study is part of the Fundamental Research Grant in 2015 from the Director General of Research and Community Service, Ministry of Research, Technology and Higher Education, Indonesia.

References

- [1] Ketaren S, Pengantar Minyak Atsiri, PN. Balai Pustaka, Jakarta, 1985.
- [2] Sastrohamidjojo H, Kimia Minyak Atsiri, Gadjah Mada University Press, 2004.
- [3] Lense O, The wild plants used as traditional medicines by indigenous people of Manokwari, West Papua, Bio Diversitas, 2012, 13 (2), 98-106.
- [4] Jantan I, Ali N. A. M, Ahmad A. R and Ahmad A. S, Chemical constituents of the essential oils of *Cinnamomum sintoc*, Blume, Pertanika J. Sci. & Techno., 1994, 2 (1), 39-45.
- [5] Iskandar Y, Supriyatna S, Chemical composition of volatile oil from *Cinnamomum sintoc* stem barks, Proceeding of The international seminar on chemistry, Jatinangor, Indonesia, 2008, 601-603
- [6] Reynertson K. A, Balick M. J, Lee R, Raynor W, Pelep Y, Kennelly E. J, A traditional method of *Cinnamomum carolinense* preparation eliminates safrole from a therapeutic Pohnpean tea, J. Ethnopharmacology, 2005, 102, 269-274.
- [7] Luo Y, LuoYo, Chen F, Liu H, Studies on the Chemical constituents in the essential oils from the leaves of *Cinnamomum bodinieri* Levi, Adv. Meterials Res., 2014, 1015, 373-376.
- [8] Kapelle I. B. D, Irawadi, T. T, Rusli M. S, Mangunwidjaja D, Mas'ud, Z. A, Synthesis of new curcumin analogues from kulitlawang oils using the conventional method and microwave, Sci. J. Chem., 2015, 3 (3): 50-56.
- [9] Sohila M. R, Sohila, H. J, Fransina E, 2013, Synthesis of 34-methylenedioxy isoamylcinamic as the sunscreen Compound from culilawang oil, Ind. J. Chem. Res., 2013, 1, 1-5.
- [10] Li H. Q, Liu Q. Z, Liu Z. L, Du Sh. Sh and Deng Z. W, Chemical composition and nematocidal activity of essential oil of *Agastacherugosa* against *Meloidogyne incognita*, Molecules, 2013, 18, 4170-4180.
- [11] Chu S. S, Cheng Fang Wang C. F, Du S. S, Liu S. L, Liu Z. L, Toxicity of the essential oil of *Illicium difengpistem* bark and its constituent compounds towards two grain storage insects, J. Insect Sci., 2011, 11 (152), 1-10.
- [12] Valente V. M. M, Jham G. N, Jardim C. M, Dhingra O. D, Ghiviriga I, Major antifungals in nutmeg essential oil against *Aspergillus flavus* and *A. ochraceus*, J. Food Res., 2015, 4 (1), 51-57.
- [13] Kamatou G. P, Vermaak I and Viljoen A. M, Eugenol- From the remote Maluku Islands to the international market place: A review of a remarkable and versatile molecule, Molecules, 2012, 17, 6953-6981.
- [14] Huang Y, Ho S. H, Lee H. C, Yap Y. L. Insecticidal properties of eugenol, isoeugenol and methyleugenol and their effect on nutrition of *Sitophilus zeamais* Motsch. (Coleoptera: Curculionidae) and *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae), J. Stored Prod. Res., 2002, 38, 403-412.
- [15] Ozturk A, Ozbek H, The Anti-inflammatory activity of *Eugenia caryophyllata* essential oil: An animal model of anti-inflammatory activity, Eur. J. Gen. Med., 2005, 2 (4), 159-163.