
Effects of wastewater irrigation on the growth of two bean spices and soil chemical properties under greenhouse conditions

Hamed Fathi¹, Mojan Mirzanejad²

¹Department of soil Sciences, college of Agriculture, Gorgan University of Agriculture Sciences and Natural Resources Gorgan, Iran

²Islamic Azad University, Rasht, Iran

Email address:

hamed_fathi_dokht@yahoo.com (H. Fathi)

To cite this article:

Hamed Fathi, Mojan Mirzanejad. Effects of Wastewater Irrigation on the Growth of Two Bean Spices and Soil Chemical Properties under Greenhouse Conditions. *American Journal of Agriculture and Forestry*. Vol. 2, No. 3, 2014, pp.88-93.

doi: 10.11648/j.ajaf.20140203.15

Abstract: This study has the objectives to evaluate the short-term effect of different dilutions of the wastewater on the chemical properties of sandy loam and sandy clay soils in two bean species under greenhouse condition. The effect of the irrigation with different wastewater dilutions on the bean crops and their chemical properties were studied. The results showed that the irrigation with wastewater reduced the pH in the sandy loam soil significantly in comparison with sandy clay soil, and this could be due to lower buffering capacity of sandy loam soil as compared to sandy clay soil registering a change in pH increasing. Available N, P and K status, OC and metallic cation content were significantly higher in sandy clay than in sandy loam soils. Generally, sewage water irrigation improved chemical properties and fertility status of both soils. The metallic cations contamination of the bean crops were observed below the maximum permissible limit. However, several folds have been built of metallic cations in soils, fronds and grain of the bean crops indicating that it is possible to increase the metallic cations contamination beyond maximum permissible limit if the waste water is continuously used for irrigating for a long time.

Keywords: Wastewater, Bean, Chemical Properties, Greenhouse Conditions

1. Introduction

The demand for water is continuously increasing in arid and semi-arid countries. Therefore, water of higher quality is preserved for domestic use while that of lower quality is recommended for irrigation. The use of industrial or municipal wastewater in agriculture is a common practice in many parts of the world (Ensink *et al.*, 2002; Kiziloglu *et al.*, 2008). Municipal wastewater is less expensive and considered an attractive source for irrigation in these countries (Al-Rashed and Sherif, 2000; Mohammad and Mazahreh, 2003), and any sources of water which might be used economically and effectively should be considered to promote further development. The total area irrigated with wastewater was conservatively estimated at around 9000 ha (Scott *et al.*, 2004). Iran, as a semi-arid country, suffers from shortages in water supply for domestic, industrial and agricultural purposes. Water resources in Iran are very limited because of naturally arid and semi-arid conditions.

Hence, limited water supplies require careful management for successful agricultural production. The interest in reusing wastewater for irrigation is rapidly growing in Iran as in most countries. Wastewater is a valuable source for plant nutrients and organic matter needed for maintaining fertility and productivity of arid soils (Horswell *et al.*, 2003). The reuse of wastewater for irrigation may potentially create environmental problems if not properly treated and managed (Bahri and Brissaud, 1996; Kiziloglu *et al.*, 2008). Application of treated wastewater for irrigation of plants and crops is gradually becoming a common practice worldwide (Angelakis *et al.*, 2003). Treatment of wastewater, as a segment of water management, usually produces a liquid effluent of suitable quality that can be used for irrigation purposes with minimum impacts on human health or the environment (Qishlaqi *et al.*, 2008; Hofstedt, 2005). In contractual wastewater treatment, noteworthy portions of heavy metals remain in the treated effluent if special advanced treatment is not conducted. Thus, long term effects of irrigation with wastewater might

include pollution of ground water and soil with heavy metals (Tyagi *et al.*, 2009). Other impacts of treated wastewater in agriculture include the health impacts of possible contamination of crops by pathogenic bacteria and heavy metals (Jiménez, 2006). Also Change in the basic physicochemical properties of soil (soil quality degradation) is another negative impact of wastewater application in agricultural lands (Qishlaqi *et al.*, 2008; Mahmoud *et al.*, 2010). This study has the objectives to evaluate the short-term effect of different dilutions of wastewater on chemical properties of sandy loam and sandy clay soils and assess the chemical build up in two bean species on pots under greenhouse conditions.

2. Materials and Methods

The experiment was conducted in the greenhouse of department of Soil Science, College of Agriculture, Shiraz University. The municipal wastewater used in this experiment was from the wastewater treatment plant of Marvdasht located near city of Shiraz. The experiment was laid out in factorial whit completely randomized design with three wastewater purification variants: fresh water irrigation (T1), primary treated wastewater and fresh water in equal proportions (T2), primary treated wastewater irrigation (T3) on two bean species (Red bean & Green bean) and two texture soil (sandy loam (SL) and sandy clay (SC) with three replicates). Primary treatment wastewater is the removal of settleable organic and inorganic solids by sedimentation and the removal of floating materials by skimming. Approximately 25–50% of the incoming biological oxygen demand (BOD₅), 50–70% of the total suspended solids, and 65% of the oil and grease are removed during the primary treatment but colloidal and dissolved constituents are not affected (Pescod, 1992; Kiziloglu *et al.*, 2008). Some characteristics of primary treated wastewater and fresh water are shown in Table 1. The primary treated wastewater and fresh water used for preparing various dilutions have total suspended solids (TSS) and pH within safe range for irrigation (Wu *et al.*, 2009) (Table 1). The soil samples were drawn prior to experimentation and at harvest of plant crops and analyzed for chemical properties following standard methods of analysis (Table 2). Soil pH and EC were measured by pH meter and EC meter from saturated soil paste extracted by vacuum pump. Organic carbon (OC) contents in soil were determined by the wet digestion method (Walkley and Black 1934), total N by the Kjeldahl method and available P according to Olsen *et al.* (1954). Available K⁺ was extracted by vacuum pump, from saturated paste and analyzed by flame emission spectrophotometer, metallic cation were extracted by DTPA and analyzed by atomic absorption spectrophotometer (Page *et al.*, 1982). The dry ashing method was used for analyzing the heavy metal contents in crop samples. Each determination was carried out by accurately measuring a sample of 1 g of a ground sample in a crucible. The crucible with its content was

placed in a muffle furnace and ashed at 550° C for 7 h. The ash was digested with 5 ml of 20% (V/V) Analar HCl solution. The residue was filtered into 50-ml volumetric flask using What man filter paper No.41 and the solution was made to the mark with deionised water. Atomic absorption spectrophotometry was used for the heavy metal determination. The sandy loam and sandy clay soils used were non-saline, mildly alkaline and calcareous in nature (Table 2). Also; the electrical conductivity (EC) was slight to moderately saline. The metallic cations content were below the maximum permissible limits as prescribed by Chen (2004). The status of available N and P was medium, while K was high in both the soils. The DTPA extractable Fe, Cu and Mn were high, while Zn was low in both the soils. Data were analyzed statistically the statistical software called SPSS (Abedi-Koupai, *et al.*, 2006).

Table 1. Characteristics of well and sewage water:

Charactries	unite	tab water	wastewater
TSS	mg L ⁻¹	4.5	85
PH	-Log[H ⁺]	7.78	7.8
EC	ds m ⁻¹	0.51	1.46
Na	me L ⁻¹	0.8	10.2
SAR	-----	1.02	6.8
BOD	mg L ⁻¹	-----	35

Table 2. chemical properties of soil prior to experimentation.

Parameter	unit	sandy loam	sandy clay
EC	ds m ⁻¹	0.57	0.72
PH	-Log[H ⁺]	7.5	8.1
CCE	%	44.5	39.8
availbale N	%	0.056	0.061
availbale p	mg kg ⁻¹	18.3	20.5
available K	mg kg ⁻¹	305	348
Zn	mg kg ⁻¹	0.68	0.73
Cu	mg kg ⁻¹	0.61	0.78
Mn	mg kg ⁻¹	4.35	5.87
Fe	mg kg ⁻¹	8.7	10.13
Pb	mg kg ⁻¹	Trace	Trace
Ni	mg kg ⁻¹	Trace	Trace
Cd	mg kg ⁻¹	Trace	Trace
OM	%	1.1	1.32

3. Results and Discussion

3.1. Effect on Soil Properties

3.1.1. Chemical Properties

The effect of soil texture, wastewater dilutions and vegetable crops on some of the chemical properties of the soils was analyzed (Tables 3A & 3B). The decrease in pH was more in the sandy loam soil as compared to sandy clay soil, and this could be due to lower buffering capacity of

sandy loam soil as compared to sandy clay soil registering a change in pH (Gelsomino, *et al.*, 2006). A decrease in pH of the soil with the use of wastewater dilutions for irrigation might be due to lower pH of wastewater being used for irrigation. Significantly highest value of EC was observed for sandy clay soil irrigated with wastewater as such as compared to rest of the treatment combinations indicating a higher build-up of salts in moderately fine textured soil. This could be ascribed to its high salt content with a mean value of 1.46 dS.m^{-1} as against 0.51 dS.m^{-1} of well water and hence contributing more dissolved salts in the soil. Similar results of pH were reported by Malla, *et al.*, 2007 and Ganeshamurthy, *et al.*, 2008) and of EC were reported by (Malla, *et al.*, 2007). On the other hand, the soils on which red bean grown soils recorded a small quantify EC than green bean grown soils. Available N, P and K status, OC and metallic cation content values were significantly higher in sandy clay than in sandy loam soils. It was further enhanced with wastewater irrigation as such or in its dilution indicating improvement in the fertility status of the soils except buildup of metallic cations in the soil. The increase in N, P and K content of soil with prolonged irrigation was also reported by (Masto, *et*

al., 2009). Sandy loam soil contained a higher value of CaCO_3 content which decreased on application of wastewater as such or in its dilution.

3.1.2. Metallic Cations Build up

The table 3B shows the Average of heavy metal concentration in soil of different irrigation treatments when vegetable crops were irrigated with waste water as such with reference to the control. Considering the mean values of the content buildup of metallic cations irrespective of the crops grown on them, a higher content buildup of Zn (2.58 ppm) and Ni (0.43 ppm) was recorded for sandy clay soil while, the buildup of Cd (1.4 ppm) was comparatively higher in sandy loam soil. Similarly, irrespective of the textural class of the soils buildup of Zn (2.58 ppm), Cu (2.48 ppm), Fe (17.78 ppm), Mn (10.93 ppm), Cd (1.4 ppm) and Ni (0.43 ppm) was relatively higher in soil where green bean was grown. These may be ascribed to differential removal of cations from the soil by the vegetable crops under study (Malla, R., & Totawat, K. L. 2006). The increase in DTPA extractable metallic cations was also reported by (Rana, *et al.*, 2010).

Table 3A. Effect of soil texture, sewage dilutions and vegetable crops on chemical properties of the soils.

species	Texture	Treatment	PH	EC ds m ⁻¹	OM	Caco3 -----%-----	N	P -----mg kg ⁻¹ -----	K
G	SL	T2	7.48	0.856	1.47	36.74	0.073	20.9	283
G	SL	T3	7.21	1.12	1.83	33.23	0.823	26.8	278.9
G	SC	T1	8.14	0.87	1.41	35.8	0.065	18.88	327.6
G	SC	T2	7.95	1.13	1.72	32.98	0.83	24.86	324.2
G	SC	T3	7.78	1.62	2.18	31.27	0.102	29.36	320.74
R	SL	T1	7.59	0.61	1.21	39.6	0.051	15.9	286.5
R	SL	T2	7.43	0.89	1.4	35.68	0.069	20	282.3
R	SL	T3	7.2	1.2	1.78	32.97	0.81	25.1	9
R	SC	T1	8.16	0.84	1.38	36.2	0.062	17.23	327
R	SC	T2	7.93	1.09	1.675	33.85	0.079	22.36	322
R	SC	T3	7.81	1.91	2.12	32.1	0.989	27.45	319

G: Green Bean R: Red Bean SL: Sandy Loam SC: Sandy Clay

Table 3B. Effect of soil texture, sewage dilutions and vegetable crops on chemical properties of the soils.

species	Texture	Treatment	Fe	Zn	Cu	Mn	Cd	Ni
G	SL	T1	6.95	0.573	0.498	4.1	T	T
G	SL	T2	10.23	1.23	1.11	6.88	0.8	0.158
G	SL	T3	12.79	1.95	1.65	8.45	1.4	0.346
G	SC	T1	9.112	0.678	0.711	5.64	T	T
G	SC	T2	14.23	1.632	1.84	9.23	0.65	0.2
G	SC	T3	17.78	2.58	2.48	10.93	1.1	0.43
R	SL	T1	6.9	0.58	0.493	3.96	T	T
R	SL	T2	9.65	1.12	0.98	6.5	0.65	0.146
R	SL	T3	11.3	1.79	1.58	8.1	1.3	0.324
R	SC	T1	8.88	0.61	0.683	5.4	T	T
R	SC	T2	14.7	1.55	1.72	9	0.48	0.187
R	SC	T3	16.88	2.43	2.42	10.88	0.91	0.394

G: Green Bean R: Red Bean SL: Sandy Loam SC: Sandy Clay T: Trace

3.2. Effect on Plant Composition

3.2.1. Metallic Cations Buildup in Grain and Fronds

The content buildup of metallic cations in the fronds and grain of crops when irrigated with waste water as such with reference to the control is given in Tables 4A and 4B, and figure 1. Considering the mean values of the content buildup of metallic cations in the fronds of crops irrespective of the type of crop grown, comparatively higher buildup of Zn (97.6 ppm), Cu (9.7 ppm), Fe (266.4 ppm) and Mn (179.9 ppm) was observed when crops were grown on sandy clay soil. Irrespective of the textural class of the soil, content buildup of Fe, Cu, Mn, and Zn was higher in the fronds of kidney bean. Similar to the content buildup of metallic cations in fronds, comparatively higher buildup of Zn (42.2 ppm), Cu (29.9 ppm), Fe (258 ppm) and Mn (75.3 ppm) was recorded in the grain of crop

grown on sandy clay soil. Irrespective of the textural class of the soils, content buildup of Fe, Cu, Mn, and Zn was higher in the fronds of kidney bean. The variations in the content buildup of metallic cations in the soils and vegetable crops may be attributed to a greater variation in the initial values of the metallic cations in the soil prior to experimentation and preferential absorbance of a particular cation by different vegetable crops under study (Sharma and Shukla, 2013). Even if the concentration of heavy metal in sewage effluents is low, the extent of build-up of metals in sewage/waste water-irrigated soils often depends on the period of its application (Malla, *et al.*, 2007). In spite of higher content buildup of metallic cations in the vegetable crops, they were all within the safe limit prescribed by (WHO, 1996).

Table 4A. Metallications build up in grain in sandy loam soil.

SPICES	Part of plant	Irrigation Treatment	N%	mg kg ⁻¹					
				P	K	Cl	Ca	Ni	
G	Grain	T1	8.46 E	0.453 F	0.451 F	2.987 E	N.D	N.D	
G	Grain	T2	9.01 ED	0.519 E	0.534 E	3.198 E	N.D	N.D	
G	Grain	T3	9.54 CD	0.567 D	0.583 D	3.421 D	N.D	N.D	
R	Grain	T1	10.2 BC	0.588 C	0.609 C	2.9 C	N.D	N.D	
R	Grain	T2	10.86 AB	0.675 B	0.723 B	3.8 B	N.D	N.D	
R	Grain	T3	11.54 A	0.732 A	0.787 A	4.7 A	N.D	N.D	
Metallic cations build up in fronds in sandy loam soil									
G	Fronds	T1	4.83 D	0.323 F	2.164 F	6.328 E	N.D	N.D	
G	Fronds	T2	5.33 C	0.386 E	2.264 D	6.249 D	N.D	N.D	
G	Fronds	T3	5.95 B	0.438 C	2.352 E	6.153 F	N.D	N.D	
R	Fronds	T1	4.08 E	0.42 D	2.921 C	6.8 C	N.D	N.D	
R	Fronds	T2	6.04 B	0.501 B	3.056 B	7.2 B	N.D	N.D	
R	Fronds	T3	7.168 A	0.576 A	3.175 A	7.6 A	N.D	N.D	

N.D: Not detection

Table 4B. Metallic cations build up in grain in sandy clay soil.

SPICES	Part of plant	Irrigation Treatment	N%	mg kg ⁻¹					
				P	K	Cl	Ca	Ni	
G	Grain	T1	9.06 F	0.49 C	0.48 C	3.4 C	N.D	N.D	
G	Grain	T2	9.69 E	0.563 B	0.57 BC	3.63 BC	N.D	N.D	
G	Grain	T3	10.3 D	0.61 B	0.62 BC	3.9 AB	N.D	N.D	
R	Grain	T1	11.4 C	0.6 B	0.8 BC	3.638 BC	N.D	N.D	
R	Grain	T2	12.6 B	0.71 A	0.85 B	3.884 AB	N.D	N.D	
R	Grain	T3	14.1 A	0.748 A	1.2 A	4.173 A	N.D	N.D	
Metallic cations build up in fronds in sandy loam soil									
G	Fronds	T1	5.2 D	0.35 D	2.3 C	7.2 A	N.D	N.D	
G	Fronds	T2	5.74 C	0.418 CD	2.40 C	7.11 A	N.D	N.D	
G	Fronds	T3	6.4 B	0.48 C	2.5 C	7 A	N.D	N.D	
R	Fronds	T1	4.3 E	0.8 B	3.6 B	7.704 A	N.D	N.D	
R	Fronds	T2	6.3 B	0.88 AB	3.9 AB	7.607 A	N.D	N.D	
R	Fronds	T3	7.2 A	0.9 A	4.2 A	7.49 A	N.D	N.D	

N.D: Not detection




Figure 1. The content builds up of metallic cations in the dry weight of fronds and grain of two beans (KB: Red Bean GB: Green Bean HT: Sandy clay LT: Sandy loam).

4. Conclusions

Generally speaking, sewage water irrigation improved chemical properties and fertility status of both soils. The metallic cations contamination of vegetable crops was observed to be below the maximum permissible limit. However, several folds buildup of metallic cations in soils, fronds and grain of vegetable crops under study indicated that there might be an increase in metallic cations contamination beyond maximum permissible limit if such waste water is continuously used for irrigation on long term. Thus timely monitoring of sewage water, soil and crops is essential to maintain soil quality for consumption of safe foods grown on them. Establishment of at least a primary sewage treatment plant is suggested where raw sewage is being diverted for irrigation.

References

- [1] Abedi-Koupai, J., Mostafazadeh-Fard, B., Afyuni, M., & Bagheri, M. R. (2006). Effect of treated wastewater on soil chemical and physical properties in an arid region. *Plant Soil and Environment*, 52(8), 335.
- [2] Al-Rashed, M. F., & Sherif, M. M. (2000). Water resources in the GCC countries: an overview. *Water Resources Management*, 14(1), 59-75.
- [3] Angelakis, A. N., Bontoux, L., & Lazarova, V. (2003). Challenges and prospectives for water recycling and reuse in EU countries. *Water Science & Technology: Water Supply*, 3(4).
- [4] Bahri, A., & Brissaud, F. (1996). Wastewater reuse in Tunisia: assessing a national policy. *Water Science and Technology*, 33(10), 87-94.
- [5] Chen, Y., Wang, C., Wang, Z., & Huang, S. (2004). Assessment of the contamination and genotoxicity of soil irrigated with wastewater. *Plant and Soil*, 261(1-2), 189-196.
- [6] Ensink, J. H., Van der Hoek, W., Matsuno, Y., Munir, S., & Aslam, M. R. (2002). Use of Untreated Wastewater in Peri-Urban Agriculture in Pakistan.
- [7] Ganeshamurthy, A. N., Varalakshmi, L. R., & Sumangala, H. P. (2008). Environmental risks associated with heavy metal contamination in soil, water and plants in urban and periurban agriculture. *J HorticulturSci*, 3(1), 1-29.
- [8] Gelsomino, A., Badalucco, L., Ambrosoli, R., Crecchio, C., Puglisi, E., & Meli, S. M. (2006). Changes in chemical and biological soil properties as induced by anthropogenic disturbance: A case study of an agricultural soil under recurrent flooding by wastewaters. *Soil Biology and Biochemistry*, 38(8), 2069-2080.
- [9] Hofstedt, C. (2005). Wastewater use in Agriculture in Andhra Pradesh, India: An evaluation of irrigation water quality in reference to associated health risks and agricultural suitability.
- [10] Horswell, J., Speir, T. W., & Van Schaik, A. P. (2003). Bio-indicators to assess impacts of heavy metals in land-applied sewage sludge. *Soil Biology and Biochemistry*, 35(11), 1501-1505.
- [11] Jiménez, B. (2006). Irrigation in developing countries using wastewater. *International Review for Environmental Strategies*, 6(2), 229-250.

- [12] Kiziloglu, F. M., Turan, M., Sahin, U., Kuslu, Y., & Dursun, A. (2008). Effects of untreated and treated wastewater irrigation on some chemical properties of cauliflower (*Brassica oleracea* L. var. botrytis) and red cabbage (*Brassica oleracea* L. var. rubra) grown on calcareous soil in Turkey. *Agricultural water management*, 95(6), 716-724.
- [13] Kiziloglu, F. M., Turan, M., Sahin, U., Kuslu, Y., & Dursun, A. (2008). Effects of untreated and treated wastewater irrigation on some chemical properties of cauliflower (*Brassica oleracea* L. var. botrytis) and red cabbage (*Brassica oleracea* L. var. rubra) grown on calcareous soil in Turkey. *Agricultural water management*, 95(6), 716-724.
- [14] Mahmoud, A., Olivier, J., Vaxelaire, J., & Hoadley, A. F. (2010). Electrical field: A historical review of its application and contributions in wastewater sludge dewatering. *Water research*, 44(8), 2381-2407.
- [15] Malla, R., & Totawat, K. L. (2006). Effect of Municipal Sewage on Soil Properties and Chemical Build-up in Vegetable Crops Grown on Haplusteps of Sub-humid Southern Plains of Rajasthan. *Journal of the Indian Society of Soil Science*, 54(2), 226-231.
- [16] Malla, R., Tanaka, Y., Mori, K., & Totawat, K. L. (2007). Effect of short-term sewage irrigation on chemical build up in soils and vegetables.
- [17] Masto, R. E., Chhonkar, P. K., Singh, D., & Patra, A. K. (2009). Changes in soil quality indicators under long-term sewage irrigation in a sub-tropical environment. *Environmental geology*, 56(6), 1237-1243.
- [18] Mohammad, M. J., & Mazahreh, N. (2003). Changes in soil fertility parameters in response to irrigation of forage crops with secondary treated wastewater. *Communications in soil science and plant analysis*, 34(9-10), 1281-1294.
- [19] Page AL (1982). Methods of soil analysis, part 2: chemical and microbiological properties. *Agronomy monograph No. 9* (part 2). ASA/SSSA, Madison, Wisconsin
- [20] Pescod, M. B. (1992). Wastewater treatment and use in agriculture.
- [21] Qishlaqi, A., Moore, F., & Forghani, G. (2008). Impact of untreated wastewater irrigation on soils and crops in Shiraz suburban area, SW Iran. *Environmental monitoring and assessment*, 141(1-3), 257-273.
- [22] Olsen, S. R., C. V. Cole, F. S. Watanabe, and L. A. Dean. 1954. Estimation of available phosphorous in soil by extraction with sodium bicarbonate. *USDA Circ. 39*, U. S. Gover. Prin. Office Washington, DC.
- [23] Rana, L., Dhankhar, R., & Chhikara, S. (2010). Soil Characteristics Affected by long term Application of Sewage Wastewater. *International Journal of Environmental Research*, 4(3).
- [24] Scott, C. A., Faruqi, N. I., & Raschid-Sally, L. (2004). Wastewater Use in Irrigated Agriculture: Management Challenges in Developing Countries.
- [25] Sharma, D., & Shukla, A. K. (2013). Analysis of heavy metals in soil and vegetables irrigated with wastewater in Meerut city region. *Int. J. Adv. Life Sci*, 6(4), 287-292.
- [26] Tyagi, R. D., Surampalli, R. Y., Yan, S., Zhang, T. C., Kao, C. M., & Lohani, B. N. (2009). Sustainable sludge management: production of value added products. *ASCE Publications*.
- [27] Walkley, A., & Black, I. A. (1934). An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil science*, 37(1), 29-38.
- [28] World Health Organization, 1996. *Water Quality Monitoring - A Practical Guide to the Design and Implementation of Freshwater Quality Studies and Monitoring Programmers*.
- [29] Wu, L., Chen, W., French, C., & Chang, A. C. (2009). Safe application of reclaimed water reuse in the southwestern United States (Vol. 8357). *UCANR Publications*.