Woody Species Richness and Diversity at Ades Dry Afromontane Forest of South Eastern Ethiopia

Muktar Reshad*, Alemayehu Beyene, Muktar Mohammed

College of Natural Resource and Environmental Science, Oda Bultum University, Chiro, Ethiopia

Email address:

r1muk@yahoo.com (M. Reshad)

*Corresponding author

To cite this article:

Muktar Reshad, Alemayehu Beyene, Muktar Mohammed. Woody Species Richness and Diversity at Ades Dry Afromontane Forest of South Eastern Ethiopia. *American Journal of Agriculture and Forestry*. Vol. 7, No. 2, 2019, pp. 44-52. doi: 10.11648/j.ajaf.20190702.12

Received: January 12, 2019; **Accepted:** March 22, 2019; **Published:** April 18, 2019

Abstract: The study was conducted at Ades Dry Afromontane Forest at 407 km Southeast of Addis Ababa to assess the diversity and composition of woody plant species. The sampling design was based on a US Forest Service model for the indigenous forests. A total of 60 circular plots each with an area of 0.017 ha were arranged in groups of four where a central plot is surrounded by three plots that are each at 36.6m from the central plot. In each plot, all woody plants that were ≥ 10 cm in DBH were sampled for floristic diversity. Biodiversity analyses were accomplished using the Shannon-Weaver's Index (H') to assess the tree species diversity and Shannon Equitability ($H'E$) was calculated to assess the evenness values of species while the Importance Value Index (IVI) of each woody species was analyzed to see the Importance of individual tree and shrub species at the site. A total of 65 trees and shrub species belonging to 38 families had been recorded in this study. Rosaceae was a family with the highest number of species comprising about 9.23 % of the total number of species. The H' ranged between 0.004 to 0.362 with the overall H' of 2.82. The $H'E$ values ranged between 0.001 to 0.087 with a mean value of 0.01. A mean $H'E$ value of 0.01 indicates that the relative homogeneity of woody plant species of the sampled plots was 1% of the maximum possible even population. The evenness values are not enough to justify uniformity in composition of tree species. The mean IVI value ranged between 0.36 to 49.06 with mean IVI value of 6.0. In this study only 15% of the recorded species were found with IVI values > 10 and the rest of 85% have IVI values < 10 . The variation in survival mechanisms of species made some species to be dominant and most species to be lower in number in a given ecosystem. Tree species with high IVI were also found to have higher H' of diversity. The spatial distribution and dominance of species can be affected both by the properties of the species themselves and the environmental factors. The diversity of woody species observed in the Ades dry afromontane is encouraging since, among other reasons, it is useful for conservation strategy.

Keywords: Shannon-Weaver, Evenness, Trees, Shrubs, Sampling Design, Important Value Index, DBH

1. Introduction

Dense and Extensive forest resources were once covered the highland area of Ethiopia [13]. Today these extensive forest resources of the country are under threats of deforestation and forest degradation. The rate of deforestation in Ethiopia has been estimated to be between 150,000 ha and 200,000 ha [12].

The continuous deforestation and degradation of forest ecosystems in Ethiopia in general and in Ades dry afromontane forest in particular, is of major concern due to the negative impacts this has on many of the ecosystem

service. The Ades dry afromontane forest, which is characterized by a variety of tree species have lost its cover due to agricultural land expansion and illegal logging activities. This degradation has affected the plant species composition and presented threats for some of the tree species.

In recognition to the above listed threats to forest biodiversity, the government of Oromia National Regional State has designed the establishment of Participatory Forest Management (PFM) as one of various strategies for

conservation of forest resources through solving the problem of open access to the forest resource and promoting sustainable forest management. Because of lack of awareness on the principles of PFM by the members of PFM living in and/or near by the forest, still there exist exploitation of the forest resource through illegal logging, cutting for fuel wood and expansion of agricultural lands. Information with regard to the diversity, composition and importance values of woody plant species in the forest is necessary for conservation and sustainable utilization of the forest resources.

This study was therefore aimed to assess the floristic composition and diversity and to analyze important value index of woody species in Ades dry afromontane forest by exploring the identity and variety of woody species that have regenerated within the forest.

2. Materials and Methods

2.1. Description of the Study Site

The study was conducted in Ades dry afromontane forest located between 1029419m E to 1030017m E and 746685m N to 746986m N. It is about 407 km to East of Addis Ababa, in Oromia National Regional State, Southeastern Ethiopia (Figure 1). The forest covers a total area of 618 ha and its altitude ranges from 2517m to 2743m a.s.l. The weather condition of the study area is characterized by coldest climate which is locally known as ‘Baddaa’ in Afan Oromo language. The mean annual temperature ranges between 17°C and 24°C and the mean annual rainfall ranges between 600mm to 1250mm [18]. Its rain fall distribution is mostly from July to September. The dry and hot season is mostly starts from the middle of December to the end of March.

Figure 1. Map of the study area.

2.2. Methods

2.2.1. Sampling Design

The layout of the plots for the assessments of woody species diversity was adopted from design developed for The Forest Inventory and Analysis Programme (FIA) [29]. A circular plots each with an area of 0.017 ha arranged in groups of four where a central plot is surrounded by three plots to form an equilateral triangle at a distance of 36.6m out from the central plot to each of the three surrounding plots were laid out (Figure 2). Pairs of plots were established and two plots had been selected at random from each group of plots, and totally of 60 plots from 30 groups were sampled in this study.

2.2.2. Measurement of Woody Vegetation

Counting the number of stems and measurement of diameter at breast height (DBH) in each plot were conducted

Figure 2. Layout of plots and sub-plots.

for each of the woody plant species. The DBH of all living woody stems (trees and shrubs ≥ 10 cm DBH) was measured using Caliper and a 5m diameter tape. The identification of species was made using a guide book Flora of Ethiopia and Eritrea [14], with the assistance of a local Para-taxonomist. For those species difficult to identify in the field, fresh specimens were collected and taken to the National Herbarium of Addis Ababa University for identification.

2.2.3. Data Analysis

- Species diversity:** The Shannon-Weaver's index (H') was used as measure of diversity [20] and calculated as $H' = -\sum P_i \ln P_i$ where: i , is the proportion of the species relative to the total number of species (p_i) and $\ln$ is a natural logarithm.
- Species richness (S):** Species richness as the number of species present in an ecosystem was calculated as: $S = \sum n$ Where: n is number of species.
- Species evenness ($H'E$):** Species evenness was assessed by Shannon's equitability index calculated as: $H'E = H'/H_{\max}$ Where: $H_{\max}$ is defined as $\ln S$
- The Importance Value Index (IVI):** The index was calculated by integrating: The Relative frequency (RF); Relative density (RD) and Relative Dominance (RB) of species [15, 25]. It is calculated as: $IVI = (RF + RD + RB)$

3. Results and Discussion

3.1. Floristic Composition

A total of 65 tree and shrub species belonging to 38 families (Table 1) were identified, out of these trees constituted 67.7% while shrubs were 32.3%. Sixteen families each with more than one species comprise 66.2 % of the total number of species recorded in the study area and the rest of 22 families each were represent only by one species (Table 2). Rosaceae comprise the highest number of species (6) followed by Oleaceae and Myrsinaceae (4 species each), Flacourtiaceae, Celastraceae, and Asteraceae (3 species each), Ulmaceae, Rubiaceae, Polygonaceae, Myrtaceae, Meliaceae, Fabaceae, Euphorbiaceae, Cupressaceae, Boraginaceae, and Araliaceae (2 species each). Habitat adaptation and favorable environmental conditions could encourage the prevalence of Rosaceae family [7, 30]. The influence of edaphic parameter (soil nutrients) in species richness and establishment in an ecosystem was also supported by other studies [28, 3].

Species that are at low level in some parts of Ethiopia such as: *Juniperus procera*, *Hagenia abyssinica*, and *Podocarpus falcatus*, are highly found in Ades dry Afromontane forest. This could be due to the establishment of Participatory Forest Management where the local communities are engaged in forest protection and management.

Table 1. Lists of trees and shrub species in Ades dry afromontane forest, South Eastern Ethiopia.

Family	Species	Life form	Family	Species	Life form
Anacardiaceae	<i>Rhus glutinosa</i>	Shrub	Myrsinaceae	<i>Myrsine africana</i>	Tree
Apocynaceae	<i>Carrisa edulis</i>	Shrub		<i>Rapanea simensis</i>	Tree
Aquifoliaceae	<i>Ilex mitis</i>	Tree	Myrtaceae	<i>Eucalyptus globulus</i>	Tree
Araliaceae	<i>Polyscias fulva</i>	Tree		<i>Sizygeum guineense</i>	Tree
	<i>Schfflera abyssinica</i>	Tree		<i>Jasminum abyssinicum</i>	Tree
	<i>Vernonia amygdalina</i>	Tree	Oleaceae	<i>Olea europaea</i>	Tree
Asteraceae	<i>Vernonia auriculifera</i>	Shrub		<i>Olea capensis</i>	Tree
	<i>Vernonia leopoldii</i>	Shrub		<i>Olinia rochetiana</i>	Shrub
Balanitaceae	<i>Balanites aegyptica</i>	Tree	Papilionaceae	<i>Erythrina brucei</i>	Tree
Boraginaceae	<i>Cordia africana</i>	Tree	Phytolacaceae	<i>Phytolacca dedocandra</i>	Shrub
	<i>Ehretia cymosa</i>	Tree	Pittosporaceae	<i>Pittosporum abyssinicum</i>	Tree
	<i>Chata edulis</i>	Shrub	Podocarpaceae	<i>Podocarpus falcatus</i>	Tree
Celastraceae	<i>Maytenus arbutifolia</i>	Tree		<i>Rumex abyssinicus</i>	Shrub
	<i>Maytenus obscura</i>	Tree	Polygonaceae	<i>Rumex nervosus</i>	Shrub
Cupressaceae	<i>Cupressus lusitanica</i>	Tree	Proteaceae	<i>Gravelea robusta</i>	Tree
	<i>Juniperus procera</i>	Tree	Rhamnaceae	<i>Rhamnus prenois</i>	Shrub
Euphorbiaceae	<i>Croton macrostachys</i>	Tree		<i>Hagenia abyssinica</i>	Tree
	<i>Macaranga kilimandscharica</i>	Tree		<i>Prunus africana</i>	Tree
Fabaceae	<i>Calpurnia subdecandra</i>	Shrub	Rosaceae	<i>Rosa abyssinica</i>	Shrub
	<i>Calpurina aura</i>	Shrub		<i>Rosmarinus offinalis</i>	Tree
	<i>Dovyalis caffra</i>	Shrub		<i>Rubus apetalus</i>	Tree
Flacourtiaceae	<i>Dovyalis abyssinica</i>	Shrub		<i>Rubus studneri</i>	Tree
	<i>Dovyalis vericosa</i>	Shrub	Rubiaceae	<i>Gardenia lutea</i>	Tree
Hypericaceae	<i>Hypericum revolutum</i>	Shrub		<i>Rytigynia neglecta</i>	Shrub
Icacinaeae	<i>Apodytes dimidiata</i>	Tree	Sapindaceae	<i>Dodonaea angustifolia</i>	Tree
Lamiaceae	<i>Ocimum lamifolium</i>	Shrub	Sapotaceae	<i>Aningeria altissima</i>	Tree
Lauraceae	<i>Ocotea kenyensis</i>	Tree	Simaroubaceae	<i>Brucea antidysenterica</i>	Shrub
Meliaceae	<i>Ekebergia capensis</i>	Tree	Solanaceae	<i>Discopodium penninervium</i>	Tree
	<i>Lepidotrichilia volkensii</i>	Tree	Sterculiaceae	<i>Dombeya torrida</i>	Tree

Family	Species	Life form	Family	Species	Life form
Melanthaceae	<i>Barsama abyssinica</i>	Tree	Ulmaceae	<i>Celtis africana</i>	Tree
Moraceae	<i>Ficus sur</i>	Tree		<i>Clerodendron myricoides</i>	Shrub
Myrsinaceae	<i>Embelia schimpri</i>	Tree	Verbenaceae	<i>Premna schimperii</i>	Tree
	<i>Maesa lanceolata</i>	Tree			

This finding was not in line with the works from Forest of Assam on vegetative assessment of tree species and shrubs where Moraceae, Meliaceae and Papilionaceae have been represented with more number of species due to their ability to produce numerous seeds which was eventually establish at suitable sites [10]. But in this study, they were represented only by one species. This may be due to the geographic variations of the two study sites and the result of this study was in line with the study conducted on tropical forests regeneration and survival of tree seedlings [17].

The number of plant species (65) recorded in Ades dry afromontane forest was higher as compared with some studies conducted in Ethiopia [1, 27, 32, 6] and elsewhere in tropical areas from Tungnath [22]. This may be due to geographic differences and variations in vegetation formation of these study sites. However, this figure is lower when

compared to the number of species recorded in Jello-Muktar dry afromontane forest under similar agro-ecology from Eastern Ethiopia [24]. Similarly, the number of plant species recorded in this study were lower when compared to the number of species recorded for forests of other ago-ecologies in Ethiopia [5, 33, 23] and some tropical countries [7, 8].

The reasons for those 22 families in the present study represented each only by one species or the poor number of species observed in these families could be attributed to diseases and browsing by herbivores which resulted in poor growth and establishment and perhaps seeds need scarification treatment before germination as study conducted on seedling populations and regeneration of woody species in dry Afromontane forests of Ethiopia [11]. The low number of species could also be attributed to anthropogenic activities which affected species growth and production [9].

Table 2. Plant Families with their number of woody plant species occurred in Ades dry afromontane forest, South Eastern Ethiopia.

Family	Species number	%	Family	Species number	%
Anacardiaceae	1	1.54	Myrsinaceae	4	6.15
Apocynaceae	1	1.54	Myrtaceae	2	3.08
Aquifoliaceae	1	1.54	Oleaceae	4	6.15
Araliaceae	2	3.08	Papilionaceae	1	1.54
Asteraceae	3	4.62	Phytolacaceae	1	1.54
Balanitaceae	1	1.54	Pittosporaceae	1	1.54
Boraginaceae	2	3.08	Podocarpaceae	1	1.54
Celastraceae	3	4.62	Polygonaceae	2	3.08
Cupressaceae	2	3.08	Proteaceae	1	1.54
Euphorbiaceae	2	3.08	Rhamnaceae	1	1.54
Fabaceae	2	3.08	Rosaceae	6	9.23
Flacourtiaceae	3	4.62	Rubiaceae	2	3.08
Hypericaceae	1	1.54	Sapindaceae	1	1.54
Icacinaeae	1	1.54	Sapotaceae	1	1.54
Lamiaceae	1	1.54	Simaroubaceae	1	1.54
Lauraceae	2	1.54	Solanaceae	1	1.54
Meliaceae	1	3.08	Sterculiaceae	1	1.54
Melanthaceae	1	1.54	Ulmaceae	2	3.08
Moraceae	1	1.54	Verbenaceae	1	1.54

The number of trees and shrub species in the present study, about 95% of which are indigenous to the area, indicates the significant role of the forest in conservation of biological diversity as reported from studies elsewhere in East Usambara Forest Reserves [16]. The high number of species richness in the study area is again attributed to the current protection measures given to the forest that via allows the regeneration of different species from soil seed bank. Some economic activities particularly, beekeeping through participatory forest management is highly encouraged in Ades dry afromontane forest by the presence of such floristic diversity.

3.2. Pattern of Distribution of Woody Species in Plant Families by Their Life Forms

With regard to the pattern of distribution of the number of woody species by life forms in each family, Rosaceae again comprise the highest number of tree species (5) and followed by Myrsinaceae (4) and Oleaceae (3) and the rest twelve families comprise each 2 tree species while 6 families were represented only by 1 tree species (Figure 3).

The distribution of shrub species in each family ranges between 1 to 3 where Flacourtiaceae comprise the highest number of shrub species (3) followed by Asteraceae,

Fabaceae and Polygonaceae (each with 2 shrub species) and the rest 12 families were represented only by one shrub species (Figure 4).

Figure 3. Distribution of tree species in the plant families.

Figure 4. Distribution of Shrub species in plant families.

3.3. Species Diversity and Evenness

The Shannon index of diversity for tree and shrub species in this study ranged between 0.004 to 0.3623 with the overall H' of 2.82 (Table 3). Species noted to have contributed to high Shannon index of diversity include: *Podocarpus falcatus* (0.3623), *Eucalyptus globules* (0.2290), *Juniperus procera* (0.2120), *Croton macrostachys* (0.074), *Cordia africana* (0.066), *Ficus sur* (0.063), *Cuprecess lusitanica* (0.063), *Embelia schimpri* (0.059), *Hagenia abyssinica* (0.054), *Ekebergia capensis* (0.053), *Ehretia cymosa* (0.050), *Hypericum revolutum* and *Carrisa edulis* (each 0.047), *Myrsine africana*, *Maytenus obscura* and *Barsama*

abyssinica (each 0.042) (Table 3). The overall H' of these species accounts about 50% of the total H' for all species recorded in the study site. Species of *Sizygeem guineense*, *Celtis africana*, *Rosmarinus offinalis*, *Rubus apetalus* and *Rubus studneri* were among tree and shrub species with the lowest Shannon index of diversity. The Shannon index of diversity recorded for *Podocarpus falcatus* (0.3623) was again higher than the overall H' of the first eighteen trees and shrub species with lowest H' . *Podocarpus falcatus* with the highest Shannon index of diversity in this study also showed the highest IVI.

Table 3. Tree and shrub species diversity indices at Ades dry afromontane forest, South Eastern Ethiopia.

Species	H'	H'E	Species	H'	H'E
<i>Aningeria altissima</i>	0.041	0.01	<i>Macaranga kilimandsharica</i>	0.039	0.009
<i>Apodytes dimidiata</i>	0.036	0.009	<i>Maesa lanceolata</i>	0.041	0.01
<i>Balanytus aegyptica</i>	0.023	0.006	<i>Maytenus arbutifolia</i>	0.037	0.009
<i>Barsama abyssinica</i>	0.042	0.01	<i>Maytenus obscura</i>	0.042	0.01
<i>Brucea antidysenterica</i>	0.034	0.008	<i>Myrsine africana</i>	0.042	0.01
<i>Calpurnia subdecandra</i>	0.035	0.008	<i>Ocimum lamifolium</i>	0.031	0.008
<i>Carrisa edulis</i>	0.047	0.011	<i>Ocotea kenyensis</i>	0.034	0.008
<i>Cupressus lusitanica</i>	0.063	0.015	<i>Olea europaea</i>	0.041	0.01
<i>Calpurina aura</i>	0.041	0.01	<i>Olea capensis</i>	0.035	0.008
<i>Celtis africana</i>	0.014	0.003	<i>Olinia rochetiana</i>	0.041	0.01
<i>Chata edulis</i>	0.03	0.007	<i>Phytolacca dedocandra</i>	0.022	0.005
<i>Clerodendron myricoides</i>	0.021	0.005	<i>Pittosporum abyssinicum</i>	0.037	0.009
<i>Croton macrostachys</i>	0.074	0.018	<i>Podocarpus falcatus</i>	0.362	0.087
<i>Cordia africana</i>	0.066	0.016	<i>Polyscias fulva</i>	0.024	0.006
<i>Discopodium penninervium</i>	0.038	0.009	<i>Prunus africana</i>	0.018	0.004
<i>Dodonaea angustifolia</i>	0.027	0.007	<i>Premna schimperi</i>	0.035	0.008
<i>Dombeya torrida</i>	0.027	0.007	<i>Rapanea simensis</i>	0.024	0.006
<i>Dovyalis caffra</i>	0.036	0.009	<i>Rhamnus prenois</i>	0.031	0.008
<i>Dovyalis abyssinica</i>	0.025	0.006	<i>Rhus glutinosa</i>	0.021	0.005
<i>Dovyalis vericosa</i>	0.023	0.006	<i>Rosa abyssinica</i>	0.022	0.005
<i>Erythrina brucei</i>	0.025	0.006	<i>Rosmarinus offinalis</i>	0.016	0.004
<i>Ehretia cymosa</i>	0.05	0.012	<i>Rubus apetalus</i>	0.016	0.004
<i>Ekebergia capensis</i>	0.053	0.013	<i>Rubus studneri</i>	0.017	0.004
<i>Embelia schimpri</i>	0.059	0.014	<i>Rumex abyssinicus</i>	0.02	0.005
<i>Eucalyptus globulus</i>	0.229	0.055	<i>Rumex nervosus</i>	0.018	0.004
<i>Ficus sur</i>	0.063	0.015	<i>Rytigynia neglecta</i>	0.02	0.005
<i>Gardenia lutea</i>	0.038	0.009	<i>Schfflera abyssinica</i>	0.023	0.006
<i>Gravelea robusta</i>	0.033	0.008	<i>Sizygeem guineense</i>	0.004	0.001
<i>Hagenia abyssinica</i>	0.054	0.013	<i>Vernonia amygdalina</i>	0.031	0.008
<i>Hypericum revolutum</i>	0.047	0.011	<i>Vernonia auriculifera</i>	0.039	0.009
<i>Ilex mitis</i>	0.041	0.01	<i>Vernonia leopoldii</i>	0.025	0.006
<i>Jasminum abyssinicum</i>	0.023	0.006			
<i>Juniperus procera</i>	0.111	0.026			
<i>Lepidotrichilia volkensii</i>	0.021	0.005			

The study revealed Shannon-Wiener Index of diversity (H') of 2.82 for the Ades dry Afromontane forest (Table 3). This index tells about species richness (number of species) and evenness (species distribution) [21]. The larger the value of H' the greater the species diversity and vice versa. An ecosystem with H' value greater than 2 has been regarded as medium to high diverse in terms of species [4]. An ecosystem having high species diversity has also a larger value of Shannon index of diversity and vice-versa [10]. The Ades dry afromontane forest is thus belongs to an area with high species diversity.

The Shannon index of diversity recorded in this study was higher than those studies conducted for some other forests types [31, 26]. The sizes of trees enumerated in those studies (i.e., within minimum DBH set at 12.7cm as opposed to 10cm in this study) could be among the factors for the observed differences in terms of Shannon index of diversity between the two sites. The high diversity of tree species observed in the Ades dry Afromontane is encouraging since, among other reasons, it is useful for conservation purposes.

The evenness values for tree species of the present study ranged between 0.001 to 0.087 with a mean value of 0.01 and overall evenness value of 0.67 (Table 3). A mean evenness value of 0.01 indicates that the relative homogeneity of tree species of the sampled forest plots was 1% of the maximum possible even population. The evenness values are not enough to justify uniformity in composition of tree species. This is expected because not all trees are equally distributed and there is some variation in the distribution of species in the study area. Tree species that showed higher evenness values were *Podocarpus falcatus* (0.087), *Eucalyptus globules* (0.055), *Juniperus procera* (0.026), *Croton macrostachys* (0.018), *Cordia africana* (0.016), *Cupressus lusitanica* and *Ficus sur* (each 0.015) and *Embelia schimpri* (0.015). These species also showed higher Shannon index of diversity (Table 3). The evenness value of tree with the highest index (*Podocarpus falcatus*) was about 87 times higher than the tree species with the lowest evenness value (*Sizygeem guineense*). This was supported by study conducted on Masha forest where the lower evenness values in forest indicate the dominance of a few species [2]. In this

study, a high evenness values for indigenous species indicates that little dominance by any single species but repeated co-existence of species over all plots in the forest as reported for some species in Northeastern Ethiopia [2]. The evenness values recorded in this study are lower than the evenness values reported from Western Kenya that ranged between 0.24 to 0.71 [19].

3.4. Importance Value Index (IVI)

In all sample plots assessed *Podocarpus falcatus* ranked first with mean IVI of 49.06, followed by *Juniperus procera* (34.15), *Eucalyptus globules* (30.6), *Cupressus lusitanica* (23), *Olea europaea* (20.17), *Hagenia abyssinica* (14.69), *Vernonia auriculifera* (13.32) and *Croton macrostachys* (12.25) (Table 4). Contrary to this, *Sisyrinchium guineense*, *Rytigynia neglecta*, *Rosmarinus officinalis*, *Rubus apetalus*, *Balanites aegyptiaca*, *Rumex abyssinicus* were the species with the lowest IVI. The mean IVI of *Hagenia abyssinica* was higher than those tree species which were officially declared endangered at the national level in Ethiopia next to *Juniperus procera*. Most of the tree species with higher IVI were indigenous although *Eucalyptus globules* and

Cupressus lusitanica species ranked first and second among species of exotic plantations.

The mean IVI value in this study ranged between 0.36 to 49.06 with mean IVI value of 6.0. It has been reported that IVI value of tree species in stands with only one species can reach a maximum of 300 [25]. In this study only 15 % of the recorded species were found with IVI values > 10 and the rest of 85% have IVI values < 10. On the other hand the first fifteen species with higher IVI values (Table 4), accounted about 64% of the total IVI for the present study which indicates that few species are dominating the area. The variation in survival mechanisms of species made some species to be dominant and most species to be lower in number in a given ecosystem [15]. The spatial distribution and dominance of species can be affected both by the properties of the species themselves and the environmental factors [28].

These results disclose that the most important species in Ades dry afro-montane forest have high diversity in the scale of Shannon-Weiner Index of Diversity. The IVI rank species in a way as to give an indication on which species come out as important element of the Ades dry afro-montane forest trees.

Table 4. Overall Mean IVI of trees and shrub (>10cm DBH) species in Ades dry afro-montane forest, South Eastern Ethiopia.

Species	RF	RD	RB	IVI	Species	RF	RD	RB	IVI
<i>Aningeria altissima</i>	1.19	0.87	1.33	3.38	<i>Macaranga kilimandscharica</i>	2.14	0.82	0.36	3.31
<i>Apodytes dimidiata</i>	1.43	0.74	0.1	2.27	<i>Maesa lanceolata</i>	4.5	4.3	0.3	9.1
<i>Balanites aegyptiaca</i>	0.59	0.42	0.12	1.14	<i>Maytenus arbutifolia</i>	3.01	0.89	1.12	5.02
<i>Barsana abyssinica</i>	1.07	0.89	0.44	2.4	<i>Maytenus obscura</i>	2.26	0.89	0.4	3.54
<i>Brucea antidysenterica</i>	1.31	0.69	0.04	2.03	<i>Myrsine africana</i>	2.38	0.89	0.32	3.58
<i>Calpurnia subdecandra</i>	1.3	1.4	2.5	5.2	<i>Ocimum lamifolium</i>	1.43	0.62	0.07	2.11
<i>Carrisa edulis</i>	1.78	1.02	0.5	3.3	<i>Ocotea kenyanensis</i>	2.14	0.69	0.26	3.1
<i>Cupressus lusitanica</i>	1.66	9	6.97	23	<i>Olea europaea</i>	6.8	0.87	12.5	20.17
<i>Calpurina aura</i>	3.2	0.87	0.4	4.47	<i>Olea capensis</i>	2.02	0.72	0.45	3.19
<i>Celtis africana</i>	0.83	0.22	1.71	2.76	<i>Olinia rochetiana</i>	2.26	0.87	0.1	3.23
<i>Chata edulis</i>	0.59	0.59	0.08	1.27	<i>Phytolacca dedocandra</i>	2.8	0.4	7.6	10.8
<i>Clerodendron myricoides</i>	0.95	0.37	0.1	1.42	<i>Pittosporum abyssinicum</i>	5.4	1.5	0.6	7.5
<i>Croton macrostachys</i>	3	1.86	7.4	12.26	<i>Podocarpus falcatus</i>	3.56	25.4	20.1	49.06
<i>Cordia africana</i>	1.78	1.61	1.23	4.62	<i>Polyscias fulva</i>	1.4	2.3	2.4	6.1
<i>Discopodium penninervium</i>	1.78	0.79	0.42	2.99	<i>Prunus africana</i>	1.07	0.32	1.09	2.49
<i>Dodonaea angustifolia</i>	1.43	0.52	0.04	1.99	<i>Premna schimperii</i>	3.8	0.72	0.6	5.12
<i>Dombeya torrida</i>	0.83	0.52	0.54	1.89	<i>Rapanea simensis</i>	1.54	0.45	0.03	2.02
<i>Dovyalis caffra</i>	0.95	0.74	0.05	1.74	<i>Rhamnus prenoideis</i>	5.2	0.62	0.01	5.82
<i>Dovyalis abyssinica</i>	1.07	0.47	0.05	1.59	<i>Rhus glutinosa</i>	3.5	1.4	3.8	8.7
<i>Dovyalis vericosa</i>	1.54	0.42	0.03	1.99	<i>Rosa abyssinica</i>	2.3	0.4	0.03	2.72
<i>Erythrina brucei</i>	0.95	0.47	0.68	2.1	<i>Rosmarinus officinalis</i>	0.59	0.27	0.04	0.9
<i>Ehretia cymosa</i>	1.66	1.11	0.68	3.46	<i>Rubus apetalus</i>	0.71	0.27	0.03	1.01
<i>Ekebergia capensis</i>	1.54	1.19	1.2	3.93	<i>Rubus studneri</i>	0.95	0.3	0.11	1.35
<i>Embelia schimpri</i>	2.26	1.36	0.96	4.58	<i>Rumex abyssinicus</i>	0.83	0.35	0.01	1.19
<i>Eucalyptus globulus</i>	8.6	12	10	30.6	<i>Rumex nervosus</i>	1.07	0.32	0.3	1.69
<i>Ficus sur</i>	1.9	1.49	4.38	7.76	<i>Rytigynia neglecta</i>	0.48	0.35	0	0.83
<i>Gardenia lutea</i>	2.49	0.79	0.21	3.5	<i>Schfflera abyssinica</i>	0.83	0.42	0.42	1.67
<i>Gravelea robusta</i>	4.5	0.9	0.8	6.2	<i>Sisyrinchium guineense</i>	0.24	0.05	0.07	0.36
<i>Hagenia abyssinica</i>	6.05	1.24	7.4	14.69	<i>Vernonia amygdalina</i>	4.65	0.87	0.6	6.12
<i>Hypericum revolutum</i>	1.9	1.02	0.04	2.95	<i>Vernonia auriculifera</i>	12	0.65	0.67	13.32
<i>Ilex mitis</i>	2.14	0.87	0.41	3.41	<i>Vernonia leopoldii</i>	1.07	0.47	0.06	1.6
<i>Jasminum abyssinicum</i>	1.19	0.42	0.03	1.64					
<i>Juniperus procera</i>	5.6	5.05	23.5	34.15					
<i>Lepidotrichilia volkensii</i>	1.07	0.37	0.33	1.77					

4. Conclusion and Recommendation

A total of 65 trees and shrub species belonging to 38 families were identified, out of these trees constituted 67.7% while shrubs were 32.3%. Sixteen families each with more than one species comprise 66.2 % of the total number of species recorded in the study area and the rest of 22 families each represent only one species. The Shannon index of diversity (H') for tree and shrub species in this study ranged from 0.004 to 0.3623 with the overall H' of 2.82 and the evenness values for tree species of the present study ranged between 0.001 to 0.087 with a mean value of 0.01 and overall evenness value of 0.67. A mean evenness value of 0.01 indicates that the relative homogeneity of tree species of the sampled forest plots was 1% of the maximum possible even population. The evenness values are not enough to justify uniformity in composition of tree species. This is expected because not all trees are equally distributed and there is some variation in the distribution of species in the study area. The mean IVI value in this study ranged between 0.36 to 49.06 with mean IVI value of 6.0. This value is lower when compared with some studies. In this study only 15 % of the recorded species were found with IVI values > 10 and the rest of 85% have IVI values < 10.

The high number of species richness in the study area is again attributed to the current protection measures given to the forest that via allows the regeneration of different species from soil seed bank. Some economic activities particularly, beekeeping through participatory forest management is highly encouraged in Ades dry afromontane forest by the presence of such floristic diversity. Further study regarding the richness and diversity of woody species to investigate the distribution of tree species for conservation and sustainable utilization is recommended.

Acknowledgements

We would like to thank Oda Bultum University for Grant funding of this Research. We appreciate and thank Office of Forestry and Wildlife Enterprise of Oromia National Regional State, Hararghe branch for providing us car during data collection. We also thank Forestry Experts of Doba District in West Hararghe Zone Administration who were involved in data collection. Finally we thank local elders who helped us in providing the local names of plant species.

References

- [1] Abera A. and Yasin A. 2018. Diversity and Abundance of Woody Plant Species of Assosa Forest Field Gene, Benishanigul Gumuz Regional State, Western Ethiopia. *International Journal of Plant Biology & Research* 6(5): 1100.
- [2] Abrham Abiyu, Vacik, H. and Glatzel, G. 2006. Population viability risk management applied to *Boswellia papyfera* (Del.) Hochst in Northeastern Ethiopia. *Journal of the Dry lands* 1 (2):98–107.
- [3] Austin M. P. & Heyligers P. C. 1989. Vegetation survey design for conservation: gradsect sampling of forests in north-eastern New South Wales. *Biological Conservation* 50:13-32.
- [4] Barbour M. D., Burk J. H. & Pitts W. D. 1987. Terrestrial plant ecology, 2nd Edition. Benjamin Cummings Inc., Menlo Park, CA, 634p.
- [5] Bekalo, T. H., S. D. Woodmatas & Z. A. Woldemariam. 2009. An ethnobotanical study of medicinal plants used by local people in the lowlands of Konta Special Woreda, southern nations, nationalities and peoples regional state, Ethiopia. *Journal of Ethnobiology and Ethnomedicine* 5:26-40.
- [6] Belaynah, A., Z. Asfaw, S. Demissew & N. F. Bussa. 2012. Medicinal plants potential and use by pastoral and agropastoral communities in Erer Valley of Babile Wereda, eastern Ethiopia. *Journal of Ethnobiology and Ethno medicine* 8:42-64.
- [7] Carlos Almazán-Núñez, R., María del Coro Arizmendi, Luis E. Eguarte and Pablo Corcuera. 2012. Changes in composition, diversity and structure of woody plants in successional stages of tropical dry forest in southwest Mexico. *Revista Mexicana de Biodiversidad* 83:1096-1109.
- [8] Cazzolla Gatti R, Vaglio Laurin G, Valentini R. 2017. Tree species diversity of three Ghanaian reserves. *iForest* 10: 362-368. – doi: 10.3832/for2056-010 [online 2017-03-07].
- [9] Chauhan DS, Dhanai CS, Bhupendra S, Chauhan S, Todaria NP, Coley PD, Barone JA. 1996. Herbivory and plant defenses in tropical forests *Ann. Rev. Ecol. Syst.* 27:305-335.
- [10] Deka J, Tripathi PO, Khan LM. 2012. High Dominance of *Shorea robusta* Gaertn. in Alluvial Plain Kamrup Sal Forest of Assam, N. E. *India Int. J. Ecosys.* 2(4):67-73.
- [11] Demel Teketay. 1997. Seedling populations and regeneration of woody species in dry Afromontane forests of Ethiopia. *Forest Ecology and Management* 98:149–165.
- [12] EFAP. 1994. Ethiopian Forestry Action Program. (EFAP), Addis Ababa.
- [13] EMA. 1988. National Atlas of Ethiopia. Ethiopian Mapping Authority, Addis Ababa.
- [14] Hedberg, I., Friis, I. and Edwards, S. 2004. Flora of Ethiopia and Eritrea. The National Herbarium, Addis Ababa University, Addis Ababa and Department of Systematic Botany, ppsala University, Uppsala.
- [15] Kent, M., and P. Coker. 1992. *Vegetation description and analysis* Belhaven Press, London.
- [16] Kessy, J. F. 1998. Conservation and Utilization of Natural Resources in the East Usambara Forest Reserves: Conventional Views and Local Perspectives. PhD thesis, 168pp.
- [17] Khan ML, Rai JPN, Tripathi RS. 1986. Regeneration and survival of tree seedlings and sprouts tropical deciduous and subtropical forests of Meghalaya, India. *Forest Ecology and Management* 14,293-304.
- [18] Kidanemariam K., Teshome S., Satish kumar B. 2015. Forest Carbon Stock in Woody Plants of Ades Forest, Western Hararghe Zone of Ethiopia and its Variation along Environmental Factors: Implication for Climate Change Mitigation. *Journal of Natural Sciences Research*. Vol. 5, No. 2.

- [19] Kindt, R. Noordin, Q., Njui, A. and Ruigu, S. 2005. Biodiversity Conservation through Agroforestry: Managing Tree Species Diversity within a Net work of Community-based Non-governmental, Governmental and Research Organizations in Western Kenya. Paper presented at 15th Annual Conference of the Eastern Africa Environmental Network on *Networking for Biodiversity*, 27-28 May, National Museum of Kenya, Nairobi.
- [20] Lou, J. 2006. Entropy and diversity. *Oikos* 113 (2): 363–375.
- [21] Magurran, A. E. 1988. Ecological Diversity and its Measurement. Princeton University Press, 192pp. Princeton
- [22] Malik ZA & Nautiyal MC. 2016. Species richness and diversity along the altitudinal gradient in Tungnath, the Himalayan benchmark site of HIMADRI. *Tropical Plant Research* 3(2): 396–407.
- [23] Mesfin Woldearegay, Zerihun Woldu and Ermias Lulekal. 2018. Species diversity, population structure and regeneration status of woody plants in Yegof dry afro-montane forest, Northeastern Ethiopia. *European Journal of Advanced Research in Biological and Life Sciences*. Vol. 6 No. 4. ISSN 2056-5984.
- [24] Muktar Reshad, Muktar Mohammed, Ahmed Mohammed and Alemayehu Beyene. 2017. Diversity of Non-Timber Forest Products (NTFPs) and its Source Plant Species: The Case of Jello-Muktar Forest, Eastern Ethiopia.
- [25] Muller-Dombois, D., H. Ellenberg. 1974. Aims and methods of vegetation ecology. John Wiley and Sons, New York. Region of Wisconsin. *Ecology* 32: 476-496.
- [26] Rogers, P. C. & Ryel, R. J. 2008. Lichen community change in response to succession in aspen forests of the Rocky Mountains, USA. *Forest Ecology and Management*, 256: 1760–1770.
- [27] Shiferaw W, Lemenih M & Gole TWM. 2018. Analysis of plant species diversity and forest structure in Arero dry Afro-montane forest of Borena zone, South Ethiopia *Tropical Plant Research* 5(2): 129–140.
- [28] Solbrig, O. T., E. Medina, and J. F. Silva. 1996. Biodiversity and Savana ecosystem process: *A Prespective Springer-Verlag* Berlin Hedelberg, Berlin.
- [29] United States Department of Agriculture Forest Service. 2007. Field methods instructions for Phase 2 (Forest Inventory) and Phase 3 (Forest Health) of the National Forest Inventory and Analysis program.
- [30] Wassie, A. & Teketay, D. 2005. Soil seed banks in Northern Ethiopia: implications for the conservation of woody plants. *Flora* 201:32–43.
- [31] Wilder, C., Brooks, T. & Lens, L. 1998. Vegetation structure and composition of the Taita Hills forest. *Journal of East African Natural History* 87:181–187.
- [32] Yineger, H., D. Yewhalaw & D. Teketay. 2008. Ethno medicinal plant knowledge and practices of the Oromo ethnic group in southwestern Ethiopia. *Journal of Ethnobiology and Ethnomedicine* 4:11.dx. doi.org/10.1186/1746-42694-11.
- [33] Zerihun Girma, George Chuyong, Paul Evangelista, and Yosef Mamo. 2018. Vascular Plant Species Composition, Relative Abundance, Distribution, and Threats in Arsi Mountains National Park, Ethiopia. *Mountain Research and Development* 38(2): 143-152.