RBFs for Integral Equations with a Weakly Singular Kernel

Jafar Biazar*, Mohammad Ali Asadi

Department of Applied Mathematics, Faculty of Mathematical Science, University of Guilan, Rasht, Iran

Email address:

biazar@guilan.ac.ir (J. Biazar), eng.asadi@gmail.com (M. A. Asadi)

To cite this article:

Jafar Biazar, Mohammad Ali Asadi. RBFs for Integral Equations with a Weakly Singular Kernel. *American Journal of Applied Mathematics*. Vol. 3, No. 6, 2015, pp. 250-255. doi: 10.11648/j.ajam.20150306.12

Abstract: In this paper a numerical method, based on collocation method and radial basis functions (RBF) is proposed for solving integral equations with a weakly singular kernel. Integrals appeared in the procedure of the solution are approximated by adaptive Lobatto quadrature rule. Illustrative examples are included to demonstrate the validity and applicability of the presented technique. In addition, the results of applying the method are compared with those of Homotopy perturbation, and Adomian decomposition methods.

Keywords: Integral Equations with a Weakly Singular Kernels, Radial Basis Functions (RBF), Adaptive Lobatto Quadrature, Homotopy Perturbation Method (HPM), Adomian Decomposition Method (ADM)

1. Introduction

Weakly singular integral equations have a major role in the fields of science and engineering. They appear in mathematical modeling of different phenomena in many disciplines such as: physics, chemistry, biology, and others. It is difficult to solve these equations analytically, hence numerical solutions are required. Singular integral equations have been approached by different methods including Collocation method [2-4], Reproducing kernel method [17], Galerkin method [5], Adomian decomposition method [1], Homotopy perturbation method [6], Radial Basis Functions [10, 11], Newton product integration method [7], and many others.

In recent years, meshless methods are used as a class of numerical methods for solving functional equations. Meshless methods just use a scattered set of collocation points, regardless any relationship information between the collocation points. This property is the main advantage of these techniques over the mesh dependent methods such as finite difference methods, finite element methods. A well-known family of meshless methods is the method of radial basis functions.

Since 1990, radial basis function method [8] is used as a meshless method to approximate the solutions of partial differential equations [9, 12-16]. These methods are developed for solving various types of linear and nonlinear functional equations such as Ordinary differential equations (ODEs). Integral and Integro-differential equations (IEs and

IDEs) are solved with the RBF method by some researchers [19-21]. In [22-24] authors used the RBF method to solve some engineering problems. Also authors of [25] solved fractional diffusion equations by RBFs.

In this paper, we will use an efficient method based on radial basis functions and collocation method to solve integral equations with a weakly singular kernel. The paper is organized as follows. In Section 2, the radial basis functions are introduced. Section 3, as the main part, presents the solution of weakly singular integral equations by RBF, via collocation method. Numerical illustrative examples are included in Section 4. A conclusion is drawn in Section 5.

2. Radial Basis Functions

Let's define the main features of the method.

2.1. Definition of Radial Basis Functions

Radial basis functions usually approximate a function as the following

$$s(x) = \sum_{i=1}^N \lambda_i \phi(\|x - x_i\|), \quad x \in \mathbb{R}.$$

Where $\phi: [0, \infty) \rightarrow \mathbb{R}$ is a fixed univariate function, the coefficients $(\lambda_i)_{i=1}^N$ are real numbers and $\|\cdot\|$, denotes the Euclidean norm.

2.2. Radial Basis Functions Interpolation

The radial basis functions approximation of a real function, say $u(x)$, is given by

$$u(x) \approx u_N(x) = \sum_{i=1}^N \lambda_i \phi(\|x - x_i\|) \\ = \sum_{i=1}^N \lambda_i \phi_i(x) = \Phi^T(x) \Lambda,$$

where

$$\phi_i(x) = \phi(\|x - x_i\|), \\ \Phi^T(x) = [\phi_1(x), \phi_2(x), \dots, \phi_N(x)], \\ \Lambda = [\lambda_1, \lambda_2, \dots, \lambda_N]^T,$$

and x_i , $i = 1, 2, \dots, N$ is a finite number of distinct points (centers) in $\mathbb{R}$. Consider N distinct support points $(x_j, u(x_j))$, $j = 1, 2, \dots, N$. One can find λ_i s by solving the following linear system

$$\mathbf{A} \Lambda = \mathbf{u}.$$

Where

$$\mathbf{A} = [\phi(\|x_j - x_i\|)]_{i,j=1}^N, \\ \Lambda = [\lambda_1, \lambda_2, \dots, \lambda_N]^T, \text{ and } \mathbf{u} = [u_1, u_2, \dots, u_N]^T.$$

Some well-known RBFs are listed in Table 1, where the Euclidian distance r is real and non-negative, and c is a positive scalar, called shape parameter.

Table 1. Some well-known RBFs.

Name of the function	Definition
Gaussian	$\phi(r) = e^{-cr^2}$
Inverse Quadric	$\phi(r) = \frac{1}{r^2 + c^2}$
Hardy Multiquadric	$\phi(r) = \sqrt{r^2 + c^2}$
Inverse Multiquadric	$\phi(r) = \frac{1}{\sqrt{r^2 + c^2}}$
Cubic	$\phi(r) = r^3$
Thin plate splines	$\phi(r) = r^2 \log(r)$
Hyperbolic secant	$\phi(r) = \text{sech}(cr)$

3. Application of RBF Method

In this paper Radial Basis Functions are used to approximate solution of integral equations with a weakly singular kernel in the following general form

$$y(x) = f(x) + \int_0^x \frac{y(t)}{\sqrt{x-t}} dt, \quad 0 \leq x \leq 1, \quad (1)$$

where $f(x)$ is a known analytic function defined on the interval $0 \leq x \leq 1$. In order to use the radial basis functions, let's consider $y(x)$ as follows,

$$y(x) \approx y_N(x) = \sum_{i=1}^N \lambda_i \phi_i(x) = \Phi^T(x) \Lambda. \quad (2)$$

Substitution in (1) from (2) leads to

$$\Phi^T(x) \Lambda = f(x) + \int_0^x \frac{\Phi^T(t) \Lambda}{\sqrt{x-t}} \Lambda dt, \quad (3)$$

To determine λ_i , $i = 1, 2, \dots, N$, from Eq. (3), let's use shifted zeros of the Legendre polynomials, $x = x_j$, $j = 1, 2, \dots, N$, as the collocation points,

$$\left(\Phi^T(x_j) - \int_0^{x_j} \frac{\Phi^T(t)}{\sqrt{x_j-t}} dt \right) \Lambda = f(x_j). \quad (4)$$

By using the Adaptive Lobatto Quadrature [18], we can approximate the integral in Eq. (4) to achieve a set of N linear equations for determining components of Λ .

4. Numerical Examples

In this section, two examples are provided to illustrate the efficiency of this approach. For the sake of comparing purposes, the same examples as [6], that have used Homotopy perturbation method, are considered.

4.1 Example

Let us consider the following weakly singular integral equation of the second kind

$$y(x) = x^2 + \frac{16}{15} x^{\frac{5}{2}} - \int_0^x \frac{y(t)}{\sqrt{x-t}} dt, \quad 0 \leq x \leq 1, \quad (5)$$

with the exact solution $y(x) = x^2$.

Let's approximate $y(x)$ as follows

$$y(x) \approx y_5(x) = \sum_{i=1}^5 \lambda_i e^{\frac{|x-x_i|^2}{100}}, \quad (6)$$

where

$$x_1 = 0.0469, \quad x_2 = 0.2308, \\ x_3 = 0.5000, \quad x_4 = 0.7692, \quad x_5 = 0.9531,$$

are shifted zeros of Legendre polynomial of degree 5. Substituting (6) into (5) gives

$$\sum_{i=1}^5 \lambda_i e^{\frac{|x-x_i|^2}{100}} = x^2 + \frac{16}{15} x^{\frac{5}{2}} - \sum_{i=1}^5 \lambda_i \int_0^x \frac{e^{\frac{|t-x_i|^2}{100}}}{\sqrt{x-t}} dt,$$

or

$$\sum_{i=1}^5 \left(e^{\frac{|x-x_i|^2}{100}} + \int_0^x \frac{e^{\frac{|t-x_i|^2}{100}}}{\sqrt{x-t}} dt \right) \lambda_i = x^2 + \frac{16}{15} x^{\frac{5}{2}}.$$

By using $x = x_j, j = 1, \dots, 5$ as collocation points, we have

$$\sum_{i=1}^5 \left(e^{\frac{|x_j-x_i|^2}{100}} + \int_0^{x_j} \frac{e^{\frac{|t-x_i|^2}{100}}}{\sqrt{x_j-t}} dt \right) \lambda_i = x_j^2 + \frac{16}{15} x_j^{\frac{5}{2}}. \quad (7)$$

Using adaptive Lobatto quadrature, integrals in Eq. (7) is computed and Eq. (7) is reduced to the following equation

$$\mathbf{A}\mathbf{\Lambda} = \mathbf{F} \quad (8)$$

where

$$\mathbf{A} = \begin{bmatrix} 1.4332 & 1.4327 & 1.4302 & 1.4256 & 1.4213 \\ 1.9603 & 1.9607 & 1.9588 & 1.9542 & 1.9494 \\ 2.4107 & 2.4130 & 2.4135 & 2.4105 & 2.4064 \\ 2.7442 & 2.7489 & 2.7525 & 2.7520 & 2.7495 \\ 2.9360 & 2.9426 & 2.9485 & 2.9503 & 2.9490 \end{bmatrix},$$

$$\mathbf{F} = \left[x_j^2 + \frac{16}{15} x_j^{\frac{5}{2}} \right]_{j=1}^5 = \begin{bmatrix} 0.0027 \\ 0.0805 \\ 0.4386 \\ 1.1453 \\ 1.8543 \end{bmatrix},$$

and $\mathbf{\Lambda} = [\lambda_1, \lambda_2, \dots, \lambda_5]^T$ is determined as the following

$$\mathbf{\Lambda} = \begin{bmatrix} 2.7558e+07 \\ -7.7979e+07 \\ 1.0093e+08 \\ -7.8189e+07 \\ 2.7683e+07 \end{bmatrix}. \quad (9)$$

Substituting from (9) into (6) yields an approximate solution as the following

$$y_5(x) = \left(2.7558e^{\frac{|x-0.0469|^2}{100}} - 7.7979e^{\frac{|x-0.2308|^2}{100}} + 10.093e^{\frac{|x-0.5000|^2}{100}} - 7.8189e^{\frac{|x-0.7692|^2}{100}} + 2.7683e^{\frac{|x-0.9531|^2}{100}} \right) \times 10^7,$$

with the maximum absolute error $7.9107e-08$.

More accurate approximations to the solution may be attained by considering additional terms. The Errors of the numerical solutions for $N=10$, and different RBFs are shown in the Table 2. Results in the Table 2 shows that all solutions which are achieved by Gaussian ($\phi(r) = \exp(-r^2/100)$), Inverse Quadric ($\phi(r) = 1/(1+r^2)$), and Multiquadric ($\phi(r) = \sqrt{1+r^2}$) RBFs are accurate enough for knowing this approach as a powerful device. Also the absolute error for this approach and Homotopy perturbation method [6] are shown in Figure 2. A Comparison which is made in Figure 2, shows that RBF method with Gaussian bases, gives more accurate solutions than the homotopy perturbation method by 18 terms. As x increases, the Error of HPM is increased rapidly, but RBF Errors are not significant yet.

Figure 1. Exact solution, $y_E(x)$, and approximate solution, $y_5(x)$, for Example 1.

Figure 2. The absolute Error of HPM, by 18 terms, and RBF, for $\phi(r) = e^{-(0.1r)^2}$.

4.2. Example

Now consider the following weakly singular integral equation

$$y(x) = \sqrt{x} + \frac{\pi}{2}x - \int_0^x \frac{y(t)}{\sqrt{x-t}} dt, \quad 0 \leq x \leq 1. \quad (10)$$

The exact solution is $y(x) = \sqrt{x}$.

First, $y(x)$ is approximated by

$$y_8(x) = \sum_{i=1}^8 \frac{\lambda_i}{1+(x-x_i)^2}, \quad (11)$$

where

$$x_1 = 0.0199, x_2 = 0.1017, x_3 = 0.2372, x_4 = 0.4083, \\ x_5 = 0.5917, x_6 = 0.7628, x_7 = 0.8983, x_8 = 0.9801,$$

are shifted zeros of Legendre polynomials of degree 8. Substituting (11) into (10) results in

$$\sum_{i=1}^8 \frac{\lambda_i}{1+(x-x_i)^2} = \sqrt{x} + \frac{\pi}{2}x - \sum_{i=1}^8 \lambda_i \int_0^x \frac{1+(t-x_i)^2}{\sqrt{x-t}} dt,$$

or

$$\sum_{i=1}^8 \left(\frac{1}{1+(x-x_i)^2} + \int_0^x \frac{dt}{(1+(t-x_i)^2)\sqrt{x-t}} \right) \lambda_i = \sqrt{x} + \frac{\pi}{2}x. \quad (12)$$

By using $x = x_j, j = 1, \dots, 8$ as collocation points, we have

$$\sum_{i=1}^8 \left(\frac{1}{1+(x_j-x_i)^2} + \int_0^{x_j} \frac{dt}{(1+(t-x_i)^2)\sqrt{x_j-t}} \right) \lambda_i = \sqrt{x_j} + \frac{\pi}{2}x_j. \quad (13)$$

Integrals in Eq. (13) is computed by adaptive Lobatto quadrature, and Eq. (13) is reduced to matrix equation

$$\mathbf{A}\mathbf{\Lambda} = \mathbf{F}, \quad (14)$$

where

$$\mathbf{F} = \left[\sqrt{x_j} + \frac{\pi}{2}x_j \right]_{j=1}^8 = \begin{bmatrix} 0.1721 \\ 0.4786 \\ 0.8597 \\ 1.2803 \\ 1.6987 \\ 2.0715 \\ 2.3589 \\ 2.5296 \end{bmatrix},$$

$$\mathbf{A} = \begin{bmatrix} 1.2818 & 1.2730 & 1.2232 & 1.1127 & 0.9647 & 0.8248 & 0.7224 & 0.6659 \\ 1.6290 & 1.6364 & 1.6014 & 1.4852 & 1.3067 & 1.1260 & 0.9893 & 0.9126 \\ 1.9063 & 1.9482 & 1.9635 & 1.8852 & 1.7072 & 1.4972 & 1.3261 & 1.2268 \\ 2.0568 & 2.1396 & 2.2295 & 2.2389 & 2.1183 & 1.9163 & 1.7260 & 1.6079 \\ 2.0799 & 2.1924 & 2.3480 & 2.4609 & 2.4477 & 2.3103 & 2.1373 & 2.0163 \\ 2.0303 & 2.1558 & 2.3482 & 2.5366 & 2.6306 & 2.5907 & 2.4732 & 2.3722 \\ 1.9662 & 2.0946 & 2.3004 & 2.5267 & 2.6904 & 2.7334 & 2.6790 & 2.6094 \\ 1.9222 & 2.0502 & 2.2591 & 2.4994 & 2.6951 & 2.7834 & 2.7700 & 2.7239 \end{bmatrix},$$

and $\mathbf{\Lambda} = [\lambda_1, \lambda_2, \dots, \lambda_8]^T$. System (14) is solved to determine components of $\mathbf{\Lambda}$ as below

$$\mathbf{\Lambda} = \begin{bmatrix} -687.02 \\ 1608.32 \\ -1719.90 \\ 1446.34 \\ -1225.73 \\ 1087.86 \\ -831.64 \\ 321.35 \end{bmatrix}. \quad (15)$$

Substituting from (15) into (11) gives an approximate solution as the following

$$y_8(x) = -\frac{687.02}{1+(x-0.0199)^2} + \frac{1608.32}{1+(x-0.1017)^2} \\ -\frac{1719.90}{1+(x-0.2372)^2} + \frac{1446.34}{1+(x-0.4083)^2} \\ -\frac{1225.73}{1+(x-0.5917)^2} + \frac{1087.86}{1+(x-0.7628)^2} \\ -\frac{831.64}{1+(x-0.8983)^2} + \frac{321.35}{1+(x-0.9801)^2},$$

with the maximum error $2.4342e-03$.

Figure 3. Exact solution, $y_E(x)$, and approximate solution, $y_8(x)$, for Example 2.

Figure 4. The absolute Error of HPM, by 18 terms, and RBF, for $\phi(r) = \sqrt{1+r^2}$.

Errors of the numerical solutions for $N=10$, and Multiquadric ($\phi(r) = \sqrt{1+r^2}$), Inverse Quadric ($\phi(r) = 1/(1+r^2)$), and Inverse Multiquadric ($\phi(r) = 1/\sqrt{1+r^2}$) RBFs are shown in the Table 3. These results show that all solutions which are achieved by these different RBFs are accurate enough for knowing RBF method as a powerful device. The absolute error for this approach and HPM [6] are shown in Figure 4. A Comparison which is made in Figure 4, shows that RBF method with Multiquadric bases

gives more accurate solutions than the homotopy perturbation method by 18 terms. As x increases, the Error of HPM is increased rapidly, but RBF Error vanishes.

5. Conclusion

This paper describes using Radial basis functions to solve integral equations, with a weakly singular kernel. The proposed method, is a meshless method, just uses a scattered set of collocation points, without any connectivity information between the collocation points. Two examples are presented to show the efficiency of the method. Numerical results at scattered points appeared in Tables 2 and 3. Exact and approximate solutions for two examples are plotted in Figures 1 and 3. Figures confirms that errors is not significant and solutions are satisfactory even by a few number of basis functions. The behavior of the errors for Homotopy perturbation method and Radial basis functions is apparent in Figures 2 and 4 for two examples. A comparison with Homotopy perturbation method shows that RBF results in more accurate solutions than Homotopy perturbation method or Adomian decomposition method which are reported in [6]. The results of two solved examples are accurate enough for knowing this approach as a powerful device. From physical point of view, these figures show that in the RBF method, as x increases, equilibrium appears.

Table 2. Solutions and Errors for Example 1.

x	y_{Exact} (y_E)	$\phi(r) = \exp(-r^2/100)$		$\phi(r) = 1/(1+r^2)$		$\phi(r) = \sqrt{1+r^2}$	
		$y_{10}(x)$	$ y_{10} - y_E $	$y_{10}(x)$	$ y_{10} - y_E $	$y_{10}(x)$	$ y_{10} - y_E $
0.0130	0.00017021	0.00017026	4.3354e-08	0.00017112	9.0317e-07	0.00017027	5.5240e-08
0.0675	0.00455197	0.00455194	2.6635e-08	0.00455079	1.1825e-06	0.00455193	3.4940e-08
0.1603	0.02569455	0.02569460	5.2464e-08	0.02569624	1.6871e-06	0.02569754	2.9851e-06
0.2833	0.08026019	0.08026009	9.6868e-08	0.08025590	4.2943e-06	0.08025913	1.0604e-06
0.4256	0.18110372	0.18110373	1.3451e-08	0.18110999	6.2725e-06	0.18110412	4.0251e-07
0.5744	0.32997806	0.32997792	1.3210e-07	0.32997235	5.7053e-06	0.32997729	7.6181e-07
0.7167	0.51365558	0.51365527	3.1387e-07	0.51366105	5.4612e-06	0.51365605	4.6475e-07
0.8397	0.70510412	0.70510411	1.1898e-08	0.70509947	4.6490e-06	0.70510354	5.7539e-07
0.9325	0.86961534	0.86961540	6.5357e-08	0.86961939	4.0569e-06	0.86961571	3.7189e-07
0.9870	0.97407674	0.97407674	2.0636e-09	0.97407264	4.0986e-06	0.97407624	4.9822e-07

Table 3. Solutions and Errors for Example 2.

x	y_{Exact} (y_E)	$\phi(r) = \sqrt{1+r^2}$		$\phi(r) = 1/(1+r^2)$		$\phi(r) = 1/\sqrt{1+r^2}$	
		$y_{10}(x)$	$ y_{10} - y_E $	$y_{10}(x)$	$ y_{10} - y_E $	$y_{10}(x)$	$ y_{10} - y_E $
0.0130	0.11422230	0.11262579	1.5965e-03	0.11261085	1.6115e-03	0.11263024	1.5921e-03
0.0675	0.25974663	0.26016436	4.1772e-04	0.26017013	4.2349e-04	0.26014565	3.9901e-04
0.1603	0.40036884	0.40101248	6.4363e-04	0.39988836	4.8048e-04	0.39993226	4.3658e-04
0.2833	0.53226149	0.53222145	4.0049e-05	0.53259710	3.3560e-04	0.53297792	7.1643e-04
0.4256	0.65235176	0.65206011	2.9166e-04	0.65214515	2.0661e-04	0.65175952	5.9224e-04
0.5744	0.75791633	0.75799302	7.6694e-05	0.75810850	1.9217e-04	0.75816263	2.4630e-04
0.7167	0.84658000	0.84641812	1.6187e-04	0.84638985	1.9014e-04	0.84637489	2.0511e-04
0.8397	0.91635407	0.91638214	2.8074e-05	0.91651646	1.6240e-04	0.91644249	8.8420e-05
0.9325	0.96567680	0.96559109	8.5710e-05	0.96555392	1.2288e-04	0.96558939	8.7405e-05
0.9870	0.99345521	0.99346109	5.8756e-06	0.99335758	9.7631e-05	0.99349627	4.1062e-05

References

- [1] Y. Mahmoudi, "A new modified Adomian decomposition method for solving a class of hypersingular integral equations of second kind", *J. Comput. Appl. Math.* Vol. 255, 2014, pp. 737-742.
- [2] P. Junghanns, K. Muller, "A collocation method for nonlinear Cauchy singular integral equations", *J. Comput. Appl. Math.* Vol. 115, 2000, pp.283-300.
- [3] P. Junghanns, R. Kaiser, "Collocation for Cauchy singular integral equations", *Linear Algebra Appl.* Vol. 439, 2013, pp. 729-770.
- [4] S. Pishbin, F. Ghoreishi, M. Hadizadeh, "The semi-explicit Volterra integral algebraic equations with weakly singular kernels: The numerical treatments", *J. Comput. Appl. Math.* Vol. 245, 2013, pp. 121-132.
- [5] K. S. Thomas, "Galerkin Methods for Singular Integral Equations", *Math. Comp.* Vol. 36, No. 153, 1981 pp. 193-205.
- [6] R. K. Pandey, O. P. Singh, V. K. Singh, "Efficient algorithms to solve singular integral equations of Abel type", *Comput. Math. Appl.* Vol. 57, 2009, pp. 664-676.
- [7] B. Babayar-Razlighi, B. Soltanalizadeh, "Numerical solution of a nonlinear singular Volterra integral system by the Newton product integration method", *Math. Comput. Model.* Vol. 58, No. 11-12, 2013, pp. 1696-1703.
- [8] R. L. Hardy, "Multiquadric equations of topography and other irregular surface", *J. Geophys. Res.* Vol. 76, 1971, pp. 1905-1915.
- [9] C. Franke, R. Schaback, "Solving partial differential equations by collocation using radial basis functions", *Appl. Math. Comput.* Vol. 93, 1997, pp. 73-82.
- [10] A. Golbabai1, S. Seifollahi, R. Gholami, "Numerical solution of weakly singular integral equations by a growing RBF network", *Int. J. Pure Appl. Math.* Vol. 46, No. 2, 2008, pp. 191-197.
- [11] K. Parand, J. A. Rad, "An approximation algorithm for the solution of the singularly perturbed Volterra integro-differential and Volterra integral equations", *Int. J. Nonlinear Sci.* Vol. 12, No. 4, 2011, pp. 430-441.
- [12] E. J. Kansa, "Multiquadrics scattered data approximation scheme with applications to computational fluid dynamics II. Solutions to hyperbolic, parabolic, and elliptic partial differential equations", *Comput. Math. Appl.* Vol.19, 1990, pp. 147-161.
- [13] R. B. Platte, T. A. Driscoll, "Computing eigenmodes of elliptic operators using radial basis functions", *Comput. Math. Appl.* Vol. 48, 2004, pp. 561-576.
- [14] E. Larsson, B. Fornberg, "A numerical study of some radial basis function based solution methods for elliptic PDEs", *Comput. Math. Appl.* Vol. 46, 2003, pp. 891-902.
- [15] M. Dehghan, "Finite difference procedures for solving a problem arising in modeling and design of certain optoelectronic devices", *Math. Comput. Simulat.* Vol. 71, 2006, pp. 16-30.
- [16] M. Dehghan, "Solution of a partial integro-differential equation arising from viscoelasticity", *Int. J. Comput. Math.* Vol. 83, 2006, pp. 123-129.
- [17] Z. Chen, W. Jiang, "The exact solution of a class of Volterra integral equation with weakly singular kernel", *Appl. Math. Comput.* Vol. 217, 2011, pp. 7515-7519.
- [18] W. Gander, W. Gautschi, "Adaptive Quadrature-Revisited", *Swets & Zeitlinger Publishers*, Vol. 40, No. 1, 2000, pp. 84-101.
- [19] P. Assari, H. Adibi, M. Dehghan, A meshless method for solving nonlinear two-dimensional integral equations of the second kind on non-rectangular domains using radial basis functions with error analysis, *J. Comput. Appl. Math.*, Vol. 239, 2013, pp. 72-92.
- [20] J. Biazar, M. A. Asadi, Indirect RBF for High-Order Integro-Differential Equations, *Brit. J. Math. Comp. Sci.*, Vol. 11, No. 1, 2015, 1-16, Article no. BJMCS 19199.
- [21] J. Biazar, M. A. Asadi, Galerkin RBF for Integro-Differential Equations, *Brit. J. Math. Comp. Sci.*, Vol. 11, No. 2, 2015, 1-9, Article no. BJMCS 19265.
- [22] A. J. M. Ferreira, E. Carrera, M. Cinefra, and C. M. C. Roque, Radial basis functions collocation for the bending and free vibration analysis of laminated plates using the Reissner-Mixed Variational Theorem, *Eur J Mech A-Solid*, Vol. 39, 2013, pp. 104-112.
- [23] M. Dehghan, V. Mohammadi, The numerical solution of Fokker-Planck equation with radial basis functions (RBFs) based on the meshless technique of Kansa's approach and Galerkin method, *Eng. Anal. Bound. Elem.*, Vol. 47, 2014, pp. 38-63.
- [24] M. Dehghan, M. Abbaszadeh, and A. Mohebbi, The numerical solution of nonlinear high dimensional generalized Benjamin-Bona-Mahony-Burgers equation via the meshless method of radial basis functions, *Comput. Math. Appl.*, Vol. 68, No. 3, 2014, pp. 212-237.
- [25] C. Piret, E. Hanert, A radial basis functions method for fractional diffusion equations, *J. Comput. Phys.*, Vol. 238, 2013, pp. 71-81.