Genetic Evaluation of Iraqi Awassi Rams According to Some Growth Traits of Their Progeny Lambs

Firas R. Al-Samarai¹, Fatten A. Mohammed², Falah H. Al-Zaydi², Nasr N. Al-Anbari³, Yehya K. Abdulrahman¹

¹Department of Veterinary, Public Health, College of Veterinary Medicine, University of Baghdad, Baghdad, Iraq

²Department of Animal Resources, Directorate of Baghdad Agriculture, Ministry of Agriculture, Baghdad, Iraq

³Department of Animal Resources, College of Agriculture, University of Baghdad, Baghdad, Iraq

Email address:

firas_rashad@yahoo.com (F. R. Al-Samarai), fatten_2000@yahoo.com (F. A. Mohammed), falah_vet@yahoo.com (F. H. Al-Zaidi), nasr_noori@yahoo.com (N. N. Al-Anbari), yehyeabulrahman@yahoo.com (Y. K. Abdulrahman)

To cite this article:

Firas R. Al-Samarai, Fatten A. Mohammed, Falah H. Al-Zaydi, Nasr N. Al-Anbari, Yehya K. Abdulrahman. Genetic Evaluation of Iraqi Awassi Rams for Some Growth Traits of Their Lambs. *American Journal of Applied Scientific Research*. Vol. 2, No. 1, 2016, pp. 1-5. doi: 10.11648/j.ajarsr.20160201.11

Abstract: This study was carried out on the research station of sheep and goat (Abo Gharib, west of Baghdad, Iraq) to evaluate the 120 Awassi rams for the birth weight (BW), weight at weaning (WW) and pre-weaning daily gains (DG) of their lambs after adjusting to the effects of parity, year of birth, sex, and type of birth (single, twin). Restricted Maximum Likelihood (REML) method in a mixed model (paternal half sib) was utilized to estimate the heritability (h^2) for above mentioned traits. Proc mixed in SAS program was used to estimate Best Linear Unbiased Prediction (BLUP) values of rams for the three studied traits. The results revealed that the means of BW, WW and DG were 3.85, 26.64 and 0.181kg respectively. The studied traits were significantly ($P < 0.01$) affected by all fixed effects except the effect of ram on WW and DG which was significant ($P < 0.05$). Heritability estimates (h^2) of BW, WW and DG were 0.23, 0.12 and 0.19 respectively. The lowest and highest BLUP values for the mentioned traits were -0.325, 0.255 kg, -1.142, 1.284 kg, and -0.103, 0.053 kg respectively. In conclusion: the results indicated that selection of rams for DG and BW could play an important role for improving the growth traits of Awassi sheep due to the high range among BLUP values along with the acceptable estimations of heritability.

Keywords: Awassi Lamb, Birth Weight, Weaning Weight, Heritability

1. Introduction

Sheep husbandry in Iraq has been a historically important component of rural development and still fulfills a sustainable role in the livelihood of farmers. The country has a tradition of the consumption of sheep products, especially lamb and mutton. The native sheep breeds in Iraq include the Karadi (Kurdi, Hamdani, Jaff and Dzaie) 20%, Awassi (Naami and Shefali) 58.2% and Arabi 21.8% (Al-Barzinji and Othman, 2013). The Awassi sheep breed has been introduced into many countries, and have been shown to have superior performance to some native breeds (Todorovski, 1988). This breed was widespread because of its good characteristics in regards to meat price and quality (Kingwell et al., 1995), milk quality (Sunderman and Johns, 1994), validity of wool for the carpet industry (Lightfoot, 1988), and its ability to cope stress of high environmental temperature (AbiSaab and Sleiman, 1995). Unfortunately, the Awassi has

taken its place among the genotypes of indigenous genetic resources requiring a protection project due to their declining numbers (Üstüner and Oğan, 2013).

Today on sheep farming, a large part of the economic income is based on meat production. Consequently, it is observed that studies aimed to increase lambs productivity and growth performance in lambs, which are the main source of meat production, have intensified (Özcan et al., 2001).

In order to devise effective breeding plans for genetic improvement of Awassi sheep, information on the extent of genetic and environmental factors on performance traits is the pre-requisite. Therefore; this study was planned to generate information on the relative importance of genetic and environmental factors on the growth performance of Awassi sheep in addition to estimate the heritability and Best Linear Unbiased Prediction (BLUP) for some growth traits.

2. Material and Methods

In this study, records of Awassi lambs bred at the research station of sheep and goats (Abu Gharib, west of Baghdad, Iraq) were utilized over three years (2007 – 2009). Data included 1318 birth weight (BW), 821 weaning weight (WW) and 821 pre-weaning daily gain (DG) records. The numbers of ewes and rams were 849 and 120 respectively.

Restricted Maximum Likelihood (REML) method in a mixed model (paternal half sib) was used to estimate the heritability (h^2) for mentioned traits. The general statistical model included fixed effects due to parity, sex, year of birth, month of birth, type of birth and age at weaning as covariate.

Proc mixed in SAS program (2010) was used to estimate Best Linear Unbiased Prediction (BLUP) values of rams for all studied traits.

SAS program Ver. 9.1 was used for data analysis. Two

Mathematical models were assumed as shown below:

Model I for BW.

$$Y_{ijklmno} = \mu + \pi_i + Y_j + S_k + T_l + M_m + F_n + e_{ijklmno}$$

Where; $Y_{ijklmno}$ = Birth weight; μ = Population mean; π_i = Effect of parity (1-3); Y_j = Year of birth, 2007, 2008, 2009; S_k = Sex, Male, female; T_l = Type of birth (Single, twin); M_m = Month of birth (January, February, and March); $e_{ijklmno}$ = random error associated with each observation. It is assumed to be normally distributed with mean zero and variance σ^2 .

Model II for WW and DG.

As weaning lambs in this station depend on weight rather than age, the WW and DG were adjusted for age at weaning.

$$Y_{ijklmno} = \mu + \pi_i + Y_j + S_k + T_l + M_m + F_n + b(X_{ijklmn}) + e_{ijklmno}$$

The notation of the second model is similar to the first model except b = regression coefficient; X_{ijklmn} = Age of lamb at weaning.

Table 1. Least square means $\pm$ SE (kg) for birth weight (BW) (kg), weaning weight (WW) and daily gain (DG) in Awassi lambs.

Sources of variation	No. of records	BW Mean $\pm$ SE	No. of records	WW Mean $\pm$ SE	DG Mean $\pm$ SE
Parity					
1 st	549	3.67 $\pm$ 0.14 ^b	239	26.82 $\pm$ 0.88 ^a	0.164 $\pm$ 0.015 ^b
2 nd	539	4.19 $\pm$ 0.11 ^a	354	28.73 $\pm$ 0.68 ^a	0.191 $\pm$ 0.012 ^a
3 rd	230	4.28 $\pm$ 0.16 ^a	228	23.48 $\pm$ 1.09 ^b	0.139 $\pm$ 0.012 ^c
Year of birth					
2007	528	3.64 $\pm$ 0.12 ^b	222	31.13 $\pm$ 0.88 ^a	0.221 $\pm$ 0.013 ^a
2008	574	3.78 $\pm$ 0.12 ^b	383	24.06 $\pm$ 0.64 ^b	0.141 $\pm$ 0.013 ^b
2009	216	4.68 $\pm$ 0.17 ^a	216	23.84 $\pm$ 1.16 ^c	0.132 $\pm$ 0.013 ^b
Month of birth					
January	326	4.25 $\pm$ 0.04 ^a	233	29.18 $\pm$ 0.36 ^a	0.202 $\pm$ 0.005 ^a
February	804	4.03 $\pm$ 0.02 ^b	483	26.20 $\pm$ 0.24 ^b	0.179 $\pm$ 0.004 ^b
March	188	3.83 $\pm$ 0.05 ^c	105	23.65 $\pm$ 0.52 ^c	0.151 $\pm$ 0.007 ^c
Type of birth					
Single	676	4.33 $\pm$ 0.03 ^a	416	28.21 $\pm$ 0.29 ^a	0.189 $\pm$ 0.004 ^a
Twin	642	3.74 $\pm$ 0.03 ^b	405	24.47 $\pm$ 0.30 ^b	0.166 $\pm$ 0.004 ^b
Sex					
Male	653	4.15 $\pm$ 0.03 ^a	398	27.77 $\pm$ 0.31 ^a	0.189 $\pm$ 0.004 ^a
Female	665	3.92 $\pm$ 0.03 ^b	423	24.92 $\pm$ 0.29 ^b	0.166 $\pm$ 0.003 ^b
Reg. on age at weaning				0.174 $\pm$ 0.019 ^{**}	0.003 $\pm$ 0.0001 ^{**}
Overall mean	1318	3.85 $\pm$ 0.02	821	26.64 $\pm$ 0.25	0.181 $\pm$ 0.003

Means of the same column with different letters differ significantly at $P < 0.01$.

** ($P < 0.01$).

Table 2. Heritability estimates and BLUP values of birth weight, weaning weight, and pre-weaning daily gain (kg) in Iraqi Awassi rams.

		Birth weight		Weaning weight		Daily gain
Heritability		0.23 $\pm$ 0.04		0.12 $\pm$ 0.04		0.19 $\pm$ 0.04
Sequence	Sire No.	BLUP	Ram No.	BLUP	Ram No.	BLUP
1	12345	-0.32520	11996.	-1.14220	11873.	-0.10330
2	11920	-0.26250	11877.	-0.90160	11877.	-0.09137
3	11975	-0.24560	11958.	-0.79060	11878.	-0.06243
4	1239	-0.15690	11873.	-0.76560	11958.	-0.03495
5	11911	-0.15100	11941.	-0.66780	11961.	-0.02903
-	-	-	-	-	-	-
116	11916	0.17260	11948.	0.57440	11914.	0.02901
117	11925	0.19210	11908.	0.63980	11931.	0.03187
118	11978	0.19350	11914.	0.77440	11989.	0.03926
119	11964	0.21480	11913.	0.77760	1748.	0.04575
120	12241	0.25520	11989.	1.28400	11913.	0.05321

3. Results and Discussion

Least square means $\pm$ SE of BW, WW and DG of the Awassi lambs in relation to parity, month and year of lambing, type of birth and sex are presented in Table 1. The overall means of BW, WW and DG were 3.85 ± 0.02 kg, 26.64 ± 0.25 kg and 0.181 ± 0.003 kg respectively.

Estimates of heritability (h^2) and BLUP values are in Table 2. The heritability of the BW, WW and DG was 0.23, 0.19 and 0.12 respectively.

The BLUP values of rams for the studied traits are shown in table 2. The lowest BLUP values for BW, WW, and DG were -0.325, -1.142 and -0.103 kg respectively. The corresponding values of highest BLUP values were 0.225, 1.284, and 0.053 kg.

The mean of the BW in current study was lower than the range of the means (4.05 to 4.52 kg) reported by several researchers for Awassi lambs (Esenbuğa and Dayioğlu, 2002; Hassen et al., 2004; Dikmen et al., 2007; Jawasreh and Khasawneh, 2007; Kridli et al., 2007; Tabbaa et al., 2008; Üstüner and Oğan, 2013). However, it is consistent with other estimates of 3.82, 3.70 and 3.67 kg reported by Al-Kass et al., (1986), Al-Wahab (2003) and Al-Khazrji et al., (2014) respectively.

The results of the current research showed that rams had significant ($P < 0.01$) effect on BW. Similar results were obtained by Al-Hilali, (1982), and Alkass et al., (1991). The parity affected the BW significantly ($P < 0.01$). The mean of BW increased with advancing parity and this could be attributed to increasing of ewe's weight. According to Křížek et al., (1983), live weight of dams significantly affected live weight of lambs at birth. Hence, balanced feeding for dam could lead to the heavier lambs at birth (Obaido, 2010). Similar finding was reported by Ghoneim et al. (1982) and Al-Khazrji et al. (2014).

The effect of year of birth on the BW was significant ($P < 0.01$). This effect reflects the variation of the availability of rainfall and pastures among different years. The month of birth also influenced BW. Lambs born early in the lambing period (January and February) surpassed those born later (March). This could be attributed to variation in quantity and quality of dam's nutrition through the gestation period particularly in the last months. Similar results were reported by Khalaf et al., (2010) who found a similar significant ($P < 0.01$) trend where BW was 3.82, 3.46 and 3.24 kg for January, February and March respectively.

The type of birth and sex affected BW significantly ($P < 0.01$). The results showed that the birth weight was higher in males compared to females and in single-born lambs compared to twins. These results were in consistent with those reported by Dikmen et al., (2007). Singles and males generally had higher birth weights than twin births and females. These results confirmed the result obtained by Üstüner and Oğan (2013).

Results revealed that the mean of the WW (26.64kg) was

lower than means of 31.29, 29.14 and 27.75 kg reported by Özcan et al. (2001), Üstüner and Oğan, (2013) and Khalaf et al., (2010) respectively. On the other hand it is higher than the value of 21.54 to 24.30 kg reported by Al- Jalili et al., (2006), Al-Wahab, (2003), Aksakal et al., (2009) and Al-Salman (2009).

WW differed significantly ($P < 0.01$) among parities where it was heaviest in the 2nd parity than those born in the 1st and 3rd parities.

This could be attributed to lamb growth rate, which is mainly affected by the dams' milk yield. High milking ewes' lambs grow faster as compared to the poor milkers (Obaido, 2010). This result was in agreement with results obtained by Al-Salman, (2009) and Al-Khazrji et al. (2014), but disagreed with others (Jawasreh and Khasawneh, 2007; Üstüner and Oğan, 2013);

There was a significant ($P < 0.01$) decreasing in the WW a cross years of birth. Year of birth within locations generally showed highly significant effects on BW, WW, and yearling weight. Fluctuations in some environmental factors prevailed; in particularly quality and quantity of the available feed stuff could be an explanation (AKF, 2006).

The month of birth also influences WW. Lambs born early in the lambing period (January and February) will gain weight better than those born late (March) due to the accessibility to the pastures in spring season. Lambs born late will not be able to use pasture in spring because of their young age besides they may have higher exposure to internal parasites which thrive in the high temperature. Thus, their weaning and yearling weights will be lower than those lambed early (Elwakil et al., 2009).

The differences in WW between the months of lambing were attributed to the yearly variation in the rain precipitation and its effect on the density, growth and availability of pastures, forage, and other feeds. Similarly, different climates have been reported to influence milk production of ewes. This indirectly affects the growth of lambs (Shaker et al., (2002). The results achieved in this study are in congruency with Al-Salman, (2009) who found that WW of lams born in November and December had significantly ($P < 0.05$) heavier WW (25.98 and 25.74kg) as compared with those born in January and February (22.63 and 20.85 kg).

In the current study, the type of birth and sex affected significantly ($P < 0.01$) the WW. The WW was higher in males compared to females and in single-born lambs compared to twins. Similar finding was obtained by Dikmen et al., (2007), Al-Salman (2009) and Üstüner and Oğan (2013).

The regression coefficient of WW on age at weaning was positive and significant ($P < 0.01$). Thus, it is an important to adjusting WW for this variation in age at weaning.

The present estimate of DG (0.181 kg) appeared to be comparable with the corresponding estimates of Awassi lambs (0.168 to 0.205 kg) reported by other researchers (Al-Salman, 2009; Üstüner and Oğan, 2013; Al-Khazrji et al., 2014).

The impact of all fixed effects on DG was significant ($P < 0.01$). The trend of changes in DG due to fixed effects is parallel to changes in WW. This could be attributed to the effect of the DG on WW.

Paternal half-sib heritability for BW, WW, and DG was given in Table 2. Their values were considered low to moderate. Heritability estimates of 0.10, 0.19, 0.18, and 0.16 for BW of Awassi lambs were reported by Kazzal, (1973), Thrift *et al.*, (1973), Aziz, (1977), Gursoy *et al.*, (1995). A higher estimates of 0.29 and 0.41 for the same breed were obtained by Al-Hilali, (1982), and Alkass *et al.*, (1991). The heritability estimate of the BW in the present study indicated that rams differed in their genetic potential.

The heritability estimate of the WW (0.12) was much lower than the estimate of the BW. This could be attributed to variation in maternal environment, mainly milk supply rather than to differences in genetic merit of the lambs. This estimate was similar to some estimations (0.10, 0.12 and 0.10) reported by Thrift *et al.*, (1973), Dzakumah *et al.* (1978) and Alkass *et al.*, (1991).

In current work, heritability of DG (0.19) was higher than 0.12 and 0.07 reported by Al-Rawi *et al.*, (1982) and, Kamber, (1987) for the same breed, while our estimate was lower than 0.51 reported by Alkass *et al.*, (1991). Differences in heritability estimates among various studies for the same trait of the same breed could be due to differences in the records number used, the correction for different non-genetic factors, the model used and the methodology for estimating heritability of the trait (Abou-Bakr, 2009).

The lowest and highest BLUP values (Table 2) for the three traits were expressed as the percentages of the overall means for each trait to identify its weighted effect. The percentages of the lowest value of BW, WW, and DG were 8.4%, 4.2% and 56% respectively, whereas, the corresponding percentages for the highest values were 6.6%, 4.8%, and 29% respectively. Hence, the selection of the rams for the DG would be effective in raising this trait of Awassi lambs. However, the h^2 of DG (0.19) was little lower than the h^2 (0.23) of BW. The selection of the rams for the BW would be also effective as the h^2 of BW was moderate. The expected genetic gain per generation under mass selection would be the product of the selection differential and heritability. On the other hand, the differences between highest and lowest percentage of BLUP values for WW was not high enough to justify selection procedures for its improvement particularly the trait has low h^2 (0.12).

Results of the year of birth effects in the present study indicated that while BW generally tends to increase along with advancing years, the WW and DG tend to decrease. Despite the breeding plan implemented in this flock, management as well as the effectiveness of the breeding plan has to be considered in order to improve mutton production. In conclusion: the results indicated that selection of rams for DG and BW could play an important role for improving the growth traits of Awassi sheep due to the high range among BLUP values along with the acceptable estimations of heritability.

References

- [1] AbiSaab, S., Sleiman, F. T. (1995). Physiological responses to stress of filial crosses compared to local Awassi sheep. *Small Rum. Res.*, 16: 55-59.
- [2] Abou-Bakr, S. (2009). Genetic and phenotypic trends of 305-day milk yield of Holstein cows raised at commercial farm in Egypt. *Egyptian J. Anim. Prod.*, 46: 85-92.
- [3] Aga Khan Foundation (AKF). (2006). Report of the Rural Support Program Activities. Salamieh, Hama (in Arabic).
- [4] Aksakal, V., Macit, M., Esenbuga, N. (2009). Effect of various ages of weaning on growth characteristics, survival rate and some body measurements of Awassi lambs. *J. Anim. Vet. Adv.*, 8: 1624-1630.
- [5] Al-Barzinji, Y. M. S., Othman, G. U. (2013). Genetic Polymorphism in *FecB* Gene in Iraqi Sheep Breeds Using RFLP-PCR Technique. *IOSR Journal of Agriculture and Veterinary Science*. 2: 46-48.
- [6] Al-Hilliali, A. H. (1982). Genetic analysis of some post-weaning productive characteristics in Awassi sheep. M. Sc. Thesis, Agricultural University Baghdad, Baghdad.
- [7] Al-Jalili, Z. F., Al-Azzawi, W. A. R., Mohammed, Q. S. (2006). The effect of type of birth of both ewe and lambs and other factors on lamb birth and weaning weights. *Egyptian J. Sheep, Goat Desert Anim. Sci.*, 1: 31-40.
- [8] Alkass, J. E., Alrawi, A. A., Al-Hillali, A. H., Al-Tayy, H. M. A. (1986). Body weight of horn-vs-polled-sired Awassi lambs. 3rd World Congress on Genetics Applied to Livestock Production. Paper 3. <http://digitalcommons.unl.edu/wcgalp/3>.
- [9] Alkass, J. E., Aziz, D. A., Hermiz, D. A. (1991). Genetic parameters of growth traits in Awassi sheep. *Emirates J. of Agric. Sci.*, 3: 152-161.
- [10] Al-Khazrji, W. J., Al-Samari, W. A., Ajil, A. M. (2014). The effect of some genetic and nongenetic factors on fertility and growth traits in Awassi sheep. *The Iraqi J. Agric. Sci.* 45: 205-211.
- [11] Al-Rawi, A. A., Badawi, F. S., Said, A. S. I., Faraj, M. S. (1982). Genetic and phenotypic Parameter estimates for growth traits in awassi sheep. (Abst.) Proceeding American Society Animal Science Meeting, 38: 35.
- [12] Al-Salman, M. H. (2009). Some factors influencing growth of Awassi lambs and their crosses from birth till weaning and weight at the age of 6 months, *Al-Anbar J. Agric. Sci.*, 7: 128-136.
- [13] Al-Wahab, R. M. H. (2003). Effects of crossing Finnrams with Awassi sheep on prolificacy and other quantitative traits, *J. Bio. Sci.*, 3: 32-42.
- [14] Aziz, D. A. (1977). Genetic, non-genetic parameters of some traits in Awassi sheep. MS. Thesis, College of Agriculture, Forestry, Univ. of Mosul (In Arabic).
- [15] Dikmen, S., Turkmen, I. I., Üstüner, H., Alpay, F., Balci, F., Petek, M., Ogan, M. M. (2007). Effect of weaning system on lamb growth and commercial milk production of Awassi dairy sheep. *Czech J. Anim. Sci.*, 52: 70-76.

- [16] Dzakuma, J. M., Nielsen, M. K., Doane, T. H. (1978). Genetic and phenotypic parameters of lamb weight. (Abst.) Proceeding American Society Animal Science Meeting, 45: 16.
- [17] ElWakil, S. L., ElSayed, M., Ahmed, A. M., Sadek, R. R., Nigm, A. A. (2009). Genetic and phenotypic parameters of birth, weaning and yearling body weights of Barki sheep raised in the North West coast of Egypt. *Egyptian J. Anim. Prod.*, 46: 43-52.
- [18] Esenbuğa, N., Dayioğlu, H. (2002). Effects of some environmental factors on growth traits of Awassi and Redkaraman lambs. *Turk. J. Vet. Anim. Sci.*, 26: 145-150.
- [19] Ghoneim, K. E., Shrode, R. R., Abdallah, R. K. (1982). Statistical analysis of birth weight in Awassi sheep in northern Iraq. *Proceedings of the Sixth International Conference on Animal and Poultry Production held at University of Zagazig, Zagazig, Egypt, 21-23 September. 2: 488-496.*
- [20] Gursoy, O., Kirk, K., Cebeci, Z., Pollot, G. E. (1995). Genetic evaluation of growth performance in Awassi sheep. In: Gabiña D. (ed.) *Strategies for sheep and goat breeding; Zaragoza: CIHEAM*, p. 193-201. (Cahiers Options Méditerranéennes; n. 11). Meeting of the joint FAO/CIHEAM Network on Sheep and Goats, Subnetwork on Animal Resources, 1995/03/26-28, Sidi-Thabet (Tunisia). <http://om.ciheam.org/om/pdf/c11/96605556.pdf>
- [21] Hassen, Y., Sölkner, J., Fuerst-Waltl, B. (2004). Body weight of Awassi and indigenous Ethiopian sheep and their crosses. *Small Rum. Res.*, 55: 51-56.
- [22] Jawasreh, K. I. Z., Khasawneh, A. Z. (2007). Studies of some economic characteristic on Awassi lambs in Jordan. *Egyptian J. of Sheep and Goats Sciences*, 2: 76-85.
- [23] Kamber, F. H. (1987). Studies on some factors affecting milk production and growth of lambs from birth to weaning in Awassi sheep. MS. Thesis, College of Agriculture, Forestry, Univ. of Mosul (In Arabic).
- [24] Khalaf, A. I., Said, S. I., Edriss, S. M. (2010). Role of some genetic and environmental factors in growth traits of Turkish Awassi, local and crossbred lambs. *The Iraqi J. Agric. Sci.*, 41: 12-22.
- [25] Kingwell, R. S., Abdil-Ghadim, K. A. (1995). Robinson, A. D.: Introducing Awassi sheep to Australia: an application of farming system models. *Agric. Systems.*, 47: 451-471.
- [26] Kridli, R. T., Abdullah, A. Y., Shaker, M. M., Al-Smadi, N. M. (2007). Reproductive performance and milk yield in Awassi ewes following crossbreeding. *Small Rum. Res.*, 71: 103-108.
- [27] Kazzal, N. T. (1973). Evaluation of some genetic and environmental factors affecting growth and development of Awassi. Ph. D. Dissertation, Univ. Tennessee, Knoxville, Tennessee, USA.
- [28] Křížek J., Louda F., Jakubec V., Řeháček E. (1983): Genetické a negenetické faktory ovlivňující mateřské vlastnosti ovcí. *Živoč. V ýr.* 28, 63-68.
- [29] Lightfoot, J. (1988). The Awassi fat tail sheep. *J. Agric., Western Australia.*, 28: 107-113.
- [30] Obaido, M. (2010). Characterization and comparative evaluation of sheep production under traditional and improved management conditions of central Syria: MSc Thesis, Dep. of Anim. Sci. Çukurova University. Turkey.
- [31] Özcan, M., Altinel, A., Yilmaz, A., Güneş, H. (2001). Studies on the Possibility of Improving Lamb Production by Two-way and Three-way Crossbreeding with German Black-Headed Mutton, Kıvrıcık and Chios Sheep Breeds 1. Fertility, Lamb Survival and Growth of Lambs. *Turk. J. Vet. Anim. Sci.*, 25: 687-694.
- [32] SAS, (2010). Statistical Analysis System. Version 9.1, SAS Institute Inc., Cary NC, USA.
- [33] Shaker, M. M., Abdullah, A. Y., Kridli, R., Šáda, I., Sovjak, R., Muwalla, M. M. (2002). Effect of crossing indigenous Awassi sheep breed with mutton and prolific sire breeds on the growth performance of lambs in a subtropical region. *Czech J. Anim. Sci.*, 47: 239-246.
- [34] Sunderman, F., Johns, M. (1994). Awassi fat tail- a chance for premium export. *J. Agric., Western Australia.*, 35: 99-105.
- [35] Tabbaa, M. J., Alnimer, M. A., Shboul, M., Titi, H. H. (2008). Reproductive characteristics of Awassi ewes mated artificially or naturally to Jordanian or Syrian Awassi rams. *Anim. Reprod.*, 5: 23-29.
- [36] Thrift, F. A., Whiteman, J. V., Kratzer, D. D. (1973). Genetic analysis of preweaning and postweaning lamb growth traits. *J. Anim. Sci.*, 36: 640-643.
- [37] Todorovski, N. (1988). The improved Awassi breed of sheep is wholly acclimatized to the natural conditions of Macedonia-Yugoslavia. *Proceedings of the Third World Congress on Sheep and Beef Cattle Breeding, Paris, France, 2: 739-742.*
- [38] Üstüner, H., Oğan, M. M. (2013). Main productive performance of Awassi sheep in the Central Anatolian Region of Turkey. *Turk. J. Vet. Anim. Sci.*, 37: 271-276.