

# The relationship between segmentation failure and spermatozoa motility characteristics

Muhammad Harfous, Hassn Hasan

Department of Anatomy and Histology and Embryology, Faculty of Medicine, Al-Andalus University for Medical Sciences, Tartus, Syria

## Email address:

dr.harfoush@hotmail.com (M. Harfous)

## To cite this article:

Muhammad Harfous, Hassn Hasan. The Relationship between Segmentation Failure and Spermatozoa Motility Characteristics. *American Journal of Biomedical and Life Sciences*. Special Issue: Spectral Imaging for Medical Diagnosis “Modern Tool for Molecular Imaging”. Vol. 3, No. 2-3, 2015, pp. 25-28. doi: 10.11648/j.ajbls.s.2015030203.15

**Abstract:** Dynamics of sperm motility (sperm velocity, percentage of motility and flagellar beat frequency) and monthly variations of semen characteristics (semen volume and osmolality and sperm concentration and motility) were studied. These criteria were used in the embryology laboratory and reproductive biology in the city of Lattakia, Syria, This study showed that sperm velocity, percentage of motility and beat frequency significantly and rapidly decreased after the activation of sperm motility. To study the dynamics of sperm motility parameters, twenty permeating males were randomly selected and electronically tagged to study the variations of semen characteristics. Specially we focused on the relationship between segmentation failure in cases treated with In Vitro Fertilization (IVF) to define the characteristics using the Computer Assisted Sperm Analysis (CASA). This study compares the reference group (normal) and a group of individuals treated with IVF, who have failed the treatment where segmentation did not occur. The role of the spermatozoon in fertilization expresses itself especially through the motility criterion. Studying the motility characteristics using CASA method seems interesting in clarifying the relationship between some of the motility characteristics and the ability to fertilize in order to determine the notification in cases of IVF. This study based on the acquired information of 85 normal voluntaries as references cases. And 45 patients with segmentation failure in the Embryology-lab of Al-Andalus University for medical Sciences between 2012- 2013. The results were clear, that significant relationship and correlation between segmentation failure and motility of the sperms.

**Keywords:** Vitro Fertilization (IVF), Spermatozoon, Computer Assisted Sperm Analysis CASA

## 1. Introduction

Knowledge on the sperm biology of a species is important for two main reasons. Percent motility and sperm velocity have been correlated to reproductive success in fish [1]. For the analysis of these indicators of sperm quality, computer-assisted sperm analysis (CASA) represents a most objective and comprehensive method of quantification [5]. CASA has therefore been employed in the determination of individual variation of sperm quality [6].

To date, very little is known about hake sperm biology. By the lateral insertion of the flagellum and the location of the centrioles outside the nuclear fossa, hake sperm belongs to the perciform type[7]

CASA device, as it shown in figure 1, consists of a microscope with high-resolution camera, in addition to an image-recording device. The device records, each 1/25 of second, the location and the weight-center of the

spermatozoon head. After converting these images into digital values, the system can draw a track for the spermatozoa head motility[8].



Figure 1. CASA test system

This method provides a sensitive means to study the spermatozoa study criteria routinely without relying on laboratories' experience and without errors[9].

## 2. Methods and Materials

### 2.1. In Vitro Fertilization - IVF

The IVF process involves:

- Stimulating multiple follicles and eggs to develop: There are several ovarian stimulation medication protocols that are used to "pump up" the ovaries to make enough follicles and eggs. Without stimulating medications, the ovaries make and release only 1 mature egg per menstrual cycle (month). In order to maximize success rates with in vitro fertilization we want a good number of high quality eggs from the woman.
- Egg retrieval to get the eggs
- Fertilizing the eggs in the laboratory
- Embryo transfer to the uterus

With ovarian stimulation for in vitro fertilization, the goal is to get approximately 8 to 15 quality eggs at the egg retrieval procedure.

Computer Assisted Sperm Analysis (CASA) was used to measure a series of parameters characterizing the motility and the sperm swimming performances.

### 2.2. Dynamics of Sperm Motility Parameters

If the sperm count or motility is extremely low, we usually assume this is the cause of the fertility issue. If the count or the motility is slightly low, it could be a contributing factor, but the sperm might not be the only fertility issue in the couple[10]. The following characteristics were studied using these computer imaging system:

- The proportion of the motile spermatozoa
- Linear speed VSL that reaches 35\_45  $\mu\text{m/s}$
- Lateral amplitude of the spermatozoon head motility ALH 3-4  $\mu\text{m}$

### 2.3. Spermatozoon Motility

Spermatozoon motility describes the ability of sperm to move properly through the female reproductive tract (internal fertilization) or through water (external fertilization) to reach the egg Sperm motility can also be thought of as the quality, which is a factor in successful conception; sperm that do not "swim" properly will not reach the egg in order to fertilize it. Normally the motility can be as Straight moving, Zig-zag moving, Vibrating, Non-motile. Whoever Human spermatozoon moves in a linear, progressive manner, in a false spiral head motility[15]. This movement results from flagellar wave progression to the proximal part towards the distal part of the flagella, on the one hand, and the cellular rotation by 180 degrees. Spermatozoon head motility is made in two types of tracks. Linear track with a change in direction every 0.5-4 seconds, mostly. Irregular movement due to change in direction.

Right after triggering the sperm activation, the wave is

present along the flagellum, but its parameters change during the motility period (Figure 2).




Figure 2. regular movement due to change in direction of Spermatozoon heads

This study showed that sperm velocity, percentage of motility and flagellar beat frequency significantly decrease after the activation of sperm motility

### 2.4. Flagella Movement

The flagellar waveform is composed of a bend with a larger angle and a bend with a smaller angle to achieve oscillation of flagellar bending, the sliding of specific microtubules and its displacement during the reverse direction of sliding are considered necessary.




Figure 3. regular movement due to change in direction of Spermatozoon

As it shown in figure 2 A possible model for the oscillatory mechanism of flagellar bending. Dyneins on two sides on the plane of the CP apparatus are regulated by signals of the RS/CP apparatus, resulting in the formation of a planar wave. The axons are fixed at the basal body near the sperm head. The base or tip of the flagellum points toward the minus or plus end of doublet microtubules, respectively. Dyneins are minus-ended motors (red and pink arrows), sliding adjacent microtubules to the plus end. The bending provides feedback and switches active dyneins, resulting in the sliding of opposite microtubules across the bend as it shown in figure 3.

In our experiment the flagellar waves start at the end of the middle-piece level which remains within the spermatozoon head longitudinal axis[16]. Bending occurs within the proximal part of the flagella, creating a lateral motility within a distance of 20\_25  $\mu\text{m}$  within the transverse level of the spermatozoon head, leading to flexing the spermatozoon head towards the other direction. The spermatozoon head rotates 180 degrees and then starts a new flagellar wave.

The ability of the spermatozoon to fertilize is linked to motility, where there is a clear relationship between motility and the ability to access inside the cervical mucus, as well as the ability to be treated with IVF.

When the lateral amplitude spermatozoon head motility decreases, the ability to access inside the cervical mucus decreases too. The comparison between the characteristics of the spermatozoon motility in men with normal fertility and men treated for procreation using CASA shows a clear lack in the criterion of lateral amplitude of the spermatozoon head motility.

Many researchers have found a relationship between the lateral amplitude of the spermatozoon head motility and the zona pellucida penetration test in rats[10].

In cases of IVF, the mobility criterion remains the most important one. According to the medical literature, it is clearly shown that the lateral amplitude of the spermatozoon is a very important criterion in determining the ability to fertilize.

### 3. Results

Eighty-five normal cases have been studied as reference objects; they met the following criteria:

- A married man who has children
- The youngest of his children is two years old
- Age between 20-45 years old.

The results were as follows:

**Table 1.** Reference group, 85 cases

	Mean value	SD
Concentration	94.4	5.6
Velocity	32.5	0.8
Linearity	5	1
Motility	64.4	1.3
ALHmax	3.54	0.07

Forty-one objects of failed segmentation were studied and the acquired data were as following

**Table 2.** Failed segmentation group, 41 cases

	Mean value	SD
Concentration	33,65u/u1	+4,54u/u1
Velocity	41,191	3+
Linearity	4,7	1,01
Motility	30,01	3,4
Alh max	3,16	0,95

The results confirmed that the motility and the lateral movement of the head remain the most important criteria in fertility.

Using only the concentration used on the failed segmentation group of progressively motile spermatozoa, we were able to achieve an overall 73 % accurate prediction of test results. Mean values and Standard deviations permitted accurate prediction of 33,65u/u1 and +4,54u/u1. The mean value of the velocity was 41,191 and the standard deviation was +3 with mean linearity factor of 4,7 and standard deviation of 1.01. However the mean value of Motility was 30,01 and standard deviation of 3,4 and the ALHmax has the

mean value of 3,16 with standard deviation of 0,95

Inclusion of the concentration of progressively motile spermatozoa into the selection gave the best prediction of results.

### 4. Conclusion

The results presented here have clearly demonstrated that the outcome of our experiments are strongly influenced by semen quality. Previous studies have reported significant correlations between the percentages of motile spermatozoa in semen samples and the outcome of segmentation failure.

We were able to identify significant relationships between sperm movement characteristics and the outcome of the segmentation failure.

In this study we found a significant difference between ALHmax in the failed segmentation and reference groups. In the failed segmentation group, the movement criterion ALHmax was 3.54 while it was 3.16 in the reference group as it shown in table 1 and 2.

Finally we can consider the movement criterion ALHmax as an indicator for segmentation in IVF.

### Acknowledgements

We thank the Faculty of medicine, Al-Andalus University for medical Sciences for help and supporting this study.

### References

- [1] Aitken, R.J., Best, F.S.M., Warner, P. & Templeton, A.A. (1984) A prospective study of the relationship between semen quality and fertility in cases of unexplained infertility. *J. Androl.* 5, 297-303.
- [2] Aitken, R.J., Sutton, M., Warner, P. & Richardson, D.W. (1985) Relationship between the movement characteristics of human spermatozoa and their ability to penetrate cervical mucus and zona-free hamster oocytes. *J. Reprod. Fert.* 73, 441-449.
- [3] Alexander, N. J. (1981) Evaluation of male infertility with an in vitro cervical mucus penetration test. *Fert. Steril.* 36, 201-208.
- [4] Beauchamp, P. J., Galle, P.C. & Blasco, L. (1984) Human sperm velocity and postinsemination cervical mucus test in the evaluation of the infertile couple. *Archs Androl.* 13, 107-112.
- [5] David, M.P., Amit, Bergman, Yedwab, G., Paz, G.F. & Homonnai, Z.T. (1979) Sperm penetration in vitro: correlations between parameters of sperm quality and the penetration capacity. *Fert. Steril.* 32, 676-680.
- [6] Feneux, D., Serres, C. & Jouannet, P. (1985) Sliding spermatozoa: a dyskinesia responsible for human infertility? *Fert. Steril.* 44, 508-511.
- [7] Gould, J.E., Overstreet, J.W., Yanagimachi, H., Yanagimachi, R., Katz, D.F. & Hanson, F.W. (1983) What functions of the sperm cell are measured by in vitro fertilization of zona-free hamster eggs? *Fert. Steril.* 40, 344-352.
- [8] Katz, D. F. & Overstreet, J.W. (1981) Sperm motility assessment by videomicrography. *Fert. Steril.* 35, 188-193.

- [9] Overstreet, J. W. (1982) Evaluation of sperm function by tests of sperm ovum interaction in vitro. In *Human Fertility Factors*, pp. 263-280. Eds A. Spira & P. Jouannet. INSERM, Paris.
- [10] Overstreet, J.W., Katz, D.F., Hanson, F.W. & Fonseca, J.R. (1979) A simple inexpensive method for objective assessment of human sperm movement characteristics. *Fert. Steril.* 31, 162-172.
- [11] Overstreet, J.W., Yanagimachi, R., Katz, D.F., Hayashi, K. & Hanson, F.W. (1980) Penetration of human spermatozoa into the human zona pellucida and the zona-free hamster egg: a study of fertile donors and infertile patients. *Fert. Steril.* 33, 534-542.
- [12] Pandya, I.J., Mortimer, D. & Sawers, R.S. (1986) A standardized approach for evaluating the penetration of human spermatozoa into cervical mucus in vitro. *Fert. Steril.* 45, 357-365.
- [13] Schats, R., Aitken, R.J., Templeton, A.A. & Djahanbakhch, O. (1984) the role of cervical mucus-semen interaction in infertility of unknown etiology. *Br. J. Obstet. Gynaecol.* 91, 371-376.
- [14] Templeton, A.A. & Mortimer, D. (1982) The development of a clinical test of sperm migration to the site of fertilization. *Fert. Steril.* 37, 410-415.
- [15] Ulstein, M. & Fjallbrant, B. (1976) In vitro tests of sperm penetration in cervical mucus. In *Human Semen and Fertility Regulation in Men*, pp. 383-388. Ed. E. S. E.
- [16] Hafez, C. V. Mosby Co., St Louis. Yanagimachi, R. (1984) Zona-free hamster eggs: Their use in assessing fertilizing capacity and examining chromosomes of human spermatozoa. *Gamete Res.* 10, 187-232.
- [17] Yanagimachi, R., Yanagimachi, H. & Rogers, B.J. (1976) The use of zona free animal ova as a test-system for the assessment of the fertilizing ability of human spermatozoa. *Biol. Reprod.* 15, 471-476.