



Potential Ex-vivo Anti-inflammatory, Cardioprotective Properties and Phytochemical Properties of Leaves of *Mussaenda Roxburghii* Hook

Farzana Akther Sumi¹, Biswajit Sikder¹, Prawej Ansari¹, Md. Reyad-ul- Ferdous^{1,2,*}, Anaytulla¹, Mustafe Khalid Mohamed¹, Md. Mustarek Uddin Meemo¹

¹Department of Pharmacy, North South University, Dhaka, Bangladesh

²Department of Pharmacy, Progati Medical Institute, Dhaka, Bangladesh

Email address:

rockyreyad@yahoo.com (Md. R. Ferdous)

To cite this article:

Farzana Akther Sumi, Biswajit Sikder, Prawej Ansari, Md. Reyad-ul- Ferdous, Anaytulla, Mustafe Khalid Mohamed, Md. Mustarek Uddin Meemo. Potential Ex-vivo Anti-inflammatory, Cardioprotective Properties and Phytochemical Properties of Leaves of *Mussaenda Roxburghii* Hook. *American Journal of Clinical and Experimental Medicine*. Special Issue: Herbal Remedies as Alternative to Future Drugs Development and Treatment. Vol. 3, No. 5-1, 2015, pp. 1-6. doi: 10.11648/j.ajcem.s.2015030501.11

Abstract: Background: *mussaenda roxburghii* hook. Belongs to family rubiaceae, is a perennial shrub moist area of valley and grows in the foothills. Objective: the present investigation was carried out to phytochemicals and pharmacological activity such anti-inflammatory, cardioprotective properties of methanolic crude extract & its fractions such as dcm extract (fraction 1), ethyl acetate extract (fraction2). Materials and methods: residual methanolic extract (fraction 3) of *mussaenda roxburghii* leaves which were obtained by solvent-solvent extraction process from crude methanolic extract of *mussaenda roxburghii*. Anti-inflammatory activity was measured by observing the mean inhibition of protein denaturation. Cardioprotective activity was evaluated by thrombolytic assay. Phytochemical screenings have done by using usual procedures. Results: an ex-vivo anti-inflammatory test demonstrated that; mean inhibition of protein denaturation of 1000/kg of crude methanolic extract of leave of *m. Roxburghii* was 17.399%. Cardioprotective properties of different extracts of *m. Roxburghii* was demonstrated which revealed that after treatment of clot with mother methanolic extract, dcm extract (fraction 1), ethyl acetate extract (fraction 2), residual methanolic extract (fraction3) of leaves and methanolic extract of root % clot lysis was 30.56%,27.61%,46.35%,26.02%,49.90 % respectively. Different tests with crude extracts showed that different types of alkaloids, carbohydrates, tannins were present in the leaves of *m. Roxburghii*. Conclusion: in these investigations demonstrates that different extracts of *m. Roxburghii* has significant pharmacological activities.

Keywords: Ex-vivo Anti-inflammatory, Cardioprotective Properties, Phytochemicals Investigation, Mean Inhibition of Protein Denaturation

1. Introduction

The goals of using plants as sources of therapeutic agents are to isolate bioactive compounds for direct use as drugs. Eg: taxol, digitoxin, morphine, digoxin, vinblastine & vincristine as well as reserpine, produce bioactive compounds of novel or known structures as lead compounds for semisynthesis to produce patentable entities of higher activity and/or lower toxicity.eg: taxotere, teniposide etc ^[1]. *Mussaenda roxburghii* hook. Belongs to rubiaceae; is a perennial shrub grows in the foothills and moist areas of valley. Roots are traditionally used in

treatment of cuts, jaundice, wounds, boils and skin diseases etc ^[2,3]. Leaves are used in the ailments of bone fracture ^[4]. The paste obtained from leaf of this plant is applied to treat boils ^[5]. Previous phytochemical investigation led to isolation of a new iridoid, shanzhiol which showed mild antibacterial activity against both staphylococcus aureus and escherichia coli with a mic of 100 µg/ml by the broth dilution method ^[6]. As part of our ongoing research with medicinal plant of bangladesh ^[9,10] the present study has been undertaken to evaluate the preliminary cytotoxicity

and antimicrobial activity of *M. Roxburghii* as well as to find out logical evidence for its folk uses and for discovery of new drug candidates.

2. Materials and Methods

2.1. Plant Materials

The leaves of *Mussaenda roxburghii* were collected from Chittagong, Bangladesh, in November 2013. A voucher specimen for this plant has been maintained in Bangladesh National Herbarium, Dhaka, Bangladesh (accession no. Mfk137. Ctg uh). The sun-dried and powdered (500 gm) leaves of *P. Cereifera* were macerated in 2.5 l of methanol for 7 days and then filtered through a cotton plug followed by Whatman filter paper number 1. All extracts were concentrated with a rotary evaporator at reduced pressure and low temperature (40-45°C). The concentrated methanolic extracts (ME) were partitioned by modified Kupchan method [7] and described by Md. Reyad-ul-Ferdous^[9] the resultant partitionates which are petroleum ether (PE), chloroform (CF), carbon tetrachloride (CTCF), and aqueous (AQSF) soluble fractions were used for the experimental processes.

2.2. Ex-vivo anti-inflammatory Activity Study

Inhibition of albumin denaturation was conducted for evaluation of ex-vivo anti-inflammatory activity. The present study was developed by certain modification of the method claimed by Mizushima Y and Kobayashi M^[8] on the basis of in-vitro determination of the protein maturation caused by the control group and comparing it with the positive and test groups. The more the inflammatory action, the more it will cause the protein maturation and the measured absorbance will be higher. The reaction mixture was consisting of test extracts at different concentrations and 1% aqueous solution of bovine albumin fraction. Reaction mixture pH was

$$\% \text{ of clot lysis} = (\text{weight of released clot} / \text{clot weight}) \times 100$$

2.4. Qualitative Screening of Phytochemicals

One gram of the methanol extract of *Mussaenda roxburghii* was dissolved in 100 ml of methanol and was subjected to preliminary phytochemical screenings for determining nature of phytoconstituents^[13-15]. A small portion of the dry extract was used for the phytochemical tests for compounds which includes carbohydrate, tannins, flavonoids, alkaloids, saponins, steroids, resins, and glycoside in accordance with the methods of with little modifications. All test procedure for chemical groups described^[16]. We identified several compounds in table-9.

(i). Test for Carbohydrate

(a). Molisch's Test

2 ml of an aqueous extract of the plant material in a test tube have to take + then have to add 2 drops of freshly prepared Molisch's reagent + mixed thoroughly + allowed 2

adjusted using small amount of 1N HCl. Aspirin was taken as standard drug. The samples were incubated at 37°C for 20 min and then heated at 57°C for 30 min. After cooling the samples, the turbidity was measured spectrophotometrically at 660 nm. The experiment was performed in triplicate. % inhibition of protein denaturation was calculated as follows: percentage inhibition = (abs control - abs sample) / abs control × 100

2.3. Cardio-protective Activity Assay

The cardio-protective activity was evaluated by thrombolytic activity assay. The thrombolytic activity of all extracts was evaluated by the method described by Dagainwala^[11] and modified by Kawsar^[112] using streptokinase (SK) as the standard as well as described by Reyad-ul-Ferdous M^[9]. 0.1 gm extract was suspended in 10 ml distilled water and the suspension was shaken vigorously on a vortex mixer. The suspension was kept overnight and decanted to remove the soluble supernatant, by using a filter. 100 µl of this aqueous preparation of herbs was added to the alpine tubes containing the clots to check the thrombolytic activity. 10 alpine tubes were taken and each eppendorf tube containing clot was properly labeled and venous blood drawn from the healthy volunteers. That was transferred in 10 different pre weighed sterile alpine tube (0.5 ml /tube) and incubated at 37°C for 45 minutes. Aspirated washout without disturbing the clot formed. Water was also added to one of the tubes containing clot and this serves as a negative thrombolytic control. All the tubes are then incubated at 37°C for 90 minutes and observed for clot lysis. After incubation, fluid obtained was removed and tubes were again weighed to observe the difference in weight after clot disruption. Difference obtained in weight taken before and after clot lysis was expressed as percentage of clot lysis. Percentage of clot lysis was determined from below formula.

ml of conc. Sulphuric acid to flow down the side of the inclined test tube. So that acid forms a layer beneath the aqueous solution. A red or reddish violet ring will form at the junction of the 2 layers. On standing or shaking a dark purple solution will form. After diluted with 5 ml of water a dull violet precipitate will form.

(b). Benedict Test

0.5 ml of an aqueous extract of the plant material + added 5 ml of Benedict's solution + boil for 5 minutes and allow cooling spontaneously. A red precipitate of cuprous oxide is formed in the presence of a reducing sugar.

(c). Fehling's Test

2 ml of an aqueous extract of the plant material + add 1 ml of a mixture of equal volume of Fehling's solutions A & B. Boil for a few minutes. A red or brick red precipitate is formed if a reducing sugar is present.

(ii). Test for Alkaloids

test name	experimentation	observation
1)mayer's test	1 ml of filtrate + few drops of mayer's reagent.	White or creamy white precipitate indicates the presence of alkaloids.
2)wagner's test	1 ml of filtrate+ few drops of wagner's reagent.	Brown or deep brown ppt indicates the presence of alkaloids.
3)hager's test	1 ml of filtrate + few drops of hager's reagent.	Yellow crystalline precipitate indicates the presence of alkaloids
4)tannic acid test	1 ml of filtrate+ few drops of tannic acid reagent.	Dirty white or black precipitate Indicates the presence of alkaloids.

(iii). Test for Glycosides

test name	experimentation	observation
General test for glycoside	A small amount of extract dissolved in small amount of water + few drops of aqueous sodium hydroxide solution	Presence of yellow colour indicates the presence of glycosides.

(iv). Test for Steroids

test name	experimentation	observation
Liebermann-burchard's test	2 ml of a chloroform extract + 2 ml of acetic anhydride + 1 ml of conc. Sulphuric acid	A greenish colour will form at the junction of 2 layers, which turns blue on standing, indicates presence of steroids.
Salkowski's test	2 ml of a chloroform extract of the plant material +1 ml of conc. Sulphuric acid from the side of the test tube.	A red color is produced in the chloroform layer if steroids are present.

(v). Test for Tannins

Test name	Experimentation	Observation
1)ferric chlorides test	5 ml of alcoholic extract solution+ 1ml 5% ferric chloride solution	A blue, blue black green or blue green colour or precipitation is produced in the presence of tannins. On addition of a few ml of dilute sulphuric acid the colour disappears followed by the formation of a yellowish brown ppt.
2)lead acetate test	5 ml of an aqueous extract of the plant material+ few drops of a 1% solution of lead acetate.	A yellow or red ppt is formed.
3)potassium dichromate test	5 ml of alcoholic extract of plant material + 1 ml 10% potassium dichromate solution	Yellow orange ppt indicates the presence of tannins.

(vi). Test for Resins

Test name	Experimentation	Observation
General test	Using gentle heat, have to dissolve a small amount of chloroformic or ethanolic extract of the plant material in 5 to 10 ml of acetic anhydride. Cool & have to add 0.05 ml of sulphuric acid.	If resin are present, a bright purplish red colour,rapidly changing to violet,is produced.

(vii). Test for Flavonoids

Test name	Experimentation	Observation
Conc. Hcl and alcoholic test	Small amount of alcoholic extract of the plant material + few drops of conc. Hcl	Immediate development of red color indicates the presence of flavonoids.

(viii). Test for Saponins

Test name	Experimentation	Observation
Shake test (aq. Solution froth test)	Boil about 0.1 gm of the powdered plant material with 10 ml of water for 3-5 minutes +filter+after cooling dilute 5 ml of the filtrate with water and shake vigorously.	Production of a persistent frothing (which remains stable on heating) indicates the presence of saponins.

2.5. Statistical Analysis

The experimental results were expressed as the mean \pm standard deviation (sd). Statistical significance of the mean mortality at each concentration was analyzed using one-way analysis of variance (anova) and compared using duncan's multiple range tests. Values of $p \leq 0.05$ were taken to be statistically significant.

3. Results and Discussion

Denaturation of proteins is a well-documented cause of inflammation. Salicylic acid, phenylbutazone, flufenamic acid (anti-inflammatory drugs) etc, have shown dose dependent ability to thermally induced protein denaturation^[8]. Ability of extracts to inhibit protein denaturation was studied to indentify mechanism of the anti-inflammatory activity. Both the extracts were effective in inhibiting heat induced albumin denaturation at different concentrations as shown in figure-1. In the present study for the in-vitro anti-inflammatory test, the crude methanolic extracts of *mussaenda roxburghii* hook. Showed mean inhibition of protein denaturation 17.399 ± 0.01937 . Whereas, for asa it was found to be 42.491 ± 0.00698 . The ability of methanol extract of leaves of *m. Roxburghii* to inhibit thermal and hypotonic protein denaturation was found to be less significant than the positive acetyl salicylic acid.

Addition Of 100 ul of streptokinase, a positive control (1500000 i.u) to the clots along with 90 minutes of incubation at 37 °c, exhibit 69.35% clot lysis. Clots when treated with 100 μ l sterile distilled water (negative control) showed only negligible clot lysis (6.23%). After treatment of clot with *m. Roxburghii* (mother methanolic extract), *m. Roxburghii* (dcm extract ,fraction 1), *m. Roxburghii* (ethyl acetate extract, fraction 2), *m. Roxburghii* (residual methanolic extract, fraction 3), *m. Roxburghii* (methanolic extract of leaves) 30.56%,27.6096%,46.35%,26.02%,49.90% clot lysis were observed respectively and mean of percentage of clot lysis was more than water. Results are shown in figure-2.

The result obtained in the present investigation phytochemicals screening of the methanol extract of leaves of *m. Roxburghii* revealed that the crude extract contained

flavonoids, saponins, steroids, tannins and triterpinoids, terpinoids, cardiac glycosides and anthraquinones (table-1). *M. Roxburghii* leaves can also have various medicinal values such as anti-inflammatory, membrane stabilizing and thrombolytic activity. The presence of flavonoids exhibit significance anticancer, cardioprotective, anti-microbial activity may demonstrate gram-positive, gram-negative as

well as fungi also. Among of the microorganisms are resistance to several drugseach though, this is only a preliminary study of the occurrence of certain properties of *m. Roxburghii* leaves an in-depth study will provide a good concerted base of all the phytochemicals and several pharmacological functions mention above.

Table 1. Chemical group test result for *Mussaenda roxburghii* leaves parts.

Examixanation	Test performed	Result for <i>m. roxburghii</i> leaves
1)Carbohydrate	1)Molisch’s test (test for Gum)	(+) ve
	2)Benedict test (test for Reducing sugar)	(-) ve
	3)Fehling’s test (test for Reducing sugar)	(-)ve
2)Alkaloids	1)Mayers test	(+)ve
	2)Wagner’s test	(+)ve
	3)Hager’s test	(+)ve
	4)Tannic acid test	(+)ve
3)Tannins	1) Ferric chloride test	(+)ve
	2)Lead acetate	(-)ve
	3)Potassium dichromate test	(+)ve
4)Resins		(-)ve
5)Flavonoids	Conc. HCl & Alcoholic test.	(-)ve
6)Saponins	Shake test(aq. Solution froth test.	(-)ve
7)Glycosides	1) General test	(-)ve
9)Steroids	1)Liebermann Barchards test	(-) ve
	2)Salkowski’s test	(-) ve

(+) = present and (-) = absence




Figure 1. Mean inhibition of protein denaturation (%MIPD) and Mean protein denaturation (%MPD) VS treatment groups. Here, ASA= Acetyl salicylic acid, MEMR= Methanolic extract of *Mussaenda roxburghii*.




Figure 2. % clotlysis of different extracts of *Mussaenda roxburghii* (Leaves).

Here, MMEMR= Mother methanolic extract of *M. roxburghii*, DCM F1 MR= DCM extract (fraction 1) of *M. roxburghii*, EA F2 MR= Ethyl acetate extract (fraction 2) of *M. roxburghii*, RME F3 MR= Residual methanolic extract (fraction 3) of *M. roxburghii*.

4. Conclusion

In the present study, we have found that most of the biologically active phytochemicals were present in the methanolic extract of *M. Roxburghii* leaves. This is only a preliminary study and to make final comment the extract should thoroughly investigated phytochemically and pharmacologically to exploit their medicinal and pharmaceutical potentialities.

Acknowledgement

We are acknowledge to north south university and progati medical institute for provide us required facilities to complete this work.

References

- [1] Farnsworth nr, akerele o, bingel as, soejarta dd, eno z. Medicinal plants in therap. World health organ 1985; 63: 965–81.
- [2] Saha j, sarkar pk, chattopadhyay s. 2011. A survey of ethnomedicinal plants of darjeeling hills for their antimicrobial and antioxidant activities. Ind j nat prod res 2: 479 - 492.
- [3] Patil sa, joshi vg. 2011. Evaluation of antibacterial and wound healing activity of leaves of *mussaenda frondosa* linn. Int j res pharmac biomed sc 2: 147 - 154.
- [4] Das hb, majumdar k, datta bk, ray d. 2009. Ethnobotanical uses of some medicinal plants by tripuri and reang tribes of tripura. Nat prod rad 8: 172 - 180.
- [5] Rahman ma. 2010. Indigenous knowledge of herbal medicines in bangladesh and treatment of skin diseases by tribal communities of the hill tracts districts. Bangladesh j bot 39: 169 – 177
- [6] Chandra du, ghosh r, chowdhury s, dinda b. 2012. New iridoid from aerial parts of *mussaenda roxburghii*. Nat prod commun 7: 1 - 2.
- [7] Van wagenen bc, larsen r, cardellina jh, ran dazzo d, lidert zc and swithenbank c: ulosantoin (1993) a potent insecticide from the sponge *ulosaruetzleri*. Journal of organic chemistry 58: 335-337.
- [8] Mizushima y and kobayashi m. Interaction of anti-inflammatory drugs with serum preteins, especially with some biologically active proteins. J of pharma pharmacol. 1968; 20:169-173.
- [9] Reyad-ul-ferdous m, alam tt, islam ma, khan mzi, tasnim f, et al (2014) *ex-vivo* cardioprotective and cytotoxic screening of fruits of *parmentiera ereifera* seem. Biol med 6: 219. Doi: 10.4172/0974-8369.1000219.
- [10] Md. Reyad-ul-ferdous, sayma akhter, md. Zahirul islam khan, md. Eshak khan, md. Atiqul islam, md. Sharif ullah. Ex-vivo anti-inflammatory and antimicrobial activities of the leaves of *bauhinia acuminata*. American journal of life sciences. Vol. 2, no. 5, 2014, pp. 267-270. Doi: 10.11648/j.ajls.20140205.13.
- [11] Dagainawala hf, prasad s, kashyap rs, deopujari jy, purohit hj and taori gm (2006) development of an in vitro model to study clot lysis activity of thrombolytic drugs. Thrombosis journal 4: 14.

- [12] Kawsar mh, sikder ma, rana ms, nimmi i and rashid ma (2011) studies of thrombolytic and cytotoxic properties of two asteraceous plants of bangladesh. Bangladesh pharmaceutical journal 14: 103-106.
- [13] Md. Reyad-ul-ferdous, nasir uddin, d.m. shamim shahjahan, mosharaf hossen, md. Saiful islam arman, ariful islam. Preliminaryin-vitro potential phytochemicals investigation of barks of ravenala madagascariensis sonnerat. Wjpps, 2014; 3(9), 1418-1423.
- [14] Egwaikhide, p.a., s.o. okeniyi and c.e. gimba, 2007. Screening for antimicrobial activity and phytochemical constituents of some nigerian medicinal plants. Adv. Biological res., 1: 155-158.
- [15] Harborne, j.b. phytochemical methods. A guide to modern techniques of plant analysis. 3rd edn., new york; chapman and hall int ed.1998.
- [16] Gani a. Methods of screening, practical phytochemistry, 2nd edn, dhaka:chaya prokashoni; 2007.p. 12-17.