Effect of Alpha-cypermethrin on morphological parameters in tomato plants (*Lycopersicon esculentum* Mill.)

Karim CHAHID¹, Amin LAGLAOUI², Said ZENTAR³, Abdeslam ENNABILI¹

¹Laboratoire LAMEC, Faculté des sciences Dhar El Mehraz, Fès

²Equipe de Recherche ERBGB, Faculté des Sciences et Techniques, Tanger

³Station d'ionisation SIBO de l'INRA, Tanger

Email address:

karimchahid@yahoo.fr (K. CHAHID)

To cite this article:

Karim CHAHID, Amin LAGLAOUI, Said ZENTAR, Abdeslam ENNABILI. Effect of Alpha-Cypermethrin on Morphological Parameters in Tomato Plants (*Lycopersicon esculentum* Mill.). *American Journal of Environmental Protection*.

Vol. 2, No. 6, 2013, pp. 149-153. doi: 10.11648/j.ajep.20130206.15

Abstract: Devastating insects are responsible of losses in quantity and quality of agricultural production. To overcome this problem, farmers use pesticides, obtained by chemical synthesis and representing the major cause of agricultural contamination of soil and groundwater. Thus, pesticides may present important risks because of their persistence, bioavailability and mobility, in spite of their correct application. This study has evaluated the effect of alphacypermethrin (pyrethroids class), largely used in tomato (*Lycopersicon esculentum* Mill.) treatment in the Northern area of Morocco. Synthetic pyrethroids are widely used as the broad-spectrum pest control agents in agricultural production because of their selective insecticidal activity, rapid biotransformation and excretion by the mammalian catabolic system and non-persistence in the environment. The effect of alpha-cypermethrin on seeds germination and seedlings growth of tomato has been studied based on morphological parameters and by using four dilutions of the normal concentration used in agriculture (100%, 75%, 50%, 25%) for germinating seeds, and only the normal concentration used in agriculture for growing tomato plants. The results indicated that alpha-cypermethrin induced a delay of germination and growth process. The germination rate of treated seeds was generally 20% lower than the control treatment. Generally the control's germination rate was around 97% in all days of measurement period. A decrease in germination rate was observed in all concentrations of α -cypermethrin; the rate was between 80% and 88.7% and it was generally constant throughout the test period. Furthermore, the length of roots and shoots in treated seeds was significantly reduced. In this regard, shoot length of the treated seedlings was 25% and 50%-reduced for the concentrations of 25% and 100%, respectively, when compared to control shoots length. A similar result was also observed in roots, the length of the treated seedlings' roots was generally 29% and 50%-reduced for the concentrations of 25% and 100%, respectively, when compared to control roots length. Concerning the growth of roots and shoots in treated plantlets, a reduction was observed when compared to the control plantlets growth. The growth delay in the treated seedlings was observed at the 2nd week of the test period. Shoot length of treated plantlets was generally around 12% reduced when compared to the control. The same result was observed in treated plants' roots which length was also 7% reduced compared to untreated seedlings. The analysis of variance (ANOVA) and the Tukey test were utilised for the Post-hoc tests. A significance level of 0.05 was used for all statistical tests.

Keywords: Insecticides, Tomato, Seed Germination, Plant Growth

1. Introduction

The tomato (*Lycopersicon esculentum* Mill.) is one of the most widely grown vegetables in the world. In recent years, competition has intensified as the world exports of tomato products from main suppliers have increased substantially.

Tomatoes growth can be inhibited by various arthropods, plant diseases and nematodes, which significantly reduce yield and quality of fruit (Oerke *et al.*, 1994).

In Northern Morocco, the most important way to protect crops is the use of chemical pesticides. Many pesticide types are used, especially: organochlorine, organophosphorus, carbamate and pyrethroids pesticides

(El Bakouri *et al.*, 2008). However, the use of these pesticides obtained by chemical synthesis represents the major cause of agricultural soil and groundwater contamination because of their persistence, bioavailability and mobility (Arias-Estevez *et al.*, 2008). The pesticide occurrence in agricultural soil of the Tangier region shows the presence of many pesticides types such as endosulfan isomers (alpha and beta), endosulfan sulfate, some DDT metabolites and alpha HCH (El Bakouri *et al.*, 2008).

2. Materials and Methods

The methodology utilized in the experimental work consisted of: plant material used to prepare the samples and a variety of germination tests; growth evaluation tests; and the different variables monitored and measured for evaluation purposes. Following the implementation of experimental design, a statistical analysis was performed to evaluate the significant level of tolerance and impact. The different experimental steps are discussed below.

The samples were prepared by selecting tomato (*Lycopersicon esculentum* Mill.) seeds from the Vilmorin commercial seeds, which were surface-sterilized in 10% commercial bleach with stirring for five minutes, followed by extensive washing in sterile-distilled water. The germination was initiated by selecting batches of 50 tomato seeds that were germinated in Petri dishes (diameter 9 cm) upon two layers of filter paper moistened with 6 ml of distilled water or insecticide solutions at the concentrations of 25%, 50%, 75% and 100%, where 100% represents the normal concentration usually used. The Petri dishes were stored in a growth chamber in darkness at 25°C for six days. At various stages of the tomato seed germination (3, 4 and 5 days), seeds of each replicate were collected for measurement. Three repetitions were

performed. Germination time was determined as the time of rupture of seed coats, and the emergence of the radicle through the seed coat.

For growth evaluation, ten seeds were sown in each plastic pot for germination and growth. Seedlings were grown at 24/20°C (day/night) and 16 hours of light, and watered each day. After growth for 30 days, the seedlings were treated with the insecticides at the normal concentrations used in agriculture (100%). The tests were realised also on the 2nd, 5th, 8th, 11th and 14th day after treatment. Each value represents the average of three replications.

The growth was evaluated by the rate of germination. The germination rate of tomato seeds was evaluated by counting the number of germinated seeds in batches of 50 seeds; each value represents the average of three repetitions. Measurements were made of the length of shoots and roots of the germinated seeds. For the elongation of growing tomato seedlings, the scaled length of shoots and roots was made; each value represents the average of three repetitions.

Following the completion of the experimental sample, the data was evaluated statistically. The data were processed by using Statistica Software (Statistica, 1997) for one-way analysis of variance (ANOVA) and the Tukey test for the Post-hoc tests. A significance level of 0.05 was used for all statistical tests.

3. Results

It was observed that germination rate of treated seeds was lower than the control one as shown in Table 1. The control's germination rate was around 97% on all days of the measurement period. A decrease in germination rate was observed in all concentrations of α -cypermethrin; the rate was between 80% and 88.7% and it was generally constant throughout the test period.

Table 1. Effect of α -cypermethrin on germination percentage of tomato seeds at a significant difference at $P \leq 0.05$, 0.01 and 0.001 levels.

	3 rd day	4 th day	5 th day
Control	97.33% $\pm$ 1.15	97.33% $\pm$ 1.15	97.33% $\pm$ 1.15
α -cypermethrin			
25%	88.66% $\pm$ 1.15***	88.66% $\pm$ 1.15***	88.66% $\pm$ 1.15***
50%	80% $\pm$ 0***	80% $\pm$ 0***	80% $\pm$ 0***
75%	80.66% $\pm$ 2.3***	82% $\pm$ 2***	82% $\pm$ 2***
100%	88% $\pm$ 0***	88% $\pm$ 0***	88% $\pm$ 0***

Note: *, ** and *** indicate significant difference at $P \leq 0.05$, 0.01 and 0.001 levels, respectively

Concerning the plantlets growth, a delay was observed for the treated seeds as shown in Table 2. Shoot length of both control and treated seed plantlets increased depending on the time, but the growth of the treated seed plantlets was

lower than the control ones. In this regard, shoot length of the treated seedlings was 25% and 50%-reduced for the concentrations of 25% and 100%, respectively.

Table 2. Effect of α -cypermethrin on shoot and root length in tomato seedlings indicate significant difference at $P \leq 0.05$, 0.01 and 0.001 levels.

Shoot length (cm)			Root length (cm)		
3 rd day	4 th day	5 th day	3 rd day	4 th day	5 th day
Control	0.81 ± 0.09	1.84 ± 0.09	2.01 ± 0.14	2.81 ± 0.17	3.59 ± 0.07
cypermethrin					
25%	0.62 ± 0.02	1.19 ± 0.08**	1.31 ± 0.08***	2.02 ± 0.14***	2.41 ± 0.15***
50%	0.49 ± 0.08**	1 ± 0.25***	1.26 ± 0.33***	1.97 ± 0.16***	1.92 ± 0.21***
75%	0.5 ± 0.05**	0.9 ± 0.15***	1.3 ± 0.22***	1.57 ± 0.12***	2.05 ± 0.21***
100%	0.44 ± 0.07***	0.93 ± 0.17***	1.1 ± 0.03***	1.79 ± 0.05***	1.95 ± 0.14***

Note: *, ** and *** indicate significant difference at $P \leq 0.05$, 0.01 and 0.001 levels respectively

A similar effect was also observed in seedlings root growth. The delay of roots growth was at the same order as the delay observed in shoot growth. Root length was around 3.6 cm in control plantlets on the 5th day and 2.41 cm in treated ones (25%- α -cypermethrin, 5th day) as shown in Table2.

Table 3. Effect of α -cypermethrin on shoot and roots length in growing tomato seedlings. *, ** and *** indicate significant difference at $P \leq 0.05$, 0.01 and 0.001 levels.

	Shoot length (cm)		Root length (cm)	
	Control	α -cypermethrin	Control	α -cypermethrine
2 nd day	18 ± 1	17.9 ± 0.65	22.8 ± 0.7	22.56 ± 0.81
5 th day	18.5 ± 0.5	18.13 ± 2.51	23.5 ± 0.58	23.3 ± 0.1
8 th day	19.23 ± 0.25	18.43 ± 0.32*	24.5 ± 0.72	23.9 ± 0.4
11 th day	20.4 ± 0.36	18.9 ± 0.6*	25.6 ± 0.28	24.13 ± 0.47**
14 th day	22.1 ± 0.62	19.6 ± 0.95*	26.7 ± 0.17	25 ± 0.55**

Note: *, ** and *** indicate significant difference at $P \leq 0.05$, 0.01 and 0.001 levels respectively

Furthermore, evaluation of length progression of growing seedlings shows a growth delay in the treated seedlings at the 2nd week of the test period. Shoot length of treated plantlets was around 19.6 cm as opposed to 22.1 cm for the control ones as shown in Table3. Root length of treated plantlets was also lower and reached 25 cm, as opposed to 26.7 cm in untreated seedlings as shown in Table3.

4. Discussion

The percentage of germination may reflect the reaction rate of plant seeds to their living environment (Li *et al.*, 2007). The results obtained in this study illustrate an inhibitory effect of germination after treatment by the α -cypermethrin at various concentrations. In literature, a decline in seed germination rate (more than 50 %) have been reported for other pesticides such as paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride) at 1.0 mg/L in *Typhalatifolia* (Moore *et al.*, 1999). A similar effect was also shown in *Triticumaestivum*, treated by using heavy metals (5–20 mg/kg of Arsenic) (Li *et al.*, 2007).

The general symptom of plants under such stresses is growth inhibition (Yang *et al.*, 2001; Wang *et al.*, 2004; Wang and Yang, 2005). It was observed that the tomato seeds plantlets and seedlings growth was significantly delayed after insecticide application, affecting the plantlets shoots as well as roots. However, many studies reported an inhibitory effect of growth after application of pesticides: (i) around 50%-decreasing of root growth in *Phaseolus vulgaris* and *Pisumsativum* after treatment with chlorsulfuron during the germination process (Fayez and Kristen, 1996), and (ii) low root growth in *Zea mays* seedlings in the presence of pesticides such as chlorsulfuron and metsulfuron methyl (Fayez *et al.*, 1994). This effect was also demonstrated with other xenobiotic types like heavy metals. Khatunet *al.* (2008) underlined a severe growth inhibition of shoots and roots in *Withaniasomnifera* because of copper contamination.

Many hypotheses could explain this delay of growth in treated plants and seedling.

Firstly, insecticides could induce damages in the meristematic cells; in this way, Fayez and Kirsten (1996) showed that chlorosulfuron has an obvious influence on the

cellular structure of root caps of *Pisumsativum*, *Phaseolus vulgaris* and *Vicia faba*, and induce a reduction of root cell division, delaying the root growth (Ray, 1982; Rost, 1984).

On the other hand, insecticides could affect the photosynthetic system by the inhibition of photo system II and chain electron transport activities, as reported for example by Mishra *et al.* (2008) in *Vigna unguiculata* when treated by dimethoate. Pesticides could also lead to a delay in photosynthetic pigments rates such as chlorophylls (Mishra *et al.*, 2008).

Moreover, in literature, the delay of growth was reported as an indicator of oxidative stress (Yang *et al.*, 2001; Wang *et al.*, 2004; Wang and Yang, 2005). Generally, the plants cells produce the ROS as a second messenger in some processes of growth and development (Grun *et al.*, 2006; Schurmann, 2003 ; Borland *et al.*, 2006). There is an interaction between activation and repression of reactive oxygen species (ROS) and phytohormones (Delledome *et al.*, 2003; Termanand Brunk, 2006; Del Rio *et al.*, 2006; Shao *et al.*, 2006, McCarty and Chory, 2000; Shao *et al.*, 2008). When the plant is under an oxidative stress, the excess of ROS production could lead to growth perturbations.

In perspective, other studies on insecticides effect in tomato are undertaken: reserves substances, enzymes degradation and oxidative stress.

5. Conclusion

To summarize, the insecticides studied showed a negative effect in both tomato seeds and growing plants. The insecticides influenced both germination and growth process by causing a decrease in germination rate and delay in length progress of treated shoots and roots in both seeds and plants. Other studies are undertaken to have a clear idea about the effect of insecticides in tomato.

References

- [1] Arias-Estevez M, Lopez-periago E, Martinez-Carballo E, Simal-Gandara J, Mejuto JC, Garcia-Riu L (2008). The mobility and degradation of pesticides in soils and the pollution of groundwater resources, *Agriculture, Ecosystems and Environment*. 123: 247-260.
- [2] Borland A, Elliott S, Patterson S (2006). Are the metabolic components of crassulacean acid metabolism up-regulated in responses to an increase in oxidative burden? *Journal of Experimental Botany*. 57 (2) : 319-328.
- [3] Del Rio LA, Sandalio LM, Corpas FJ, Palma JM, Barroso JB (2006). Reactive oxygen species and reactive nitrogen species in peroxisomes. Production, scavenging, and role in cell signaling. *Plant Physiology*. 141: 330-335.
- [4] Delledome M, Polverari A, Murgia I (2003). The functions of nitric oxide-mediated signaling and changes in gene expression during the hypersensitive response. *Antioxidants and Redox Signaling*. 5 (1): 33-41.
- [5] El Bakouri H, Aassiri A, Morillo J, Usero J, Khaddor M, Ouassini A (2008). Pesticides and lipids occurrence in Tangier agricultural soil (northern Morocco). *Applied Geochemistry*. 23: 3487-3497
- [6] Fayez KA, Kristen U (1996). The influence of herbicides on the growth and proline content of primary roots and on the ultrastructure of root caps. *Environmental and Experimental Botany*. 36 (1): 71-81.
- [7] Fayez KA, Gerken I, Kristen U (1994). Ultrastructural responses of root caps to the herbicides chlorsulfuron and metsulfuron methyl. *Plant and Soil*. 167(1): 127-134.
- [8] Grun S, Lindermayr C, Sell S, (2006). Nitric oxide and gene regulation in plants. *Journal of Experimental Botany*. 57 (3): 507-516.
- [9] Khatun S, Babar Ali M, Hahn E-J, Paek K-Y (2008). Copper toxicity in *Withaniasomnifera*: Growth and antioxidant enzymes responses of *in vitro* grown plants. *Environmental and Experimental Botany*. 64: 279-285
- [10] Li CX, Feng SL, Shao Y, Jiang LN, Lu XY, Hou XL (2007). Effects of arsenic on seed germination and physiological activities of wheat seedlings. *Journal of Environmental Sciences*. 19(6): 725-732
- [11] McCarty DR, Chory J, (2000). Conservation and innovation in plant signaling pathways. *Cell*. 103: 201-209.
- [12] Mishra V, Srivastava G, Prasad SM, Abraham G (2008). Growth, photosynthetic pigments and photosynthetic activity during seedling stage of cowpea (*Vigna unguiculata*) in response to UV-B and dimethoate. *Pesticide Biochemistry and Physiology*. 92: 30-37.
- [13] Moore MT, Huggett DB, Huddleston GM, Rodgers JH, Cooper CM (1999). Herbicide effects on *Typha latifolia* (Linnaeus) germination and root and shoot development. *Chemosphere*. 38 (15): 3637-3647.
- [14] Oerke EC, Dehne DW, Schonbeck F, Weber A (1994). Crop Production and Crop Protection: Estimated Losses in Major Food and Cash Crops. Elsevier Hardcover, Amsterdam, 830pp.
- [15] Ray TB (1982). The mode of action of chlorsulfuron: a new herbicide for cereals, in Fayez KA, Kristen U (1996). The influence of herbicides on the growth and proline content of primary roots and on the ultrastructure of root caps. *Environmental and Experimental Botany*. 36 (1): 71-81.
- [16] Rost TL (1984). The comparative cell cycle and metabolic effects of chemical treatments on root tip meristem. III. Chlorsulfuron, in Fayez KA, Kristen U (1996). The influence of herbicides on the growth and proline content of primary roots and on the ultrastructure of root caps. *Environmental and Experimental Botany*. 36 (1): 71-81.
- [17] Schurmann P, (2003). Redox signaling in the chloroplast: the ferredoxin/ thioredoxin system, *Antioxidants Redox Signaling*. 5 (1), 69-78.
- [18] Shao HB, Chen XY, Chu LY, Zhao XN, Wu G, Yuan YB, Zhao CX, Hu ZM, (2006). Investigation on the relationship of Proline with wheat anti-drought under soil water deficits. *Biointerfaces*. 53 (2), 113-119.
- [19] Shao HB, Chu LY, Cheruth AJ, Zhao CX, (2008). Water-deficit ,stress-induced anatomical changes in higher plants.

Comptes Rendus de Biologies 331, 215-225.

- [20] Terman A, Brunk UT, (2006). Oxidative stress, accumulation of biological 'Garbage', and aging, *Antioxidants Redox Signaling*. 8 (1-2), 197-204.
- [21] Wang YS, Yang ZM (2005). Nitric oxide reduces aluminum toxicity by preventing oxidative stress in the roots of *Cassia tora* L. *Plant Cell Physiol*. 46(12): 1915–1923.
- [22] Wang YS, Wang J, Yang ZM, Lu B, Wang QY, Li SQ, Lu YP, Wang SH, Sun X (2004). Salicylic acid modulates aluminium induced oxidative stress in roots of *Cassia tora* L. *Acta Bot. Sin.* 46: 819–828.
- [23] Yang H, Wong JWC, Yang ZM, Zhou LX (2001). The ability of *Agropyronelongatum* to accumulate the single metal of cadmium, copper, nickel and lead and root exudation of organic acids. *J Environ Sci*, 13(3): 368–75.
- [24] Yücel E, Hatıpoğlu A, Sözen E, Güner ST (2008). The effects of the lead (PbCl) on mitotic cell division of Anatolian Black Pine (*Pinus nigra* ssp. *pallasiana*). *Biodicon* 1(2): 124129.