Review Article

Antimicrobial Activity of Mn Complexes Incorporating Schiff Bases: A Short Review

Sharif Uddin¹, Md. Saddam Hossain², Md. Abdul Latif², Md. Rabiul karim¹,
Ranjan K. Mohapatra³, Md. Kudrat-E-Zahan^{1,*}

¹Department of Chemistry, Rajshahi University, Rajshahi, Bangladesh

²Department of Chemistry, Begum Rokeya University, Rangpur, Bangladesh

³Department of Chemistry, Government College of Engineering, Keonjhar, Odisha, India

Email address:

kudrat.chem@ru.ac.bd (Md. Kudrat-E-Zahan)

*Corresponding author

To cite this article:

Sharif Uddin, Md. Saddam Hossain, Md. Abdul Latif, Md. Rabiul karim, Ranjan K. Mohapatra, Md. Kudrat-E-Zahan. Antimicrobial Activity of mn Complexes Incorporating Schiff Bases: A Short Review. *American Journal of Heterocyclic Chemistry*. Vol. 5, No. 2, 2019, pp. 27-36. doi: 10.11648/j.ajhc.20190502.12

Received: May 8, 2019; Accepted: June 12, 2019; Published: July 1, 2019

Abstract: Antibiotic resistance has been growing at an anxious rate and as a result the activity of antibiotics against Gram-negative and Gram-positive bacteria has dropped dramatically day by day. Metal ions play many critical functions in humans. Deficiency of some metal ions can lead to disease like pernicious anemia, growth retardation, heart disease in infants etc. In this sense there is a strong need to synthesis new substances that not only have good spectrum of activity, but having new mechanisms of action. Inorganic compounds particularly Mn metal complexes have played an important role in the development of new metal based drugs. Over the past few years, there have been many reports on their applications in homogeneous and heterogeneous catalysis. The activity is usually increased by complexation therefore to understand the properties of both ligands and metal can lead to the synthesis of highly active compounds. The influence of certain metals on the biological activity of these compounds and their intrinsic chemical interest as multidentate ligands has prompted a considerable increase in the study of their coordination behavior. Development of a new chemotherapeutic Schiff bases and their metal complexes is now attracting the attention of medicinal chemists. In this review we have focused on research of Mn complexes incorporation of Schiff bases carry out over the past few decades which has sought to possess preclinical pharmacological screenings like anti-bacterial, anti-fungal, anti-tuberculosis, anti-inflammatory, anti-cancer, DNA- interaction and anti-tumor action of synthetic metal complexes.

Keywords: Schiff Bases, Mn Complexes, Metal Based Drugs, Antibacterial, Anti-Inflammatory

1. Introduction

Interest in the study of Schiff base hydrazones has been growing because of their antimicrobial, anti-tuberculosis, and anti-tumour activity [1-12]. Schiff bases play an important role in inorganic chemistry, as they easily form stable complexes with most transition metal ions. The development of the field of bioinorganic chemistry has increased the interest in Schiff base complexes, since it has been recognized that many of these complexes may serve as models for biologically important species [1-5]. Coordination compounds

derived from aroyl hydrazones have been reported to act as enzyme inhibitors and are useful due to their pharmacological applications [6-12]. Schiff bases are a class of compounds having azomethine linkage (C=N) and they are widely used in the field of coordination chemistry due to their ability to form stable complexes [13]. Manganese Schiff base complexes have subsequently been found to play an important role in many biological processes such as trans amination, decarboxylation, condensation and -elimination

and racemization [14] gave complexes of various metals with Schiff bases derived from thiazoles and their carbohydrazones having antimicrobial activity. Seleem reported stability constants and thermodynamic parameters of some complexes with Schiff base hydrazones containing pyrimidine moiety [15]. They are among the most expedient and versatile ligands which are easily synthesized in a one-step classical condensation reaction through common aldehydes with amines or ketones usually in quantitative yields. Schiff base ligands of heterocyclic compounds and their transition metal complexes are of great interest as simple structural models of biological system due to the presence of hetero atoms such nitrogen, oxygen and sulphur groups [13]. Schiff bases have gained importance in medicinal and pharmaceutical fields due to a broad spectrum of biological activities like anti-inflammatory [16-20], analgesic [21-23], antimicrobial [24, 25], anticonvulsant [26], antitubercular [27], anticancer [28, 29], antioxidant [30], antihelmintic [31], and so forth. The wide ranges of applications of these compounds have continued to attract interest among inorganic chemists. Many of them are also thermally stable and have synthetic flexibility. Some of these compounds are found to be useful both as clinical diagnostic agents and in chemotherapeutic application. In view of the properties mentioned above, a large number of Schiff bases have been synthesized and extensively studied the literature [32]. The design and synthesis of organic-inorganic hybrid complexes based on strong coordinate bonds and multiple weak non-covalent forces has become a rapidly growing field of research [32]. Microbes encounter a variety of metal ions in the environment and interact with them, which is sometimes beneficial or detrimental depending on the chemical/physical nature and oxidation state of the metal ion. The microbes have the ability to bind to metal ions present in the external environment at the cell surface and to transport them into the cell for various intracellular functions [33]. In this review we explored the overview application of antimicrobial activities of Mn complexes incorporating Schiff bases.

2. Antimicrobial Activity of Schiff Base Manganese Complexes

The antimicrobial activity of the Schiff base ligand and their metal complexes at low concentrations were evaluated against Gram positive and Gram negative bacteria as well as fungal species [34]. Antimicrobial activity of Schiff base metal complexes of Mn(II) derived from 2-amino pyridine have been synthesized by Rehab K Al-Shemary and *et al.* The in-vitro antimicrobial activity of the scrutinised ligand and its metal complexes was tested against the bacteria *Escherichia coli* and *Staphylococcus* *usurios* by disc diffusion method using nutrient agar as a medium. The stock solution was prepared by dissolving the compounds in DMSO. In a typical procedure, a well was made on agar medium inoculated with

the microorganism. The well was filled with the test solution using a micropipette and the plate was incubated 24 h for bacteria at 37°C and 72 h for fungi at 30°C. After incubation, the diameter of the clear zone of inhibition surrounding the sample is taken as a measure of the inhibitory power of the sample against the particular test organism. This may be due to that the chelation considerably reduces the polarity of the metal ion mainly because of partial sharing of its positive charge with the donor groups and possible electron delocalization over the whole created ring such, chelation could also enhance the lipophilic character of the central metal atom, which subsequently favors its permeation through the lipid layer of the cell membrane [35]. Schiff base metal complexes of Mn derived from Ofloxacin have been synthesized by Rabia Arshad and *et al.* All metal complexes except Ni complex showed enhanced antimicrobial activity while the Mn complex being more active. Similarly against *E. coli*, metal complexes showed somewhat comparable or enhanced activity than the parent drug except Ni complex which has just lower activity than the drug. The order of antibacterial activity is as follows; Metal Complexes of ligands > Ligands > Parent Drug. Ofloxacin is an antibacterial drug so it does not show any activity against fungi. In case of antifungal activities, we know that some other quinolones show very low activities against any fungi. Besides very poor antifungal records of quinolones, we managed to measure the antifungal properties of our prepared complexes in order to know any effect on certain fungi. The prepared complexes were used on two fungal strains *i.e.* *A. flavus* and *A. niger*. Activities of ligands are not better enough to be declared as good antifungal agents. Metal complexes of these ligands also exhibited very slight increase in antifungal activities [36].

Figure 1. Structural formula of Ofloxacin.

G. kumar and *et al* were synthesized some M(III) complexes of Cr, Mn and Fe (figure 3) with a Schiff base derived from 2-amino-4-ethyl-5-hydroxybenzaldehyde and thiocarbohydrazide (figure 2). The antibacterial and antifungal activities of the newly synthesized compounds were evaluated by the agar well diffusion method. The Mn(III) complex exhibited moderate activity against *P. aeruginosa* and *B. megaterium*, was highly effective against *K. fragilis*, *C. albicans* and *T. reesei*, but showed no activity against *E. coli*, *S. aureus* and *R. rubra*.

Figure 2. Synthesis of the Schiff base ligand.

Figure 3. Synthesis of the M(III) Schiff base complexes.

Sangita Sharma and et al were investigated Antimicrobial Activity of Some Transition Metal Complexes (Mn, Co, Zn, Ni) derived from L-Proline and Kojic Acid. The antimicrobial effect of metal complexes at 2×10^{-3} M concentration on the growth of bacteria (*Escherichia coli* and *Bacillus subtilis*), yeast (*Sacharomyces cerevisiae*) and fungi (*Aspergillus niger*) was investigated in liquid media. The highest activity was reported with Ni(II) complexes with 92.65%, 88.09%, 85.77% and 45.1% growth inhibition for *S. cerevisiae*, *E. coli*, *B. subtilis* and *A. niger* respectively. The second highest antimicrobial activity was exerted by cobalt complex with 80.39%, 71.38%, 70.51% and 52.62% growth inhibitory effect respectively for *B. subtilis*, *S. cerevisiae*, *E. coli*, and *A. niger*. The Ni(II) complexes exerted the highest activity amongst all test organisms under study except for *A. Niger* which was best with Zn(II) complexes. The least activity has been reported for the Mn(II) complexes. This may be due to fact that manganese is an essential trace nutrient in living cell life and a wide group of enzymes have manganese as cofactor [37]. Schiff base, (E)-N'-(-1-(1-ethyl-4-hydroxy-2-oxo-1,2-dihydroquinolin-3-yl) ethylidene)nicotinohydrazide and its Cu(II), Ni(II), Co(III), Mn(II) and Fe(III) complexes have been prepared by Trimbak K Chondhekar et al (figure 4). The ligand and its metal complexes have been screened in vitro for their antibacterial and antifungal activities by agar- cup method using DMSO as a solvent and poison- plate method using potato dextrose agar medium in DMSO respectively. All the metal complexes do not show any activity against Gram – ve bacteria, *E. coli* and *S. typhi*, except Mn(II) complex which show highest activity against *E. coli* comparable to the standard, penicillium. On the other hand, the ligand show considerable activity against Gram +ve bacteria, *S. aureus* and

remains inactive against *B. subtilis*. The most active is the Ni(II) complex which shows better inhibitory action against both gram positive bacterial species. Mn(II) and Cu(II) complexes also inhibit the growth. However Co(III) and Fe(III) complexes do not have significant effect against both the species. All these active metal complexes exhibited a more inhibitory effect compared to the ligand. This is because the chelation reduces the polarity of the metal ion because of partial sharing of its positive charge with the donor groups which results in to the increase of lipophilic nature of the complexes which in turn favors its permeation to the lipid layer of the membrane. Moreover, other factors, like stability constants, molar conductivity, solubility and magnetic moments also affect the antimicrobial activity of the complexes [38].

Figure 4. Proposed structure for the complexes.

Novel Schiff base and its 3d transition metal complexes of Mn(II), Fe(III) and V(IV) has been designed and synthesized from 4-aminoantipyrine and Ethyl 4- methyl – oxo-6-phenylhexahydro pyrimidine-5-carboxylate by Mohanambal D. and Arul Antony S. In vitro antibacterial

(*E. coli*, *Staphylococcus aureus*, *Bacillus spp.*, *Salmonella spp.*) and invitro antifungal (*Candida*, *Aspergillus niger*,) tested by Mueller Hinton Agar solid media (MHA) and Sabouraud's dextrose 9SDA media. Agar diffusion assay was carried out to evaluate the antimicrobial activity of ligand and its metal complexes synthesized. The plates were incubated at 37°C for 24hr, during which activity was evidenced by the presence of azone of inhibition surrounding the well and antibacterial and antifungal activity was expressed as the mean diameter of inhibition zones (mm) produced by the synthesized compounds [39]. Aishatu Salihu Mohammed and et al were investigated Antifungal Activity of Manganese (II) Complex with Schiff Base Derived from Acetylacetone and Leucine. The antifungal activities of the prepared compounds were studied using *Candida albican* and *Sacchro myces cerevisia et al.* The manganese (II) complex showed strong activity on the fungal isolates at medium and high concentrations. All organisms were found to be resistant to the Schiff base at all concentrations. The diameter of inhibition zone for each treatment ranged from 7 mm to 38 mm [40]. Complexes of Mn(II) and Fe(III) with trimethoprim Schiff base ligands, HL¹ (figure 5) and HL² (figure 6) derived from salicylaldehyde and benzaldehyde were synthesized and investigated by Naomi P. Ndahi and et al. The results obtained on antimicrobial studies, the ligand (HL¹) and its complexes did not show much effect against the bacterial isolates as well as the fungus *Candida albicans* tested as compared with the standard trimethoprim drug at all concentrations with the exception of Fe(HL¹) Cl]Cl which indicated susceptibility with zone of 2 2 inhibition of 30.00 mm (30 µg/ml), 24.70 mm (20 µg/ml) and 20.30 mm (10 µg/ml) against *Corynae* bacterium species compared to 23.30 mm (30 µg/ml), 18.7 mm (20 µg/ml) and 15.00 mm (10 µg/ml) shown by the parent drug trimethoprim. In contrast, HL² complexes showed greater activity against *Streptococcus pyogenes*, *Corynae* bacterium species and *Shigella dysenteriae* at all concentrations compared to the trimethoprim standard. Mn(HL₂)Cl₂ complex (Figure 8) showed low to moderate activity against all the tested organisms. [Fe(HL₂) Cl]Cl did not show any visible inhibition against *Bacillus subtilis*. Antimicrobial resistance against some promising antibiotics drugs such as trimethoprim and others was commonly observed in many countries worldwide during in vivo analysis of pathogens [41].

Figure 5. Proposed structure of HL¹.

Figure 6. Proposed structure of HL².

Figure 7. Proposed structure of [Mn (HL¹)₂Cl₂].

Figure 8. Proposed structure of [Mn(HL²)₂Cl₂]Cl.

Mohammad Muzammil Y. Kuddushi and et al were Synthesis And Characterization of Schiff Base Aniline With 5-Bromo -2- Hydroxyl Benzaldehyde and Their Metal Complexes. The ligands and the metal complexes show very good antimicrobial properties against The gram-positive bacteria-*B. megaterium* and gram-negative-*Escherichia coli* [42]. Manganese(III) complexes derived from the bis_Schiff bases N, N' bis (5 fluorosalicylidene)-1, 2 diaminoethane (H2La) and 3, 4-bis(2-hydroxybenzylideneamino)pyridine (H2Lb), respectively, have been prepared and characterized by Q. R. Liu, L. W. Xue and G. Q. Zhao. Qualitative determination of antimicrobial activity was done using the disk diffusion method [24, 25]. In general complex I shows greater antibacterial and antifungal activities against *Staphylococcus aureus*, *Escherichia coli*, and *Candida albicans* compared to complex II and it is obvious that H² L^a has also greater antibacterial and antifungal activities against *Staphylococcus aureus*, *Escherichia coli*, and *Candida albicans* compared to H² L^b [43]. Carbazoles containing two

new Schiff bases (Z,Z)-N,N-bis[(9-ethyl-9H-carbazole-3-yl)methylene]propane-1, 3-diamine (L1) and (Z,Z)-N,N-bis[(9-ethyl-9H-carbazole-3-yl)methylene]-2, 2-dimethylpropane-1, 3-diamine (L2) and their Co(II) and Mn(II) complexes were synthesized and their antimicrobial properties investigated by Selma Bal and Sedat Salih Bal. The synthesized Mn and Co compounds were effective for almost all of the microorganisms; in particular, for the microorganisms *K. pneumoniae*, *E. aerogenes*, *S. faecalis*, and *S. aureus*. The title ligands brought about bigger inhibition zones. Generally, they were most effective, both antimicrobial and antifungal. Among the fungi, they gave the biggest inhibition zones for *C. Utilis* with 27mm and 25mm zones, respectively. Among the coordination compounds, the higher values, that are both antimicrobial and antifungal, were recorded for the manganese (II) complex compounds [44]. M (III) complexes of Cr, Mn and Fe with a Schiff base derived from 2-amino-4-ethyl-5-hydroxybenzaldehyde and thiocarbonylhydrazide were synthesized and characterized by G. Kumar and et al. The Cr (III) complex showed high antibacterial and antifungal activity against *B. megaterium*, *K. fragilis*, *R. rubra* and *C. albicans*, moderate activity against *E. coli*, *S. aureus* and *P. aeruginosa*, but no activity against *T. reesei*. The Mn(III) complex exhibited moderate activity against *P. aeruginosa* and *B. megaterium* and highly effective against *K. fragilis*, *C. albicans* and *T. reesei*, but showed no activity against *E. coli*, *S. aureus* and *R. rubra*. The Fe (III) complex exhibited moderate activity against *S. aureus*, *P. aeruginosa*, *B. megaterium* and *T. reesei*, had a higher effect against *K. fragilis* and *C. albicans*, but no activity against *E. coli* and *R. rubra* microorganisms [45]. A new series of transition metal complexes of Cu (II), Ni (II), Co (II), Mn (II), Zn (II), VO (IV), Hg (II) and Cd (II) have been synthesized from the Schiff base (L) derived from 4-aminoantipyrine, 3-hydroxy-4-nitrobenzaldehyde and o-phenylenediamine by N Raman and et al. The in vitro biological screening effects of the investigated compounds were tested against the bacteria: *Salmonellatyphi*, *Staphylococcus aureus*, *Escherichiacoli*, and *Bacillus subtilis*. The antifungal activities of the compounds were evaluated by the well-diffusion method against the fungi viz., *Aspergillusniger*, *Aspergillus flavus* and *Rhizoctonia bataicola* [46]. Metal complexes of Mn(II), Cu(II) and Ni(II) with Schiff base ligand (HL) derived from condensation of propionaldehyde with ethylenediamine were synthesized and investigated their antimicrobial activity by Wakil IM and et al. The antimicrobial activity of the Schiff base ligand and their metal complexes [Mn(II), Cu(II) and Ni(II) complexes] at concentrations of 10, 20 and 30 µg/ml were evaluated against Gram positive (*Staphylococcus aureus*, *Streptococcus pyogenes*) and Gram negative(*Escherichia coli*, *Salmonella typhi*)bacteria as well as fungal specie – *Candida albicans*. The Schiff base ligand (HL1) did not show efficacy against all the microorganisms tested. But, on complexation with Mn(L1-L1)X₂ showed increased activity (P<0.001) against all the organisms that were previously resistant to the parent ligand. *Albicans* which was resistant to the Mn(II) complex. At concentration of 30 µg/ml, it was observed that all the test

organisms were susceptible on Cu(L1- L1)X₂ complex. Ni(L¹-L¹)X₂ was found to be resistant to *E. coli* while *S. typhi*, *S. aureus*, *S. pyogenes* and *C. Albicans* were found to be susceptible to Ni(II) complex. Enhanced activity (P<0.001) was observed for Cu(II) complex on comparison with Mn(II) and Ni(II) [47].

Figure 9. Propionaldehyde.

Figure 10. Proposed structure of Schiff base (HL).

Figure 11. Proposed structure of the metal complexes.

Two new Schiff base ligands (L1, L2) have been prepared from the reaction of 2, 6 diacetylpyridine and 2-pyridinecarboxyaldehyde with 4-amino-2, 3-dimethyl-1-phenyl-3-pyrozolin-5-on, and their Co(II), Cu(II), Ni(II), Mn(II), and Cr(III) metal complexes have also been prepared by E. Ispir and et al. The antibacterial and antifungal activities of the new compounds against eight bacteria, namely, *Escherichia coli*, *Staphylococcus aureus*, *Klebsiellapneumoniae*, *Mycobacteriums megmatiss*, *Pseudomonas aeruginosa*, *Enterococcus cloacae*, *Bacillus megaterium*, and *Micrococcusluteus*, and five fungi, namely, *Kluyveromyces fragilis*, *Rhodotorularubra*, *Candida albicans*, *Saccharomycescerevisiae*, and *Trichodermareesei*. Cefodizime, Cefuroxime, Cephalothin, Oxacillin and Nystatin are taken as the standard reference antibiotics for antibacterial and antifungal activities. The antibacterial and antifungal

activities of standard antibiotics gave 45 and 8 mm inhibition zones for fungi and bacteria respectively [49].

Figure 12. The proposed structure of the new Schiff base ligands.

Figure 13. The proposed structures of the L1 (a) and L2 (b) complexes.

Metal complexes of Mn(II), Fe(II), Co(II) and Cd(II) ions (Figure 15) with Schiff base ligand 4-[(pyridin-2-ylimino)methyl]phenol (figure 14) derived from condensation of 2-amino pyridine with 4-hydroxybenzaldehyde was prepared and investigated their antimicrobial properties by M. S. Hossain and et al. The free Schiff base ligand and their metal complexes were screened for their antibacterial activity against the strains the *Escherichia coli*, *Pseudomonas aereuginosa*, *Acetobacteraceti*. The susceptibility zones were the clear zones around the discs killing the bacteria. All the Schiff base and metal complexes individually exhibited varying degrees of inhibitory effects on the growth of tested bacterial species. Most of the metal complexes showed more antibacterial activity than Schiff base ligand [50].

Figure 14. Structure of Schiff base $L = [C_{12}H_{10}N_2O]$.

Figure 15. Structure of metal complexes.

The new Schiff base ligand Salicylaldehyde-4-Chlorobenzoylhydrazone (HL) (figure 16) and its Mn(II), Co(II) and Ni(II) metal complexes (figure 17) were synthesized and characterized by A. H Manik shete et al. The anticancer activity of the ligand (HL) and its Mn(II) complexes was determined by sulforhodamine -B colorimetric

assay. Mn(II) complex shows moderate activity for breast (MCF7) cell line with IC₅₀ value of 11.1 μ M and resistant to the colon (HT29) cell line with IC₅₀ value of 84.9 μ M. Mn(ii)

complexes showed significant anticancer activity. Ligand and its Mn(II), complexes were active against MCF7 than HT29 cell line [51].

Figure 16. Synthesis of Schiff base.

Figure 17. Synthesis of metal complexes.

Figure 18. SOVPTH metal complexes ($M=Mn^{2+}$).

Metal complexes of a novel Schiff base derived from condensation of p-toluic hydrazide and o-Vanillin are reported and characterized by S. Kondaiah and *et al.* The biological activity of the Schiff base ligand and their metal complexes and streptomycin (as a standard compound) were tested against bacteria because bacteria can achieve resistance to antibiotics through biochemical and morphological modifications. The organisms used in the present

investigations included *Staphylococcus aureus* and *Bacillus subtilis* (as gram positive bacteria) and *Pseudomonas aeruginosa* and *Escherichia coli* (as gram negative bacteria). The data obtained reflect that the Schiff base ligand shows moderate activity in comparison with *Staphylococcus aureus*, *Bacillus subtilis* and less active in comparison with *Pseudomonas aeruginosa*, *Escherichia coli* [52].

A series of manganese(II) and copper(II) complexes with reduced Schiff bases derived from o-phenylenediamine has been prepared and characterized by S. Belaid *et al.* Mn(II) complexes (figure 20) present a good activity against most of the dermatophytes tested and fungi of the genus *Scedosporium*, especially for [MnL1(H₂O)₂] which possesses the more simple structure. In addition, their activity against *Scedosporium* species, i.e., *S. apiospermum* which ranks the second among the filamentous fungi colonizing the airways of patients with cystic fibrosis and *S. prolificans* which causes severe disseminated infections in leukemic patients. [53].

Figure 19. Formula of the ligands: $R=H$: H_2L^1 , $R=CH_3$: H_2L^2 , $R=Cl$: H_2L^3 .

Figure 20. Structural scheme of the complexes (b) $[MnL^1(H_2O)_2]$, $[MnL^3(H_2O)_2]$.

Manganese (II) complexes of bishydrazones were prepared and investigated their antimicrobial activities by P. Mittal and V. Uma. The results of antibacterial and antifungal activity show that these compounds exhibit greater antibacterial and antifungal activity against the *E. Coli*, *Klebsiella*, *Enterococci*, *S. Aureus* *Candida Albicans* and *Candida Krusei* [54].

3. Conclusions

In this work, the antimicrobial activity of Mn complexes has been reviewed. With the improvement in the field of medicinal chemistry, the role of transition metal complexes especially Mn complexes as therapeutic compounds is becoming increasingly significant. Recent advances in medicinal chemistry have made possible formation of number of transition metal complexes with organic ligand of interest, which can be used as therapeutic agent. Significant progress in the synthesis of platinum based anti-cancer drugs like cisplatin has been made. These drugs have proven to be highly effective chemotherapeutic agents for treating various types of cancers. The use of transition metal complexes as therapeutic compounds has become more and more pronounced. These complexes offer a great diversity in their action; they do not only have anti-cancer properties but have also been used as anti-inflammatory, anti-infective and anti-diabetic compounds. Development of Mn complexes as drugs is not an easy task; considerable effort is required to get a compound of interest. Besides all these limitations and side effects, Mn complexes

would be widely used as antimicrobial agents and make a large contribution to medicinal therapeutics.

Acknowledgements

The authors are thankful to the Chairman, Department of Chemistry, Rajshahi University, Bangladesh for the laboratory facilities.

References

- [1] Chohan, Z. H.; Sheazi, S. K. A. Synthesis and characterization of some Co (II), Cu (II) and Ni (II) complexes with nicotinyldiazine derivatives and their biological role of metals and anions on the antibacterial properties. *Synth. React. Inorg. Met.-Org. Chem.* 1999, 29, 105-118.
- [2] Jayabalakrishnan, C.; Natarajan. K. Synthesis, characterization and biological activities of ruthenium (II) carbonyl complexes containing bifunctional tridentate Schiff bases. *Synth. React. Inorg. Met.-Org. Chem.* 2001, 31, 983-995.
- [3] Jeeworth, T.; Wah, H. L. K.; Bhowon, M. G.; Ghoorhoo. D.; Babooram, K. Synthesis and antibacterial/catalytic properties of schiff bases and Schiff base metal complexes derived from 2, 3-diaminopyridine. *Synth. React. Inorg. Met.-Org. Chem.* 2000, 30, 1023-1038.
- [4] Dharmaraj, N.; Viswanalhamurthi, P.; Natarajan. K. Ruthenium (II) complexes containing bidentate Schiff bases and their antifungal activity. *Transit. Metal Chem.* 2001, 26, 105-109.
- [5] Collins, C. H.; Lyne, P. M. In *Microbiol Methods*; University Park Press: Baltimore, 1970.
- [6] Savanini, L.; Chiasserini, L.; Gaeta, A.; C. Pellerano. Synthesis and anti-tubercular evaluation of 4-quinolyldiazones. *Biorg. Med. Chem.* 2002, 10, 2193-2198.
- [7] Ochiai, Ei-ichiro. *Bioinorganic Chemistry*; Allyn and Bacon: Boston, 1977.
- [8] Ferrari, M. B.; Capacchi, S.; Pelosi, G.; Reffo, G.; Tarasconi, P.; Alberlini, R.; Pinelli, S.; Lunghi, P. Synthesis, structural characterization and biological activity of helicon thiosemicarbazone monohydrate and a copper (II) complex of salicylaldehyde thiosemicarbazone. *Inorg. Chim. Acta* 1999, 286, 134-141.
- [9] Agarwal, R. K.; Singh, L.; Sharma, D. K.; Singh, R. Synthesis, spectral and thermal investigation of some oxovanadium (IV) complexes of isonicotinic acid hydrazide. *Turk J. Chem.* 2005, 29, 309-310.
- [10] Elo, H.; Sunila, I.; Lumme, P. Studies on the acute toxicity of antineoplastic metal chelate. *Transbis (salicyldoximate) copper(II) in rats.* *Inorg. Chim. Acta* 1987, 136, 61-63.
- [11] Elo, H.; Lumme, P. *Trans-bis (salicyldoximate) copper (II) and its derivatives as antiproliferative and antineoplastic agents.* *Inorg. Chim. Acta* 1987, 136, 149-153.
- [12] Ali, M. A.; Kabir, M. H.; Nazimuddin. M.; Majumder, S. M. H.; Tarafder, M. T. H.; Akhair, M. Synthesis, characterization and antifungal properties of some four-coordinate nickel (II) and four and five-coordinate copper (II) complexes containing tridentate thiosemicarbazones and heterocyclic bases. *Indian J. Chem.* 1988, 27A, 1064-1067.

- [13] Mohammad S, Summaiya H, Mohammed S, Owais M, Mohammad A, Saud, IA (2016). Pharmacophore hybrid Approach of New Modulated bis-diimine CuII/ZnII Complexes based on 5-chloro Isatin Schiff base Derivatives Synthesis, Spectral Studies and Comparative Biological Assessment. *Journal of Photochemistry & Photobiology*. 157, 39–56.
- [14] S. D. Angadi, R. K. Shikkargoi, N. N. Millikarjuna, K. Siddapa and V. H. Kulkarni, E. *Journal Chem*. 2003, 5, 566.
- [15] H. S. Seleem, Stability Constants and Thermodynamic Parameters of $Mn^{2+}Co^{2+}$, Ni^{2+} Complexes with Some Schiff Base Hydrazones Containing the Pyrimidine Moiety, *Ann. Chem.*, (2003) pp. 305-314.
- [16] Sathe BS, Jaychandran E, Jagtap VA, Sreenivasa GM. Synthesis characterization and anti-inflammatory evaluation of new fluorobenzothiazoleschiff's bases. *International Journal of Pharmaceutical Research and Development*. 2011; 3 (3):164-169.
- [17] Sondhi SM, Singh N, Kumar A, Lozach O, Meijer L. Synthesis, anti-inflammatory, analgesic and kinase (CDK-1, CDK-5 and GSK-3) inhibition activity evaluation of benzimidazole/benzoxazole derivatives and some Schiff's bases, *Bioorganic and Medicinal Chemistry*. 2006; 14 (11):3758-3765.
- [18] Pandey A, Dewangan D, Verma S, Mishra A, Dubey RD. Synthesis of schiff bases of 2-amino-5-aryl-1, 3, 4-thiadiazole and its analgesic, anti-inflammatory, antibacterial and antitubercular activity. *International Journal of ChemTech Research*. 2011; 3 (1):178-184.
- [19] Chandramouli C, Shivanand MR, Nayanbhai TB, Bheemachari B, Udupi RH. Synthesis and biological screening of certain new triazole schiff bases and their derivatives bearing substituted benzothiazole moiety, *Journal of Chemical and Pharmaceutical Research*. 2012; 4 (2):1151-1159.
- [20] Chinnasamy RP, Sundararajan R, Govindaraj S. Synthesis, characterization, and analgesic activity of novel schiff base of isatin derivatives. *Journal of Advanced Pharmaceutical Technology and Research*. 2010; 1 (3):342-347.
- [21] Mounika K, Anupama B, Pragathi J, Gyanakumari C. Synthesis, characterization and biological activity of a Schiff base derived from 3-ethoxy salicylaldehyde and 2-amino benzoic acid and its transition metal complexes. *Journal of Scientific Research*. 2010; 2 (3):513-524.
- [22] Venkatesh P. Synthesis, characterization and antimicrobial activity of various Schiff bases complexes of Zn (II) and Cu (II) ions. *Asian Journal of Pharmaceutical and Health Sciences*. 2011; 1 (1):8-11.
- [23] Chaubey AK, Pandeya SN. Synthesis & anticonvulsant activity (Chemo Shock) of Schiff and Mannich bases of Isatin derivatives with 2-Amino pyridine (mechanism of action), *International Journal of Pharm Tech Research*. 2012; 4 (4):590-598.
- [24] Aboul-Fadl T, Mohammed FA, Hassan EA. Synthesis, antitubercular activity and pharmacokinetic studies of some Schiff bases derived from 1-alkylisatin and isonicotinic acid hydrazide (INH), *Archives of Pharmacal Research*. 2003; 26(1):778-784.
- [25] Miri R, Razzaghi-asl N, Mohammadi MK. QM study and conformational analysis of an isatin Schiff base as a potential cytotoxic agent. *Journal of Molecular Modeling*. 2013; 19 (2):727-735.
- [26] Ali SMM, Abul Kalam Azad M, Jesmin M. In vivo anticancer activity of Vanillin semicarbazone, *Asian Pacific Journal of Tropical Biomedicine*. 2012; 2 (6):438-442.
- [27] Wei D, Li N, Lu G, Yao K. Synthesis, catalytic and biological activity of novel dinuclear copper complex with Schiff base. *Science in China B*. 2006; 49 (3):225-229.
- [28] Avaji PG, Vinod Kumar CH, Patil SA, Shivananda KN, Nagaraju C. Synthesis, spectral characterization, in-vitro microbiological evaluation and cytotoxic activities of novel macrocyclic bis hydrazone. *European Journal of Medicinal Chemistry*. 2009; 49 (9):3552-3559.
- [29] Venugopala KN, Jayashree BS. Synthesis of carboxamides of 2-amino-4-(6-bromo-3-coumarinyl)thiazole as analgesic and anti-inflammatory agents, *Indian Journal of Heterocyclic Chemistry*. 2003; 12 (4):307-310.
- [30] Emayavaramban M, Kumar K, Mani P, Prabhakaran B, Muthuvel A. Synthesis, complexation, spectral and antimicrobial study of some novel 5 bromofluorobenzaldehydeoxime and semicarbazone under ultrasonic irradiation. *International Journal of Advanced Chemistry*. 2014; 2 (1):20-23.
- [31] Salman A Khan, Abdullah M Asiri, Khalid Al-Amry, Maqsood Ahmad Malik. Synthesis, Characterization, Electrochemical Studies, and In vitro Antibacterial. *Activity of Novel Thiosemicarbazone and Its Cu (II), Ni (II), and Co (II) Complexes*. Hindawi Publishing Corporation Scientific World Journal, 2014, Article ID 592375:9.
- [32] Hina Z, Anis A, Asad UK, and Tahir AK. Synthesis, Characterization and Antimicrobial Studies of Schiff base Complexes. *Journal of Molecular Structure*, 2015, 1097, 129–135.
- [33] Sayed M. Abdallah, Gehad G. Mohamed, M. A. Zayed, Mohsen S. Abou El-Ela. Spectroscopic study of molecular structures of novel Schiff base derived from o-phthalaldehyde and 2-aminophenol and its coordination compounds together with their biological activity. *Spectrochimica Acta Part A*, 2009, 73, 833–840.
- [34] Md. Saddam Hossain, C. M. Zakaria, Md. Kudrat-E-Zahan. Metal Complexes as Potential Antimicrobial Agent: A Review. *American Journal of Heterocyclic Chemistry*. 2018, 4 (1), 1-21.
- [35] Rehab K Al-Shemary, Wurood A Jaafar, and Ali N Niseaf. Structure, Diagnosis, and in the Vitro Antimicrobial evaluation of 2-amino pyridine-derived Ligand Schiff base and its complexes with Cu (II), Hg (II), Ni (II), Mn (II) and Co (II). *RJPBCS*, 2017, 8 (3), 174-184.
- [36] Rabia Arshad, Iftikhar Hussain Bukhari, Fareeha Anum, Zartasha Aftab, Zahida Noreen. Synthesis, Spectral and Biological studies of transition metal complexes of schiff base derived from Ofloxacin. *International Journal of Advanced Research in Biological Sciences*, 2016. 3 (4), 99-108.
- [37] Sangita Sharma, Jayesh Ramani, Jasmin Bhalodia, Neha Patel, Khushbu Thakkar and Rajesh Patel. Synthesis, Characterization and Antimicrobial Activity of Some Transition Metal Complexes (Mn, Co, Zn, Ni) With L-Proline and Kojic Acid. *Advances in Applied Science Research*, 2011, 2 (4), 374-382.

- [39] Sandeep A. Kumbalpur, Archana A. Kachare, Sunil G. Shankarwar and Trimbak K. Chondhekar. Synthesis, characterization and antimicrobial activities of Schiff base ligand and its metal complexes. *Der Pharma Chemica*, 2015, 7 (8), 88-93.
- [40] Mohanambal D. and Arul Antony S. Synthesis, Characterization and Antimicrobial activity of some novel Schiff Base 3d Transition Metal Complexes Derived from Dihydropyrimidinone and 4-Aminoantipyrine. *Research Journal of Chemical Sciences*, 2014, 4 (7), 11-17.
- [41] Aishatu Salihu Mohammed, Fatima Abubakar, Abigail Ebi Aye. Synthesis, Characterization and Antifungal Activity of Manganese (II) Complex with Schiff Base Derived from Acetylacetone and Leucine. *American Journal of Nano Research and Applications*, 2017, 5 (6), 110-113.
- [42] Naomi P. Ndahi, Hussein Garba, Ibrahim Waziri, Aliu A. Osunlaja, and Habila A. N. Putaya. Complexes of Mn (II) and Fe (III) with Schiff bases Derived from Trimethoprim with Salicylaldehyde and Benzaldehyde as Potential Antimicrobial Agents. *Nigerian Journal of Pharmaceutical and Biomedical Research*, 2018; 3 (1):53.
- [43] Mohammad Muzammil Y. Kuddushi., Mohammed Abrar H. Malek., Vinod L. Patidar, Mihir S. Patel., Roma K. Patel and Rohit H. Dave. Synthesis and Characterization of Schiff Base Aniline With 5-Bromo-2-Hydroxyl Benzaldehyde and Their Metal Complexes. *International Journal of Recent Scientific Research*, 2018, 9 (4), 26026-26030.
- [44] Q. R. Liu, L. W. Xue, and G. Q. Zhao. Manganese (III) Complexes Derived from bis-Schiff Bases: Synthesis, Structures, and Antimicrobial Activity. *Russian Journal of Coordination Chemistry*, 2014, 40, (10), 757-763.
- [45] Selma Bal and Sedat Salih Bal. Cobalt (II) and Manganese (II) Complexes of Novel Schiff Bases, Synthesis, Characterization, and Thermal, Antimicrobial, Electronic, and Catalytic Features. *Advances in Chemistry*, 2014, 1-14.
- [46] Gajendra Kumar, Dharmendra Kumar, C. P. Singh, Amit Kumar and V. B. Rana. Synthesis, Physical Characterization and Antimicrobial Activity of Trivalent Metal Schiff Base Complexes. *J. Serb. Chem. Soc.* 2010, 75 (5), 629-637.
- [47] N. Raman, J. Dhavethu Raja and A. Sakthivel. Synthesis, Spectral Characterization of Schiff Base Transition Metal complexes: DNA cleavage and antimicrobial activity studies. *J. Chem. Sci.*, 2007, 119 (4), 303-310.
- [48] I. M. Wakil, N. P. Ndahi, M. B. Abubakar, I. Waziri. Synthesis, Characterization, and Antimicrobial Studies of Propionaldehyde Schiff base Metal (II) Complexes. *Chemistry Research Journal*, 2017, 2 (2), 39-45.
- [49] E. Ispir, S. Toroglu, A. Kayral, Z. Synthesis, characterization, antimicrobial and genotoxic activities of new Schiff bases and their complexes. *Transition Met Chem*, 2008, 33, 953-960.
- [50] M. S. Hossain, C. M. Zakaria, M. K. Zahan. Synthesis and Characterization with Antimicrobial Activity Studies on some Transition Metal Complexes of N, O Donor Novel Schiff Base Ligand. *J. Sci. Res.* 2017, 9 (2), 209-218.
- [51] A. H. Manikshete, M. M. Awatade, S. K. Sarsamkar, M. R. Asabe. Synthesis, characterization, antimicrobial, anticancer and antidiabetic activity of new Manganese (II), Nickel (II) and Cobalt (II) complexes with Salicylaldehyde-4-chlorobenzoylhydrazone. *International Journal of Engineering Science Invention*, 2015, 4 (1), 22-29.
- [52] S. S. Kondaiah, G. Naga Raja Reddy, D. Rajesh, Jacob Joseph. Synthesis, Characterization, and Antibacterial Activity of the Schiff Base Derived from P-Toluic Hydrazide and o-Vanillin (OVPTH Ligand) and its Mn (II), Co (II), Ni (II) and Cu (II) Complexes. *Indian Journal of Advances in Chemical Science*, 2013, 1 (4), 228-235.
- [53] Sabrina Belaid, Anne Landreau, Safia Djebbar, Ouassini Benali-Baitich, Gilles Bouet, Jean-Philippe Bouchara. Synthesis, characterization and antifungal activity of a series of manganese (II) and copper (II) complexes with ligands derived from reduced N, N'-O-phenylenebis (salicylideneimine), *Journal of Inorganic Biochemistry*, 2008, 102, 63-69.
- [54] P. Mittal and V. Uma. Coordination Behavior and Biological Activity of Unsymmetrical Bis-Hydrazone Towards Manganese (II) Ion. *Int. J. Chem.* 2008, 6 (2), 1050-1060.