Isolation and Identification of *Enterobacteriaceae* from Patients with Community Acquired Urinary Tract Infection

Abd Elrahman Mustafa Abd Elrahman Osman^{*}, Shingray Osman Hashim,
Mohammed Abdall Musa, Omer Mohammed Tahir

Medical Laboratory Sciences Division, Port Sudan Ahlia College, Port Sudan, Sudan

Email address:

abdo_rahman96@hotmail.com (A. E. M. A. E. Osman)

^{*}Corresponding author

To cite this article:

Abd Elrahman Mustafa Abd Elrahman Osman, Shingray Osman Hashim, Mohammed Abdall Musa, Omer Mohammed Tahir. Isolation and Identification of *Enterobacteriaceae* from Patients with Community Acquired Urinary Tract Infection. *American Journal of Health Research*. Vol. 6, No. 1, 2018, pp. 25-31. doi: 10.11648/j.ajhr.20180601.14

Received: October 28, 2017; **Accepted:** December 4, 2017; **Published:** January 22, 2018

Abstract: The study was carried out in red sea state during the period from November 2013 to March 2014 to investigate the *Enterobacteriaceae* from patients suffering from community-acquired urinary tract infections and then do sensitivity test for each of isolates. In this study out of 100 specimens 52 *Enterobacteriaceae* species were isolated from different clinics and hospitals in Port Sudan city. The specimens were cultured on CLED media (cystine lactose electrolyte deficiency). Identification was done by gram's stain and conventional biochemical reactions. Then the anti-microbial sensitivity tests were done as follows: (Ampicillin-Sulbactam "AS", Co. trimoxazole "BA", Ceftriaxone "CI", Chloramphenicol "CH", Cephalaxin "PR", Tetracycline "TE", Ciprofloxacin "CP", Amikacin "AK", Sparfloxacin "SC", Gatifloxacin "GF", Norfloxacin "NX") by Kirby-Bauer disc diffusion method. The study revealed that the most part of strains are sensitive to Chloramphenicol and Amikacin and resistant to Ampicillin-Sulbactam (AS). The identified *Enterobacteriaceae* were as follows; *Escherichia coli* 34 (65%), *Klebsiella pneumoniae* 10 (19%), *Klebsiella oxytoca* 3 (6%), *Salmonella Paratyphi A* 3 (6%), *Proteus mirabilis* 1 (2%), *Citrobacter* 1 (2%).

Keywords: Enterobacteriaceae, Urinary Tract Infections, Escherichia Coli, Red Sea State

1. Introduction

Urinary tract infection represents one of the commonest bacterial infections. The *Enterobacteriaceae* are the most frequent pathogen detected, causing most of the urinary tract infection, *Enterobacteriaceae* are a group of largest, most heterogeneous collection of medically important gram-negative bacilli, being found worldwide in soil, water and are part of the normal flora in the intestine of most animal and human. This family includes many genera such as (*Escherichia coli*, *Klebsiella*, *Salmonella* ...etc). These days the main cause for concern among *Enterobacteriaceae* especially that cause community acquired infection is *Escherichia coli*. [1].

A urinary tract infection (UTI) is a condition in which one or more parts of the urinary system (the kidneys, bladder, and urethra) become infected. Although some cases of UTI are due to fungus or a virus, most are caused by one of several

types of bacteria. Most cases of UTIs are caused by bacteria that typically multiply at the opening of the urethra and travel up to the bladder. Much less often, bacteria spread to the kidney from the bloodstream [1].

The urinary system helps maintain proper water and salt balance throughout the body and also expels urine from the body. The symptoms of urinary tract infections result from the presence and growth of bacteria or other microorganisms in the urinary tract. The urinary tract is normally a sterile environment. The bacteria, in some cases, can get flushed up into the kidneys and therefore can cause a kidney infection. Both kidney and bladder infections are more common in women because their urethras are shorter than men, making it easier for organisms to get from outside into the bladder. Most typically, a woman develops a UTI if she has been

sexually active, (hence the moniker "honeymoon cystitis"), or has been careless with her hygiene habits (for example, wiping from back to front after a bowel movement) [2].

Early recognition of bacteraemic UTI and prompt, appropriate treatment are critical in reducing the mortality. A diagnosis of a urinary tract infection can easily be missed or delayed in the elderly. This is because some symptoms, such as fatigue and weakness, may not be noticed or might be associated with aging. Also the extensive and inappropriate use of antimicrobial agents has invariably resulted in the development of antibiotic resistance which, in recent years, has become a major problem worldwide. Family physicians can provide empirical treatment without the benefit of a pre-therapy urine culture. The extensive and inappropriate use of antimicrobial agents has invariably resulted in the development of antibiotic resistance which, in recent years, has become a major problem worldwide. For example, Quinolones which are one of the most widely used antibiotics in the community for the treatment of UTI, the unfortunate excessive use of this agent that has led to a considerable and worrying increase in the rate of *E. coli* resistant isolates in many countries, and this is just a start because as more patients use antimicrobial drugs without pre-therapy culture, more resistant bacteria will gain. [2].

Urologists have tended to ignore the clinical importance and urologic realities of community-acquired urinary tract infections (UTIs) despite their significant prevalence, cost, morbidity, and increasing management problems. This is primarily because of our perception that uncomplicated UTIs are common but not a serious problem (patients do not die from uncomplicated UTIs), easy to diagnose (simple midstream urine culture), and simple to treat (short course of antibiotics). Nevertheless, data on increasing prevalence, cost, morbidity, antibiotic resistance, recurrence, and relapse suggest that the urological community needs to have another look at community-acquired UTIs. [3].

This study aimed to determine the frequency of isolated *Enterobacteriaceae* and antimicrobial susceptibility of uropathogens in culture-positive community-acquired UTIs over a 3 months period, community-acquired UTIs over a four months period in order to. Understanding the pathogenesis of UTI which may lead to better methods of prevention and treatment. [3].

2. Materials and Method

2.1. Study Approach

Study approach is to identify and isolate the possible causative organisms and the characterization of isolated strains using bio typing and anti bio gram.

2.1.1. Study Type and Design

Cross-sectional descriptive study.

2.1.2. Study Area

The study area is different hospitals and clinics in Red Sea state.

2.1.3. Study Period

During the period from November 2013 to March 2014

2.2. Methodology

2.2.1. Sample Size and Collection

According to standard method all specimens were examined to detect, and identify pathogens or their products using: microscopic examination of specimen to detect their motility, morphology and staining reaction. Also culture techniques to isolate pathogens in pure form and to identify them then test their antibiotic sensitivity and biochemical.

a. Sampling

Samples for community acquired UTI to investigate and diagnosis of microbial diseases (urine specimen).

b. Urine Specimens (100 Samples)

Urine specimens were collected in case of urinary tract infection as following. A mid stream urine is obtained in a sterile container after cleaning the external genitalia with tap water and drying. Samples should reach the laboratory within one hour after voiding or kept refrigerated at 4°C to avoid multiplication of bacteria in urine then we do appropriate test to identify and isolate the causative organisms.

2.2.2. Cultivation of Specimens

Urine specimens were inoculated onto CLED medium by using sterile loop, then incubated aerobically at 37°C overnight.

2.2.3. Examination of Bacterial Growth

The primary culture on CLED medium that showed significant growth was examined for fermentation. The morphological character, size, shape, colour were observed and recorded.

2.2.4. Interpretation of Culture Growth

The culture growth obtained was interpreted as significant ($>10^5$ CFU/ml). Cultured of less than (10^3 CFU/ml) of urine was considered insignificant, while culture with no growth were considered negative. Significant cultures were further investigated.

2.2.5. Purification of Bacterial Growth

The isolates were streaked onto Nutrient agar and incubated overnight at 37°C. The resultant growth was checked for purity and stored in Bijiou bottle for further investigation.

2.2.6. Identification of the Isolated Bacteria

I. Colonial Morphology

Colonial characteristics were observed on CLED medium after overnight incubation isolated organisms were grown on CLED medium.

II. Biochemical Tests (Conventional Test)

These are called biochemical tests because they are tests which identify the bacteria on the basis of the presence of certain enzymes and other biochemical properties.

III. Kliglar Iron Agar

this medium was originally designed as a multi-test medium. It provides a low degree of sensitivity for H₂S production (often required when differentiating members of the (Enterobacteriaceae). The medium is now used principally as a standard test for H₂S. One disadvantage of multitest media is that chemical interaction-in this case acid production from fermentable sucrose-may inhibit blackening of the iron indicator. Some Citrobacter and Proteus species have this ability. KIA agar should be used in conjunction with a urease test to eliminate Proteus spp when screening for Salmonellae.

IV. Citrate Utilization Test

This is for the ability of on organism to utilize citrate as the sole carbon and energy source for growth and an ammonium salt as sole source of nitrogen.

V. Indole Test

This test demonstrates the ability of certain bacteria to decompose the amino acid tryptophan to indole, which accumulates in the medium. Indole is then tested for by a colorimetric reaction with p- dimethylaminobenzaldehyde. Add 0.5ml Kovacs reagent to see result.

VI. Urease Test

Bacteria, particularly those growing naturally in an environment exposed to urine, may decompose urea by means of the enzyme urease. The occurrence of this enzyme can be tested for by growing the organism in the presence of urea and testing for alkali production by means of a suitable pH indicator. An alternative method is to test for the production of ammonia from urea by means of Nessler's reagent.

VII. Motility Test

In semi solid agar media, motile bacteria (Swarm) give a diffuse spreading growth that is easily recognized by the naked eye. Motility may thus be detected more easily than by the microscopical method.

VIII. Antimicrobial Sensitivity Test

All isolated microorganisms were subjected to antimicrobial sensitivity test using Modified Kirby-Bauer disc diffusion method [10].

3. Results

One hundred urine specimens were collected from patients suffering from UTI during the period from November 2013 to March 2014. Out of 100 patients (52%) were positive, while (48%) were negative. Among the positive growth 4 of them were NLF organisms (8%), and 48 (92%) were L. F organisms (Table 1).

The most frequent patients were female (75%), 38 of them were positive culture and the male were less frequent (25%) only 14 of them were positive culture (Figure 1).

The isolated Enterobacteriaceae were as follows: *Escherichia coli* 34 (65.4%), *Klebsiella pneumoniae* 10 (19.2%), *Klebsiella oxytoca* 3 (5.7%), *Salmonella Paratyphi A* 3 (5.7%), *Proteus mirabilis* 1 (2%), *Citrobacter* 1 (2%) (Figure 2).

For female patients the age ranged from (one year to 10 years) one patient showed positive culture, 4 urine cultures were positive from the ages (11 to 20 years). Seven urine cultures were positive from the ages of (21 to 30 years). From the age of 31 to 40 years old 13 of them were positive urine culture and this is the most predominant age infected with Enterobacteriaceae, from (41 years old to 50 years) 9 patients were positive urine culture and more than (50 years) only 4 patients were positive urine culture (Figure 3). For male patients the ages ranged from (one year to 10 years) one patient was positive urine culture, from the ages of (21 to 30 years) 4 patients were positive for urine culture, from (41 years to 50 years) 4 patients were positive for urine culture. More than (50 years) 4 patients represent the most frequent infection in male (Figure 4).

The result of antimicrobial sensitivity tests revealed in (Table 2) and (Figure 5).

Figure 1. Distribution of specimen according to gender and the number of positive culture for each gender.

Table 1. Significant and insignificant growth.

Total Number of specimens	Significant growth (%)		Insignificant growth(%)
100 specimens	52%	LF(%)	48%
	NLF(%)	48 specimens92.3%	
	4 specimens8%

Figure 2. Distribution of specimen according to gender and the number of positive culture for each gender. **Figure 3.** Percent of each isolated Enterobacteriaceae.

Figure 4. Distribution of female specimens according to ages and positive culture.

Figure 5. Distribution of male specimens according to ages and positive culture.

Table 2. Number and percentage of antimicrobial sensitivity tests (S=sensitive, R=resistance).

Organisms	AS	BA	CI	CH	PR	TE	CP	Ak	SC	GF	NX	
S	<i>E.coli</i>	1	13	22	34	18	7	21	33	14	29	19
		3%	38%	65%	100%	53%	20%	62%	97%	41%	85%	55%
	<i>k.pnuemonie</i>	10	4	4	9	1	7	6	7	4	8	5
		100%	40%	40%	90%	10%	70%	60%	70%	40%	80%	50%
	<i>k.oytoca</i>	0	0	0	2	0	1	2	3	2	3	2
		0%	0%	0%	67%	0%	33%	67%	100%	67%	100%	67%
	<i>S.para A</i>	0	3	3	3	2	3	3	3	2	2	3
		0%	100%	100%	100%	67%	100%	100%	100%	67%	67%	100%
	<i>Proteus</i>	0	0	1	1	1	1	0	1	0	1	0
		0%	0%	100%	100%	100%	100%	0%	100%	0%	100%	0%
R	<i>ciyrobacter</i>	0	0	0	1	0	0	1	1	1	1	1
		0%	0%	0%	100%	0%	0%	100%	100%	100%	100%	100%
	<i>E.coli</i>	33	21	12	0	16	27	13	1	20	5	15
		97%	62%	35%	0%	57%	80%	38%	3%	59%	15%	45%
	<i>k.pnuemonie</i>	0	6	6	1	1	3	4	3	6	2	5
		0%	60%	60%	10%	90%	30%	40%	30%	60%	20%	50%
	<i>k.oytoca</i>	3	3	3	1	3	2	1	0	1	0	1
		100%	100%	100%	33%	100%	67%	33%	0%	33%	0%	33%
	<i>S.para A</i>	3	0	0	0	1	0	0	0	1	1	0
		100%	0%	0%	0%	33%	0%	0%	0%	33%	33%	0%
	<i>Proteus</i>	1	1	0	0	0	0	1	0	1	0	1
		100%	100%	0%	0%	0%	0%	100%	0%	100%	0%	100%
	<i>ciyrobacter</i>	1	1	1	0	1	1	0	0	0	0	0
		100%	100%	100%	0%	100%	100%	0%	0%	0%	0%	0%

Figure 6. Percentage of each antimicrobial used for UTI infections.

4. Discussion

This study was conducted to determine the main *Enterobacteriaceae* caused infection in community acquired UTI patient and the antimicrobials sensitivity test. The results revealed that the lactose ferment *Enterobacteriaceae* type represent (92%), while non lactose fermented organisms represent (8%) and this result agreed with the result obtained by Panahi Y., *et al.* [4]. The ratio between male to female is 1:3, 25% --75% respectively this result is nearly similar to result 1:2 report by Sharifian *et al.* [5].

The result showed that UTI cases among female of age group (31-40 years) were found more susceptible to UTI (33.4%), that agree with results obtained by Orret and Shurland [6]. In the present study the most prevalent organisms were *E.coli* (65.4%) of all *Enterobacteriaceae* isolated, this result agree that obtained by Ekweozor and Onyemenen [7], who showed that *E.coli* account (62%), the followed organism is *Klebsiella Spp* this result is similar to that obtained by Gupta K., *et al.* [8]. In our study the elder males (over 50 years) are more susceptible to UTI this result agree with a result obtained by Torkaman M., *et al* [9]. In the present study the antimicrobial sensitivity test demonstrated that Ampicillin-Sulbactam (21.2%) and Tetracycline (36.5%) are the lowest active agent, While Chloramphenicol (96.2%) and Amikacin (92.3), are the highest active agent this result confirmed the result obtained by Modarres *et al.* [7].

On other hand, the result showed that *E.coli* was more susceptible to Chloramphenicol (100%), Amikacin (97%) and Gatifloxacin (97%), respectively, this results disagree with a result obtained by (Sharifian *et al* [5]. who found that *E. coli* was most sensitive to Ceftizoxime.

5. Conclusions and Recommendations

5.1. Conclusions

- I. The most prevalent organism in UTI infections is *E.coli*.
- II. Female are more susceptible to UTI than male.
- III. Ages from 31-40 are the most infected among female.
- IV. Men whom ages more than 50 are more susceptible to UTI than other ages.
- V. Chloramphenicol and Amikacin are the highest active agents in most strains.
- VI. Ampicillin-Sulbactam is the lowest active antimicrobial agents due to high resistant rate.
- VII. Resistance rate of *Enterobacteriaceae* increased to commonly used antimicrobial agents.

5.2. Recommendations

- a. Urine specimen should be investigated for culture and susceptibility test before giving the patient any therapy to decrease the resistant rate among organisms causing community acquired UTI.
- b. Increase awareness about the hazards of using antibiotic for treatment without pre-therapy culture and how bacteria gain resistant among people with UTI infection.
- c. Uses of API 20 E are recommended for investigation of all *Enterobacteriaceae* in Microbiology lab for more accuracy.
- d. Establishment of antimicrobial policies and treatment guidelines.
- e. Uses of Chloramphenicol and Amikacin as routine treatment for community acquired UTI infections.
- f. In depth studies by using modified technique (molecular methods) for precision and accuracy is recommended.

References

- [1] Okesola. A. O and Makanjuola. O. (2009). Resistance to Third-generation Cephalosporins and other Antibiotics by Enterobacteriaceae in Western Nigeria, *American Journal of Infection disease*, 5(1):17-20.
- [2] Murray Patrick. R, Rosenthal. Kens, Kabayashi George S and Pfaaller. Michael A. (2002). *Medical Microbiology*, 4th edition, Mosby. P266-279.
- [3] Rayan. Kenneth. J and George. C. Ray. (2004). *Sherris Medical Microbiology* 4th edition, McGraw-HILL, New York, Chicago. P. 343-367.
- [4] Panahi Y, Beiraghdar F, Moharamzad Y, Matinzadeh ZK, Einollahi B. The incidence of urinary tract infections in febrile children during a two-year period in Tehran, Iran. *Trop Doct*. 2008; 38(4):247-9.
- [5] Sharifian M, Karimi A, Tabatabaei SR, Anvaripour N. Microbial sensitivity pattern in urinary tract infections in children: a single center experience of 1,177 urine cultures. *Jpn J Infect Dis*. 2006;59(6):380-2.
- [6] Orret and Shurland, 1998. Bacterial etiologic agents of urinary tract infection in Hospitalized patients Nigeria, Abuja.
- [7] Modarres S, Oskoi NN. Bacterial etiologic agents of urinary tract infection in children in the Islamic Republic of Iran. *East Mediterr Health*. 1997;3:290-5.
- [8] Gupta K. Emerging antibiotic resistance in urinary tract pathogens *Infect Dis Clin North Am*. 2003;17(2):243-59.
- [9] Torkaman M, Afsharpaiman SH, Hoseini MJ, Moradi M, Mazraati A, Amirjalali S, et al. Platelet count and neonatal sepsis: a high prevalence of Enterobacter spp. *Singapore Med J*. 2009; 50(5):482-512
- [10] Collee, J. G; Morison, B. P.; Fraser, A. G. Simmons, A. (1996). *Maki and McCartney Practical Medical Microbiology* (14th edition). Churchill Livingstone New York.