Serum Metabolomics Reveals Cholic Acid, Chenodeoxycholic Acid, and Taurochenodeoxycholic Acid as Potential Biomarkers for Hypothyroid Rats

Hidegori Nagao^{1,*}, Masanori Suzuki², Hironori Aoki¹, Kouichi Minato¹

¹Pharmacokinetics Research Department of ASKA Pharmaceutical Co., Ltd., Kawasaki, Japan

²Department of Analytical Research, ASKA Pharma Medical Co., Ltd., Kawasaki, Japan

Email address:

nagao-h@aska-pharma.co.jp (H. Nagao)

To cite this article:

Hidegori Nagao, Masanori Suzuki, Hironori Aoki, Kouichi Minato. Serum Metabolomics Reveals Cholic Acid, Chenodeoxycholic Acid, and Taurochenodeoxycholic Acid as Potential Biomarkers for Hypothyroid Rats. *American Journal of Life Sciences*.

Vol. 3, No. 4, 2015, pp. 295-305. doi: 10.11648/j.ajls.20150304.17

Abstract: Hypothyroidism decreases energy metabolism including carbohydrate and lipid metabolism and protein synthesis, due to reduced serum levels of the thyroid hormones thyroxine (T_4) and triiodothyronine (T_3). Although many endogenous serum metabolites are influenced by hypothyroidism, serum metabolomic profiling has rarely been applied to the study of hypothyroidism. In the present study, we investigated potential biomarkers for hypothyroidism using serum metabolomics, and then measured serum levels of these endogenous metabolites using an analytical method: ultra-performance liquid chromatography coupled with quadrupole time-of-flight mass spectrometry. There was a significant difference in the metabolic profiles of thyroidectomized (Tx) and normal rats. We found that many bile acid (BA) levels were significantly changed in serum of Tx rats. Simultaneous measurement of 12 different BAs in serum revealed that cholic acid (CA), chenodeoxycholic acid (CDCA), and taurochenodeoxycholic acid (TCDCA) levels significantly increased in Tx rats by approximately 25-fold, 11-fold, and 3-fold, respectively, compared with those of control rats. In Tx rats with active hormone T_3 replacement, serum T_3 levels were returned to physiological levels. However, these changes in BA levels were maintained at a high level. These results indicate that T_3 replacement does not normalize the thyroid hormonal milieu. Thus, increased CA, CDCA, and TCDCA levels in serum after Tx may be a homeostatic response to not only T_3 but also pro-thyroid hormone T_4 deficiency. This study is the first to report that CA, CDCA, and TCDCA may be potential biomarkers for hypothyroidism and the efficacy of thyroid hormone replacement therapy in hypothyroidism.

Keywords: Metabolomics, Biomarker, Bile Acid, Hypothyroidism, Thyroid Hormone Replacement Therapy

1. Introduction

Thyroid hormones (THs) play an important role in both energy homeostasis and regulation and also during the development and differentiation of cells. THs secreted from the thyroid gland are thyroxine (T_4) and triiodothyronine (T_3). T_4 has little biological activity and is considered a pro-hormone, as activation occurs through deiodination of T_4 at the 5'-position of the phenolic ring to form T_3 , the active form of thyroid hormone [1]. Most of the actions of TH seem to be dependent on binding of T_3 to a nuclear thyroid hormone receptor (TR). There are three major forms of the TR, designated $\alpha 1$, $\beta 1$, and $\beta 2$. TR $\alpha 1$ is the predominant subtype in bone, the gastrointestinal tract, cardiac and skeletal muscle, and the central nervous system;

TR $\beta 1$ is most abundant in the liver and kidney; and TR $\beta 2$ is more discretely expressed in the hypothalamus, pituitary, cochlea, and retina [2-5]. Thus, THs act on a wide variety of tissues, and hypothyroidism results in multiple physiological responses. Some studies have shown that TH deficiency due to hypothyroidism leads to a decreased rate of metabolism of lipids, proteins, and carbohydrates [6,7].

Metabolomics, which can be defined as measurement of all cellular metabolites, is the comprehensive assessment of endogenous metabolites and quantification of metabolites from a biological sample [8]. In contrast to genomics, transcriptomics, and proteomics, which involve macromolecules with similar chemical properties, such as

DNA, RNA, and proteins, metabolomics analysis deals with diverse properties of low molecular weight compounds. This technique can be widely applied to a number of fields including pharmacology, toxicology, and food science [9,10]. As metabolomics involves the downstream products of genomics and proteomics, it is also regarded as a complement to other “omics” for interpretation of gene function [11,12]. Metabolomics is a powerful tool for discovering new small molecule biomarkers. Following further validation, the identified markers can help to diagnose and evaluate prognosis for various diseases, such as diabetes mellitus, liver injury, and acute renal toxicity [13–15].

In this study, we used serum metabolomics profiling in rats with hypothyroidism induced by surgical thyroidectomy using ultra-performance liquid chromatography coupled with quadrupole time-of-flight mass spectrometry (UPLC-QTOF/MS). First, we analyzed the metabolites in serum, and detected differences in bile acids (BAs). These metabolites were analyzed using supervised orthogonal partial least squared discriminant analysis (OPLS-DA) and online metabolite data bases search. Then, concentrations of the detected BAs in serum were quantified using a simultaneous analytical method [16]. We investigated sequential changes in individual serum BA levels in rats after thyroidectomy, and then investigated the effects of replacement of the active hormone T_3 on individual serum BA levels. Our results suggest that cholic acid (CA), chenodeoxycholic acid (CDCA), and taurochenodeoxycholic acid (TCDCA) may be potential biomarkers for hypothyroidism and the efficacy of TH replacement therapy in hypothyroidism.

2. Materials and Methods

2.1. Materials

T_3 was purchased from Sigma-Aldrich Co. (St. Louis, MO, USA). CA and CDCA were purchased from Wako Pure Chemical Industries, Ltd. (Osaka, Japan). Glycocholic acid (GCA), tauroursodeoxycholic acid (TUDCA), TCDCA, and taurodeoxycholic acid (TDCA) were purchased from Merck (Darmstadt, Germany). Taurocholic acid (TCA), taurine- β -muricholic acid (β TMCA), glycochenodeoxycholic acid (GCDCA), tauroolithocholic acid (TLCA), taurohyodeoxycholic acid (THDCA), and taurohyocholic acid (THCA) were purchased from Steraloids UK Ltd. (London, England). The internal standards, CA-2,2,3,4,4- d_5 and TCA-2,2,4,4- d_4 , were purchased from Toronto Research Chemicals Inc. (Tronto, Canada), and GCA-2,2,4,4- d_4 was from C/D/N isotopes Inc. (Quebec, Canada). Deionized water used for UPLC mobile phase was prepared on Milli-Q advantage A10 water purification system (Millipore, Bedford, MA, USA). Methanol and acetonitrile were of HPLC grade and obtained from Kanto Chemical Inc. (Tokyo, Japan) and Wako Pure Chemical Industries, Ltd. (Osaka, Japan), respectively. Formic acid

and ammonium acetate were from Sigma Chemical (St. Louis, MO, USA).

2.2. Animals

Seven-week-old male Sprague-Dawley rats were obtained from Charles River Laboratories Japan (Kanagawa, Japan). Animals were fed a commercial diet (FR-2, Funabashi Farm Co., Chiba, Japan) and filtered tap water *ad libitum*. The cages were located in a light (0800 - 2000 h lights on), temperature ($23 \pm 5^\circ\text{C}$), and humidity ($60 \pm 20\%$) controlled room. The rats were allowed to acclimatize for 1 week before starting the experiments.

All experimental procedures were approved by the Animal Research Committee of ASKA Pharmaceutical Co., in accordance with the Basic Guidelines for Proper Conduct of Animal Testing and Related Activities in the Research Institutions under the Jurisdiction of the Ministry of Health, Labour and Welfare of Japan.

2.3. Thyroidectomy

Rats were made hypothyroid by surgical thyroidectomy as described previously in detail [17]. After the surgery, serum thyroid stimulating hormone (TSH) rapidly increased, and body weight gain achieved complete stasis. Complete resection of the thyroid in the thyroidectomized (Tx) rats was confirmed at the end of the experiment by macroscopic observation at necropsy. The Tx rats drank 1% calcium lactate aqueous solution *ad libitum* for 1 week after Tx. Normal group received sham operation as control.

2.4. Experiment 1

Rats were divided into two groups ($n=5$), the Tx and normal (control) groups. Blood were collected from tail vein for 3 weeks. All blood samples were obtained between 0900 and 1100 h to minimize fluctuations in THs [18]. Serum samples were separated by centrifugation at $16000 \times g$ for 5 min at 4°C , and stored at -20°C until analyzed.

2.5. Experiment 2

Osmotic minipumps (Alzet, model 2ML4, DURECT, CA, USA), delivering $1.5 \mu\text{g } T_3/\text{head/day}$ (Tx+ T_3 group; $n=6$) were implanted subcutaneously in rats on day 0 under isoflurane anesthesia at 2 weeks after Tx. The doses of T_3 were determined by our preliminary experiment and previous literature [19]. The vehicle was saline, in which T_3 (final concentration; $25 \mu\text{g/mL}$ in $0.5 \text{ mmol/L NaOH/saline}$) was diluted after NaOH solubilization ($5 \text{ mg } T_3/20 \mu\text{L}$ of 1 mol/L NaOH). Six normal and Tx rats (control and Tx group) received pumps with $0.5 \text{ mmol/L NaOH/saline}$ only. On the day before pump implantation (day -1) and after implantation (day 14), blood were collected from tail vein between 0900 and 1100 h. Serum samples were separated by centrifugation at $16000 \times g$ for 5 min at 4°C , and stored at -20°C until analyzed.

2.6. Metabolomics Analysis

2.6.1. UPLC-QTOF/MS

Serum samples which were collected on the final day (3 weeks after Tx) of the experiment 1, were prepared for UPLC-QTOF/MS analysis by acetonitrile protein precipitation. The serum samples (40 μ L) were mixed with cold acetonitrile (80 μ L) for 30 sec and centrifuged at $16000 \times g$ for 5 min at 10°C . An aliquot of the supernatant transferred to a clean tube. An equivalent 0.1% formic acid was added to the supernatant. Metabolite extracts (2 μ L) were injected onto a reverse-phase ACQUITY UPLC HSS T3 column (2.1×100 mm i.d., 1.8 μ m particles, Waters, Milford, MA, USA) using an ACQUITY UPLC system (Waters). The flow rate of the mobile phase was 400 μ L/min. Mobile phase A consisted of 0.1% formic acid in water, while mobile phase B was acetonitrile. The gradient was initiated with 99% A and maintained for 2 min, and then linearly decreased to 5% within 3 min, and maintained for 4 min. The column eluent was introduced directly into the mass spectrometer. Mass spectrometry was performed on a Q-TOF instrument (Xevo QToF MS, Waters), operating in negative electrospray ionization (ESI⁻) mode from m/z 50 to 1000 in MS scanning. Capillary voltage was 2.5 kV, sample cone was 25 V, desolvation temperature 500°C , source temperature 150°C , and desolvation gas flow 1000 L/h. Leucine enkephalin (m/z 556.2771) was used as the lock-mass; a solution of 0.2 ng/mL (50:50 acetonitrile: water containing 0.1% formic acid) was infused into the instrument at 5 μ L/min via an auxiliary sprayer. MassLynx software (v.4.1, Waters) was used to acquire the chromatogram and mass spectrometric data in centroid format. In the MS^E (Waters) was performed on all samples, where data were collected at both low (5.0 eV) and high (20.0 to 35.0 eV) collision energies, in order to obtain fragmentation data simultaneously.

2.6.2. Data Processing and Analysis

The raw data of chromatographic peaks obtained from the UPLC-TOF/MS system were recognized and matched by MarkerLynx XS software (Waters). A list of the detected peaks intensities was generated for the first chromatogram, and the retention time and m/z value pairs were used as identifiers. The ion intensities for each detected peak were then normalized to the sum of the peak intensities in that sample. There was no significant correlation between the total intensities used for normalization and the sample groups being compared in the study. The multivariate data matrix was analyzed by EZ-info software (Waters). The supervised orthogonal partial least squares discriminant analysis (OPLS-DA) was used to concentrate group discrimination into the first component with remaining unrelated variation contained in subsequent components. Candidate markers were extracted from S-plots constructed following analysis with OPLS-DA, and markers were chosen based on their contribution to the variation and correlation within data set.

2.6.3. Identification of Candidate Markers

The elemental compositions were generated by MarkerLynx based on the accurate mass of the high-contribution-score metabolites. MassFragment application manager (Waters) was used to facilitate the MS/MS fragment ion analysis. The markers were matched based on retention time, mass assignment, and online database query. The accurate mass and structure information of the markers obtained from KEGG (www.genome.jp/kegg/), HMDB (www.hmdb.ca/), and ChemSpider (www.chemspider.com/) databases.

2.7. Quantitation of BA Concentrations in Serum

Serum BA concentrations were determined using the slightly modified method [16]. In brief, serum samples (50 μ L) were mixed with 50 μ L methanol and 50 μ L of isotopic standards mixture (2000 ng/mL) in an ice-cold water bath. After being mixed thoroughly for 1 min, the mixture stood at -20°C for 20 min. The mixture was centrifuged at $16000 \times g$ for 10 min at 4°C , and then 2 μ L of the supernatant was injected to the UPLC-TOF/MS system. The UPLC system was coupled online to a Xevo Q-TOF mass spectrometer (Waters) in negative electrospray mode with a scan range of m/z 50 to 1000. The chromatographic separation of the analytes was performed using an Acquity UPLC BEH C18 column (100×2.1 mm i.d., 1.7 μ m particles, Waters). Details of the LC and MS conditions including gradient elution and the selected reaction monitoring transitions for all the compounds as well as the corresponding internal standard for each BA were described in a previous report [16]. The calibration curve was linear for all BAs over a range of 10 to 25 ng/mL (Min.) ~ 1000 to 10000 ng/mL (Max.).

2.8. Serum TH Concentrations

Serum T_4 and T_3 concentrations were determined by the method using online solid-phase extraction liquid chromatography-mass spectrometry/mass spectrometry (online SPE LC-MS/MS) [17]. In brief, an API5000 triple-quadrupole mass spectrometer (AB SCIEX, CA, U.S.A.) equipped with a TurboIonSpray source and Shimadzu HPLC system was employed to perform the analysis using isotope dilution with deuterium labeled internal standard, [$^2\text{H}_5$] T_4 . A 20 μ L of rat serum was mixed with internal standard acetonitrile solution for deproteinization, and was diluted with 0.1% formic acid. After centrifugation, the supernatant was injected into the LC-MS/MS system. Online extraction and chromatographic separation of the analytes were performed using respectively a Shim pack MAYI-ODS column (10 mm $\times$ 2.0 mm i.d., 50 μ m particles, Shimadzu, Kyoto, Japan) and a Synergi Polar-RP 80A column (50 mm $\times$ 2.0 mm i.d., 4 μ m particles, Phenomenex, Utrecht, the Netherlands). Details of the LC conditions including gradient elution and the selected reaction monitoring transitions are described in a previous report [17]. Linear calibration curves of T_4 and T_3 were obtained in the concentration range of 0.1 to 100 ng/mL, with a lower limit of quantitation of 0.1 ng/mL.

2.9. Serum TSH Concentrations

Serum TSH concentrations were determined by Enzyme-Linked Immunosorbent Assay using TSH ELISA Test Kit (Endocrine Technology, Newark, CA, U.S.A.).

2.10. Calculations and Statistical Analysis

Total bile acid (total BA) concentration was calculated as the sum of the 10 bile acids: CA, CDCA, GCA, GCDCA, TCA, TCDCA, TDCA, THDCA, TUDCA, and β TMCA. TLCA and THCA were not included as only trace amounts were detected. All data of serum BA concentrations were expressed as mean $\pm$ standard deviation (S.D.). Statistical significance of serum BA concentrations between control and Tx group in the experiment 1 was determined by an unpaired Student's *t*-test. Differences between three groups in the experiment 2 were assessed by one way analysis of variance. Post hoc testing was performed for inter-group comparisons

using the Tukey's test. In all cases, the results were considered significant at $p < 0.05$.

3. Results

3.1. Serum Metabolomic Profiling

Serum samples were collected on the final day (3 weeks after Tx) of the experiment 1, and analyzed using UPLC-QTOF/MS. As more metabolites were detected in negative ion mode than in positive ion mode, negative ion mode was used for this study. Representative base peak ion (BPI) chromatograms of the serum of control and Tx rats are presented in Fig. 1. The variations in *m/z* values and retention times of selected peaks were less than 5 mDa and 0.05 min, respectively. After peak alignment, the serum datasets contained 4456 metabolite ions. The pre-processed UPLC-QTOF/MS data were further investigated using multivariate statistical analysis.

Fig. 1. The UPLC-QTOF/MS BPI profile of serum samples in (A) control group and (B) Tx group.

3.2. Multivariate Data Analysis

The supervised OPLS-DA was applied to obtain better discrimination between the control and Tx groups. Good separation between the two groups was observed, and there

was no variation among the rats within each group and no outliers on the vertical axis (Fig. 2A). The goodness-of-fit parameter R^2 and the predictive ability parameter Q^2 were 0.999 and 0.895, respectively. These results indicate that this

model fits and is useful for discriminating between the two groups.

The S plots from the OPLS-DA model were created using the loading profile of the first component (p) and the correlation of p with the first component ($p(\text{corr})$), which represents the reliability of p to the first component (Fig. 2B). Each plot represents one chromatogram of a metabolite with a specific retention time. We used cut-off values of $p \geq |0.05|$ and $p(\text{corr}) \geq |0.8|$, and 38 plots located inside the squares in Fig. 2B were selected; 28 plots were more increased in the Tx group compared with the control group, and 10 plots were more decreased in the Tx group compared with the control

group. According to an identity check based on raw data and features of the chromatographic peaks, the candidate metabolites detected in the profiling process were identified by MS/MS analysis or by searching their accurate masses using the online metabolite databases, KEGG, HMDB, and ChemSpider. Several metabolites of pyrimidine, phenylalanine, and the amino acid metabolism pathway were matched (data not shown). Many BAs and their metabolites, CA, CDCA, GCA, GCDCA, CA glucuronide, and CDCA sulfate, were matched as markers (Table 1). Among these, the metabolite that suggested the largest change in Fig. 2B (retention time 4.79) was identified as CA.

Fig. 2. OPLS-DA plots of serum samples in control group and Tx group. (A) scores plot, (B) loadings S-plot.

Table 1. List of bile acid and their metabolites searched by online metabolite database.

Retention time (min)	Mass (<i>m/z</i>)	Metabolite	Mass accuracy (mDa)	Change ^a
4.79	407.2784	Cholic acid	-1.3	↑
4.84	391.2841	Chenodeoxycholic acid	-0.7	↑
4.47	464.3010	Glycocholic acid	0.1	↑
4.46	448.3052	Glycochenodeoxycholic acid	1.0	↑
4.38	583.3093	Cholic acid glucuronide	2.5	↓
4.50	471.2421	Chenodeoxycholic acid sulfate	-0.5	↓

^a Relative level in serum of thyroidectomized group in comparison to the control group

3.3. Changes in Serum BA Concentrations After Tx

Metabolomic analysis using UPLC-QTOF/MS revealed that serum BAs, and especially CA, levels markedly elevated in the Tx group compared with controls at the endpoint of the experiment. Serum concentrations of primary BAs including CA, CDCA, GCA, and GCDCA were quantified using a simultaneous analytical method [16]. Sequential changes in individual serum BA levels after Tx were also analyzed.

Serum CA, CDCA, and TCDCA levels increased rapidly more 7 days after Tx, and significantly increased to 25-fold, 11-fold, and 3-fold of the control group, respectively, 21 days after Tx (Fig. 3). CA and CDCA constituted approximately

80% of total BA content (CA: 62% and CDCA: 18%). Serum TDCA levels increased immediately after Tx, and significantly increased to 21-fold the level of controls on day 21 after Tx. THDCA levels did not change until 14 days after Tx, but significantly increased to 21-fold of controls on day 21. Serum GCA, GCDCA, and TCA levels were also increased in Tx rats (2-fold, 3-fold, and 2-fold, respectively, of controls on day 21), but these differences were not statistically significant. Serum TUDCA and β TMCA levels did not significantly differ between the control and Tx groups. Serum TLCA and THCA were not analyzed as only trace amounts were detected.

Fig. 3. Sequential changes in individual serum BA concentrations of control and Tx rats.

All data represent the mean $\pm$ S.D. of five animals. * $p < 0.05$ and ** $p < 0.01$ compared with the controls. Undefined abbreviations are: CA, cholic acid; CD, chenodeoxy; G, glycol; T, tauro; D, deoxy; H, hyo; HD, hyodeoxy; UD, ursodeoxy; M, muri.

3.4. Serum Concentrations of TH

Serum T₃ and T₄ concentrations were determined before

(day -1) and after (day 14) subcutaneous implantation of osmotic minipumps, delivering vehicle (Control and Tx), 1.5 μ g T₃/head/day (Tx+T₃) to control or Tx rats (Fig. 4). In the

all Tx groups before T_3 replacement, serum T_4 and T_3 levels significantly decreased compared with controls (T_4 : 39.0 ± 5.0 ng/mL for control, 2.4 ± 0.9 ng/mL for Tx, 2.6 ± 0.6 ng/mL for Tx+T3 and T_3 : 0.61 ± 13 ng/mL for control, 0.10 ± 0.06 ng/mL for Tx, 0.10 ± 0.05 ng/mL for Tx+T3). These levels in Tx group did not significantly change between before and

after vehicle infusion. In the Tx+T3 group after T_3 replacement, serum T_3 levels significantly increased compared with that of before replacement, and were approximately 92% of control group (0.49 ± 0.09 ng/mL for control, 0.45 ± 0.06 ng/mL for Tx+T3).

Fig. 4. Serum T_4 (A) and T_3 (B) concentrations before and after subcutaneous infusion of vehicle or T_3 in control or Tx rats.

Serum samples were collected before (day -1) and after (day 14) subcutaneous implantation of osmotic minipumps, delivering vehicle (control and Tx) or $1.5 \mu\text{g}$ T_3 /head/day (Tx+T3) to control or Tx rats. All data represent the mean $\pm$ S.D. of six animals. * $p < 0.05$ and ** $p < 0.01$ compared with the controls. # $p < 0.05$ and ## $p < 0.01$ compared with the before replacements.

3.5. Serum TSH Concentrations

Table 2. Serum TSH concentrations in control and Tx rats.

Group	Concentration (ng/mL)	
	Before ^a	After ^b
Control	1.47 ± 0.671	$3.91 \pm 1.26^{##}$
Tx	$10.8 \pm 3.26^{**}$	$26.9 \pm 10.3^{\#}$
Tx+T3	$11.8 \pm 1.03^{**}$	16.7 ± 6.50

The control and Tx groups were subcutaneous implanted osmotic minipumps, delivering vehicle. The Tx+T3 group was subcutaneous implanted osmotic minipumps, delivering T_3 ($1.5 \mu\text{g}$ T_3 /head/day).

Data are expressed as the mean values $\pm$ S.D. (n = 6).

* $p < 0.05$ and ** $p < 0.01$ compared with the controls.

[#] $p < 0.05$ and ^{##} $p < 0.01$ compared with the before replacements.

^a, the day before replacement; ^b, 14th day after replacement.

Serum TSH concentrations were determined before (day -1) and after (day 14) subcutaneous implantation of osmotic minipumps, delivering vehicle (Control and Tx), $1.5 \mu\text{g}$

T_3 /head/day (Tx+T3) to control or Tx rats (Table 2). In the all Tx groups before T_3 replacement, serum TSH levels significantly increased compared with controls. In the Tx

group after vehicle infusion, serum TSH levels significantly elevated more than those of before infusion. However, in the Tx+T3 group, serum TSH levels did not significantly change between before and after T₃ replacement. Serum TSH levels in Tx rats did not return to normal by T₃ replacement.

3.6. Effects of T₃ Replacement on Serum BA Concentrations in Tx Rats

To investigate the effects of T₃ replacement on serum BA concentrations in Tx rats, serum total BA concentrations were determined before (day -1) and after (day 14) subcutaneous implantation of osmotic minipumps, delivering vehicle (control and Tx groups) or 1.5 µg T₃/head/day (Tx+T3 group) to normal or Tx rats (Table 3). Serum total BA concentrations in the Tx and Tx+T3 groups before replacement were

significantly increased compared with those of the control group (Tx: 25.3 ± 14.8 nmol/mL, Tx+T3: 20.7 ± 10.7 nmol/mL, control: 1.55 ± 0.365 nmol/mL). However, Serum total BA concentrations in Tx rats did not change after T₃ replacement. Next, the composition ratio of each BA in serum was investigated before and after T₃ replacement in Tx rats (Fig. 5). Although the composition ratios of serum GCA and GCDCA in the Tx+T3 group were significantly increased after T₃ replacement (GCA: 1.2% vs. 7.5% and GCDCA: 0.03% vs. 0.77%, respectively), the composition ratios of most bile acids did not change markedly (CA: 67.7% vs. 61.5%, CDCA: 20.3% vs. 20.8%, TCA: 6.2% vs. 5.5%, TCDCA: 1.6% vs. 1.4%, TDCA: 1.0% vs. 0.6%, THDCA: 1.3% vs. 0.6%, TUDCA: 0.04% vs. 0.13%, and βTMCA: 0.6% vs 1.2%).

Fig. 5. Effects of T₃ replacement on serum BA composition in Tx rats.

All data represent the mean of six animals. Before, the day before replacement; After, 14th day after replacement. Undefined abbreviations are: CA, cholic acid; CD, chenodeoxy; G, glycol; T, tauro; D, deoxy; H, hyo; HD, hyodeoxy; UD, ursodeoxy; M, muri.

Table 3. Effects of triiodothyronine replacement on serum total bile acid concentrations in Tx rats.

Group	Total BA Concentration (nmol/mL)	
	Before ^a	After ^b
Control	1.55 ± 0.365	1.83 ± 0.636
Tx	25.3 ± 14.8**	32.8 ± 20.2**
Tx+T3	20.7 ± 10.7**	20.4 ± 9.27**

The control and Tx groups were subcutaneous implanted osmotic minipumps, delivering vehicle. The Tx+T3 group was subcutaneous implanted osmotic minipumps, delivering T₃ (1.5 µg T₃/head/day).

Data are expressed as the mean values ± S.D. (n = 6).

***p* < 0.01 compared with the controls.

^a, the day before replacement; ^b, 14th day after replacement.

4. Discussion

Hypothyroidism is caused by thyroid gland disorders, and leads to a decrease in production and secretion of T₄ and T₃. Tx rats are a well-established hypothyroidism model. In a previous study, we reported that Tx induced a prolongation of the half-life of T₄ in serum, and influenced T₄ metabolism and elimination [17]. We speculated that many endogenous metabolites were altered by Tx.

In the present study, we performed metabolomics analysis of serum to comprehensively analyze changes in several metabolic pathways in Tx rats and to identify corresponding biochemical perturbations. We could detect more than 4000 metabolite ions in each rat, providing a very comprehensive

serum metabolome for Tx. Using OPLS-DA followed by an online metabolite database search, metabolites that differed in Tx rats were matched, including CA, CDCA, GCA, and GCDCA, which were all increased in the serum of the Tx group. These results suggest that BAs are candidate markers for hypothyroidism.

Metabolomics is very useful for biomarker investigation, but it is a qualitative analytical method. Thus, we also measured serum concentrations of 12 BAs including CA, CDCA, GCA, and GCDCA, and investigated sequential changes in individual serum BA levels after Tx. Serum concentrations of all BAs except TLCA, THCA, TUDCA, and β TMCA, increased after Tx. There was a difference in sequential changes of each BA after Tx. BAs are synthesized from cholesterol in the liver. There are many steps, and multiple intermediates involved in the formation of primary BAs, which are CA and CDCA. These BAs were dehydroxylated by intestinal bacteria to form secondary BAs deoxycholic acid and lithocholic acid. Moreover, they undergo amidation with taurine or glycine to form conjugated BAs. Thus, we considered that the difference in sequential changes of each BA after Tx would be caused by the metabolic process of BA. Among these, serum concentrations of CA, CDCA, and TCDCA were significantly increased than 7 days after Tx. In a previous study, we have demonstrated that serum concentrations of T_4 and T_3 decrease rapidly after Tx, and are maintained at very low levels than 7 days after treatment [17]. Thus, these results suggest that the increased CA, CDCA, and TCDCA levels in serum after Tx are a homeostatic response to serum thyroid hormone deficiency.

Although BAs have been known to play an essential role in dietary lipid absorption and cholesterol metabolism, in recent years BAs have been shown to act as signaling molecules through activation of a variety of nuclear and cell surface receptors [20]. Membrane-type bile acid receptor (M-BAR)/TGR5 is a member of the G-protein coupled receptor (GPCR) superfamily, and has been found to be ubiquitously expressed including in endocrine glands, brown adipose tissue (BAT), muscles, immune organs, and the enteric nervous system in humans and animals [21]. TGR5 induces the production of intracellular cyclic adenosine monophosphate (cAMP) upon interaction with BAs [21,22]. Watanabe *et al.* reported that deiodinase type 2 (D2) is activated by increased cAMP production that stems from the binding of BAs with TGR5 in human brown adipocytes and skeletal myocytes, and then activated D2 accelerates to convert the pro-hormone T_4 to active hormone T_3 , a major component involved in cellular basal metabolism [23]. Intracellular T_4 and T_3 levels decrease after Tx [24]. Moreover, we previously reported that activity and mRNA expression of D2 in brain, BAT, and muscle were increased after Tx in rats [25]. Thus, our findings suggest that the BAs (CA, CDCA, and TCDCA) act as a signaling molecule to increase intracellular T_3 levels in local tissues by accelerating the above mechanisms (the BA–TGR5–cAMP–D2– T_3 pathway).

In hypothyroid patients, T_4 and T_3 levels in both blood and

tissues are markedly decreased but return to physiological levels with T_4 replacement therapy [26]. T_4 is considered a pro-hormone, as activation occurs through deiodination of T_4 to form T_3 , the active form of TH. Thus, T_3 replacement therapy in hypothyroid patients may be more effective than T_4 replacement. In this study, we investigated the effects of T_3 replacement on serum BA concentrations in hypothyroid rats. Serum T_3 levels in Tx rats were returned to physiological levels with T_3 replacement. However, serum TSH levels were maintained at very high levels. These results were consistent with a previously study by Escobar-Morreale *et al.* [27]. They also reported that T_3 levels in many tissues of Tx rats were distinctly lower than the controls even if serum T_3 levels were returned to physiological levels with T_3 replacement. Serum concentrations of total BA in Tx rats after T_3 replacement showed little change compared with before replacement, and the composition ratio of BAs also showed no significant difference before and after T_3 replacement. Intracellular T_3 levels are known to be maintained by direct T_3 uptake from blood and by intracellular selective conversion of T_4 to T_3 by D2 [28,29]. Intracellular T_3 concentrations in BAT and brain are supplied at a ratio of 50:50 and 20:80 from these two pathways, respectively [30,31]. Moreover, we have previously reported that activity and mRNA expression of D2 in local tissues of Tx rats were maintained at high levels after T_3 replacement [25]. These results indicate that serum BA levels were maintained at a high level because intracellular TH concentrations in Tx rats were not normalized to physiological levels by T_3 replacement.

TH replacement therapy in hypothyroid patients is estimated by serum levels of THs and TSH. In general, TH in the blood is quickly taken up into cells in target tissues via specific plasma membrane transporters, members of the organic anion transporting polypeptide family [32], and the monocarboxylate transporter family [33]. However, in humans and rodents who exhibit inactivating mutations in the genes encoding these transporters, THs in the blood cannot be taken up into cells even after TH replacement, and biological function is not fully restored [34–36]. Thus, we suggest that TH levels in local tissues are a more important indicator of the efficacy of hypothyroidism therapy than TH levels in blood. However, it is unfeasible to measure TH concentrations in human patients' organ. The results of study indicate that serum BA levels are negatively correlated with intracellular TH concentrations in target tissues. Thus, serum CA, CDCA, and TCDCA may be potential biomarkers for hypothyroidism and the efficacy of TH replacement therapy in hypothyroid patients. To identify this hypothesis, further studies are needed to investigate the mechanism by which Tx increases serum BA levels in rats, and to determine if the same results can be seen in human subjects.

5. Conclusion

Serum BA levels were shown to increase markedly in rats after Tx, as indicated by metabolomics analysis. Moreover, we investigated sequential changes in serum levels of 12 BAs

in rats after Tx. We observed an increase in serum CA, CDCA, and TCDCA after Tx, and believe that this is a homeostatic response to intracellular TH deficiency in local tissues. To the best of our knowledge, this is the first report indicating that serum CA, CDCA, and TCDCA may be potential biomarkers for hypothyroidism and the efficacy of TH replacement therapy in hypothyroidism.

Acknowledgements

We thank to Ms. Etsuko Suzuki and Mr. Kazunobu Maruyama (Waters) for technical assistance.

References

- [1] T.J. Visser, "Pathways of thyroid hormone metabolism," *Acta Med. Austriaca*, vol. 23, pp. 10-16, 1996.
- [2] S.Y. Cheng, "Multiple mechanisms for regulation of the transcriptional activity of thyroid hormone receptors," *Rev. Endocr. Metab. Disord.*, vol. 1, pp. 9-18, 2000.
- [3] G.R. Williams, "Cloning and characterization of two novel thyroid hormone receptor β isoforms," *Mol. Cell. Biol.*, vol. 20, pp. 8329-8342, 2000.
- [4] H. Ying, H. Suzuki, L. Zhao, M.C. Willingham, P. Meltzer and S.Y. Cheng, "Mutant thyroid hormone receptor β represses the expression and transcriptional activity of peroxisome proliferator-activated receptor γ during thyroid carcinogenesis," *Cancer Res.*, vol. 63, pp. 5274-5280, 2003.
- [5] I. Jones, L. Ng, H. Liu and D. Forrest, "An intron control region differentially regulates expression of thyroid hormone receptor $\beta 2$ in the cochlea, pituitary, and cone photoreceptors," *Mol. Endocrinol.*, vol. 21, pp. 1108-1119, 2007.
- [6] A.R. Cappola and P.W. Ladenson, "Hypothyroidism and atherosclerosis," *J. Clin. Endocrinol. Metab.*, vol. 88, pp. 2438-2444, 2003.
- [7] P. Cettour-Rose, C. Theander-Carrillo, C. Asensio, M. Klein, T.J. Visser, A.G. Burger, C.A. Meier and F. Rohner-Jeanrenaud, "Hypothyroidism in rats decreases peripheral glucose utilisation, a defect partially corrected by central leptin infusion," *Diabetologia*, vol. 48, pp. 624-633, 2005.
- [8] A. Zhang, H. Sun, G. Yan, Y. Han, Y. Ye and X. Wang, "Urinary metabolic profiling identifies a key role for glycocholic acid in human liver cancer by ultra-performance liquid-chromatography coupled with high-definition mass spectrometry," *Clin. Chim. Acta*, vol. 418, pp. 86-90, 2013.
- [9] N. Aranibar, M. Borys, N.A. Mackin, V. Ly, N. Abu-Absi, S. Abu-Absi, M. Niemitz, B. Schilling, Z.J. Li, B. Brock, R.J. Russell 2nd, A. Tymiak and M.D. Reily, "NMR-based metabolomics of mammalian cell and tissue cultures," *J. Biomol. NMR*, vol. 49, pp. 195-206, 2011.
- [10] A. Lodi and S.M. Ronen, "Magnetic resonance spectroscopy detectable metabolomics fingerprint of response to antineoplastic treatment," *PLoS ONE*, 6:e26155, 2011.
- [11] L.M. Raamsdonk, B. Teusink, D. Broadhurst, N.S. Zhang, A. Hayes, M.C. Walsh, J.A. Berden, K.M. Brindle, D.B. Kell, J.J. Rowland, H.V. Westerhoff, K. van Dam and S.G. Oliver, "A functional genomics strategy that uses metabolome data to reveal the phenotype of silent mutations," *Nat. Biotechnol.*, vol. 19, pp. 45-50, 2001.
- [12] O. Fiehn, J. Kopka, P. Dormann, T. Altmann, R.N. Trethewey and L. Willmitzer, "Metabolite profiling for plant functional genomics," *Nat. Biotechnol.*, vol. 18, pp. 1157-1161, 2000.
- [13] L.D. Roberts, A. Koulman and J.L. Griffin, "Towards metabolic biomarkers of insulin resistance and type 2 diabetes: progress from the metabolome," *Lancet Diabetes Endocrinol.*, vol. 2, pp. 65-75, 2014.
- [14] T. Soga, M. Sugimoto, M. Honma, M. Mori, K. Igarashi, K. Kashikura, S. Ikeda, A. Hirayama, T. Yamamoto, H. Yoshida, M. Otsuka, S. Tsuji, Y. Yatomi, T. Sakuragawa, H. Watanabe, K. Nihei, T. Saito, S. Kawata, H. Suzuki, M. Tomita and M. Suematsu, "Serum metabolomics reveals γ -glutamyl dipeptides as biomarkers for discrimination among different forms of liver disease," *J. Hepatol.*, vol. 55, pp. 896-905, 2011.
- [15] T.H. Kim, M.Y. Ahn, H.J. Lim, Y.J. Lee, Y.J. Shin, U. De, J. Lee, B.M. Lee, S. Kim and H.S. Kim, "Evaluation of metabolomic profiling against renal toxicity in Sprague-Dawley rats treated with melamine and cyanuric acid," *Arch. Toxicol.*, vol. 86, pp. 1885-1897, 2012.
- [16] E.J. Want, M. Coen, P. Masson, H.C. Keun, J.T. Pearce, M.D. Reily, D.G. Robertson, C.M. Rohde, E. Holmes, J.C. Lindon, R.S. Plumb and J.K. Nicholson, "Ultra performance liquid chromatography-mass spectrometry profiling of bile acid metabolites in biofluids: application to experimental toxicology studies," *Anal. Chem.*, vol. 82, pp. 5282-5289, 2010.
- [17] H. Nagao, T. Imazu, H. Hayashi, K. Takahashi and K. Minato, "Influence of thyroidectomy on thyroxine metabolism and turnover in rats," *J. Endocrinol.*, vol. 210, pp. 117-123, 2011.
- [18] J.T. Herlihy, C. Stacy and H.A. Bertrand, "Long-term food restriction depresses serum thyroid hormone concentrations in the rat," *Mech. Ageing Dev.*, vol. 53, pp. 9-16, 1990.
- [19] T.T. Nguyen, J.J. Di Stefano, H. Yamada and Y.M. Yen, "Steady state organ distribution and metabolism of thyroxine and 3,5,3'-triiodothyronine in intestine, liver, kidneys, blood and residual carcass of the rat in vivo," *Endocrinology*, vol. 133, pp. 2973-2983, 1993.
- [20] S. Fiorucci, S. Cipriani, F. Baldelli and A. Mencarelli, "Bile acid-activated receptors in the treatment of dyslipidemia and related disorders," *Prog. Lipid Res.*, vol. 49, pp. 171-185, 2010.
- [21] Y. Kawamata, R. Fujii, M. Hosoya, M. Harada, H. Yoshida, M. Miwa, S. Fukusumi, Y. Habata, T. Itoh, Y. Shintani, S. Hinuma, Y. Fujisawa and M. Fujino, "A G protein-coupled receptor responsive to bile acids," *J. Biol. Chem.*, vol. 278, pp. 9435-9440, 2003.
- [22] T. Maruyama, Y. Miyamoto, T. Nakamura, Y. Tamai, H. Okada, E. Sugiyama, T. Nakamura, H. Itadani and K. Tanaka, "Identification of membrane-type receptor for bile acids (M-BAR)," *Biochem. Biophys. Res. Commun.*, vol. 298, pp. 714-719, 2002.
- [23] M. Watanabe, S.M. Houten, C. Matak, M.A. Christoffolete, B.W. Kim, H. Sato, N. Messaddeq, J.W. Harney, O. Ezaki, T. Kodama, K. Schoonjans, A.C. Bianco and J. Auwerx, "Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation," *Nature*, vol. 439, pp. 484-489, 2006.

- [24] H.F. Escobar-Morreale, M.J. Obregon, F. Escobar Del Rey and G. Morreale De Escobar, "Replacement therapy for hypothyroidism with thyroxine alone does not ensure euthyroidism in all tissues, as studied in thyroidectomized rats," *J. Clin. Invest.*, vol. 96, pp. 2828-2838, 1995.
- [25] H. Nagao, M. Sasaki, T. Imazu, K. Takahashi, H. Aoki and K. Minato, "Effects of triiodothyronine on turnover rate and metabolizing enzymes for thyroxine in thyroidectomized rats," *Life Sci.*, vol. 116, pp. 74-82, 2014.
- [26] A.D. Toft, "Thyroxine therapy," *N. Engl. J. Med.*, vol. 331, pp. 174-180, 1994.
- [27] H.F. Escobar-Morreale, M.J. Obregón, F. Escobar del Rey and G. Morreale de Escobar, "Tissue-specific patterns of changes in 3,5,3'-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?," *Biochimie*, vol. 81, pp. 453-462, 1999.
- [28] G.R. Williams and J.H. Bassett, "Deiodinases: the balance of thyroid hormone: local control of thyroid hormone action: role of type 2 deiodinase," *J. Endocrinol.*, vol. 209, pp. 261-272, 2011.
- [29] A.C. Bianco, D. Salvatore, B. Gereben, M.J. Berry and P.R. Larsen, "Biochemistry, cellular and molecular biology, and physiological roles of the iodothyronine selenodeiodinases," *Endocr. Rev.*, vol. 23, pp. 38-89, 2002.
- [30] A.C. Bianco and J.E. Silva, "Nuclear 3,5,3'-triiodothyronine (T3) in brown adipose tissue: receptor occupancy and sources of T3 as determined by in vivo techniques," *Endocrinology*, vol. 120, pp. 55-62, 1987.
- [31] J. van Doorn, F. Roelfsema and D. van der Heide, "Conversion of thyroxine to 3,5,3'-triiodothyronine in several rat tissues in vivo: the effect of hypothyroidism," *Acta Endocrinol.*, vol. 113, pp. 59-64, 1986.
- [32] B. Hagenbuch, "Cellular entry of thyroid hormones by organic anion transporting polypeptides," *Best Pract. Res. Clin. Endocrinol. Metab.*, vol. 21, pp. 209-221, 2007.
- [33] W.E. Visser, E.C. Friesema, J. Jansen and T.J. Visser, "Thyroid hormone transport by monocarboxylate transporters," *Best Pract. Res. Clin. Endocrinol. Metab.*, vol. 21, pp. 223-236, 2007.
- [34] E.C. Friesema, A. Grueters, H. Biebermann, H. Krude, A. von Moers, M. Reeser, T.G. Barrett, E.E. Mancilla, J. Svensson, M.H. Kester, G.G. Kuiper, S. Balkassmi, A.G. Uitterlinden, J. Koehle, P. Rodien, A.P. Halestrap and T.J. Visser, "Association between mutations in a thyroid hormone transporter and severe X-linked psychomotor retardation," *Lancet*, vol. 364, pp. 1435-1437, 2004.
- [35] L.M. Roberts, K. Woodford, M. Zhou, D.S. Black, J.E. Haggerty, E.H. Tate, K.K. Grindstaff, W. Mengesha, C. Raman and N. Zerangue, "Expression of the thyroid hormone transporters monocarboxylate transporter-8 (SLC16A2) and organic ion transporter-14 (SLCO1C1) at the blood-brain barrier," *Endocrinology*, vol. 149, pp. 6251-6261, 2008.
- [36] S. Mayerl, T.J. Visser, V.M. Darras, S. Horn and H. Heuer, "Impact of Oatp1c1 deficiency on thyroid hormone metabolism and action in the mouse brain," *Endocrinology*, vol. 153, pp. 1528-1537, 2012.