

Heavy Metals Status in Some Commercially Important Fishes of Meghna River Adjacent to Narsingdi District, Bangladesh: Health Risk Assessment

Md. Simul Bhuyan^{1,*}, Muhammad Abu Bakar², Aysha Akhtar¹, Md. Shafiqul Islam¹

¹Institute of Marine Sciences and Fisheries, University of Chittagong, Chittagong, Bangladesh

²Bangladesh Council of Scientific and Industrial Research, Chittagong, Bangladesh

Email address:

simulbhuyan@gmail.com (Md. S. Bhuyan)

*Corresponding author

To cite this article:

Md. Simul Bhuyan, Muhammad Abu Bakar, Md. Shafiqul Islam, Aysha Akhtar. Heavy Metals Status in Some Commercially Important Fishes of Meghna River Adjacent to Narsingdi District, Bangladesh: Health Risk Assessment. *American Journal of Life Sciences*. Vol. 4, No. 2, 2016, pp. 60-70. doi: 10.11648/j.ajls.20160402.17

Received: April 27, 2016; Accepted: May 6, 2016; Published: May 30, 2016

Abstract: The present study was carried out to determine the heavy metals (Cd, Pb, Zn, Al, Cu, Ni, Fe, Mn, Cr, Co) in the muscles of 32 fish species for 3 seasons in Meghna River. The heavy metals were analyzed by Atomic Absorption Spectrophotometer. The estimated concentrations of all metals in the present study were lower than the limits permitted by FAO, WHO, EU, United States Food and Drug Administration (USFDA), US Environment Protection Agency (US/EPA) and England Guidelines except the concentrations of Pb & Zn that were found above the allowable ranges in different fishes namely *Amblypharyngodon mola*, *Colisa lalia*, *Tetraodon cutcutia*, *Barbodes sarana*, *Labeo calbasu*, *Puntius sarana*, *Ompok pabda*, *Aila coila*, *Mastacembelus armatus*, *Glossogobius giuris*, *Nandus nandu*, *Tenualosa ilisha*, *Lepidocephalichthys guntea*, *Xenentodon cancila*, *Stinging catfish*. Multivariate statistical analysis such as principal component analysis and correlation matrix showed significant anthropogenic intrusions of Zn, Al, Cd, Pb, Cu, Ni, Fe, Mn, Cr, Co in fishes. There was significant positive correlation between Cd vs Co (0.733), moderate positive correlation between Fe vs Al (0.568), Ni vs Co (0.482), Mn vs Co (0.395) which indicate that their common origin especially from industrial effluents, municipal wastes and agricultural inputs.

Keywords: Heavy Metal, Multivariate Analysis, Fish, Atomic Absorption Spectrophotometer, Meghna River

1. Introduction

Industrialization is a term associated with socio-economic activities (Richard 2005; Jaillon and Poon 2009, Thanoon et al. 2003) that alter the society infrastructure (Abdullah et al. 2009) through the huge production (Abdullah et al. 2009; Thanoon et al. 2003). Most of these industries discharge untreated wastes that contribute heavy metal hugely. Industrial and agricultural as well as natural activities are the leading responsible sources of metal contamination in aquatic environment (Wilson and Pyatt 2007; Khan et al. 2008; Tarra-Wahlberg et al. 2001; Akif et al. 2002; Jordao et al. 2002; Sekabira et al. 2010; Zhang et al. 2011; Bai et al. 2011; Grigoratos et al. 2014; Martin et al. 2015). Heavy metal pollution is a great concern because of their long persistence,

bioaccumulation and biomagnifications in the food chain (Rahman et al. 2013; Sharma et al. 2007; Sankar et al. 2006; Papagiannis et al. 2004; Zhou et al. 2004; Sun et al. 2001; Zhou 1995) ultimately poses toxicity both in human and aquatic animals (Islam et al. 2015a; Ahmed et al. 2015a, b; Fang et al. 2014; Alhashemi et al. 2012; Pan and Wang 2012; Yi et al. 2011; Vieira et al. 2011; Forti et al. 2011; Banerjee et al. 2011; Tuzen 2009; Sanchez-Chardi et al. 2007; McCluggage 1991; WHO 1995). Most of the literature conducted on bioaccumulation and toxicity of heavy metal (Rainbow et al. 2000; Shuhaimi-Othmana and Pascoe 2007) as the heavy metal pollution is global concern (Islam et al. 2014). This increasing pattern of heavy metals have adverse health effects for invertebrates, fish, and humans (Yi et al. 2011; Islam et al. 2014; Martin et al. 2015; Islam et al.

2015b,d; Ahmed et al. 2015c). Recently, in many developing countries like Bangladesh are continuously being contaminated with heavy metal (Islam et al. 2015c) due to disposal of untreated effluents from different industries, domestic wastes and agrochemicals in the open water bodies and rivers deteriorating water quality (Khadse et al. 2008; Venugopal et al. 2009; Islam et al. 2015a,c). Heavy metal in aquatic ecosystems measured by monitoring concentrations in water, sediments and biota (Camusso et al. 1995) but considerable amount was found in fishes (Rashed 2001) and sediments (Namminga and Wilhm 1976). Several studies have been conducted in rivers and lakes giving special preference to environment during the last decade (Özmen et al. 2004; Begüm et al. 2005; Fernandes et al. 2008; Öztürk et al. 2008; Pote et al. 2008 and Praveena et al. 2008). Meghna River near Narsingdi is highly polluted with industrial and domestic sewage that contribute huge amount of Zn, Al, Cd, Pb, Cu, Ni, Fe, Mn, Cr, Co. But unfortunately, there was no

scientific research on heavy metal pollution in the concerned area was found so far which compelled us to conduct this research with aim (1) to determine the heavy metal concentrations in fish and (2) to compare the concentrations of heavy metal in different seasons.

2. Materials and Methods

2.1. Sampling Sites

Fish samples were collected from two points (Effluent discharge area and far from the discharge area) of Meghna River near Narsingdi District (23°55'28.52"N and 90°45'12.06"E). Sampling procedures were performed in three phases: firstly, September, 2015 (Rainy season); secondly, January, 2016 (Winter season) and thirdly, March, 2016 (Pre-monsoon).




Figure 1. Map showing sampling station of Meghna River.

2.2. Sample Collection and Preservation

A total of 32 fish species were collected from fishermen for individual season and then identified according to Rahman et al. (2009), Roy et al. (2007), Quddus et al. (1988), Quddus and Shafi (1983). After collection the fish were placed immediately in poly-ethylene bags and then kept into isolated container of polystyrene icebox. Finally, the samples were transferred to the Bangladesh Council of Scientific and Industrial Research (BCSIR) in ice box (Irwandi and Farida 2009; Ismail and Saleh 2012) where the fish were first washed with deionized water sealed in poly-ethylene bags subsequently kept in a freezer at -20°C until analysis (Elnabris et al. 2012).

2.3. Heavy Metal Determination

The heavy metal contents were determined by AAS using standard analytical procedure. Sample collection is important stage for metal analysis. Samples were generally carefully handled to avoid contamination. Glassware was properly cleaned, and the reagents were of analytical grade. Distilled water was used throughout the study. Reagents blank determinations were used to correct the instrument readings. The techniques for samples preparation, standard Preparation, analysis for metal analyses have been briefly described below.

2.4. Sample Preparation (Dry Ashing Method)

This procedure was also used for destruction of organic

matter. Precaution was to be taken to avoid losses by volatilization of elements. At first samples were homogenized. Then the samples were weighed accurately a suitable quantity (10 to 20 g) of the homogenized samples in a tared silica dish. After that the samples were dried at 100°C in a laboratory oven. These dishes were then placed in the muffle furnace at ambient temperature and slowly raised temperature to 450°C at a rate of no more than 50°C/h. The samples were ignited in a Muffle furnace at 450°C for at least 8 hrs. After ashing was completed and cool, then the dishes were removed from furnace. Then the ashes were dissolved in diluted nitric acid (Aftan *et al.* 2000). The solutions were returned to a hot plate and continued heating, adding additional acid as necessary until digestion was completed. Then the samples were filtrated into a 100 ml volumetric flask using Whatman No. 44 filter paper and washed the residue. Each sample solution was made up to the mark with distilled water.

2.5. Standard Preparation

Every metal standard solution was prepared for calibration the instrument for each element being determined on the same day as the analyses were performed due to possible deterioration of standard with time. All samples were prepared from chemicals of analytical grade with distilled water. 1 gm of metal Cadmium, Copper, Lead, Nickel were dissolved in HNO₃ solution; 1 g of Cobalt, Iron, Manganese, Zinc, Aluminum were dissolved in HCl solution; 2.8289 g K₂Cr₂O₇ (=1g Chromium) was dissolved in water and made up to 1 liter in volumetric flask with distilled water, thus stock solution of 1000 mg/l of Cd, Cu, Pb, Ni, Co, Fe, Mn, Zn, Al and Cr were prepared. (Cantle, J. E. 1982). Then 100 ml of 0.1, 0.25, 0.5, 0.75, 1.0 and 2.0 mg/l of working standards of each metal except iron were prepared from these stock using micropipettes in 5ml of 2N nitric acid. 100 ml of 2.0, 2.5, 5.0, 10.0 and 20.0 mg/l of working standards of iron metal were prepared from iron stock solution. Reagent blank was prepared in the same manners of sample preparation without sample to avoid reagents contamination.

2.6. Analysis of Sample

Finally, the atomic absorption instrument was set up carefully. At the meantime, flame condition and absorbance were optimized for the analyte. Then blanks (deionized water), standards, sample blank and samples were aspirated into the flame in AAS (Model- iCE 3300, Thermo Scientific, Designed in UK, Made by China)). The calibration curves were found for concentration vs. absorbance. Data were statistically analyzed using fitting of straight line by least square method. For more accuracy, a blank reading was also taken and necessary corrections were made during the calculation of various elements concentration.

2.7. Statistical Analysis

One Way Analysis of Variance (ANOVA) was done to

show the variations in concentration of heavy metal in terms of seasons and fish. GGraph was used for graphical presentation of heavy metal against seasons. According to Dreher (2003), Principal Component Analysis (PCA) was performed on the original data set (without any weighting or standardization). Pearson's product moment correlation matrix was done to identify the relation among metals to make the result validate obtained from multivariate analysis.

3. Results

A total 32 fish species were analyzed for heavy metals (Zn, Al, Cd, Pb, Cu, Ni, Fe, Mn, Cr, Co) detection. In rainy season, the highest concentration of Zn was recorded in *Amblypharyngodon mola* (42.45mg/kg) and the lowest was found in *Ctenopharyngodon idella* (10.27mg/kg) (Table 1). The maximum concentration of Al recorded in *Gudusiachapra* (117.55mg/kg) and minimum was found in *Ctenopharyngodon idella* (1.85mg/kg) (Table 1). The highest concentration of Mn was recorded in *Tetraodon cutcutia* (19.07mg/kg) and lowest was found in *Ctenopharyngodon idella* (0.96mg/kg) (Table 1). The highest concentration of Cd in *Ompok pabda* (0.12mg/kg); Pb in *Colisa lalia* (5.87); Cu in *Barbodes sarana* (32.44mg/kg); Ni in *Amblypharyngodon mola* (0.76mg/kg); Cr in *Anabus testudineus* (1.75mg/kg) and Co in *Mystus bleekeri* (0.43mg/kg) whereas the lowest concentration for these metals were below detection limit (Table 1).

During winter season, the highest value of Zn was recorded in *Amblypharyngodon mola* (44.48mg/kg) and the lowest value was found in *Johnius coitor* (9.6mg/kg) (Table 2). The highest concentration of Al was detected in *Labeo rohita* (106.70mg/kg) and the lowest value was documented in *Nandus nandus* (3.24mg/kg). The maximum value of Cu was measured in *Glossogobius giuris* (8.19mg/kg) and the minimum value was recorded in *Macragnathus aculeatus* (0.20mg/kg) (Table 2). The maximum amount of Fe was detected in *Mystus bleekeri* (93.16mg/kg) and the lowest amount was found in *Ctenopharyngodon idella* (9.86mg/kg). The highest concentration of Mn was found in *Colisa lalia* (19.87mg/kg) and the lowest concentration was measured in *Mastacembelus armatus* (1.65mg/kg). Moreover, the maximum amount was recorded for Cd in *Ompok pabda* (0.21mg/kg); Pb in *Colisa lalia* (6.75mg/kg); Ni in *Amblypharyngodon mola* (0.986mg/kg); Cr in *Stinging catfish* (3.01mg/kg) and Co in *Labeo rohita* (0.70mg/kg) while the lowest concentration of these metals were recorded below detection limit (Table 2).

In Pre-monsoon, the highest value of Zn was recorded in *Amblypharyngodon mola* (43.67mg/kg) and the lowest value was documented in *Johnius coitor* (8.65mg/kg) (Table 3). The greatest amount of Al was measured in *Labeo rohita* (104.87mg/kg) and the lowest amount was found in *Ctenopharyngodon idella* (1.78mg/kg). The supreme concentration of Cu was detected in *Barbodes sarana* (26.67mg/kg) and the lowest concentration was measured in *Macragnathus aculeatus* (0.21mg/kg). The maximum value of Fe was recorded in *Gudusiachapra* (77.66mg/kg) and the

minimum value was recorded in *Xenentodon cancila* (7.85mg/kg). The greatest amount of Mn was measured in *Tetraodon cutcutia* (20.01mg/kg) and the lowest value was found in *Ctenopharyngodon idella* (1.01mg/kg). The highest concentration was recorded for Cd in *Ompok pabda* (0.23mg/kg); Pb in *Colisa lalia* (6.85mg/kg); Ni in *Amblypharyngodon mola* (0.98mg/kg); Cr in *Aila coila* (8.18mg/kg) and Co in *Channa punctatus* (0.51mg/kg) whereas the lowest concentration for these metals was documented below detection limit (Table 3).

3.1. Analysis of Variance (ANOVA) in Fish Species

There were significant variations in the concentrations of Zinc, aluminium, lead, copper, nickel, iron, cobalt, manganese found across the various fish species as the significance level ($p < 0.05$). However, there were no significant variations ($p > 0.05$) in cadmium and Chromium levels across the fish species. Moreover there was no prevalent variation ($p > 0.05$) in the metal concentration in terms of seasons except for lead and manganese ($p < 0.05$).

Table 1. Heavy metal concentrations (mg/kg) in fish during Rainy season.

Sl. No.	Scientific Name	Sample ID	Zn (mg/kg)	Al (mg/kg)	Cd (mg/kg)	Pb (mg/kg)	Cu (mg/kg)	Ni (mg/kg)	Fe (mg/kg)	Mn (mg/kg)	Cr (mg/kg)	Co (mg/kg)
01	<i>Aila coila</i>	FR-01	23.85	9.70	BDL	0.04	0.19	0.35	18.62	3.45	BDL	BDL
02	<i>Aorichthys aor</i>	FR-02	19.54	6.20	0.027	BDL	0.61	0.192	18.61	2.02	BDL	0.24
03	<i>Mastacembelus armatus</i>	FR-03	15.38	2.10	BDL	BDL	1.41	BDL	24.34	1.76	BDL	BDL
04	<i>Labeo bata</i>	FR-04	15.67	12.99	BDL	BDL	0.93	0.08	19.62	3.56	BDL	BDL
05	<i>Glossogobius giuris</i>	FR-05	18.54	5.32	BDL	BDL	7.19	0.18	13.07	2.98	BDL	BDL
06	<i>Nandus nandus</i>	FR-06	19.70	3.51	0.015	BDL	1.02	0.138	17.23	4.8	BDL	0.27
07	<i>Colisa lalia</i>	FR-07	32.88	31.34	BDL	5.87	3.00	BDL	37.67	18.99	BDL	BDL
08	<i>Gudusiachapra</i>	FR-08	17.7	117.55	BDL	BDL	1.01	0.22	76.21	14.21	BDL	BDL
09	<i>Chitala chitala</i>	FR-09	15.40	13.85	BDL	0.035	0.72	BDL	45.34	2.30	0.19	BDL
10	<i>Colisa chuna</i>	FR-10	14.52	12.55	BDL	0.09	1.40	BDL	35.76	3.75	0.78	BDL
11	<i>Barbodes sarana</i>	FR-11	24.41	45.52	0.099	0.47	32.44	0.193	39.97	4.46	BDL	0.28
12	<i>Tenualosa ilisha</i>	FR-12	11.31	3.86	0.092	0.67	1.21	0.084	27.54	2.39	0.05	BDL
13	<i>Notopterus notopterus</i>	FR-13	18.62	13.21	BDL	BDL	0.03	0.01	19.23	3.67	BDL	BDL
14	<i>Tetraodon cutcutia</i>	FR-14	34.21	33.01	BDL	BDL	0.99	0.31	39.91	19.07	BDL	BDL
15	<i>Ctenopharyngodon idella</i>	FR-15	10.27	1.85	BDL	BDL	0.69	0.064	9.11	0.96	0.11	BDL
16	<i>Labeo calbasu</i>	FR-16	29.21	42.66	BDL	1.71	1.28	BDL	18.21	4.11	0.99	BDL
17	<i>Mystus cavasius</i>	FR-17	15.82	27.41	BDL	BDL	0.84	0.272	35.74	1.50	BDL	BDL
18	<i>Lepidocephalichthys guntea</i>	FR-18	24.56	7.88	BDL	3.21	2.30	0.29	14.87	3.77	BDL	0.03
19	<i>Puntius sarana</i>	FR-19	29.65	44.28	0.035	BDL	0.71	0.210	66.71	9.44	BDL	BDL
20	<i>Xenentodon cancila</i>	FR-20	20.13	5.45	BDL	0.38	1.21	0.187	18.71	2.56	BDL	0.27
21	<i>Clupisoma garua</i>	FR-21	23.01	21.44	BDL	BDL	0.38	0.12	42.42	10.10	BDL	BDL
22	<i>Anabus testudineus</i>	FR-22	26.44	17.72	BDL	BDL	1.54	0.185	65.43	7.10	1.75	BDL
23	<i>Amblypharyngodon mola</i>	FR-23	42.45	31.11	BDL	BDL	2.05	0.76	28.04	5.43	BDL	BDL
24	<i>Ompok pabda</i>	FR-24	39.50	4.5	0.121	BDL	0.64	0.362	11.16	1.92	BDL	0.41
25	<i>Johnius coitor</i>	FR-25	17.34	9.32	BDL	BDL	0.50	0.10	31.36	3.10	0.21	0.30
26	<i>Labeo rohita</i>	FR-26	28.40	103.13	0.013	BDL	1.98	0.150	64.56	9.76	0.21	0.37
27	<i>Stinging catfish</i>	FR-27	12.83	15.37	0.063	1.35	1.02	BDL	93.83	2.54	0.19	0.24
28	<i>Channa striatus</i>	FR-28	16.42	2.45	BDL	BDL	0.61	0.084	39.21	9.10	0.08	0.15
29	<i>Channa punctatus</i>	FR-29	15.59	6.10	0.011	0.040	1.27	0.12	147.77	9.55	BDL	0.29
30	<i>Macrognathus aculeatus</i>	FR-30	20.32	4.89	BDL	BDL	0.10	0.11	18.32	2.67	BDL	BDL
31	<i>Mystus bleekeri</i>	FR-31	28.15	10.90	0.039	BDL	1.41	BDL	78.35	2.77	BDL	0.43
32	<i>Silonia silondia</i>	FR-32	21.92	3.53	BDL	BDL	0.48	0.178	28.13	1.76	BDL	BDL

Table 2. Heavy metal concentrations (mg/kg) in fish during Winter season.

Sl. No.	Scientific Name	Sample ID	Zn (mg/kg)	Al (mg/kg)	Cd (mg/kg)	Pb (mg/kg)	Cu (mg/kg)	Ni (mg/kg)	Fe (mg/kg)	Mn (mg/kg)	Cr (mg/kg)	Co (mg/kg)
01	<i>Aila coila</i>	FW-01	25.77	10.69	BDL	0.15	0.24	0.456	23.63	3.51	BDL	BDL
02	<i>Aorichthys aor</i>	FW-02	14.96	43.19	BDL	BDL	0.40	0.235	54.77	2.40	0.20	BDL
03	<i>Mastacembelus armatus</i>	FW-03	14.99	6.15	BDL	BDL	1.98	0.141	20.43	1.65	BDL	BDL
04	<i>Labeo bata</i>	FW-04	16.75	14.35	BDL	BDL	1.10	0.174	21.92	4.34	BDL	BDL
05	<i>Glossogobius giuris</i>	FW-05	19.63	5.52	BDL	BDL	8.19	0.179	15.05	3.22	BDL	BDL
06	<i>Nandus nandus</i>	FW-06	17.85	3.24	BDL	BDL	0.91	0.112	19.12	10.41	BDL	BDL
07	<i>Colisa lalia</i>	FW-07	36.2	32.54	BDL	6.75	3.98	BDL	37.6	19.87	BDL	BDL
08	<i>Gudusiachapra</i>	FW-08	18.73	120.4	BDL	BDL	1.21	0.297	78.62	14.34	BDL	BDL
09	<i>Chitala chitala</i>	FW-09	11.47	5.90	BDL	BDL	0.45	0.093	13.50	2.65	BDL	BDL
10	<i>Colisa chuna</i>	FW-10	15.99	13.79	BDL	0.11	2.11	BDL	38.67	4.87	1.6	BDL
11	<i>Barbodes sarana</i>	FW-11	16.10	10.53	0.030	BDL	6.52	0.129	14.10	3.54	BDL	BDL
12	<i>Tenualosa ilisha</i>	FW-12	11.31	3.86	0.092	0.67	1.21	0.084	27.54	2.39	0.05	BDL
13	<i>Notopterus notopterus</i>	FW-13	19.47	13.50	BDL	BDL	0.33	0.130	23.07	4.91	BDL	BDL
14	<i>Tetraodon cutcutia</i>	FW-14	35.95	34.51	BDL	BDL	1.47	0.364	42.16	19.58	BDL	BDL
15	<i>Ctenopharyngodon idella</i>	FW-15	11.21	1.98	BDL	BDL	0.87	0.12	9.86	1.21	0.21	BDL

Sl. No.	Scientific Name	Sample ID	Zn (mg/kg)	Al (mg/kg)	Cd (mg/kg)	Pb (mg/kg)	Cu (mg/kg)	Ni (mg/kg)	Fe (mg/kg)	Mn (mg/kg)	Cr (mg/kg)	Co (mg/kg)
16	<i>Labeo calbasu</i>	FW-16	31.21	43.87	BDL	1.91	1.76	BDL	20.01	4.79	1.12	BDL
17	<i>Mystus cavasius</i>	FW-17	17.21	27.69	BDL	BDL	1.02	0.31	36.74	1.81	BDL	BDL
18	<i>Lepidocephalichthys guntea</i>	FW-18	25.99	9.01	BDL	4.21	2.45	0.41	15.89	4.65	BDL	0.067
19	<i>Puntius sarena</i>	FW-19	39.65	12.28	BDL	BDL	1.23	0.596	33.96	15.99	BDL	BDL
20	<i>Xenentodon cancila</i>	FW-20	19.05	6.92	BDL	BDL	0.39	0.307	13.94	3.40	BDL	BDL
21	<i>Clupisoma garua</i>	FW-21	24.12	22.24	BDL	BDL	0.52	0.227	44.40	11.43	BDL	BDL
22	<i>Anabus testudineus</i>	FW-22	15.61	3.32	BDL	BDL	0.82	0.092	27.10	6.97	0.19	BDL
23	<i>Amblypharyngodon mola</i>	FW-23	44.48	32.0	BDL	BDL	2.26	0.986	30.05	5.74	BDL	BDL
24	<i>Ompok pabda</i>	FW-24	39.98	5.01	0.21	BDL	0.69	0.38	12.01	2.05	BDL	0.51
25	<i>Johnius coitor</i>	FW-25	9.60	29.10	BDL	BDL	0.34	0.157	23.92	2.85	0.057	BDL
26	<i>Labeo rohita</i>	FW-26	29.40	106.70	0.04	BDL	2.01	0.17	66.98	11.30	0.57	0.70
27	<i>Stinging catfish</i>	FW-27	13.01	11.42	BDL	1.56	1.76	0.45	25.94	3.89	3.01	BDL
28	<i>Channa striatus</i>	FW-28	16.67	3.42	BDL	BDL	1.01	0.10	40.01	10.21	0.10	0.22
29	<i>Channa punctatus</i>	FW-29	12.40	3.22	BDL	BDL	0.32	0.087	12.93	8.67	BDL	BDL
30	<i>Macrognathus aculeatus</i>	FW-30	22.55	5.57	BDL	BDL	0.20	0.243	19.38	3.58	BDL	BDL
31	<i>Mystus bleekeri</i>	FW-31	25.47	93.65	BDL	BDL	0.41	0.766	93.16	7.80	BDL	BDL
32	<i>Silonia silondia</i>	FW-32	23.24	4.21	BDL	BDL	0.54	0.18	29.92	1.86	BDL	BDL

Table 3. Heavy metal concentrations (mg/kg) in fish during Pre-monsoon season.

Sl. No.	Scientific Name	Sample ID	Zn (mg/kg)	Al (mg/kg)	Cd (mg/kg)	Pb (mg/kg)	Cu (mg/kg)	Ni (mg/kg)	Fe (mg/kg)	Mn (mg/kg)	Cr (mg/kg)	Co (mg/kg)
01	<i>Aila coila</i>	FPM-01	26.84	9.43	BDL	3.28	1.41	0.182	17.93	3.65	8.18	BDL
02	<i>Aorichthys aor</i>	FPM-02	14.96	43.19	BDL	BDL	0.40	0.235	54.77	2.40	0.20	BDL
03	<i>Mastacembelus armatus</i>	FPM-03	11.59	5.10	BDL	1.34	1.26	0.04	12.36	2.86	1.94	0.17
04	<i>Labeo bata</i>	FPM-04	15.00	14.21	BDL	BDL	1.08	0.16	20.41	4.07	BDL	BDL
05	<i>Glossogobius giuris</i>	FPM-05	18.90	17.05	BDL	2.70	1.30	0.121	13.50	4.26	BDL	0.12
06	<i>Nandus nandus</i>	FPM-06	14.61	5.67	BDL	0.86	1.26	0.045	11.40	6.27	0.19	0.05
07	<i>Colisa lalia</i>	FPM-07	35.49	32.48	BDL	6.85	3.03	BDL	38.5	19.67	BDL	BDL
08	<i>Gudusia chapra</i>	FPM-08	18.01	119.3	BDL	BDL	1.06	0.28	77.66	14.84	BDL	BDL
09	<i>Chitala chitala</i>	FPM-09	10.98	5.72	BDL	BDL	0.46	0.08	12.78	2.45	BDL	BDL
10	<i>Colisa chuna</i>	FPM-10	15.77	12.98	BDL	0.12	1.56	BDL	36.21	3.90	0.40	BDL
11	<i>Barbodes sarana</i>	FPM-11	33.69	35.15	BDL	6.05	26.67	0.366	18.56	5.96	BDL	BDL
12	<i>Tenualosa ilisha</i>	FPM-12	11.01	3.91	0.07	0.59	1.11	0.01	27.03	2.21	0.07	BDL
13	<i>Notopterus notopterus</i>	FPM-13	19.15	9.00	BDL	2.54	0.78	0.182	19.08	6.22	BDL	BDL
14	<i>Tetraodon cutcutia</i>	FPM-14	34.21	34.01	BDL	BDL	1.40	0.31	41.87	20.01	BDL	BDL
15	<i>Ctenopharyngodon idella</i>	FPM-15	10.69	1.78	BDL	BDL	0.82	0.11	9.75	1.01	0.15	BDL
16	<i>Labeo calbasu</i>	FPM-16	30.80	43.57	BDL	1.89	1.65	BDL	19.84	4.50	1.04	BDL
17	<i>Mystus cavasius</i>	FPM-17	17.56	28.21	BDL	BDL	1.00	0.21	35.76	1.75	BDL	BDL
18	<i>Lepidocephalichthys guntea</i>	FPM-18	25.64	8.05	BDL	3.88	2.38	0.352	15.96	4.09	BDL	0.077
19	<i>Puntius sarena</i>	FPM-19	30.68	20.06	BDL	2.12	1.59	0.176	19.90	8.52	BDL	BDL
20	<i>Xenentodon cancila</i>	FPM-20	28.87	4.66	BDL	2.71	1.02	0.099	7.85	2.01	2.17	0.15
21	<i>Clupisoma garua</i>	FPM-21	24.32	20.22	BDL	BDL	0.41	0.21	41.65	10.98	BDL	BDL
22	<i>Anabus testudineus</i>	FPM-22	15.74	32.50	BDL	3.09	0.985	0.135	32.71	4.69	2.29	0.28
23	<i>Amblypharyngodon mola</i>	FPM-23	43.67	31.55	BDL	BDL	2.43	0.98	30.10	4.32	BDL	BDL
24	<i>Ompok pabda</i>	FPM-24	38.45	4.50	0.23	BDL	0.63	0.28	11.47	2.10	BDL	0.42
25	<i>Johnius coitor</i>	FPM-25	8.65	27.70	BDL	BDL	0.32	0.16	22.47	2.67	0.04	BDL
26	<i>Labeo rohita</i>	FPM-26	29.20	104.87	0.03	BDL	1.99	0.17	66.22	9.88	0.31	0.45
27	<i>Stinging catfish</i>	FPM-27	10.84	10.80	BDL	1.66	1.22	0.073	24.84	2.79	1.97	BDL
28	<i>Channa striatus</i>	FPM-28	15.76	3.21	BDL	BDL	1.12	0.08	39.31	10.13	0.09	0.04
29	<i>Channa punctatus</i>	FPM-29	16.70	66.50	BDL	5.86	0.94	BDL	54.34	8.96	3.62	0.51
30	<i>Macrognathus aculeatus</i>	FPM-30	21.45	5.07	BDL	BDL	0.21	0.21	18.32	3.01	BDL	BDL
31	<i>Mystus bleekeri</i>	FPM-31	26.54	69.53	BDL	3.00	2.19	BDL	37.62	3.71	1.23	BDL
32	<i>Silonia silondia</i>	FPM-32	22.53	4.01	BDL	BDL	0.32	0.173	29.13	1.74	BDL	BDL

FR – Fish Rainy Season

FW – Fish Winter Season

FPM- Fish Pre-monsoon

BDL – Below Detection Limit

3.2. Correlation Matrix

In aquatic environment, the inter relationship among metals in fishes provide significant information sources and

pathways of variables. The result of correlations between heavy metals acquiesced with the results obtained by PCA and CA that confirm some new associations between parameters. There was significant positive correlation between Cd vs Co (0.733), moderate positive correlation

between Fe vs Al (0.568), Ni vs Co (0.482), Mn vs Co (0.395) and weak correlation exist between Cr vs Co (0.351), Pb vs Cr (0.283). The strong and moderate correlation indicates their sources of origin are similar especially from

industrial effluents, municipal wastes and agricultural inputs. On the other hand, weak negative correlations were found between Cd vs. Zn (-0.241), Pb vs. Ni (-.155), Cd vs. Fe (-0.147) in river fish.

Table 4. Correlation matrix of heavy metals in fish.

	Zn	Al	Cd	Pb	Cu	Ni	Fe	Mn	Cr	Co
Correlation	Zn	1.000								
	Al	.217	1.000							
	Cd	-.241	-.029	1.000						
	Pb	.247	.071	-.054	1.000					
	Cu	.165	.098	-.039	.261	1.000				
	Ni	.181	.091	.721	-.155	.013	1.000			
	Fe	.070	.568	-.147	-.087	-.029	-.075	1.000		
	Mn	.187	.356	.559	.191	-.010	.446	.253	1.000	
	Cr	-.125	.017	.400	.283	-.059	.215	-.100	.157	1.000
	Co	-.054	.119	.733	-.027	.012	.482	.066	.395	.351

3.3. Principal Component Analysis

The extraction method was used to find out the principal components in PCA analysis that was Eigen values. The components were taken as principal components whose Eigen values was greater than 0.6 were taken into account. 6 PCs were extracted by using correlation matrix which reflects the processes influencing the heavy metals composition having 88.16 % of total sample variance (Table 4). The total variance of the PCs were 29.0%, 18.81% and 14.48%,

10.91%, 8.74%, 6.22% for PC 1, PC 2, PC 3, PC 4, PC 5, PC 6 respectively. PC 1 is strongly correlated with Cd, Co, Ni, Mn and PC 2 with Al, Fe, Zn. Moreover, there was strong correlation of PC 3 with Pb, PC 5 with Cu and there was no strong correlation for PC 4 and PC 6. The source of PC 1 and PC 2 can be considered as mixed source from anthropogenic inputs particularly from industrial effluents and agricultural activities in the study area. Whereas PC 3, PC 4, PC 5 and PC 6 can be considered as different source from both lithogenic and anthropogenic inputs.

Table 5. Component matrix of six factors model with strong to moderate loadings in fish.

Parameters	Component					
	PC 1	PC 2	PC 3	PC 4	PC 5	PC 6
Cd	.910	-.315			.101	-.141
Co	.804				.178	.121
Ni	.778			-.453		.165
Mn	.717	.381			-.172	-.469
Al	.234	.784	-.230	.185		.162
Fe		.697	-.494	.280	.120	
Zn		.567	.352	-.510	-.418	.263
Pb		.289	.793	.345	-.146	-.250
Cr	.494	-.201	.335	.578	-.135	.423
Cu		.277	.529	-.236	.749	
Eigen value	2.9	1.88	1.45	1.10	0.87	0.62
% Total variance	29.0	18.81	14.48	10.91	8.74	6.22
Cumulative %	29.0	47.81	62.29	73.20	81.94	88.16




Figure 2. Concentration of different heavy metals in fishes during Rainy season.




Figure 3. Concentration of different heavy metals in fishes during Winter season.




Figure 4. Concentration of different heavy metals in fishes during Pre-monsoon.

4. Discussion

The concentration of toxic metals in 32 fish species collected from Meghna River were found to be below the WHO permissible concentrations given for seafood (WHO 1972, 1987). Zinc is an important element in human diet but lower concentration can cause serious threat to human health (Agency for Toxic Substances and Disease Registry 2004). In the recent study the concentrations of Zn varied between (8.65-44.48)mg/kg. The concentration of Zn was lower than FAO maximum guideline of 30 mg/kg (FAO/WHO 1989; FAO 1983) except in *Amblypharyngnodon mola*, *Colisa lalia*, *Tetraodon cutcutia*, *Barbodes sarana*, *Labeo calbasu*, *Puntius sarena*, *Ompok pabda* for 3 consecutive seasons. The concentrations up to 40 mg/kg for Zn may induce toxicity, characterized by symptoms of irritability, muscular stiffness and pain, loss of appetite, and nausea (NAS-NRC 1974) though the toxicity of Zn rare.

In the present study the amount of Cd found in *Ompok pabda*, *Barbodes sarana*, *Tenualosa ilisha*, *Puntius sarena*, *Stinging catfish* exceeded the limit (0.02 mg/kg) set by the (EU 2006) but in most fishes the concentration was below detection limit. Cadmium can accumulate in the human body that may causes prostate cancer and breast cancer (Saha and Zaman 2012), kidney dysfunction, skeletal damage and reproductive deficiencies in human (Commission of the European Communities 2001).

The concentration of Pb ranged between (BDL-6.85) mg/kg that lower than the limit set by the EU is 0.3 mg/kg (EU 2006) except in *Colisa lalia*, *Aila coila*, *Mastacembelus armatus*, *Glossogobius giurus*, *Nandus nandu*, *Barbodes sarana*, *Tenualosa ilisha*, *Labeo calbasu*, *Lepidocephalichthys guntea*, *Xenentodon cancila*, *Stinging catfish* for 3 seasons. More or less similar result was found by Staniskiene *et al.* (2006); Copat *et al.* (2012). Lead is highly responsible for the reduced cognitive development and intellectual performance in children and increased blood pressure and cardio vascular disease in adults (Commission of the European Communities 2001).

FAO/WHO (1989) established limits for Cu in fish as 30.0 mg/kg for human health. The value of Cu in the present study was varied between (0.03-32.44) mg/kg far below the permissible limit except in *Barbodes sarana* exceeded the permissible limit. But Cu is necessary for the synthesis of haemoglobin but can cause harm at high concentrations (McCluggage 1991) such as liver and kidney damage (Agency for Toxic Substances and Disease Registry 2004).

Ni was found in very low concentration (BDL-0.986) mg/kg in fish samples. The maximum concentration was set by (USFDA 1993) for Ni is 70–80 mg/kg. The present

concentration were far below the stipulated limit. Concentrations crossing the set limit may cause cancer of the lung and nasal cavity (USFDA 1993).

Cr is important element that helps the body use sugar, protein, and fat but excess amount have adverse effects on fish and wildlife (Akan *et al.* 2009) and sometimes Cr is carcinogenic (Institute of Medicine 2002). The concentration of Cr is below the limit (12–13) mg/kg set by the United States Food and Drug Administration (USFDA 1993). Moreover, the deficiency of Cr can affect the glucose, lipid and protein metabolism and impaired growth (Akoto *et al.* 2014).

Fe concentration varied between (7.85-147.77) mg/kg and highest concentration 147.77 mg/kg in *Channa punctatus*. Fe is essential element necessary for the production of hemoglobin, myoglobin and certain enzymes (Akoto *et al.* 2014). Fe deficiency causes anemia in human (Anderson and Fitzgerald 2010), weakness, inability to concentrate and susceptibility to infection (Akoto *et al.* 2014).

Mn in small amount needed for growth and prevention of cardiac arrest, heart attack, and stroke (Akoto *et al.* 2014; Ikem and Egiebor 2005). Acute toxicity causes psychologic and neurologic disorder (Saha and Zaman 2012). In present study the concentration of Mn in the fish muscles varied from 0.96-20.01 mg/kg. Similar result found by (Akoto *et al.* 2014) and this result was lower than Begum *et al.* (2005) in Dhamondi Lake, Dhaka, Bangladesh (8.8–23.5 $\mu\text{g/g}$) and 0.59–11.74 $\mu\text{g/g}$ in Lake Tanganyika, Tanzania (Chale 2002).

Co concentrations was measured in all fish samples ranged from BDL-0.7 mg/kg wet weight. Co is beneficial to health but high level of Co may cause lung and heart effects and dermatitis (Agency for Toxic Substances and Disease Registry 2004).

The concentrations of Al ranged from 1.78-120.4 mg/kg in the fish muscles lower than the criteria for the maximum concentrations and criterion continuous concentration values of the US EPA (EPA 2002) water quality criteria. Aluminium frequently occur in the biosphere (Poleo 1995; Gromysz-Kalkowska and Szubartowska 1999; Weng *et al.* 2002) and the in concentration of Al in rivers and lakes may cause fish death (Reitz *et al.* 1996; Alloway and Ayres 1997).

Ikem and Egiebor (2005) stated that some factors (the duration of exposure of fish to contaminants in water, feeding habit of fish, concentrations of contaminants in water column, water chemistry, contamination of fish during handling and processing) influenced the contamination of fish.

5. Conclusion

The findings of the present study indicate the fish community that are consumed regularly by the local people is being contaminated by different heavy metals discharged

from the various industrial, urban as well as the agricultural activities occurred in the vicinity of the Meghna River. The elevated level of heavy metals found in most edible fishes ultimately will harm the human health. Therefore, the present research recommends that the point sources of heavy metals in the vicinity of the Meghna River should be strictly monitored for protecting the health of riverine ecosystem along with fish community.

Acknowledgements

The authors are grateful to the Bangladesh Council of Scientific and Industrial Research (BCSIR), Chittagong. The proposed research is a major contribution of Biodiversity, Environment and Climate Change Research Laboratory, Institute of Marine Sciences and Fisheries, University of Chittagong.

References

- [1] Abdullah, M. R., Mohd Kamar, K. A., Mohd Naw, M. N., Tarmizi Haron, A., and Arif, M. et al. (2009). "Industrialized building system: A definition and concept." Proc. of ARCOM Conference 2009, Nottingham, UK.
- [2] Abua Ikem and Nosa O. Egiebor, (2005). Assessment of trace elements in canned fishes (mackerel, tuna, salmon, sardines and herrings) marketed in Georgia and Alabama (United States of America), *Journal of Food Composition and Analysis* 18 (2005) 771–787. www.elsevier.com/locate/jfca
- [3] Agency for Toxic Substances and Disease Registry, (2004). Agency for Toxic Substances and Disease Registry, Division of Toxicology, Clifton Road, NE, Atlanta, GA, available at: <http://www.atsdr.cdc.gov/toxprofiles/>.
- [4] Ahmed, M. K., Baki, M. A., Islam, M. S., Kundu, G. K., Sarkar, S. K., Hossain, M. M. et al. (2015a). Human health risk assessment of heavy metals in tropical fish and shell fish collected from the river Buriganga, Bangladesh. *Environ. Sci. Pollut. Res.*, <http://dx.doi.org/10.1007/s11356-015-4813-z>.
- [5] Ahmed, M. K., Shaheen, N., Islam, M. S., Al-Mamun, M. H., Islam, S., Banu, C. P. et al. (2015b). Trace elements in two staple cereals (rice and wheat) and associated health risk implications in Bangladesh. *Environ. Monit. Assess.* 187, 326–336.
- [6] Ahmed, M. K., Shaheen, N., Islam, M. S., Al-Mamun, M. H., Islam, S., Mohiduzzaman, M., Bhattacharjee, L. et al. (2015c). Dietary intake of trace elements from highly consumed cultured fish (*Labeorohita*, *Pangasius pangasius* and *Oreochromis mossambicus*) and human health risk implications in Bangladesh. *Chemosphere* 128, 284–292.
- [7] Akoto, O, Bismark Eshun, F, Darko, G. and Adei, E. et al. (2014). Concentrations and Health Risk Assessments of Heavy Metals in Fish from the Fosu Lagoon, *Int. J. Environ. Res.*, 8(2):403-410, Spring 201
- [8] Akan, J. C., Abdul-Rahman, F. I., Sodipo, O. A. and Akandu, P. I. et al. (2009). Bioaccumulation of some heavy metals of six fresh water fishes caught from Lake Chad in Doron Buhari, Borno State, Nigeria. *J. Appl. Sci. in Environ. Sanit*, 4 (2), 103-114.
- [9] Akif, M., Khan, A. R., Sok, K., Min, K. S., Hussain, Z. and Maal-Abrar, M. et al. (2002). Textile effluents and their contribution towards aquatic pollution in the Kabul River (Pakistan). *Journal of Chemical Society of Pakistan* 24(2): 106-111.
- [10] Alhashemi AH, Sekhavatjou MS, Kiabi BH. et al. (2012). Bioaccumulation of trace elements in water, sediment, and six fish species from a freshwater wetland, Iran. *Microchem J* 104:1–6.
- [11] Alloway, B. J., Ayres, D. C. et al. (1997). *Chemical Principles of Environmental Pollution*. Second Edition. Blackie Academic & Professional, London SE1 8HN, UK.
- [12] Anawar HM, Akai J, Mostofa KM. et al. (2002). Arsenic poisoning in groundwater: Health risk and geochemical sources in Bangladesh. *Environ Internat* 27: 597–4.
- [13] Bai, J., Xiao, R., Cui, B., Zhang, K., Wang, Q., Liu, X., Gao, H., Huang, L. et al. (2011). Assessment of heavy metal pollution in wetland soils from the young and old reclaimed regions in the Pearl River Estuary, South China. *Environ. Pollut.* 159, 817–824.
- [14] Banerjee N, Nandy S, Kearns JK. et al. (2011). Polymorphisms in the TNF- α and IL10 gene promoters and risk of arsenic-induced skin lesions and other non-dermatological health effects. *Toxicol Sci* 121(1): 132–39
- [15] Begum, A., Amin, M. N., Kaneco, S. and Ohta, K. et al. (2005). Selected elemental composition of fish, *Tilapia nilotica*, *Cirrhina mrigala* and *Clarius batrachus* from the fresh water Dhanmondi Lake in Bangladesh. *Food Chemistry*, 93, 439–443.
- [16] Camusso, M., Vigano, L., Baitstrini, R. et al. (1995). Bioaccumulation of trace metals in rainbow trout. *Ecotox. Environ. Safe.* 31: 133–141.
- [17] Copat, C., Bella, F., Castaing, M., Fallico, R., Sciacca, S. and Ferrante, M. et al. (2012). Heavy Metals Concentrations in Fish from Sicily (Mediterranean Sea) and Evaluation of Possible Health Risks to Consumers. *Bull. Environ. Contam. Toxicol.*, 88, 78–83.
- [18] Commission of the European Communities, (2001). Commission Regulation (EC) No. 221/2002 of 6 February 2002 amending regulation (EC) No. 466/2002 setting maximum levels for certain contaminants in foodstuffs. Official Journal of the European Communities, Brussels, 6 February 2002.
- [19] Dreher T. (2003). Evaluation of graphical and multivariate methods for classification of water chemistry data, *Hydrogeology Journal*, 11, 605–606.
- [20] European Union, (2006). Setting maximum levels for certain contaminants in foodstuffs, Commission Regulation (EC) No 1881/2006; OJ L 364, 20.12.2006, p. 5
- [21] European Union, (2002). Heavy Metals in Wastes, European Commission on Environment. Available at http://ec.europa.eu/environment/waste/studies/pdf/heavymetal_sreport.pdf
- [22] Fang Y, Sun X, Yang W. et al. (2014). Concentrations and health risks of lead, cadmium, arsenic, and mercury in rice and edible mushrooms in China. *Food Chem* 147: 147–51.

- [23] FAO/WHO, (1989). Evaluation of certain food additives and the contaminants mercury, lead and cadmium. WHO Technical Report, Series No. 505.
- [24] Fernandes, C., Fontainhas-Fernandes, A., Cabral, D., Salgado, M. A. et al. (2008). Heavy metals in water, sediment and tissues of *Liza saliens* from Esmoriz-Paramos lagoon, Portugal. *Environ. Monit. Assess.* 136: 267–275.
- [25] Forti E, Salovaara S, Cetin Y. et al. (2011). In vitro evaluation of the toxicity induced by nickel soluble and particulate forms in human airway epithelial cells. *Toxicol in Vitro* 25:454–61
- [26] Grigoratos, T., Samara, C., Voutsas, D., Manoli, E., Kouras, A. et al. (2014). Chemical composition and mass closure of ambient coarse particles at traffic and urban background sites in Thessaloniki, Greece. *Environ. Sci. Pollut. Res.* 21, 7708–7722.
- [27] Gromysz-Kalkowska, K., Szubartowska, E. (1999). Aluminium - its ecological role and toxicity for animals. *Medycyna Weterynaryjna* 55(4), 229-223
- [28] G. Afthan; G. Cumont; H. P. Dypdahl; K. Gadd; G. N. Havre; K. Julshamn; K. Kåverud; B. Lind; J. Loimaranta; M. Merseburg; A. Olsson; S. Piepponen; B. Sundström; B.J. Uppstad; T. Waaler; L. Winnerstam. et al. (2000). Determination of Metals in Foods by Atomic Absorption Spectrometry after Dry Ashing: NMKLL Collaborative Study. LARS JORHEM. National Food Administration, Chemistry Division 2, Box 622, S-751 26 Uppsala, Sweden. Collaborators: *Journal of AOAC International* Vol. 83, No. 5, 2000 1207.
- [29] Institute of Medicine, (2002). Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. Institute of Medicine of the National Academies, The National Academy Press, 2101 Constitution Avenue, NW, Washington, DC, p. 773.
- [30] Irwandi, J. and Farida, O. (2009). Mineral and heavy metal contents of marine fin fish in Langkawi Island, Malaysia, *International Food Research Journal* 16: 105-112 (2009).
- [31] Islam, M. S., Ahmed, M. K., Habibullah-Al-Mamun, M., Islam, K.N., Ibrahim, M., Masunaga, S. et al. (2014). Arsenic and lead in foods: a potential threat to human health in Bangladesh. *Food Addit. Contam. Part A* 31 (12), 1982–1992.
- [32] Islam, M. S., Ahmed, M. K., Habibullah-Al-Mamun, M., Hoque, M. F. et al. (2015a). Preliminary assessment of heavy metal contamination in surface sediments from a river in Bangladesh. *Environ. Earth Sci.* 73, 1837–1848.
- [33] Islam, M. S., Ahmed, M. K., Raknuzzaman, M., Habibullah-Al-Mamun, M., Masunaga, S. et al. (2015b). Metal speciation in sediment and their bioaccumulation in fish species of three urban rivers in Bangladesh. *Arch. Environ. Contam. Toxicol.* 68, 92–106.
- [34] Islam, M. S., Ahmed, M. K., Raknuzzaman, M., Habibullah-Al-Mamun, M., Islam, M. K. et al. (2015c). Heavy metal pollution in surface water and sediment: a preliminary assessment of an urban river in a developing country. *Ecol. Indic.* 48, 282–291.
- [35] Ismaniza Ismail, Idaliza Mat Saleh, (2012). Analysis of heavy metals in water and fish (*Tilapia* sp.) samples from Tasik Mutlara, Puchong. *The Malaysian Journal of Analytical Sciences*, Vol 16 No 3 (2012): 346-352.
- [36] Jaillon, L. and Poon, C. S. (2009). “The evolution of prefabricated residential building systems in hong kong: A review of the public and the private sector.” *Automation in Construction*, Vol. 18, No. 3, pp. 239-248.
- [37] Jordao, C. P., Pereira, M. G., Bellato, C. R., Pereira, J. L. and Matos, A. T. et al. (2002). Assessment of water systems for contaminants from domestic and industrial sewages. *Environmental Monitoring Assessment* 79(1): 75-100.
- [38] John Edward Cantle, (1982). Atomic Absorption Spectrometry. Elsevier Scientific publishing Company, New York, 1982, pp -159-160.
- [39] Kamal J. Elnabris, Shareef K. Muzyed, Nizam M. El-Ashgar. et al. (2012). Heavy metal concentrations in some commercially important fishes and their contribution to heavy metals exposure in Palestinian people of Gaza Strip (Palestine), *Journal of the Association of Arab Universities for Basic and Applied Sciences*(2013) 13, 44–51. <http://dx.doi.org/10.1016/j.jaubas.2012.06.0>
- [40] Khan, S., Cao, Q., Zheng, Y. M., Huang, Y. Z., Zhu, Y. G. et al. (2008). Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China. *Environ. Pollut.* 152, 686–692.
- [41] Martin, J. A. R., Arana, C. D., Ramos-Miras, J. J., Gil, C., Boluda, R. et al. (2015). Impact of 70 years urban growth associated with heavy metal pollution. *Environ. Pollut.* 196, 156–163.
- [42] McCluggage D. (1991). Heavy metal poisoning, NCS Magazine. The Bird Hospital, CO, USA. Available at www.cockatiels.org/articles/Diseases/metals.html
- [43] Md Saiful Islam, Md Kawser Ahmed & Md Habibullah-Al-Mamun. et al. (2014). Determination of Heavy Metals in Fish and Vegetables in Bangladesh and Health Implications, Human and Ecological Risk Assessment: An International Journal, DOI: 10.1080/10807039.2014.950172
- [44] Namminga, H. N., Wilhm, J. (1976). Effects of high discharge and an oil refinery cleanup operation on heavy metals in water and sediments in Skeleton Creek. *Proceedings of the Oklahoma Academy of Science*, 56: 133–138.
- [45] NAS-NRC, (1982). National Drinking Water and Health, Academy of Sciences-National Research Council National Academic Press, Washington D.C.
- [46] Özmen, H., Kùlahçı, F., Çukurovalı, A., and Doğru, M. et al. (2004). Concentrations of heavy metal and radioactivity in surface water and sediment of Hazar Lake (Elazığ, Turkey). *Chemosphere*, 55: 401–408.
- [47] Öztürk, M., Özözen, G., Minareci, O., and Minareci, E. et al. (2008). Determination of heavy metals in of fishes, water and sediment from the Demirköprü Dam Lake (Turkey). *Journal of Applied Biological Sciences*, 2(3): 99–104.
- [48] Pan K and Wang WX. (2012). Trace metal contamination in estuarine and coastal environments in China. *Sci Tot Environ* 421: 3–16.
- [49] Papagiannis I, Kagaloub I, Leonardos J. et al. (2004). Copper and zinc in four freshwater fish species from Lake Pamvotis (Greece). *Environ Internat* 30: 357–62.
- [50] Pote, J., Haller, L., Loizeau, J. L., Bravo, A. G., Sastre, V., and Wildi, W. et al. (2008). Effects of a sewage treatment plant outlet pipe extension on the distribution of contaminants in the sediments of the Bay of Vidy, Lake Geneva, Switzerland. *Bioresource Technol.*, 99: 7122–7131.

- [51] Poleo, A. B. S. (1995). Aluminium polymerization -- a mechanism of acute toxicity of aqueous aluminium to fish. *Aquatic Toxicology* 31(4), 347-356.
- [52] Praveena, S. M., Radojevic, M., Abdullah, M. H., Aris, A. Z. et al. (2008). Application of sediment quality guidelines in the assessment of mangrove surface sediment in Mengkabong lagoon, Sabah, Malaysia. *Iran. J. Environ. Health. Sci. Eng.*, 5 (1): 35-42.
- [53] Quddus, M. M. A., M. N., Sarker, A. K., Banerjee, et al. (1988). *Studies of the Chondrichthyes Fauna (Sharks, Skates and Rays) of the Bay of Bengal*. The Journal of NOAMI. Vol. 5, No. 1 & 2. Pp. 19-39.
- [54] Quddus, M. M. A. and Shafi, M. (1983). *Bangapa sagarer Matsya Sampad (The Fisheries Resources of the Bay of Bengal)*. Kabir Publications, 38/3, Bangla Bazar, Dhaka, Bangladesh. 535 p.
- [55] Rahman MM, Asaduzzaman M, and Naidu R. et al. (2013). Consumption of arsenic and other elements from vegetables and drinking water from an arsenic-contaminated area of Bangladesh. *J Hazard Mater* 262:1056-63.
- [56] Rahman, A. K. A., Kabir, S. M. H., Ahmad, M., Ahmed, A. T. A., Ahmed, Z. U., Begum, Z. N. T., Hassan, M. A. and Khondker, M. et al. (2009). *Encyclopedia of Flora and Fauna of Bangladesh*, Vol. 24. *Marine Fishes*. Asiatic Society of Bangladesh, Dhaka. pp. 2-57.
- [57] Rashed, M. N. (2001). Monitoring of environmental heavy metals in fish from Nasser Lake. *Environ. Int.*, 27: 27-33.
- [58] Rainbow, P. S., Amiard-Triquet, C., Amiard, J. C., Smith, B. D., Langston, W. J. et al. (2000). Observations on the interaction of zinc and cadmium uptake rates in crustaceans (amphipods and crabs) from coastal sites in UK and France differentially enriched with trace metals. *Aquatic Toxicology* 50, 189,204.
- [59] Reitz, B., Heydorn, K., Pritzl, G. et al. (1996). Determination of aluminium in fish tissues by means of INAA and ICP-MS. *Journal of Radioanalytical and Nuclear Chemistry* 216 (1), 113-116.
- [60] Richard, R. B. (2005). "Industrialized building systems: Reproduction before automation and robotics." *Automation in Construction*, Vol. 14, No. 4, pp. 442-451.
- [61] Roy, B. J., M. P. Dey, M. F. Alam and N. K. Singha, et al. (2007). *Present Status of shark fishing in the marine water of Bangladesh*. UNEP/CMS/MS/Inf/10.17p. http://www.cms.int/bodies/meetings/regional/sharks/pdf_docs/Inf_10_Bangladesh_Presentation_on_Shark_Fishing.pdf: Accessed 15 March, 2009.
- [62] Sanchez-Chardi, A., Lopez-Fuster, M. J., Nadal, J. et al. (2007). Bioaccumulation of lead, mercury, and cadmium in the greater white-toothed shrew, *Crocidura russula*, from the Ebro Delta (NE Spain): Sex- and age-dependent variation. *Environ. Pollut.* 145, 7-14.
- [63] Saha, N and Zaman, M. R. (2012). Evaluation of possible health risks of heavy metals by consumption of foodstuffs available in the central market of Rajshahi City, Bangladesh; *Environ. Monit. Assess.* 185, 38673878.
- [64] Sankar TV, Zynudheen AA, Anandan R. et al. (2006). Distribution of organochlorine pesticides and heavy metal residues in fish and shellfish from Calicut region, Kerala, India. *Chemosphere* 65: 583-90.
- [65] Sekabira, K., Oryem Origa, H., Basamba, T. A., Mutumba, G., Kakudidi, E. et al. (2010). Assessment of heavy metal pollution in the urban stream sediments and its tributaries. *Int. J. Environ. Sci. Technol.* 7 (4), 435-446.
- [66] Sharma RK, Agrawal M, and Marshall FM. et al. (2007). Heavy metals contamination of soil and vegetables in suburban areas of Varanasi, India. *Ecotoxicol Environ Saf* 66: 258-66.
- [67] Shuhaimi-Othmana, M., Pascoe, D. (2007). Bioconcentration and depuration of copper, cadmium, and zinc mixtures by the freshwater amphipod *Hyalella azteca*. *Ecotoxicology and Environmental Safety* 66 (1), 29-35.
- [68] Sun, T. H., Zhou, Q. X., Li, P. J. et al. (2001). *Pollution Ecology*. Science Press, Beijing, China, pp. 160,194.
- [69] Staniskiene, B., Matusevicius, P., Budreckiene, R. and Skibniewska, K. A. et al. (2006). Distribution of Heavy Metals in Tissues of Freshwater Fish in Lithuania; *Polish J. of Environ. Stud.*, 15, 585-591.
- [70] Tarra-Wahlberg, N. H., Flachierm A., Lane, S. N. and Sangfors, O. et al. (2001). Environmental impacts and metal exposure of aquatic ecosystems in rivers contaminated by small scale gold mining: The Puyango River Basin, Southern Ecuador. *Science of the Total Environment* 278 (1-3): 239-261.
- [71] Thanoon, W. A., Wah Peng, L., Abdul Kadir M. R., Jaafar, M. S., and Salit, M. S. et al. (2003). "The essential characteristics of industrialised building system." *Proc. of International Conference on Industrialized Building Systems*, Kuala Lumpur, Malaysia, pp. 283-292.
- [72] Tuzen M. (2009). Toxic and essential trace elemental contents in fish species from the Black Sea, Turkey. *Food Chem Toxicol* 47, 1785-90.
- [73] USFDA, (1993). United States Food and drug administration, Guidance document for chromium in shellfish. *DHHS/PHS/FDA/CFSAN/Office of seafood*, Washington D. C.
- [74] Vieira C, Morais S, Ramos S. et al. (2011). Mercury, cadmium, lead and arsenic levels in three pelagic fish species from the Atlantic Ocean: Intra- and inter-specific variability and human health risks for consumption. *Food Chem Toxicol* 49(4): 923-32.
- [75] Weng, L., Temminghoff, E. J. M., Van Riemsdijk, W. et al. (2002). Aluminium speciation in natural waters: measurement using Donnan membrane technique and modelling using NICA-Donnan. *Water Research* 36(17), 4215-4226.
- [76] Wilson, B., Pyatt, F.B. (2007). Heavy metal dispersion persistence, and bioaccumulation around an ancient copper mine situated in Anglesey, UK. *Ecotoxicol. Environ. Saf.* 66, 224-231.
- [77] WHO (World Health Organization), (1995). *Lead*. Geneva: Environmental Health Criteria, Geneva, Switzerland.
- [78] WHO (World Health Organisation), (1972). Evaluation of certain food additives and the contaminants mercury, lead, cadmium. 16 th Report of the Joint FAO/WHO Expert Committee on Food Additives. Technical Report Series 505, Geneva.

- [79] WHO (World Health Organisation), (1987). Evaluation of certain food additives and contaminants. 33rd Report of the joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series 776, Geneva, 80 pp.
- [80] Yi, Y., Yang, Z., Zhang, S. et al. (2011). Ecological risk assessment of heavy metals in sediment and human health risk assessment of heavy metals in fishes in the middle and lower reaches of the Yangtze river basin. *Environ. Pollut.* 159, 2575–2585.
- [81] Zhang, C., Qiao, Q., Piper, J. D. A., Huang, B. et al. (2011). Assessment of heavy metal pollution from a Fe-smelting plant in urban river sediments using environmental magnetic and geochemical methods. *Environ. Pollut.* 159, 3057–3070.
- [82] Zhou, Q. X., Kong, F. X., Zhu, L. et al. (2004). *Ecotoxicology: Principles and Methods*. Science Press, Beijing, pp. 161-217.
- [83] Zhou, Q. X. (1995). *Ecology of Combined Pollution*. China Environmental Science Press, Beijing, pp. 1-29.