
Biological Potentials of Ginger Associated *Streptomyces* Compared with Ginger Essential Oil

Herivony Onja Andriambelason¹, Rado Rasolomampianina¹, Rahanira Ralambondrahety¹, Rigobert Andrianantenaina¹, Marson Raherimandimby², Fidèle Randriamiharisoa³

¹National Centre of Environmental Research, Antananarivo, Madagascar

²Department of Fundamental and Applied Biochemistry, Faculty of Sciences, University of Antananarivo, Antananarivo, Madagascar

³Ministry of Health, Antananarivo, Madagascar

Email address:

herionja@ymail.com (H. O. Andriambelason)

To cite this article:

Herivony Onja Andriambelason, Rado Rasolomampianina, Rahanira Ralambondrahety, Rigobert Andrianantenaina, Marson Raherimandimby, Fidèle Randriamiharisoa. Biological Potentials of Ginger Associated *Streptomyces* Compared with Ginger Essential Oil. *American Journal of Life Sciences*. Vol. 4, No. 6, 2016, pp. 152-163. doi: 10.11648/j.ajls.20160406.13

Received: October 11, 2016; **Accepted:** October 25, 2016; **Published:** December 2, 2016

Abstract: Medicinal plants and associated microorganisms are recognized to have beneficial relationship. These two organisms are well known for their ability to produce bioactive secondary metabolites which the similarity has been demonstrated in a few works. This study had for objective to assess biological potentials of actinomycetes isolated from ginger rhizomes and its rhizospheric soil and to determine their similarity and efficiency with ginger essential oil. Among the 63 actinomycetes strains isolated from the rhizomes and rhizospheric soils of two ginger countries plantations of Soavinandriana Itasy-Madagascar, biological activity tests showed that 16 strains (2 endophytes and 14 from rhizospheric soils of ginger) exhibited antimicrobial activity against at least one germ. The strains are more active against Gram+ bacteria and fungi than Gram- bacteria. Only, one strain isolated from ginger rhizospheric soil of the site n°2 (AHO 18) inhibited the development of all tests germs. The tests conducted on six representative strains selected on the basis of antimicrobial assay showed that extracts from the isolates AHO 3 and AHO 43 have strong antiproliferative activity on cells HT-20 (colon cancer) with IC₅₀ values of 5µg/ml and 2,2µg/ml, respectively; strong antimalaria activity against the chloroquino-resistant *Plasmodium falciparum* strain (IC₅₀=1,25µg/ml for AHO 3 extract and 2,5<IC₅₀<5µg/ml for AHO 43 extract) and antioxidant activity (IC₅₀=15mg/ml for AHO 3 extract and 10,6mg/ml for AHO 43 extract). The 2 isolates based on phenotypic and molecular characterization using their 16S rRNA gene were identified as *Streptomyces chrysomallus* (isolate AHO 3) and *Streptomyces sp* (isolate AHO 43). Moreover, the two essential oils of ginger tested showed antimicrobial activity against all tests germs used and antioxidant activity. Only, ginger essential oil from the site n°2 exhibited moderate antiproliferative potential (IC₅₀=14µg/ml) on colon cancer cells and high antiplasmodial activity (2,5<IC₅₀<5µg/ml). *Streptomyces sp* showed similar and strong biological activities than those of ginger essential oil from the site n°2. Chemical screening of the *Streptomyces sp* extract and the essential oil H2 revealed the common presence of terpens and phenolic compounds.

Keywords: Actinomycetes, Endophytes, Rhizospheric Soil, Rhizome, Ginger, Essential Oil, Antimicrobial, Antioxidant, Antimalaria, Antiproliferative

1. Introduction

Medicinal plants are well-known for their richness in natural bioactive substances. They have been used since the first civilization and the therapy has been continued to resort to these vegetal resources nowadays in two ways: the extraction of pure natural substances frequently designed for

major therapeutic indications or in nature with simple or innovative family medication forms (extract, powder...), generally used in minor pathology or in appoint therapy. In some cases, the different steps of bioactive molecules extraction, the diverse screening and the clinical assays impose an important quantity of plant materials. The over-harvesting of these natural resources could make, thus, some

species in danger. One alternative for the production of bioactive compounds similar to those of plants and for the preservation of vegetal biodiversity is the valorization of microorganisms associated with the plants. Among them, endophytic fungi are well-known. *Fusarium solani* and *Entrophospora infrequens* isolated from *Apodytes dimidiata* tree and *Nothapodytes foetida* twigs, respectively, produced the same anticancer substances as the two plants: the camptothecin [1, 2]. The cajanol, an anticancer produced by *Hypocrea lixii* isolated from *Cajanus cajan* roots [3]. Another endophytic fungus (*Mucor fragilis*) isolated from *Sinopodophyllum hexandrum* rhizomes produced similar anticancer substances as the plant: the podophyllotoxin and the kaempferol [4]. However, other endophytic microorganisms belonging to actinomycetes group have been demonstrated important sources of natural bioactive substances showing comparable biological activity and producing similar bioactive substances than host plant. Caruso *et al.* [5] were isolated from different organs of *Taxus baccata* and *Taxus brevifolia* endophytic fungi and actinomycetes producing similar anticancer compounds (the taxol) than host plants. Recently, Akshatha *et al.* [6] were isolated from leaves and stems of two antidiabetic plants (*Leucas ciliata* and *Rauwolfia densiflora*) two actinomycetes (*Streptomyces longisporoflavus* and *Streptomyces sp.*) which extracts exhibited antidiabetic potential.

Actinomycetes are prokaryote, Gram positive and filamentous bacteria which provide 70% of actual drugs and anti-infectious [7]. Secondary metabolites produced by these microorganisms represent a wide source of compounds with broad structural diversity and are endowed of important biological potential. Then, they are used in many domains as human therapeutic and veterinary, food industry and agriculture to fight against certain pathogens and toxinogens for human, animals and vegetal. Actinomycetes are ubiquitous and can be isolated from different natural habitats as soils [8, 9], plants [10, 11], waters [12], marine organisms [13] and even extreme sites (arid area, polar site...) [14]. The present work selects ginger as plant material for its therapeutic and aromatic properties well-known for millennium, its medicinal use as ubiquitous as its culinary use and its essential oil (1 to 3% of rhizomes) rich in active compounds with different properties [15, 16].

In recent years, there has been renewed interest in ginger as a source of bioactive natural products. Research works carried out on ginger have been focused on the effects of ginger consumption on health [17, 18, 19], the isolation and characterization of ginger bioactive compounds [15, 20] and the isolation of ginger endophytes [21, 22]. In order to find natural substances with comparable biological activity as active extract of the plant, actinomycetes were isolated from rhizomes and rhizospheric soil of ginger. This study was undertaken with a view to test the potential of endophytic and telluric actinomycetes from ginger as producers of natural substances showing comparable biological activity as ginger essential oil. A chemical screening of the extracts from target isolates has been reported and compared with the chemical

compounds of ginger essential oil.

2. Materials and Methods

2.1. Plant and Soil Sampling

Fresh ginger rhizomes and rhizospheric soil samples were collected from two ginger rural plantations: Ampamaha (site n°1, 19°11'S46°23'E; 1132m Alt.) and Andrefaniviny (site n°2, 19°10'S46°26'E; 1335m Alt.) located in the district of Soavinandriana in Itasy Region, Madagascar [23].

2.2. Isolation of Actinomycetes Strains

Isolation of actinomycetes strains from ginger rhizomes was conducted according to Fischer *et al.*'s methods [24] with modifications [25-26, 23] while actinomycetes isolation from rhizospheric soil samples was performed by soil dilution and heat treatment techniques as reported previously [23].

2.3. Biological Activities of Actinomycetes

2.3.1. Antimicrobial Activity of the Isolates

Actinomycetes isolates were evaluated *in vitro* for their antimicrobial activity against human pathogens and phytopathogenic microorganisms: *Klebsiella oxytoca* ATCC 8724, *Escherichia coli* ATCC 25922, *Bacillus cereus* ATCC 13061, *Staphylococcus aureus* ATCC 11632, *Candida albicans* and *Fusarium sp.*

Tests microorganisms were inoculated onto Mueller Hinton agar (for bacteria) and on Potato Dextrose Agar (PDA) (for yeast and fungus). Antibacterial activity was investigated as described by Acar and Goldstein [27] with modifications. Disks (6mm in diameter) of Mueller Hinton agar were taken aseptically and substituted by disks of mature actinomycetes grown on starch casein agar (SCA). Antifungal activity was, however, performed according to Loqman *et al.*'s techniques [28] by aseptic transfer of mature actinomycetes disks to the inoculated PDA plates with fungal mycelial disk (6mm of diameter) in the center.

Plates were kept in a refrigerator + 4°C for at least 4 h to allow the diffusion of any antibiotics produced, then incubated at 37°C for human pathogens and at 30°C for *Fusarium sp.*

The inhibition of pathogens growth was appreciated by the measure of the diameter of inhibition zone around the actinomycetes colonies, after 1 to 2 days of incubation for human pathogens and after 4 days for phytopathogenic fungus. Only isolates showing an inhibition zone greater than 8 mm were considered as active isolates.

2.3.2. Antioxidant Activity

i. Extraction of secondary metabolites

Actinomycetes isolates showing large spectrum of antimicrobial activity were selected to extract secondary metabolites. SCA media were inoculated with pure cultures of selected isolates and incubated for 8 to 14 days at 30°C. Thirty milliliters of ethanol (95°) were, then, poured into inoculated SCA. The mixture was settled at room temperature for two

hours and filtered with a steriflip. The obtained filtrat constitutes the extract which was dried by speedvac.

ii. DPPH radical scavenging activity

In vitro antioxidant capacity of actinomycetes extracts was assessed by the measure of DPPH (1,1-diphényl-2-picrylhydrazyl) radical scavenging according to Leitao *et al.* [29]. Two milliliters of different concentrations of extract methanolic solution from 16mg/ml to 0,5mg/ml were prepared and mixed with DPPH methanolic solution (0,002%). The inhibition of DPPH radical was estimated by spectrophotometer at 517nm after incubation for 30min in dark at room temperature. Antioxidant capacity of the extracts was estimated in comparison with a natural antioxidant, the ascorbic acid. The extracts were tested in triplicate and the inhibition of free DPPH radical in percentage (I %) was calculated as follows:

$I \% = \frac{A_{\text{blank}} - A_{\text{sample}}}{A_{\text{blank}}} \times 100$ [30]; A blank: Absorbance of the blank (DPPH in methanol), A sample: Absorbance of the extract.

The IC₅₀ value (concentration of required antioxidant to reduce initial concentration of DPPH to 50%) was calculated by linear extrapolation of the data that lay on the either side of the 50% inhibition level.

2.3.3. Antiproliferative Activity

In vitro antiproliferative activity of extracts from actinomycetes of ginger rhizomes and rhizospheric soils was assayed on human A-2780 ovarian and HT-20 colon cancer cells at Virginia Polytechnic Institute and State University, USA according to Cao *et al.*'s methods [31].

2.3.4. Antiplasmodial Activity

This assay was also performed at Virginia Polytechnic Institute and State University, USA according to Harinantenaina *et al.*'s methods [32] on chloroquine-resistant *Plasmodium falciparum* Dd2 strain.

2.4. Biological Activities of Ginger Rhizomes Essential Oil

In view of the various properties attributed to ginger and in the purpose to compare biological activities of ginger essential oil and those of associated actinomycetes extracts, previous tests were performed with ginger essential oil. The same methods as described above were used for antioxidant, antiproliferative and antiplasmodial tests whereas antimicrobial activity was tested using aromatogram method [23].

For all activities, the results are expressed according to the average (\pm SE) of three replicate determinations.

2.5. Identification of Active Isolates

Isolates exhibiting activities were characterized phenotypically and identified by 16S rRNA gene sequencing as reported by Andriambelosen *et al.* [23].

2.6. Chemical Screening of *Streptomyces sp* Extract

As the biological activity of an organism is due to a specific chemical compound, this work aims to determine the chemical families of *Streptomyces sp* extract and the

chemical compounds of ginger essential oil.

The screening method used for the determination of *Streptomyces sp* extract chemical families was the same as used in chemical phytoscreening which is a qualitative analysis based on coloration or precipitation reaction of the compounds of the chemical families in the crude extract [33]. The tested families are: the terpenoides, the tannins, the leucoanthocyanes, the flavonoides, the alkaloids, the saponosides, the anthraquinones, the polysaccharides and the polyphenols.

2.7. Determination of the Chemical Composition of Ginger Essential Oil

Chemical compounds of the essential oil were analyzed qualitatively and quantitatively by Gas Chromatography at Virginia Polytechnic Institute and State University, USA.

3. Results and Discussions

3.1. Actinomycetes Isolation

From the 2 ginger rhizomes samples and the 2 ginger rhizospheric soil samples collected in two ginger plantations localized in the middle-west region of Madagascar, a total of 63 actinomycetes strains were isolated on SCA medium. Among them, 8 strains were obtained from ginger rhizomes samples (AHO 1- AHO 8) and 55 from the rhizospheric soils (AHO 9- AHO 63) (figure 1). Cultural characters of the isolates are summarized in the table 1 and some varieties of isolated strains are shown in the figure 2. According to these results, the number of isolated endophytes is largely lower than actinomycetes from rhizospheric soils'number. In our knowledge, any comparative work of the number of endophytic actinomycetes and rhizospheric soil actinomycetes isolated from the same plant has been cited in the literature. However, our results confirm those obtained by Intra *et al.* [34] who reported that the rhizospheric soil is an important source of actinomycetes. Crawford *et al.* [35] showed also that soils associated with the rhizosphere contain twice as many actinomycetes as soils non- associated with the rhizosphere.




Figure 1. Number of actinomycetes strains isolated according to the plantation sites.

For plants, many works have been demonstrated that many

endophytic actinomycetes could be isolated from a plant. However, they are especially abundant in the root [36, 37,

38] confirming, thus, their isolation from the rhizomes in the present work.




Figure 2. Varieties of actinomycetes isolates.

Table 1. Cultural characters of actinomycetes isolates on SCA medium.

Characteristics	Actinomycetes isolates
Aspects of the colonies	Opaque, powdery, embedded in agar
Color of substrate mycelia	White, grey, yellow, brown
Color of aerial mycelium	White, grey, yellow, brown, orange
Diffusible pigment	Brown, yellow
Size of the colonies	1mm-5mm

3.2. Biological Activities

3.2.1. Antimicrobial Activity of Actinomycetes Isolates

Modified method used for antimicrobial test of the isolates leads to good results, the diameters of inhibition zone in Acar and Goldstein's method were low than those obtained in modified method (figure 3). Out of the 63 isolates, 16 actinomycetes strains were active against one or more of the tested pathogens, especially against Gram positive bacteria

(10 isolates) and fungi (13 isolates). These results were in agreement with those of some works demonstrating that antagonistic reaction of actinomycetes against Gram positive bacteria was much higher than Gram negative bacteria [39, 8, 40]. It could be explained by the difference in the composition of the cell membrane which is complex in Gram negative bacteria offering their resistance to diverse metabolites [41]. One isolate (AHO 18) from ginger rhizospheric soil of Andrefanivany showed inhibition of all tested microorganisms growth. Nevertheless, this strain was less active than nalidixic acid (30µg) against Gram negative bacteria but more active than fusidic acid (10µg) and nystatin (100UI) against *Staphylococcus aureus* and *Candida albicans*, respectively (table 2). In addition, other isolates as AHO 12, AHO 14 and AHO 43 exhibited high inhibition of pathogens growth than standard antibiotics used.


Table 2. Antimicrobial activity of active actinomycetes isolates.

Diameter of the inhibition zone (mm)							
Strains	Sites	<i>Escherichia coli</i> ATCC 25922	<i>Klebsiella oxytoca</i> ATCC 8724	<i>Bacillus cereus</i> ATCC 13061	<i>Staphylococcus aureus</i> ATCC 11632	<i>Candida albicans</i>	<i>Fusarium sp.</i>
AHO 1	2	-	-	-	-	-	13,5 ± 0,7 ^d
AHO 3	2	-	-	-	-	-	17,5 ± 0,7 ^{bc}
AHO 12	1	-	-	*19± 1,4 ^b	25 ± 0,0 ^b	10 ± 0,0 ^{ef}	21 ± 1,4 ^a
AHO 13	2	-	-	-	-	14 ± 1,4 ^{de}	-
AHO 14	2	-	-	-	-	43,5 ± 2,1 ^b	20 ± 0,0 ^{ab}
AHO 16	1	-	-	20± 0,0 ^b	19 ± 1,4 ^c	-	0
AHO 18	2	12,5 ± 0,0 ^a	12± 0,0 ^a	12,5± 0,7 ^c	28,5 ± 2,1 ^b	59 ± 2,9 ^a	18 ± 0,0 ^{bc}
AHO 24	1	-	-	20± 0,0 ^b	19,5± 0,7 ^c	14,5 ± 0,7 ^d	17± 1,4 ^c
AHO 31	1	-	-	12± 0,0 ^c	18 ± 1,4 ^c	10 ± 0,0 ^{ef}	-
AHO 32	1	-	-	9± 0,0 ^{ed}	-	-	-
AHO 36	1	-	-	-	-	-	10,5 ± 0,7 ^{ef}
AHO 38	2	-	-	-	-	18 ± 1,4 ^d	-
AHO 41	1	-	-	9,5± 0,7 ^{ed}	-	-	13 ± 1,4 ^{de}
AHO 43	2	-	-	52,5 ± 3,5 ^a	38 ± 2,8 ^a	33,5 ± 2,1 ^c	11 ± 0,0 ^{def}
AHO 44	1	-	-	13 ± 0,7 ^c	-	-	-
AHO 51	1	-	-	14 ± 0,7 ^c	-	-	10 ± 0,0 ^f
NA 30	-	18 (S)	23 (S)	-	-	-	-
NET 30	-	-	-	27 (S)	-	-	-
FA 10	-	-	-	-	25 (S)	-	-
NY 100	-	-	-	-	-	25 (S)	-

(-): no inhibition; (S): sensitive, NA: Nalidixic acid, NET: Netilmicin, FA: Fusidic acid, NY: Nystatin

* The data in the same column followed by the same letter don't show significant difference according to Anova test (P<0,05)

For the 8 endophytic actinomycetes, 2 isolates were active against one of the 2 tested fungi, *Fusarium sp* (table 2). This result concurs with the finding of Taechowisan and Lumyong [21] who reported that endophytic actinomycetes isolated from *Zingiber officinale* and *Alpinia galanga* exhibited fungicide activity against phytopathogens (*Colletotrichum musae* and *Fusarium oxysporum*). Furthermore, previous works showed antagonistic effect of endophytic actinomycetes against many varieties of fungi such as *Alternaria*, *Rhizoctonia*, *Verticillium*, *Fusarium*, *Phytophthora* and *Phytium spp* [42, 43, 44, 45, 11, 46].



Effect of AHO 43 on *Candida albicans*



Effect of AHO 12 on *Staphylococcus aureus*

a: Acar and Goldstein's method; b: modified method

Figure 3. Inhibition of pathogens growth by actinomycetes strains.

3.2.2. Antioxidant Activity of Selected Actinomycetes Extracts

Six actinomycetes isolates (AHO 1, AHO 3, AHO 12, AHO 18, AHO 24 and AHO 43) with large spectrum of antimicrobial activity were kept for metabolites extraction. The extracts were, then, screened for their antioxidant, antiplasmodial and antiproliferative activities. Among the 6 extracts, 2 extracts from the isolates AHO 3 and AHO 43 showed antioxidant activity with IC₅₀ values of 15±1,5mg/ml and 10,6±0,7mg/ml, respectively. It was reported that the majority of actinomycetes extracts possessed weak antioxidant activity than ascorbic acid [47, 48]. This result is consistent with our findings in which antioxidant activity of the isolates AHO 3 and AHO 43 was 1,53 times and 1,08 times, respectively, lower than ascorbic acid activity which IC₅₀ value is 9,8±0,7mg/ml.

3.2.3. Antiproliferative of Actinomycetes Extracts

Of the 6 actinomycetes extracts screened for their antiproliferative activity, the two same extracts from the isolates AHO 3 and AHO 43 exhibited a strong cytotoxic activity on colon cancer cells HT-20 with IC₅₀ values of 5±0,5µg/ml and 2,2±0,2µg/ml, respectively according to the criteria of National Cancer Institute [49]. These results corroborate those of many studies which indicate that actinomycetes constitute an important source of antiproliferative compounds [5, 50, 51]. The two extracts

were, yet, less active than taxol (standard) which IC₅₀ value is 0,0082±0,003µg/ml).

3.2.4. Antiplasmodial Activity of Actinomycetes Extracts

Even though, the study on the evaluation of actinomycetes antiplasmodial activity is yet scarce, the 2 extracts from the isolates AHO 3 and AHO 43 showed very strong activity against *Plasmodium falciparum* Dd2 chloroquino-resistant. The IC₅₀ values were 1,25±0,04µg/ml for the extract AHO 3 and 2,5<IC₅₀<5µg/ml for the extract AHO 43. The two extracts were also less active than artemisinin (standard) which IC₅₀ value is 0,7±0,02µg/ml.

3.3. Biological Activities of Ginger Essential Oil

3.3.1. Aromatogram

The results of aromatogram assay showed that all tests-pathogens were sensitive to both essential oils extracted from ginger rhizomes of Ampamaha and Andrefaniviny plantations (figure 4). *Bacillus cereus* is very sensitive to both essential oils H1 and H2 while *Candida albicans* is extremely sensitive to essential oil H1 and very sensitive to essential oil H2 according to the criteria of Ponce *et al.* [52]. These results were in agreement with Oussalah *et al.* [53] who demonstrated in their work on a large range of essential oils screened for their potential antimicrobial that ginger essential oil is one of the most efficient essential oils against fungi.




Figure 4. Antimicrobial activity of essential oils extracted from ginger rhizomes samples.

The bar indicates the mean standard error for each essential oil

H1: essential oil extracted from Ampamaha ginger rhizomes

H2: essential oil extracted from Andrefaniviny ginger rhizomes

3.3.2. Antioxidant Activity of Ginger Essential Oil

The two essential oils H1 and H2 showed antioxidant activity with IC₅₀ values of 19,5mg/ml and 19mg/ml, respectively. It emphasizes what Kikuzaki and Nakatani [54] reported, about forty antioxidant compounds were identified in ginger.

3.3.3. Antiproliferative and Antiplasmodial Activities of Ginger Essential Oil

Among the two essential oils screened for their antiproliferative activity, only the essential oil H2 exhibited a moderate activity on colon cancer cells HT-20 with IC₅₀ value of 14±4,0µg/ml. In spite of this, anticancer potential of ginger essential oil has been demonstrated in several works [55, 56]. In addition, this essential oil showed a very strong antiplasmodial activity with IC₅₀ value included between 2,5 and 5µg/ml. In our knowledge, antiplasmodial potential of ginger was demonstrated for the first time.

From these results, it could be deduced that the endophytic actinomycete AHO 3 and the ginger rhizospheric soil actinomycetes AHO 43 showed comparable biological

activities as the essential oil H2 of ginger rhizomes. However, the isolate AHO 43 possessed the closest biological activities as ginger essential oil. Moreover, the biological activities of these 2 isolates were more accentuated than those of ginger essential oil. For the antimicrobial activity, the diameters of inhibition zone of the isolate AHO 43 were 2 or 3 times higher than those of ginger essential oil. For the antioxidant, antiproliferative and antiplasmodial activities, all the IC₅₀ values of the two actinomycetes extracts were higher than those of ginger essential oil (table 3). Zhao *et al.* [3] obtained a comparable result in which cytotoxicity level of the cajanol produced by the endophytic fungus (*Hypocrea lixii*) of *Cajanus cajan* roots against lung cancer cells A-549 was higher than the cajanol produced by the host plant.

Compared to the standards used for each activity (ascorbic acid, taxol and artemisinin), the activity of the two extracts from the isolates AHO 3 and AHO 43 were low. Nevertheless, it is important to emphasize that the tested extracts were yet crude extracts whereas all standards used were pure compounds.

Table 3. Recapitulation of the biological activities of actinomycetes isolates (AHO 3, AHO 43) and ginger essential oil H2 of Andrefaniviny plantation.

Isolates/Extracts	Antimicrobial activity (diameters of inhibition zone in mm)						Antioxidant activity (IC ₅₀ mg/ml)	Antimalaria activity (IC ₅₀ µg/ml)	Antiproliferative activity (IC ₅₀ µg/ml)
	<i>E. cloacae</i>	<i>K. oxytoca</i>	<i>B. cereus</i>	<i>S. aureus</i>	<i>C. albicans</i>	<i>Fusarium sp.</i>			
AHO 3 (site 2)	6	6	6	6	6	17,5	15	1,25	5
AHO 43 (site 2)	6	6	38	52,5	33,5	11	10,6	1,25<x<2,5	2,2
H2	9	10	15,5	14,5	16,5	9	19,00	2,5<x<5	14

E: *Enterobacter*, K: *Klebsiella*, S: *Staphylococcus*, B: *Bacillus*, C: *Candida*, -: no activity

For the case of Ampamaha ginger plantation; none of endophytes or telluric actinomycetes isolated showed comparable biological activities to the essential oil H1. This could be due to the isolation method where the diversity of actinomycetes isolated is widely dependent on [57].

3.4. Identification of Active Isolates

Phenotypic and molecular characterization using 16S rRNA gene sequence of the active isolates showed that they belong to the genus *Streptomyces* as reported previously [23]. Thus, the isolate AHO 3 was identified as *Streptomyces chrysomallus* and the isolate AHO 43 as *Streptomyces sp* (figure 5).




Figure 5. Neighbor-joining tree based on partial 16S rRNA gene sequences of active isolates.

3.5. Chemical Screening of *Streptomyces sp* Extract

As the essential oil H2 and the ethanolic extract of *Streptomyces sp* were closely comparable in term of biological activity, this work suggested determining the chemical family of the active compounds in the extract. Thus, the results revealed the presence of sterols, triterpens, tannins, leucoanthocyanes, flavonoids and alkaloids (table 4). According to several studies, these compounds have been demonstrated to possess important biological potentials. The alkaloids exhibit particular pharmacological activities as

antitumor and antiplasmodial [58, 59, 60]. The triterpens have been demonstrated to have antiplasmodial and anticancer activities [61, 62]. The sterols are endowed of antitumoral properties [63].

Tannins and flavonoids are phenolic compounds with various biological activities. The flavonoids display antiplasmodial activity [64] whereas the tannins present antibacterial, antifungal, antioxidant and antitumor activities [65, 66, 67]. Concerning the leucoanthocyanes, these compounds have been demonstrated to have bactericide activity [68].

Table 4. Chemical profile of *Streptomyces sp* extract.

Chemical families	Tests	Observations	Results
ALKALOIDS	Mayer	Abundant precipitation	++
	Wagner	Weak precipitation	+
	Dragendorff	Weak precipitation	+
FLAVONOIDES	Willstätter	Red purple coloration	+
	Willstätter modified	Red purple coloration	+
LEUCOANTHOCYANES	Bate-Smith	Red purplish coloration	+
TERPENOIDES	Liebermann Burchard	Pink coloration	++
STEROIDES	Salkowski	Red partition ring	++
	Badjet-Kedde	Green coloration	-
ANTHRAQUINONES	Borntrager	No change of coloration	-
SAPONOSIDES	Height of the foam	At t=0 (h=5cm); at t=30min (h=2cm)	-
POLYPHENOLS	Gelatine 1%	No precipitation	-
TANNINS	Salted gelatine	Formation of precipitate	++
	FeCl ₃	Black blue solution	++
POLYSACCHARIDES	-	No precipitation	-

-: absence, +: presence, ++: abundance

3.6. Chemical Composition of the Essential Oil H2

The results from Gas Chromatography revealed that 96,28% of the compounds of the essential oil H2 were identified: 60,88% are terpenic compounds and 35,40% oxygenated compounds. The terpenic compounds are zingiberene (14,48%), β -phellandrene (8,36%), camphene (7,98%) and β -sesquiphellandrene (6,45%). However, the

majority compounds of oxygenated compounds are geranial or citral a (9,35%), neral or citral b (6,23%), eucalyptol (3,57%) and borneol (2,09%). The percentage of α -terpineol, elemol and zingiberenol is 1-2%. Unknown products are 3,72% and don't show any majority products (table 5).

Table 5. Components of ginger rhizomes essential oil from Andrefaniviny by Gas chromatography.

Compounds	Retention time (min)	% (w/w)	Compounds	Retention time (min)	% (w/w)
2-Heptanone	4,498	0,01	Cyclosativene	13,879	0,15
2-n-Butyl furan	4,539	0,01	Geranyl acetate	14,025	0,58
2-Heptanol	4,622	0,07	α -Copaene	14,038	0,19
Tricyclene	5,103	0,15	iso- β -Elemene	14,181	0,02
α -Thujene	5,157	0,01	cis- β -Elemene	14,312	0,67
α -Pinene	5,3	2,79	7-epi-Sesquithujene	14,507	0,12
Camphene	5,588	7,98	β -Caryophyllen	14,882	0,05
Cosmene	5,829	0,02	γ -Elemene	15,061	0,17
Sabinene	6,011	0,17	Sesquisabinene A (RI 1454)	15,212	0,02
β -Pinene	6,096	0,38	(E) Isoeugenol	15,309	0,07
6-Methyl-5-heptene-2-one	6,193	0,21	(E)- β -Farnesene	15,387	0,22
Myrcene	6,284	1,21	Sesquisabinene B (RI 1467.3)	15,441	0,13
Octanal	6,505	0,08	α -Humulene	15,496	0,01
α -Phellandrene	6,594	0,54	allo-Aromadendrene	15,632	0,18
3-Carene	6,717	0,03	Selina-4,11-diene	15,862	0,20
α -Terpinene	6,831	0,02	Ar-Curcumene	15,926	3,88
p-Cymene	6,987	0,14	Germacrene D	15,984	1,16
Limonene	7,115	0,87	Tridecan-2-one	16,08	0,01
β -Phellandrene	7,112	8,26	Zingiberene	16,171	14,48
Eucalyptol	7,148	3,57	α -Selinene	16,228	0,70
2-Heptanol, acetate	7,246	0,03	α -Farnesene	16,32	3,60
cis- β -Ocimene	7,407	0,00	β -Bisabolene	16,381	3,30
Melonal	7,528	0,01	γ -Cadinene	16,53	0,25
(E)-2-Octen-1-al	7,607	0,03	epi-Cubebol	16,574	0,20
γ -Terpinene	7,661	0,04	β -Sesquiphellandrene	16,664	6,45
trans-Linalool oxide	7,943	0,00	(E)- γ -Bisabolene	16,791	0,20
Terpinolene	8,271	0,32	Unknown	17,005	0,21
2-Nonanone	8,277	0,20	Elemol	17,117	1,43
Inconnu	8,399	0,04	trans-Sesquisabinene hydrate	17,142	0,38
2-Nonanol	8,432	0,26	trans-Nerolidol	17,262	0,82
Linalool	8,447	0,92	Germacrene B	17,331	0,38
Perillene = Furan, 3-(4-methyl-3-pentenyl)	8,49	0,01	Germacrene D-4-ol	17,608	0,11
(E)-4,8-Dimethylnona-1,3,7-triene	8,788	0,06	cis-Sesquisabinene hydrate	17,769	0,96
Bicyclo [2.2.1] heptane, 2-methoxy-1, 7,7-trimethyl	8,814	0,05	Zingiberenol	18,16	1,45
cis-2-Menthenol	8,952	0,09	epi- γ -Eudesmol	18,376	0,21
Inconnu	8,984	0,04	Zingiberenol isomer	18,433	0,67
trans-2-Menthenol	9,311	0,04	Unknown	18,499	0,41
Camphor	9,475	0,09	γ -Eudesmol	18,537	0,11
Citronellal	9,543	0,39	α -Cadinol	18,685	0,06
Isobornéol	9,71	0,05	β -Eudesmol	18,868	0,73
cis- β -Terpineol	9,81	0,03	Unknown	18,983	0,08
Borneol	9,89	2,09	Intermedeol	19,03	0,08
Terpinen-4-ol	10,107	0,21	β -Bisabolol	19,08	0,08
p-Cymen-8-ol	10,23	0,03	Unknown	19,296	0,08
Cryptone	10,31	0,03	Unknown	19,361	1,24
α -Terpineol	10,365	1,22	Unknown	19,416	0,09

Compounds	Retention time (min)	% (w/w)	Compounds	Retention time (min)	% (w/w)
cis-Piperitol	10,466	0,01	Unknown	19,495	0,45
Myrtenol	10,493	0,09	Unknown	19,525	0,48
n-Decanal	10,592	0,08	Unknown	19,756	0,17
trans-Piperitol	10,694	0,04	Unknown	20,01	0,10
Citronellol	11,033	0,64	trans, trans-Farnesal	20,164	0,16
Nerol	11,064	0,11	Xanthorrhizol	20,291	0,06
3-Methyl-3-(4-methylpent-3-en-1-yl) oxirane-2-carbaldehyde	11,18	0,04	Unknown	20,355	0,22
Citral b=Neral	11,364	6,23	Unknown	20,904	0,05
Geraniol	11,564	1,09	Unknown	20,956	0,02
2-Decenal, (2E)-	11,722	0,06	Unknown	21,191	0,16
Citral a=Geranial	11,949	9,35	Unknown	22,081	0,04
Borneol, acetate	12,276	0,17	Unknown	22,42	0,02
2-Undecanone	12,328	0,48	Geranyl-p-cymene	23,304	0,02
2-Undecanol	12,451	0,05	[6]-Gingerone	26,931	0,01
Myrtenyl acetate	13,013	0,00	[6]-Shogaol	27,661	0,04
δ -Elemene	13,278	0,10	[6]-Gingerol	28,805	0,00
Citronellyl acetate	13,452	0,08	[8]-Shogaol	30,086	0,00
Neryl acetate	13,669	0,00			

These different constituents have been also demonstrated to show a large biological activity spectrum. For examples, β -sesquiphellandrene is antitumor [69]; the citrals (geranial and neral) and camphene possess bactericide and antioxidant effects [70, 71]. Moreover, it would be emphasized that biological activity of the essential oils are not only from their majority compounds. Other minority compounds can develop an efficient system with synergic or antagonistic reaction against target strains [72].

Thus, biological activities of *Streptomyces sp* extract and the essential oil H2 which were at the same time antimicrobial, antioxidant, antiplasmodial and antiproliferative, are linked to the chemical components of their secondary metabolites. One or several chemical components of the extract and the essential oil can display, therefore, one or several activities demonstrated above. Accordingly, it was observed that *Streptomyces sp* extract and the essential oil H2 contained together as common chemical components the terpenes and the phenolic compounds.

4. Conclusion

From this work, it would be concluded that two actinomycetes associated to ginger plant (*Streptomyces chrysomallus*, endophyte and *Streptomyces sp*, from ginger rhizospheric soil) present high pharmacological potential and great therapeutic value as the active extract (essential oil) of the medicinal plant. These bacteria showed accentuated antimicrobial, antioxidant, antiplasmodial and antiproliferative activities than ginger essential oil which prove that actinomycetes constitute important sources of secondary metabolites with wide biological and structural diversity. Further investigation is however, required for isolation, identification, *in vitro* and *in vivo* assays of the active compounds produced by the two isolates. Likewise, this work reports the presence of rhizospheric soil actinomycetes showing comparable biological activities as

those of plant active extract. Anyway, these findings provide evidence that plant associated actinomycetes could be taken as sustainable alternatives for drugs production and for plant preservation.

References

- [1] Amna T., Puri S. C., Verma V., Sharma J. P., Khajuria R. K., Musarrat T. J., Spitteller M. and Qazi G. N. (2006). Bioreactor studies on the endophytic fungus *Entrophospora*
- [2] Shweta S., Zuehlke S., Ramesha B. T., Priti V., Mohana Kumar P., Ravikanth G., Spitteller M., Vasudeva R. and Uma Shaanker R. (2010). Endophytic fungal strains of *Fusarium solani*, from *Apodytes dimidiata* E. Mey. Ex Arn (Icacinaeace) produce camptothecin, 10-hydroxycamptothecin and 9-methoxycamptothecin. *Phytochemistry*, 71: 117-22.
- [3] Zhao J., Li C., Wang W., Zhao C., Luo M., Mu F. *et al.* (2013). *Hypocrea lixii*, novel endophytic fungi producing anticancer agent cajanol, isolated from pigeon pea (*Cajanus cajan* [L] Mill sp.). *J. Appl. Microb.*, 115: 102-113.
- [4] Huang J. X., Zhang J., Zhang X. R., Zhang K., Zhang X. *et al.* (2014). *Mucorfragilis* as a novel source of the key of pharmaceutical agents podophyllotoxin and kaempferol. *PharmBiol.*, 52: 1237- 1243.
- [5] Caruso M., Colombo A. L., Fedeli L., Pavesi A., Quaroni S., Saracchi M., and Ventrella G. (2000). Isolation of endophytic fungi and actinomycetes taxane producers. *Ann. Microbiol.* 50: 1-13.
- [6] Akshatha V. J., Nalini M. S., D' Souza, Prakash H. S. (2014). Streptomycete endophytes from anti-diabetic medicinal plants of the Western Ghats inhibit alpha-amylase and promote glucose uptake. *Lett Appl Microbiol.*, 58: 433-439.
- [7] Newman D. J., Cragg G. M. and Snader K. M. (2003). Natural Products as sources of new drugs over the period 1981-2002. *J. Nat. Prod.*, 66 (7): 1022-1037.

- [8] Oskay M., Tamer A. U. and Azeri C. (2004). Antibacterial activity of some actinomycetes isolated from farming soils of Turkey. *African Journal of Biotechnology*, 3 (9): 441-446.
- [9] Jeffrey L. S. H. (2008). Isolation, characterization and identification of actinomycetes from agriculture soils at Semongok, Sarawak. *African Journal of Biotechnology*, 7 (20): 3697-3702.
- [10] Igarashi Y., Iida T., Sasaki T., Saito N., Yoshida R., and Furumai T. (2002). Isolation of actinomycetes from live plants and evaluation of anti-phytopathogenic activity of their metabolites. *Actinomycetologica*, 16 (1): 9-13.
- [11] Inderiati S. and Franco C. M. M. (2008). Isolation and Identification of Endophytic Actinomycetes and their Antifungal Activity. *Journal of Biotechnology Research in Tropical Region*, 1: 1-6.
- [12] Boughachiche F., Reghioia S., Oulmi L., Zerizer H., Kitouni M., Boudemagh A., Boulahrouf A. (2005). Isolement d'actinomycétales productrices de substances antimicrobiennes à partir de la Sebkhia de Ain Mlila. *Sciences & Technologie C*, 5-10.
- [13] Kumar S. M. and Selvam. K. (2011). Prospective efficacy of bioactive extract of novel actinomycetes against malignant cell and multidrug resistant bacteria. *Pharmacology on line*, 1: 1262-1290.
- [14] Zerizer H., Oulmi L., Boughachiche F., Reghioia S., Boudemagh A., Kitouni M. and Boulahrouf A. (2006). Identification d'une actinomycétale, productrice d'antibactériens, isolée de sols arides de la région de Biskra. *Sciences & Technologie*, 24: 17-22.
- [15] Butt M. S and Sultan M. T. (2011). Ginger and its health claims: molecular aspects. *Crit Rev Food Sci Nutr*, 51 (5): 383-393.
- [16] Katzer G. (2012). Ginger (*Zingiberofficinalerose*). gernet-katzers-spice-pages.com.
- [17] Chaiyakunapruk N., Kitikannakorn N. et al. (2006). The efficacy of ginger for the prevention of postoperative nausea and vomiting: a meta-analysis. *Am J Obstet Gynecol.*, 194 (1): 95-99.
- [18] Apariman S., Ratchanon S., Wiriyasirivej B. (2006). Effectiveness of ginger for prevention of nausea and vomiting after gynecological laparoscopy. *J Med Assoc Thai*. 89 (12): 2003-2009.
- [19] Jolad S. D., Lantz R. C. et al. (2005). Commercially processed dry ginger (*Zingiber officinale*): composition and effects on LPS-stimulated PGE2 production. *Phytochemistry*, 66 (13): 1614-1635.
- [20] Karna P., Chagani S., Gundala S. R., Rida P. C., Asif G., Sharma V., Gupta M. V. and Aneja R. (2012). Benefits of whole ginger extract in prostate cancer. *Br J Nutr.*, 107 (4): 473-484.
- [21] Taechowisan T. and Lumyong S. (2003). Activity of endophytic actinomycetes from roots of *Zingiber officinale* and *Alpinia galanga* against phytopathogenic fungi. *Annals of Microbiology*, 53 (3): 291-298.
- [22] Jasim B., Aswathy Agnes Joseph, Jimtha John C., Mathew J., Radhakrishnan E. K. (2013). Isolation and characterization of plant growth promoting endophytic bacteria from the rhizome of *Zingiber officinale*. *3 Biotech.*, 8p.
- [23] Andriambeloson O., Rasolomampianina R. and Rahehimandimby M. (2014). Selection and characterization of bioactive actinomycetes associated with the medicinal plant Ginger (*Zingiber officinale*). *Journal of International Academic Research for Multidisciplinary*. 2 (9): 30-45.
- [24] Fisher P. J., Petrini O. and Lappin-Scott H. M. (1992). The distribution of some fungal and bacteria endophytes in maize (*Zea mays* L.) *New Phytol.* 122: 299-305.
- [25] Pereira J. O., Azevedo J. L. and Petrini O. (1993). Endophytic fungi from stylosantes: a preliminary report. *Mycologia*. 85: 362-364.
- [26] Pereira J. O., Carneiro-Vieira M. L. and Azevedo J. L. (1999). Endophytic fungi from *Musa acuminata* and their reintroduction in axenic plants. *World J. Microbiol. Biotechnol.* 15: 47-51.
- [27] Acar J. F. and Goldstein F. W. (1996). Disk Susceptibility Test. In: V. Lorian. "Antibiotics in Laboratory Medicine". Baltimore: William and Wilkins Co. 4th Edt: 1-51.
- [28] Loqman S., Barka E. A., Clément C. and Ouhdouch Y. (2009). Antagonistic actinomycetes from Moroccan soil to control the grapevine gray mold. *World Journal of Microbiology and Biotechnology*, 28: 81-91.
- [29] Leitao G. G., Leitao S. G. and Vigelac W. (2002). Quick preparative separation of natural naphthopyranones with antioxidant activity by high-speed counter-current chromatography. *Z Naturforsch.* 57: 1051-1055.
- [30] Sharififar F., Moshafi M. H., Mansouri S. H., Khodashenas M. and Khoshnoodi M. (2007). *In vitro* evaluation of antibacterial and antioxidant activities of the essential oil and methanol extract of endemic *Zataria multiflora* Boiss. *Food Control*, 18: 800-805.
- [31] Cao S., Brodie P. J., Miller J. S., Randrianaivo R., Ratovoson F., Birkinshaw C., Andriantsiferana R., Rasamison V. E., and Kingston D. G. I. (2007). Antiproliferative Xanthones of *Terminalia calcicola* from the Madagascar Rain Forest. *J. Nat. Prod.*, 70: 679-681.
- [32] Harinantenaina L., Bowman J. D., Brodie P. J., Slebodnick C., Callmender M. W., Rakotobe E., Randrianaivo R., Rasamison V. E., Gorka A., Roepe D., Cassera M. B. and Kingston D. G. I. (2012). Antiproliferative and Antiplasmodial Dimeric Phloroglucinols from *Mallotus oppositifolius* from the Madagascar Dry Forest. *Journal of natural products*, A-F.
- [33] Fong H. H. S., Tin W. A. M. and Farnsworth N. R. (1977). Phytochemical screening. Review. Chicago: University of Illinois, 73-126.
- [34] Intra B., Mungsuntisuk I., Nihira T., Igarashi Y. and Panbangred W. (2011). Identification of actinomycetes from plant rhizospheric soils with inhibitory activity against *Colletotrichum. spp*, the causative agent of anthracnose disease. *BMC Research Notes*, 4: 98.
- [35] Crawford D. L., Lynch J. M. and Ousley M. A. (1993). Isolation and characterisation of actinomycetes antagonists of a fungal root pathogen. *Appl. Environm. Microbiol.* 59 (11): 899-905.

- [36] Zinniel D. K., Lambrecht P., Harris N. B., Feng Z., Kuczumski D., Higley P., Ishimaru C. A., Arunakumari A., Barletta R. G. and Vidaver A. K. (2002). Isolation and characterization of endophytic colonizing bacteria from agronomic crops and prairie plants. *Appl Environ Microbiol*, 68: 2198-2208.
- [37] Hasegawa S., Meguro A., Shimizu M., Nishimura T. and Kunoh H. (2006). Endophytic actinomycetes and their interactions with host plants. *Actinomycetologica*, 20: 72-81.
- [38] Gangwar M., Dogra S., Gupta U. P. and Kharwar R. N. (2014). Diversity and biopotential of endophytic actinomycetes from three medicinal plants in India. *African Journal of Microbiology Research*, 8 (2): 184-191.
- [39] Basilio A., González I., Vicente M. F., Gorrochategui J., Cabello A., González A. and Genilloud O. (2003). Patterns of antimicrobial activities from soil actinomycetes isolated under different conditions of pH and salinity. *J. Appl. Microbiol.*, 95: 814-823.
- [40] Sacramento D. R., Coelho R. R. R., Wigg M. D., Linhares L. F. T. L., Santos M. G. M., Semêdo L. T. A. S. and Silva A. J. R. (2004). Antimicrobial and antiviral activities of an actinomycete (*Streptomyces* sp.) isolated from a Brazilian tropical forest soil. *World J. Microbiol. Biotechnol.*, 20: 225-229.
- [41] Scherrer R. and Gerhardt, P. (1971). Molecular sieving by the *Bacillum megaterium* cell wall and protoplast. *Journal of Bacteriology*, 107: 718-735.
- [42] Yuan, W. M. and Crawford D. L. (1995). Characterization of *Streptomyces lydicus* WYEC 108 as a Potential Biocontrol Agent against Fungal Root and Seed Rots. *Appl. Environ. Microbiol.*, 61: 3119-3128.
- [43] El-Tarabily K. A., Hardy G. E. St. J., Sivasithamparam K., Hussein A. M. and Kurtboke D. I. (1997). The Potential for the Biological Control of Cavity-Spot Disease of Carrots, Caused by *Pythium coloratum*, by Streptomyces and NonStreptomyces Actinomycetes. *New Phytol.*, 137: 495-507.
- [44] Getha K. and Vikineswary. S. (2002). Antagonistic Effects of *Streptomyces violaceusniger* Strain G10 an *Fusarium oxysporum* f. sp. cubense Race 4: Indirect Evidence for the Role of Antibiotics in the Antagonistic Process. *J. Industrial Microbiol. Biotech.*, 28: 303-310.
- [45] Aghighi S., Bonjar G. H. S., Rawashdeh R., Batayneh S. and Saadoun I. (2004). First Report of Antifungal Spectra of Activity of Iranian Actinomycetes Strains against *Alternaria solani*, *Alternata alternate*, *Fusarium solani*, *Phytophthora megasperma*, *Verticillium dahliae* and *Saccharomyces cerevisia*. *Asian J. Plant Sci.*, 3: 463-471.
- [46] Verma V. C.; Gond S. K.; Kumar A.; Mishra A.; Kharwar R. N. and Gange A. C. (2009). Endophytic actinomycetes from *Azadirachta indica* A. Juss.: Isolation, diversity, and antimicrobial activity. *Microbial Ecol.*, 57: 749-756.
- [47] Zhong K., Gao X. L., Xu Z. J., Gao H., Fan S., Yamaguchi I., Li L. H. and Chen R. J. (2011). Antioxidant activity of a novel *Streptomyces* strain Eri 12 isolated from the Rhizosphere of *Rhizoma curcumae longae*. *Cur Res Bacteriol*, 4: 63-72.
- [48] Thenmozhi M. and Kannabiran. (2012). Antimicrobial and antioxidant properties of marine actinomycetes *Streptomyces* sp VITSTK7. *Oxid Antioxid Med Sci.*, 1 (1): 51-57.
- [49] Boyd M. R. (1997). Preclinical Screening, Clinical Trials, and Approval in: Teicher, B. A. *Anticancer Drug Development Guide*. Totowa: Ed. Humana Press, 23-42.
- [50] Kwon H. C., Kauffman C. A. and Jensen P. R. (2006). Marinomycins a-d, antitumor antibiotics of a new structure class from a marine actinomycete of the recently discovered genus *Marinospora*. *Journal of the American Chemical Society*, 128: 1622-1632.
- [51] Malet-Gason L., Romero F., Espliego-Vazquez F., Gravalos D. and Fernandez-Puentes J. L. (2009). IB-00208, a new cytotoxic polycyclic xanthone produced by a marine-derived *Actinomadura*. *The Journal of Antibiotics*, 56: 219-225.
- [52] Ponce A. G., Fritz R., Del Valle C. and Roura S. I. (2003). Antimicrobial activity of essential oils on the native microflora of organic Swiss chard. *Lebensmittel- Wissenschaft und Technologie*, 36: 679-684.
- [53] Oussalah M., Caillet S., Saucier L. and Lacroix M. (2007). Inhibitory effects of selected plant essential oils on four pathogen bacteria growth: *E. coli* O157: H7, *Salmonella typhimurium*, *Staphylococcus aureus* and *Listeria monocytogenes*. *Food Control.*, 18 (5): 414-420.
- [54] Kikuzaki H. and Nakatani N. (1996). Cyclic diarylheptanoids from rhizomes of zingiber officinale. *Phytochemistry*, 43 (1): 273-277.
- [55] Kim E. C., Min J. K. *et al.* (2005). [6]-Gingerol, a pungent ingredient of ginger, inhibits angiogenesis *in vitro* and *in vivo*. *Biochem Biophys Res Commun.*, 23 (2): 300-308.
- [56] Aggarwal B. B. and Shishodia S. (2006). Molecular targets of dietary agents for prevention and therapy of cancer. *Biochem Pharmacol.*, 71 (10): 1397-1421.
- [57] Takahashi Y. and Omura S. (2003). Isolation of new actinomycete strains for the screening of new bioactive compounds. *Journal of Genetic Applied Microbiology*, 49: 141-154.
- [58] Wall M. E., Wani M. C. and Taylor H. (1987). Plant antitumor agents. Isolation, structure, and structure activity relationships of alkaloids from *Fagara macrophylla*. *Journal of Natural Products*, 50: 1095-1099.
- [59] Basco L. K., Mitaku S., Skaltsounis A. L., Ravelomanantsoa N., Tillequin F., Koch M. and Le Bras J. (1994). *In vitro* activities of furoquinoline and acridone alkaloids against *Plasmodium falciparum*. *Antimicrobial Agents and Chemotherapy*, 38: 1169-1171.
- [60] Gakunju D. M., Mberu E. K., Dossaji S. F., Gray A. I., Waigh R. D., Waterman P. G. and Watkins W. M. (1995). Potent antimalarial activity of the alkaloid nitidine, isolated from a Kenyan herbal remedy. *Antimicrobial Agents and Chemotherapy*, 39: 2606-2609.
- [61] Namukobe J., Kiremire B. T., Byamukama R., Kasenene J. M., Dumontet V., Guéritte F., Krief S., Florent I and Kabasa J. D. (2013). Nouveaux triterpènes à activité antiplasmodiale isolés des feuilles de *Neoboutonia macrocalyx* L., une plante consommée par les chimpanzés du parc national de Kibale (Ouganda). *Revue de primatologie*, 5.
- [62] Konoshima T., Takasaki M., Tokuda H., Masuda K., Arai Y., Shiojima K. and Ageta H. (1996). Anti-tumor-promoting activities of triterpenoids from ferns. *Biological and pharmaceutical bulletin*, 19 (7): 962-965.

- [63] Bok J. W., Lermer L., Chilton J., Klingeman H. G and Towers Neil G. H. (1999). Antitumor sterols from the mycelia of *Cordyceps sinensis*. *Phytochemistry*, 51: 891-898.
- [64] Chen M., Theander T. G., Christensen S. B., Hviid L., Zhai L. and Kharazmi A. (1994). Licochalcone A, a New Antimalarial Agent, Inhibits *In Vitro* Growth of the Human Malaria Parasite *Plasmodium falciparum* and Protects Mice from *P. yoelii* Infection. *Antimicrobial Agents and Chemotherapy*, 1470-1475.
- [65] Kolodzie J., Kayser O., Latte K P., Ferreira D. (1999). Evaluation of the antimicrobial potency of tannins and related compounds using the microdilution both method. *Planta medica*, 65 (5): 444-446.
- [66] Bouchet N., Barrier L. and Fauconneau B. (1998). Radical scavenging activity and antioxydant proprieties of tannins from *Guiera senegalensis* (Combretaceae). *Phytotherapyresearch*, 12 (3): 159-162.
- [67] Yoshida T. Hatano T., Miyamoto K., Okuda T., Brouillard R. and Jay M. (1995). Antitumor and related activities of ellagitannin oligomers. Les colloques: Polyphenols 94, Palma de Mallorca, 69: 123-132.
- [68] Velomalala N. M., Raherimandimby M., Ramamonjisoa D. (2013). Contribution à l'étude des propriétés biologiques des extraits de feuilles de *Senecio fujasioïdes* (Asteraceae). *Journal International de santé au travail*, 1: 33-40.
- [69] Ahn B. Z. and Lee. J. H. (1989). Cytotoxic and cytotoxicity-potentiating effects of the *Curcuma* root on L1210 cell. *Korean Journal of Pharmacognosy*, 20: 223-226.
- [70] Edris A. E. (2007). Pharmaceutical and therapeutic potentials of essential oils and their individual volatile constituents. *A reviewPhytother Res*, 21: 308-323.
- [71] Encheqroun H. K. B., Ghanmi M., Satrani B., Aafi A. and Chaouch A. (2012). Activité antimicrobienne des huiles essentielles d'*Artemisia esatlantica*, plante endémique du Maroc. *Bulletin de la Société Royale des Sciences de Liège*, 81: 4-21.
- [72] Sing R., Marimuthu P., De Heluani C. S and Catalan Ceser A. N. (2006). Antioxidant and biocidal Activities of *Carum nigrum* (seed) Essential oil, Oleoresin, and their selected components. *Journal of Agricultural and Food Chemistry*, 54: 174-181.