



Irradiation and Storage Temperature Influence the Physiological Changes and Shelf Life of Mango (*Mangifera indica* L.)

Manoj Kumar Yadav^{1, 5, *}, Kirtivardhan², Paramveer Singh³, Pushpraj Singh⁴

¹Department of Horticulture, Navsari Agricultural University, Navsari, India

²Department of Basic Science, Navsari Agricultural University, Navsari, India

³Bihar Agriculture College, Bihar Agricultural University, Sabour, India

⁴International Crop Research Institute for Semi-Arid Tropics, Patancheru, India

⁵Department of Horticulture and Food Processing, Office of the DDH, Civil Lines, Satna, India

Email address:

manoj_kadali11@hotmail.com (M. K. Yadav), kirtivardhan@nau.in (Kirtivardhan), shekhawatdeep@rediffmail.com (P. Singh), pushprajsngh9@gmail.com (P. Singh)

*Corresponding author

To cite this article:

Manoj Kumar Yadav, Kirtivardhan, Paramveer Singh, Pushpraj Singh. Irradiation and Storage Temperature Influence the Physiological Changes and Shelf Life of Mango (*Mangifera indica* L.). *American Journal of Plant Biology*. Vol. 2, No. 1, 2017, pp. 5-10.

doi: 10.11648/j.ajpb.20170201.12

Received: October 17, 2016; **Accepted:** November 23, 2016; **Published:** January 3, 2017

Abstract: In spite of being the leading producer of mango in the world, India registers the highest post harvest losses and minuscule exports. The reasons for this paradox are manifold. However, issues related to quarantine, shelf life and quality are the major stumbling blocks to trade, both national and international. Therefore, a balance between the required effective dose and tolerance of fruit to irradiation has to be investigated under various storage temperatures. The experiment was arranged with sixteen treatment combinations of irradiation doses (0.00kGy-0.60kGy) and similar to storage temperature (Ambient at 27±2°C and 60-70% RH, 9°C and 90% RH, 12°C and 90% RH and Control atmospheric (CA) storage (12°C, O₂ 2%, CO₂ 3% and RH 90%) from the year 2008-2010. The fruits were exposed to gamma radiation from the source of ⁶⁰Co. The data indicated that the fruits irradiated with 0.40kGy gamma rays recorded significantly minimum per cent reduction in physiological loss in weight, reduced ripening per cent, increased marketability of fruits and maximum average days to ripening and shelf life of fruits. Similar pattern were noticed when fruits kept at 9°C storage temperature and in combined effect of irradiation and storage conditions. Suggestions are made for maximizing storage potential by use of irradiation and adequate storage facilities for qualitative and hygiene produces.

Keywords: Alphonso Mango, Gamma Irradiation, Marketability, Ripening, Shelf Life, Storage Temperature

1. Introduction

Mangoes (*Mangifera indica* L.; family Anacardiaceae) are still judged as luxuries and expensive items in the markets of many industrialized countries. Asia accounts for 77% of global mango production and the Americas and Africa account for 13 and 19%, respectively [14]. India is the global leader in mango production [28]. The high cost of mangoes in importing countries is due to high air freight charges. Mangoes are classified as climacteric fruits and ripen rapidly after harvest. Mango is generally harvested when

physiologically mature and is allowed to ripen under suitable conditions of temperature and humidity. Therefore, if freshly harvested fruit is allowed to ripen at normal ambient conditions, ripening processes increase rapidly within the week [30]. As fruit, it's also remain alive after harvesting it is necessary to restrain their respiration and metabolic activity in order to maintain cell life and optimize post harvest quality by delayed ripening and senescence. For overcome these metabolic activities, innovation in irradiation and cold

storage is the new tools for the enhancement of physiological changes and health promoting components in most of climacteric fruits. Mango is susceptible to chilling injury and an optimum temperature of 12-13°C is generally recommended [7], [31]. Irradiation of fruits has been successfully shown to delay ripening in Papaya [16]. Irradiation is a physical process for the treatment of foods akin to conventional process like heating or freezing. It prevents food poisoning, reduces wastage to contamination and at the same time preserves quality [9]. In spite of being the leading producer of mango in the world, India registers the highest post harvest losses and minuscule exports. The reasons for this paradox are manifold. However, issues related to quarantine, shelf life and quality are the major stumbling blocks to trade, both national and international [30]. Therefore, a balance between the required effective dose and tolerance of fruit to irradiation has to be investigated under various storage temperatures. So, irradiation can be used in combination with low temperature for to assess the effects of different doses of gamma irradiation and storage temperature on reduction of the physiological loss in weight and ripening phenomena of fruit. The loss in weight of fruits is likely to reduce the marketability and quality of fruit drastically.

2. Materials and Methods

The experiment was at conducted at Department of Horticulture, Navinchandra Mafatlal College of Agriculture, Navsari agricultural University, Navsari, Gujarat from the year 2008-2010. Export grade mangoes of cv. Alphonso were harvested from University orchard. The selected mangoes from class I as per the quality parameters specified and described in "post harvest manual for mangoes" published by Agricultural Production and Export Development Authority. These fruits sorted by uniformity in size, maturity and freedom from defects. The fruits were kept in plastic crates with cushioned material and transported to cold storage of Post Harvest Technology Unit, Navsari Agricultural University, Navsari (Gujarat) India. Than after, fruits were again sorted to remove those with spotty and having bad appearance. The individual fruit weight was from 250-350g. The selected fruits wash with chlorine water and after drying fruits packed in corrugated fibre board boxes cushioned (CFB) with tissue paper. The dimension of CFB box was 370 X 275 X 90mm and gross weight of box with fruits was 3.0 kg. One box having nine fruits for each treatment and each treatment replicated thrice as per experimental design. The packed boxes kept in cold storage at 12°C for 8 hours for pre-cooling treatment. The time gap between harvesting and pre-cooling was not more than 6 hours.

2.1. Irradiation Treatment

After pre-cooling, fruits transport to irradiation treatment in air conditioned vehicle. It was carried out at Board of Radiation and Isotope Technology, Bhabha Atomic Research Centre, Mumbai (India). The fruits were exposed to gamma

radiation for different doses from the source radio isotope ^{60}Co with energy 1.33MeV. There were four irradiation doses *i.e.* I₁.0.00kGy (Unirradiated), I₂.0.20kGy, I₃.0.40kGy and I₄. 0.60kGy. The time gap from pre-cooling to irradiation was not more than 9 hours. After irradiation, fruits immediately transported to cold storage of university in air conditioned vehicle.

2.2. Storage Conditions

The boxes were kept in cold storage at different temperature as per storage temperature treatments *viz.*, Ambient at 27±2°C temperature and 65±5% relative humidity (S₁), 9°C temperature and 90% relative humidity (S₂), 12°C temperature and 90% relative humidity (S₁) and Control atmospheric storage at 12°C temperature, O₂ 2%, CO₂ 3% and 90% relative humidity (S₁). Post harvest biochemical changes of these fruits were studied by measuring the total soluble solids, sugars, acidity and ascorbic acid content of fruits.

2.3. Measurement Protocols

2.3.1. Determination of Physiological Parameters

i. Physiological Loss in Weight (Per Cent)

Four fruits from each treatment were weighted on first day of treatment and subsequently their weight was recorded from forth day to at six day interval up to the end of shelf life. The Physiological loss in weight was expressed in percentage and calculated as follows. $PLW \% = \frac{W_1 - W_2}{W_1} \times 100$, Where, W₁=initial weight and W₂=final weight [27]

ii. Ripening Per Cent

Ripening was measured by the number of fruits having change in colour from greenish to yellow and soft in texture were counted from four day of storage to at six day intervals up to the eating ripeness and expressed in percentage over total number of fruits taken for study.

2.3.2. Marketable Fruits Per Cent

The number of good quality and visibly sound fruits that can be marketed were counted and expressed as percentage over the total number of fruits at prescribed interval up to 90 per cent fruits has marketability.

2.3.3. Average Number of Days Taken to Ripening

The number of days taken by mango fruits in each treatment to attain eating ripeness by more than 80 per cent fruits of treatment was considered as days taken for ripening.

2.3.4. Shelf Life of Fruits

The number of days taken in each treatment to attain the maximum eating ripeness by all the fruits was computed considering the weighted mean. Each fruit was thoroughly scrutinized for any visible symptoms of spoilage and the end of shelf life was considered when the 90 per cent fruits were over ripened.

2.4. Statistical Analysis

Statistically the data obtained from experiment was analyzed using ANOVA for completely randomized design with factorial concept. Significance differences among treatments were compared using the Fisher's analysis of variance at the 5% probability level [15]. The data were subjected to appropriate transformation (arcsine) to meet the assumptions of normality.

3. Results and Discussion

3.1. Physiological Loss in Weight

The data indicated that the physiological loss in weight of fruits increased with the advancement of storage period and significantly influenced by irradiation and storage temperature. It is evident from Table 1, that the minimum reduction in physiological loss in weight was recorded in the fruits exposed with 0.40kGy irradiation and stored at 9°C

(I₃S₂) i.e. 0.932% at 6th day, 1.780% at 12th day, 2.760% at 18th day, 3.480% at 24th day, 4.730% at 30th day and 5.670% at 36th day of storage. The physiological loss of weight in fruits was possibly on account of loss of moisture through transpiration and utilization of some reserve food materials in the process of respiration [12]. The irradiation significantly reduced physiological loss in weight during storage period over control which might be attributed to reduction in utilization of reserve food material in the process of respiration [18]. The delay in respiration rate as a result of irradiation is also reported by [19] in guava. Similar findings were also observed by [29], [17] [5] in mango. Lower physiological loss in weight was noted in 9°C and 12°C and in CA (12°C) storage temperature which might be due to lesser water vapour deficit compared to ambient condition and the low temperature which had slowed down the metabolic activities like respiration and transpiration [27]. The observations accordance with the results of [23] in mango; [14] in banana and [8] in guava.

Table 1. Effect of Irradiation and storage conditions on physiological loss in weight of mango Cv. Alphonso during storage.

Physiological loss in weight (%) days after storage															
	4					10					16				
Source	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean
S ₁	3.493	2.783	2.633	2.743	2.913	12.197	7.463	7.170	7.243	8.518	0.000	10.310	10.820	11.490	8.150
S ₂	0.853	0.683	0.473	0.733	0.686	2.357	1.867	1.417	2.037	1.919	3.180	2.960	2.340	3.090	2.890
S ₃	0.873	0.753	0.613	0.837	0.769	2.557	1.990	1.640	2.217	2.108	3.870	3.090	2.570	3.370	3.220
S ₄	2.683	2.303	1.700	2.367	2.263	7.110	3.520	3.387	3.610	4.407	10.100	5.270	5.220	5.470	6.510
Mean	1.976	1.630	1.355	1.670		6.055	3.710	3.403	3.777		4.290	5.410	5.250	5.850	
Source	I	S		I X S		I	S		I X S		I	S		I X S	
S. Em ±	0.0035	0.003		0.007		0.005	0.005		0.010		0.010	0.010		0.021	
CD (P≤0.05)	0.010	0.010		0.020		0.015	0.015		0.030		0.031	0.031		0.062	

Table 1. Cont.

Physiological loss in weight (%) days after storage															
	22					28					34				
Source	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean
S ₁	0.000	13.470	12.650	13.810	9.980	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
S ₂	4.650	4.230	3.230	3.360	3.870	6.410	5.440	4.440	5.760	5.520	7.930	6.990	5.500	7.090	6.880
S ₃	4.420	4.370	3.430	4.420	4.160	6.940	5.690	4.610	5.970	5.800	0.000	7.230	5.510	7.520	5.060
S ₄	13.320	7.320	6.940	7.990	8.890	0.000	9.300	9.200	5.490	5.990	0.000	0.000	0.000	0.000	0.000
Mean	5.600	7.350	6.850	7.100		3.340	5.110	4.560	4.300		1.980	3.550	2.750	3.650	
Source	I	S		I X S		I	S		I X S		I	S		I X S	
S. Em ±	0.005	0.005		0.010		0.005	0.005		0.011		0.005	0.005		0.010	
CD (P≤0.05)	0.16	0.16		0.031		0.016	0.016		0.033		0.015	0.015		0.030	

Where I=irradiation, S=Storage temperature, CA=Control atmospheric storage

3.2. Ripening Per Cent

The Table 2, indicated that irradiated fruits significantly delayed the ripening process over unirradiated fruits irrespective of storage condition and not fully ripen up to 30th day of storage at 9°C. The fruits exposed to gamma rays (0.40kGy) and stored at 9°C and 12°C were remained unripe (I₃S₂ and I₃S₃) on 30th day of storage, the gamma rays @ 0.40kGy (I₃) exposed fruits showed 50.10 and 52.00 per cent ripening, respectively at 9°C and 12°C (S₃) storage. Rest of the treatments has high ripening or discarded due to complete of their shelf life. The unirradiated mangoes had early ripeness whereas; gamma rays exposed mangoes had a

significantly delayed ripening. The possible mechanisms that have been postulated include: a) irradiations results in decreased sensitivity to ripening action of ethylene and b) alteration in carbohydrates metabolism by regulating certain key enzymes, which interfere with production of ATP which is required for various synthetic processes during ripening [22]. Same findings noted by [26] and [6] in mango and [1] in banana. The lower and delayed ripening was noted at 9°C and 12°C and in CA (12°C) storage as compared to ambient temperature. The decrease of ripening per cent and increase in days for ripening at low temperature may be due to desirable inhibition of enzymatic activities leading reduction in the respiration and ethylene production. These results are

supported by [11] in mango and [2] in banana. The minimum and delayed ripening in fruits due to exposed to gamma rays and stored at low temperature might be due to the joint balancing effect of irradiation and low temperature [26].

Table 2. Effect of Irradiation and storage conditions on ripening of mango Cv. Alphonso during storage.

Ripening (%) days after storage															
Source	10					16					22				
	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean
S ₁	93.88	0.000	0.000	0.000	23.470	0.000*	71.630	69.900	72.920	53.610	0.00*	0.00*	0.00*	0.00*	0.00
S ₂	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.00	0.00	0.00	0.00	0.00
S ₃	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.00	0.00	0.00	0.00	0.00
S ₄	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	77.15	69.36	64.94	73.17	71.15
Mean	23.470	0.000	0.000	0.000		0.000	17.910	17.470	18.230		19.29	17.34	16.23	18.29	
Source	I	S		I X S		I	S		I X S		I	S		I X S	
S. Em ±	0.005	0.005		0.010		0.013	0.013		0.026		0.01	0.01		0.02	
CD ($P \leq 0.05$)	0.014	0.014		0.029		0.038	0.038		0.075		0.03	0.03		0.07	

Table 2. Cont.

Ripening (%) days after storage										
Source	28					34				
	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean
S ₁	0.00*	0.00*	0.00*	0.00*	0.00	0.00*	0.00*	0.00*	0.00*	0.00
S ₂	65.23	29.52	28.25	47.08	42.52	0.00*	86.06	84.13	97.76	66.98
S ₃	74.66	33.08	28.93	57.90	48.65	0.00*	97.01	96.36	98.10	72.87
S ₄	0.00*	0.00*	0.00*	0.00*	0.00	0.00*	0.00*	0.00*	0.00*	0.00
Mean	34.97	15.65	14.29	26.24		0.00	45.77	45.12	48.96	
Source	I	S		I X S		I	S		I X S	
S. Em ±	0.01	0.01		0.02		0.02	0.02		0.04	
CD ($P \leq 0.05$)	0.04	0.04		0.08		0.06	0.06		0.12	

Note: 1.-* indicate fruits completely discarded

2. Where I=irradiation, S=Storage temperature, CA=Control atmospheric storage

3.3. Marketable Fruits Per Cent

The marketable fruits percentage were influenced with the advancement of storage period and significantly affected by irradiation, storage temperature and their interaction (Table 3). During storage, few treatments had 100 per cent values for marketability and few had 0.00 per cent marketability due to induction of senescence. On 12th day of storage only unirradiated ambient stored fruits showed 98.53% marketability, rests are cent per cent marketable but up to 30th day of storage all treatments showed lower marketability

compared to cent per cent in treatments I₂S₃ and I₃S₃. The gamma rays irradiated fruits stored at 9°C and 12°C and in CA (12°C) had significantly higher marketable fruits compared to unirradiated (0.00kGy) fruits. The possible reasons may be that irradiation maintains water content in the fruit and low temperature coupled with high humidity in cold storage maintains the health of the fruits. These results are in conformity with the findings of [27] and [5] with respect to irradiation and [10] with respect to low temperature in mango.

Table 3. Effect of Irradiation and storage conditions on marketable fruits of mango Cv. Alphonso during storage.

Marketable fruits (%) days after storage										
Source	10					16				
	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean
S ₁	97.69	100	100	100	99.42	0.00	97.56	100	97.28	73.71
S ₂	100	100	100	100	100	100	100	100	100	100
S ₃	100	100	100	100	100	100	100	100	100	100
S ₄	100	100	100	100	100	100	100	100	100	100
Mean	100	100	100	100		75.00	99.39	100	99.32	
Source	I	S		I X S		I	S		I X S	
S. Em ±	0.05		0.05	0.10		0.11	0.11		0.22	
CD ($P \leq 0.05$)	0.15		0.15	0.30		0.32	0.32		0.66	

Table 3. Cont.

Marketable fruits (%) days after storage										
	22					28				
Source	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean
S ₁	0.00	88.82	89.37	86.00	66.05	0.00	0.00	0.00	0.00	0.00
S ₂	100	100	100	100	100	84.57	95.97	100	94.11	93.66
S ₃	100	100	100	100	100	98.47	100	100	100	99.62
S ₄	94.07	100	100	96.22	97.57	59.49	71.08	84.34	69.32	71.06
Mean	73.52	97.20	97.34	95.55		60.63	141.76	71.08	65.86	
Source	I	S	I X S			I	S	I X S		
S. Em ±	0.11	0.11		0.22		0.08	0.08		0.16	
CD (P≤0.05)	0.32	0.32		0.64		0.24	0.24		0.48	

Note: Where I=irradiation, S=Storage temperature, CA=Control atmospheric storage

3.4. Average Number of Days Taken to Ripening

Significantly the maximum number of days (38.40) to ripening was recorded in fruits exposed to 0.40kGy gamma rays and stored at 9°C storage temperature (I₃S₂) followed by 37.41 days when fruits exposed with 0.20kGy gamma rays and stored at same situations (I₂S₂). It can be observed that medium dose of irradiation and low temperature significantly enhanced the days taken to ripening of fruits and helped in extending the ripening process (Table 4).

3.5. Shelf Life of Fruits

The data on shelf life of fruits are presented in Table 4. It is cleared from the data that shelf life of fruits significantly influenced by irradiation, storage temperature and their interaction. Significantly the maximum shelf life (41.78 days) was recorded in fruits exposed to 0.40kGy gamma rays

and stored at 9°C storage (I₃S₂) followed by I₂S₂ (40.46 days) when fruits exposed to 0.20kGy gamma rays and stored at same storage temperature. The minimum shelf life (14.11 days) was recorded in unirradiated fruits at ambient storage temperature (I₁S₁). The higher shelf life in irradiated fruits might be due to delaying the ripening process and senescence [16]. Same findings noted by [13] and [3] in mango; [32] in banana and [21] in guava. The fruits stored The shelf life increased in low temperature might be due to the delayed ripening as a result of inhibition of enzymatic activities and reducing the respiration and ethylene production [10]. The present findings confirmed with the findings of [4] in mango and [2] in banana. In combine the irradiated fruits stored at low temperature showed maximum shelf life compared to unirradiated fruits stored at ambient temperature which may be due to combined effect of irradiation and low temperature.

Table 4. Effect of Irradiation and storage conditions on average days taken to ripening and shelf life of mango Cv. Alphonso.

Source	Average days taken to ripening					Shelf life (days)				
	I ₁	I ₂	I ₃	I ₄	Mean	I ₁	I ₂	I ₃	I ₄	Mean
S ₁	12.00	21.18	21.90	20.92	18.99	14.13	24.16	24.80	24.10	21.79
S ₂	35.13	37.51	38.07	35.87	36.65	37.36	40.47	41.79	38.26	39.47
S ₃	30.90	23.27	35.91	34.90	31.24	34.02	38.27	38.95	37.14	37.09
S ₄	25.00	27.01	27.07	26.27	26.34	26.13	28.58	30.26	28.13	28.27
Mean	25.72	27.15	30.68	29.44		27.91	32.87	33.95	31.90	
Source	I	S	I X S			I	S	I X S		
S. Em ±	0.01	0.01	0.03			0.01	0.01		0.02	
CD (P≤0.05)	0.04	0.04	0.09			0.03	0.03		0.06	

4. Conclusion

The fruits of Alphonso mango subjected to 0.40kGy gamma rays irradiation subsequently stored at 9°C temperature delayed the ripening process which maintained lower percentage of physiological loss in weight and ripening per cent, higher percentage of marketable fruits, and increase the shelf life for longer period. This was followed by the fruits treated with 0.40kGy gamma irradiation subsequently stored at 12°C temperature.

References

- [1] Aina, J. O., Adesiji, O. F. and Ferris, S. R. B. (1999). Effect of gamma irradiation on post harvest ripening of plantain fruit (*Musa paradisiaca* L.). Cultivars. *Journal of Science, Food and Agriculture*, 79 (5), 653-656.
- [2] Deka, B. C., Choudhury, A., Bhattacharyya, K. H., Begum and Neog, M. (2006). Postharvest treatment for shelf life extension of banana under different storage environments. *Acta Horticulturae*, 2, 841-849.

- [3] Dhakar, S. D., Savagraon K., A., Srirangarajan, A. N., and Sreenivasan, A. (1966). Irradiation of mangoes. I. Radiation-induced delay in ripening of Alphonso mangoes. *Journal of Food Science*, 31 (6), 863-869.
- [4] Dhemre, J. K. and Waskar, D. P. (2004). Effects of post harvest treatments on shelf life quality of Kesar mango fruits during storage. *Journal of Maharashtra Agricultural University*, 29 (2), 159-163.
- [5] El-Salhy, F. T. A., Khafagy, S. A. A. and Haggay, L. F. (2006). The changes that occur in mango fruits treated by irradiation and hot water during cold storage. *Journal of Applied Research*, 2 (11), 864-868.
- [6] Farzana, Panhwar (2005). Post harvest technology of mango fruits, its development, physiology, pathology and marketing in Pakistan. Digital Verlag GmbH Pub., Germany.
- [7] Gomez-Lim, M. R. (1993). Mango fruit ripening: Physiological and Molecular biology. In: Forth international mango symposium, *Acta Horticulturae*, 341 (Eds. Bruce Schaffer) Tropical Research and Education Center, Florida. pp. 484-496.
- [8] Gutierrez, A. O., Nieto, A. D., Martinez, D., Dominguez, A. M. T., Delgadillo, S. and Qutierrez, A. J. G. (2002). Low temperature plastic film, maturity stage and shelf life of guava fruits. *Revista Chapingo Serie Horticulture*, 8 (2), 283-301.
- [9] Mahindru, S. N. (2009). Food preservation and irradiation. APH publishing corporation, New Delhi.
- [10] Mane, S. R. and Patel, B. N. (2010). Effect of maturity indices, postharvest treatments and storage temperatures on physiological changes of mango (*Mangifera indica* L.) cv. Kesar. In: *National seminar on precision farming in horticulture during Dec., 28-29 held at CoHF, Jhalawar, India* (p. 291).
- [11] Mann, S. S. and Singh, R. N. (1975). Studies on cold storage of mango fruits (*Mangifera indica* L.) cv. Langra. *Indian Journal of Horticulture*, 32 (1), 7-14.
- [12] Mayer, B. S., Anderson, D. S. and Bhing, R. H. (1960). *Introduction to plant physiology*. D. Van Nastrand Co. Ltd., London.
- [13] Moreno, M., Castell-Perez, M. E., Gomes, C., Da silva, P. F. and Moreira, R. G. (2006). Effects of electron beam irradiation on physical, textural and micro-structural properties of Tommy Atkins mangoes (*Mangifera indica* L.). *Journal of Food Science*, 71 (2), 521-524.
- [14] Nagaraju, C. G. and Reddy, T. V. (1995). Deferral of banana fruit ripening by cool chamber storage. *Advances in Horticultural Science*, 9 (4), 162-166.
- [15] Panse, V. G. and Sukhatme, P. V. (1967). *Statistical Methods for Agricultural workers*. ICAR: New Delhi.
- [16] Pimentel, R. M. de Azeveda and Walder, J. M. M. (2004). Gamma irradiation in papaya harvested at three stages of maturation. *Scientia Agriculture*, 61 (2), 140-150.
- [17] Prasadini, P. P., Khan, M. A. and Reddy, P. G. (2008). Effect of irradiation on shelf life and microbiological quality of mangoes (*Mangifera indica* L.). *Journal of Research ANGRU*, 36 (4), 14-23.
- [18] Purohit, A. K., Rawat, T. S. and Kumar, A. (2004). Shelf life and quality of ber fruit cv. Umran in response to post harvest application of ultra violet radiation and paclobutrazole. *Plant Foods for Human Nutrition*, 58 (3), 1-7.
- [19] Roy, S. K. and Joshi, G. D. (1989). An approach to integrated post harvest handling of mango. *Acta Horticulturae*, 231, 469-661.
- [20] Singh, S. P. and Pal, R. K. (2009). Ionizing radiation treatment to improve postharvest life and maintain quality of fresh guava fruit. *Radiation Physics and Chemistry*, 78, 135-140.
- [21] Singh, S. P. and Pal, P. K. (2008). Post harvest fruit fly disinfestations strategies in rainy season guava crop. *Acta Horticulturae*, 375, 591-596.
- [22] Udipi, S. A. and Ghurge, P. S. (2010). Applications of food irradiation. In: *Food irradiation* (eds. Udipi, S. A. and Ghugre, P. S.). Agrotech Publishing academy, Udaipur, pp. 40-71.
- [23] Waskar, D. P. and Masalkar, S. D. (1997). Effect of hydrocooling and bavistin dip on the shelf life and quality of mango during storage under various environments. *Acta Horticulturae*, 455, 687-695.
- [24] Yadav, M. K. and Parmar, B. R., (2014). Response to γ radiation and storage temperature on quarantine pests of Alphonso mango. *Journal of Horticulture and Forestry*, 6 (5): 50-52.
- [25] Yadav, M. K. and Patel, N. L., (2013). Effect of Gamma Radiation and Storage Temperature on Post Harvest Rotting Management of Kesar Mango, *Journal of Mycology and Plant Pathology*, 42 (2): 201-204.
- [26] Yadav, M. K. and Patel, N. L., (2014). Optimization of irradiation and storage temperature for delaying ripening process and maintaining quality of Alphonso mango fruit (*Mangifera indica* L.). *African Journal of Agricultural Research*, 9 (5): 562-571.
- [27] Yadav, M. K., N. L. Patel, B. R. Parmar and Dileswar Nayak., (2013^a). Evaluation of physiological and sensory changes of Kesar mango (*Mangifera indica* L.) influenced by ionizing radiation and storage temperature. *SAARC Agricultural Journal*, 11 (2): 69-80.
- [28] Yadav, M. K., Patel, N. L. & Patel, S. R., (2013^b). Effect of irradiation and storage temperature on quality parameters of Kesar mango (*Mangifera indica* L.). *Indian Journal of Plant Physiology*, 18 (3): 313-317.
- [29] Yadav, M. K., Patel, N. L., Chaudhary, A. D. and Parmar, B. R., (2013^c). Gamma irradiation and storage temperature affects the physiological weight loss and ripening of Kesar mango. *International Journal of Agricultural and Medicinal Plants Research*, 1 (3), 6-12
- [30] Yadav, M. K., Patel, N. L. and Patel, S. R. (2013). Effect of irradiation and storage temperature on quality parameters of Kesar mango (*Mangifera indica* L.). *Indian Journal of Plant Physiology*, 18 (3), 313-317.
- [31] Yimyong, S., Datsenka, T. U. and Handa, A. K., (2011). Hot water treatment delays ripening associated metabolic shift in 'Okrong' mango fruit during storage. *Journal American Society for Horticultural Science*, 136 (6), 441-451.
- [32] Zaman, W., Paul, D., Alam, K., Ibrahim, M. and Hassan, P. (2007). Shelf life extension of banana (*Musa sapientum*) by gamma radiation. *Journal of Bio-Science*, 15, 47-53.