
The Demographic Characteristics and the Risk Factors of Dementia in SAUDI Elderly

Muneerah Albugami^{1,*}, Najeeb Qadi², Fahed Almugbel¹, Alaa Mohammed¹, Alawi Alttas², Abdelazeim Elamin¹, Mumin Siddiquee¹, Usama El Alem¹, Yasmin Al Twaijri³

¹Medicine Department, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia

²Neuroscience Department, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia

³Department of Biostatistics, Epidemiology and Scientific Computing, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia

Email address:

Drmb99@yahoo.com (M. Albugami), nqadi@kfshrc.edu.sa (N. Qadi), fahd2525@hotmail.com (F. Almugbel), dr.alaibrahim@gmail.com (A. Mohammed), alawiattas@yahoo.com (A. Alttas), abdelazeim@hotmail.com (A. Elamin), msiddiquee@kfshrc.edu.sa (M. Siddiquee), uelalem82@kfshrc.edu.sa (U. El Alem), yasmin@kfshrc.edu.sa (Y. Al Twaijri)

*Corresponding author

To cite this article:

Muneerah Albugami, Najeeb Qadi, Almugbel, Alaa Mohammed, Alawi Alttas, Abdelazeim Elamin, Mumin Siddiquee, Usama El Alem, Yasmin Al Twaijri. The Demographic Characteristics and the Risk Factors of Dementia in SAUDI Elderly. *American Journal of Psychiatry and Neuroscience*. Vol. 6, No. 1, 2018, pp. 1-8. doi: 10.11648/j.ajpn.20180601.11

Received: November 30, 2017; **Accepted:** December 8, 2017; **Published:** January 10, 2018

Abstract: There is a little information about dementia in Saudis. This is a retrospective chart review study from 1995 -2010. to describe the demographic characteristics and the risk factors of dementia, the prevalence of different types of dementia, and the current clinical practice of dementia in Saudi tertiary care hospital. A total of 418 demented patients (236 males, 182 females) their mean age was 78.8. Prevalence of diabetes 32%, hypertension 71.53%, dyslipidemia 30.05% and depression 24.41%. Clinically 64.37% of patients had memory impairment, 54.25% had confusion and 34.63% had personality changing. The commonest type of dementia was mixed dementia 18.37% followed by Alzheimer disease 15.87%. 16.10% of patients had received cholinesterase inhibitor and 9.78% had received memantine. Infection was the commonest cause of frequent admission (40%) Mortality rate was 77.99%. The commonest cause of death was infection (38.34%) followed by cardiovascular causes like stroke (23.34%) and cardiac diseases (17.48%). Conclusion: (1) Mixed dementia is the commonest type of dementia in Saudis due to high prevalence of cardiovascular diseases risk factors. (2) High prevalence of depression among demented Saudi patients. It requires early recognition and treatment. (3) Demented patients have frequent admissions and long stay in hospital which makes the economic cost is very high. (4) Mortality rate among demented patients is high and the outcome of dementia is expected to be poor. The underlying message of this study is to increase awareness of the public and health system about the impact of dementia in Saudis and the need for prevention strategies, trained physicians and more research.

Keywords: Dementia, Elderly, Alzheimer's Disease, Mixed Dementia, Depression

1. Introduction

Dementia is characterized by the impairment of memory and learning and at least one other cognitive domain (i.e., aphasia, apraxia, agnosia or executive function), representing a highly severe functional deterioration that interferes with the patient's daily functional abilities and independence [1].

Dementia is a syndrome caused by different etiologies and

it has a substantial medical and social impact [2]. Dementia prevalence has increased continuously over the past decades due to aging populations. The World Health Organization (WHO) estimated that the number of persons living with dementia worldwide (36 million in 2010) will double over the next 20 years [3].

Saudi Arabia is facing the challenge of an ageing population. According to the last Saudi census in 2011 the

life expectancy at birth is 74. Increased in elderly population will have an impact on health care and social services. There is a little information about dementia in Saudis [4]. We have scanty data regarding the demographics characterizing this population or the etiologic diagnoses among those affected in Saudi Arabia.

The objective of this study is to describe the demographic characteristics and the risk factors of dementia, the prevalence of different types of dementia, and the current clinical practice of dementia in Saudi Arabia at tertiary care hospital.

2. Methods

It is a retrospective, cohort study utilized chart review of patients have dementia who received regular care at King Faisal Specialist Hospital and Research Centre (KFSH&RC) in Saudi Arabia from 1995 -2010. Exclusion criteria include depression without dementia and delirium due to acute illness or medications.

3. Statistical Analysis

For this study, all the statistical analysis of data was done by using the software package SAS version 9.4 (SAS

Institute Inc., Cary, NC, USA). Descriptive statistics for the continuous variables are reported as mean ± standard deviation and categorical variables are summarized as frequencies and percentages. The categorical variables are compared by Chi-square test. The level of statistical significance is set at $p < 0.05$.

4. Results

A total of 418 demented patients (236 males, 182 females). The mean age was 78.8 and 63.41% of patients diagnosed to have dementia at age >65. Their basic characteristics and clinical presentation are shown in table 1 & 2.

261 patients (59.32%) had diabetes, 314 patients (71.53%) had hypertension, 131 patient (30.05%) had dyslipidemia, 104 patients (24.41%) had depression, 90 patients (30.72%) were on medication for depression, 81 patients (18.93%) had seizure, 280 patients (64.37%) had memory impairment, 236 patient (54.25%) had confusion, 151 patients (34.63%) had personality changing. Laboratory and radiological investigations are shown in table 3. Neurologists had evaluated 277 patients, internists had evaluated 220 patients, and both had evaluated 130 patients, psychiatrists had evaluated 91 patients.

Table 1. Basic characteristics of dementia patients.

Total no. = 418	Frequency	Percent (%)	Mean
Current age			78.8
Age on diagnosis <65 y	92	22.44	
Age on diagnosis >65y	260	63.41	
not recorded/missing	66	14.15	
Body mass index on presentation (BMI)			26.4
BMI 7 years after dementia diagnosis			25.5
Sex			
Male	236	56.46	
Female	182	43.54	
Education level			
Collage	14	3.18	
Elementary school	7	1.58	
High school	1	0.23	
Illiterate	26	5.91	
Not recorded/missing	370	89.1	
Marital status:			
-Divorced	9	2.01	
-Married	232	52.73	
-single	20	4.55	
-Widow	57	7.98	
-Not recorded/missing	100	32.73	
Comorbid conditions			
Diabetes mellitus	261	59.32	
Hypertension	314	71.53	
Dyslipidemia	131	30.05	
Seizure	81	18.93	
Depression	104	24.41	
patients on medication for depression	90	30.72	
Cigarette smoking			
Yes	22	5.02	
No	110	25.11	
Not recorded/missing	286	69.87	
Alcohol			
Yes	3	0.68	
No	103	23.41	

Total no. = 418	Frequency	Percent (%)	Mean
Not recorded/missing	312	75.91	
Family history of dementia			
Yes	8	1.82	
No	97	22.04	
Not recorded/missing	313	76.14	
Stroke	46	15.03	
Parkinson disease	17	5.55	
Chronic kidney disease	113	36.93	
Hypothyroidism	12	3.92	
HIV	8	2.61	
Brain Malignancy	10	3.27	
Ischemic heart disease	16	5.23	
Atrial fibrillation	8	2.61	
Hepatitis B virus	1	0.33	
Hepatitis C virus	3	0.98	
Liver cirrhosis	3	0.98	
Tuberculosis	3	0.98	
Normal pressure hydrocephalus have shunt	4	1.31	
Multiple sclerosis	2	0.65	
Neurosyphilis	1	0.33	
Vasculitis	2	0.65	
More than 3 Comorbid condition	121	27.44	
Feeding			
Oral feeding	93	21.14	
Nasogastric tube (NGT)	37	8.41	
Gastrostomy tube (GT)	134	30.45	
Jejunostomy tube (JGT)	6	1.36	
Not recorded/missing	148	38.64	

Table 2. Clinical presentation.

Clinical presentation	Frequency	Percent (%)
Memory impairment	280	64.37
Confusion	236	54.25
Personality change	151	34.63
Visual hallucination	9	12.5
Agitation	8	11.11
Urine incontinence	6	1.96
Poor oral intake	1	1.39
More than 3 symptoms	122	27.66

Table 3. Laboratory and radiological investigations.

Laboratory Investigations Results	Frequency	Percent (%)	Radiological Investigation	Frequency	Percent (%)
Vitamin B12			CT brain		
Low	11	2.63	Normal	23	5.11
Normal	126	30.14	Atrophy	129	28.67
High	76	18.19	Vascular changes	163	36.22
not recorded/missing	205	49.04	Evidence of stroke		
			White matter changes	119	26.44
				16	3.56
Folate			MRI brain		
Low	3	0.72	Normal	4	0.85
Normal	86	20.57	Atrophy	62	13.20
High	74	17.70	Vascular changes	61	12.98
not recorded/missing	255	61	Evidence of stroke	130	27.66
			White matter changes	213	45.31
Thyroid stimulating hormone (TSH)			Doppler carotid US		
Low	14	3.35	Normal	35	8.38
Normal	246	58.85			
High	55	13.16	Stenosis	21	5.02
Not done	15	3.59		362	86.60
not recorded/missing	88	21.05	not recorded/missing		
Syphilis			EEG		
Negative	84	20.11	Abnormal	70	16.76
Positive	12	2.88	Normal	12	2.87
not recorded/missing	322	77.01	not recorded/missing	336	80.37

Laboratory Investigations Results	Frequency	Percent (%)	Radiological Investigation	Frequency	Percent (%)
HIV			CSF analysis		
Negative	76	18.2	Abnormal	16	3.83
Positive	8	1.9	Normal	48	11.48
not recorded/missing	334	79.9	Not done	325	77.75
			not recorded/missing	29	6.94
Creatinine level					
Low	10	2.39			
Normal	290	69.38			
High	113	27.03			
not recorded/missing	5	1.20			
ALT level in hepatic profile					
Low	8	1.91			
Normal	375	89.71			
High	30	7.18			
not recorded/missing	5	1.20			
Hemoglobin					
Low	147	35.17			
Normal	213	50.96			
High	9	2.15			
not recorded/missing	49	11.72			
WBC					
Low	7	1.67			
Normal	359	85.89			
High	50	11.96			
not recorded/missing	2	0.48			
Platelet					
Low	22	5.3			
Normal	361	86.4			
High	28	6.7			
not recorded/missing	7	1.6			
Total cholesterol					
Low	4	0.96			
Normal	177	42.34			
High	134	32.06			
not recorded/missing	103	24.64			
LDL					
Low	2	0.48			
Normal	179	42.82			
High	233	55.74			
not recorded/missing	4	0.96			
Triglyceride					
Low	1	0.24			
Normal	199	47.61			
High	35	8.37			
not recorded/missing	183	43.78			
HDL					
Low	17	4.06			
Normal	205	49.04			
High	4	0.96			
not recorded/missing	192	45.94			
HBA1C					
Normal	116	27.75			
High	202	48.33			
not recorded/missing	100	23.92			
Anti DNA profile					
Normal	155	37.08			
High	17	4.07			
not recorded/missing	246	58.85			

81 patients (18.37%) had mixed dementia, 70 patients (15.87%) had Alzheimer disease, 34 patients (7.71%) had vascular dementia, 17 patients (3.85%) had Parkinson dementia, 10 patients (2.27%) had Lewis body dementia, 6 patients (1.36%) had frontotemporal dementia, only 110 patients (25.88%) had received treatment of dementia, 71 patients (16.10%) had received cholinesterase inhibitor, 39

patients (9.78%) had received memantine, 21.14% patients on oral feeding, 8.41% on nasogastric tube (NGT), 30.45% on gastrostomy tube (GT) & 1.36% on jejunostomy tube (JGT). 40% of the causes of frequent admission to hospital were infection like aspiration pneumonia (27%), urinary tract infection (10.24%), bed sores infection (2.76%), followed by cardiac causes (24.52%) like heart failure 11.06%, ischemic

heart diseases 13.46%, family members had taken care of 210 patients (50.24%), housemaid of 6 patients (1.44%), private nurse of 6 patients (1.44%), home health care (HHC) of 68 patients (16.27%) and 128 patients (30.62%) not documented who was looking after them.

We found 51 patients (11.78%) were stay in the hospital for long nursing care, 92 patients alive (22.01%), 326 died (77.99%). The mean age at death was 77.4. Gender and causes of death are shown in table 4.

Table 4. Causes of death.

	Frequency	Percent%	Mean
Current status			
Alive	92	22.01	
Male	50		
Female	42		
Died	326	77.99	
Male	218		
Female	108		
Age at death			77.4
Cause of death			
Infection	125	38.34	
Pneumonia	85	68	
Urinary tract infection	25	20	
Bed sores	15	12	
Cardiac causes	57	17.48	
Heart failure	34	59.65	
MI	14	24.56	
Other cardiac causes	9	15.79	
Malignancy	19	5.83	
Stroke	76	23.31	
pulmonary embolism	6	1.84	
Non specify cause of death (No autopsy is available)	43	13.2	

5. Discussion

78.8 is the mean age of patients and 63.41% of them diagnosed above age 65years. Older age is strongly associated with dementia and has been reported globally [5, 6]. Dominant gender in this study is male 56.46%. Female sex is a dementia risk factor in many international studies [5]. In the Rotterdam Study the incidence of dementia is higher in women than men; age-specific incidence rates are very similar up to the age of 85years. Whereas in women it continued to rise above age 85, the difference could be explained by women living longer than men. Framingham study estimates of lifetime risks for dementia 18.4% and 31.8% for 75-year-old men and women, respectively [7].

The prevalence of cardiovascular diseases (CVD) risk factors like hypertension, dyslipidemia and diabetes mellitus (DM) are high in the current study. This may be responsible for the high prevalence of stroke and chronic renal diseases. It is reported that demented patients have a higher number of comorbidities and the two most frequent comorbidities both for men and women are hypertension and DM [8]. Patients with dementia usually have on average 2 to 8 chronic diseases (comorbidities). In our study 27.44% of patients have more than 3 risk factors comorbidities [9, 10].

The prevalence of DM in general population in Saudi

Arabia is 25.4% [11]. Another study showed the overall prevalence of DM in Saudis is 23.1%, obesity 31%, hypertension and coronary artery disease 25.1% in women, 36.5% in men [12].

Studies finding an association between obesity and incident dementia have usually measured body mass index (BMI) or adiposity in mid rather than late life. Most of these studies found that mid-life obesity increases the risk of dementia later in life. In this study there is no difference in mean BMI on presentation and 7 years later on [13, 14]. Low BMI in later life is associated with the development of dementia, because weight loss is an early manifestation of the disease rather than a true risk factor [15].

Once Alzheimer's disease (AD) develops patients with higher education or occupational levels appear to experience more rapid cognitive decline as compared with those with less education [16, 17]. Unfortunately the education level was not documented in 89.1% of patients.

The prevalence of depression in dementia is reported to be 20–60% [18]. The relationship between depression and dementia is complex [19]. The prevalence of depression in general Saudi population is 49.9% [20]. The prevalence in this study is 24.41% and 30.72% of them on treatment for depression. In cohort study depressive symptoms have an association with cognitive decline that is independent of the neuropathologic hallmarks of dementia [21]. Vascular and mixed subtypes of dementia have a higher prevalence of depression as compared to AD [22]. Treatment for depression can significantly improve quality of life.

Prevalence of seizure among patients is 18.93%. Epileptic events in dementia are frequent and under recognized. Epileptic seizures after stroke are independent predictors of new-onset dementia. The preexisting vascular pathologies that may predispose to both epileptic seizures and new-onset dementia could be white matter changes, silent infarcts, or microbleeds. This could be due to an underlying preclinical degenerative disorder such as Alzheimer's disease [23]. A seizure in dementia; particularly AD is established in many studies. Mechanisms underlying seizure pathogenesis are unresolved, but recent studies raise the possibility, that seizures are related to the same pathogenetic processes responsible for cognitive decline, Treatment of seizures in dementia remains empirical [24].

HIV prevalence in Saudi Arabia is very low 0.02% [25]. HIV associated neurocognitive disorder (HAND) encompass a hierarchy of progressively more severe patterns of neurological involvement. It can range from asymptomatic neurocognitive impairment (ANI) to minor neurocognitive disorder (MND) to more severe HIV-associated dementia (HAD) also called AIDS dementia complex (ADC) [26]. The incidence of ADC has dropped by antiretroviral therapy. In this study 8 patients are reported to have HIV and they had dementia when their age <65.

Dementia symptoms are cognitive changes and behavioral/psychological Symptoms of dementia (BPSD). The commonest symptoms among demented patients in this study are memory impairment, confusion and personality changing.

27.66% of patients have more than 3 symptoms on presentation.

The difficulty in caring for patients with AD is more closely linked to the existence and intensity of the BPSD than the actual cognitive decline. Current treatment for these BPSD is difficult but is generally approached by medications and/or psycho-social support therapies. Several studies have estimated the prevalence of the neuropsychiatric symptoms of dementia to affect 50% to 80% of demented patients in the course of the disease [27]. Systematic reviews of the prevalence of BPSD in community-dwelling older people with dementia give widely ranging results, depending on which tools are used and the length of the observation period [28].

Dementia diagnosis is based on structured history, neurological examination and neuropsychological assessment. Neuroimaging will support for but did not change the clinical diagnosis. Laboratory investigations may help to rule out secondary causes of dementia.

The commonest type of dementia among patients is mixed dementia followed by AD. High prevalence of mixed dementia could be related to high prevalence of CVD risk factors like hypertension, dyslipidemia and diabetes mellitus among Saudis. Stroke is reported in 15.03% of our patients and stroke is well known to double the risk of developing dementia [29]. Dementia medications can temporarily improve symptoms and slow the progression of the disease process. 25.88% of patients have received treatment for dementia, 16.10% have received cholinesterase inhibitor, 5.22% have received memantine. Duration of treatment in 10.79% of patients is <5 years and in 8.63% of patients is > 5 years.

Feeding can be difficult in demented patients. 30.45% of patients on gastrostomy tube (GT) feeding. In observational studies, tube feeding has not been shown to prevent aspiration, heal pressure wounds, improve nutritional status, or decrease mortality in persons with advanced dementia. Previous study in Saudi patients reported that 38.1% of demented patients are on tube feeding and 51% of them have stroke. Demented patients who have difficulty swallowing or reduced food intake often receive feeding tubes [30]. Demented patients are admitted to hospital two to three times more often than people of the same age without dementia. In the USA, admissions to hospital for people over the age of 85 years with dementia increased from 700 000 in 2000 to 1.2 million in 2008. Prevalence of demented patients' admission is 40 - 43% in the UK, Italy and Switzerland [31]. The commonest cause of frequent admission among patients is infection (aspiration pneumonia, urinary tract infection and bed sores infection). Infection is the commonest cause of frequent admission as reported in the literature. Frequent admissions and long stay in hospital make the economic cost is high [32]. The caregivers in 50.24% of patients are family members. Caregivers of demented patients are faced with a number of challenges that can affect their own health and wellbeing which needs further research. 11.78% of demented patients stay at hospital for long nursing care which make the cost is high because no nursing home in Saudi Arabia.

Dementia is a major risk factor for death in advanced age.

Moderate and severe dementia was associated with an increased mortality risk even after appropriate control of comorbid conditions [33]. Mortality rate among our patients is high. The mean age at death is 77.4. The mean age at death of HIV patients with dementia is 45.5.

The commonest cause of death is infection followed by cardiovascular causes like stroke and cardiac diseases. No matter what care is offered, the outcome of dementia is likely to be poor. Half of people with moderate dementia admitted with acute illness such as hip fracture or pneumonia will die within 6 months [34]. There is an apparent difference in survival between AD patients with onset of illness before 75 years and those after 75 years: the younger patients have a longer life expectancy. However, there are conflicting data on survival (in years) comparing male and female patients and comparing patients of different ethnicities [35]. Dementia increases the mortality risk at ten years in the NEDICES Study as in other cohort studies. Age and co-morbidity are associated with higher mortality in dementia patients. One third of deaths in persons over 85 years-old could be attributable to dementia [36].

6. Conclusion

This study has some limitations. It is a retrospective chart review where some missing data are expected and poor documentation is common. Since this study was performed at tertiary care hospital, generalizability may be limited due to sample size. No autopsy is available. Despite these limitations, up to our knowledge this study is the first data study in Saudi Arabia that highlights the following:

(1) Mixed dementia is the commonest type of dementia in Saudis due to high prevalence of cardiovascular diseases risk factors. Prevent and control these risk factors may help to prevent vascular dementia. (2) High prevalence of depression among demented Saudi patients. It requires early recognition and treatment. (3) Despite lack of evidence that feeding tubes benefit patients with dementia, patients who have difficulty swallowing or reduced food intake often receive feeding tubes, the use of feeding tubes are expected to increase. (4) Demented patients have frequent admissions and long stay in hospital which makes the economic cost is very high. There is an urgent need to have national strategies to open nursing homes in Saudi Arabia. (5) Mortality rate among demented patients is high.

The underlying message of this study is to increase awareness of the public and health system about the impact of dementia in Saudis and the need for prevention strategies, trained physicians and more research.

Conflict of Interest

The authors declared that there is no conflict of interest

Acknowledgements

The authors are grateful to Mrs Nada Alhumidan and Mrs

Nada Bawayan for their effort in data entry and Mr. Abdelmoneim Eldali from Research Centre at KFSH&RC for his effort in the statistical analysis of our study.

References

- [1] Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders. 5th edition. Arlington, VA: American Psychiatric Association; 2013.
- [2] Qiu C, von SE, et al: Twenty-year changes in dementia occurrence suggest decreasing incidence in central Stockholm, Sweden. *Neurology* 2013, 80: 1888-1894.
- [3] World Health Organization and Alzheimer's disease International (ADI): Dementia: A Public Health Priority. Geneva: World Health Organization; 2012.
- [4] Ogunniyi A, Daif AK, et al: Dementia in Saudi Arabia: experience from a university hospital. *Acta Neurol Scand* 1998; 98: 116120.
- [5] Borenstein AR, Copenhaver CI, et al. Early-life risk factors for Alzheimer disease. *Alzheimer Dis Assoc Disord* 2006; 20: 63-72.
- [6] Alewijn Ott, Incidence and Risk of Dementia: The Rotterdam study. *Am. J. Epidemiol.* (1998) 147 (6): 574-580.
- [7] Seshadri S, Wolf PA, et al. Lifetime risk Of dementia and Alzheimer's disease: estimates from The Framingha M Study. Presented at the 48th Annual Meeting Of the American Academy of Neurology, San Francisco, California, March 1996.
- [8] Beatriz Poblador-Plou. Comorbidity of dementia: a cross-sectional study of primary care older patients. *BMC Psychiatry* 2014, 14: 84.
- [9] Schubert CC, Boustani M, et al.: Comorbidity profile of dementia patients in primary care: are they sicker? *J Am Geriatr Soc* 2006, 54: 104-109.
- [10] Sanderson M, Wang J, et al.: Co-morbidity associated with dementia. *Am J Alzheimers Dis Other Demen* 2002, 17: 73-78.
- [11] Al-Rubeaan K, Al-Manaa HA. Epidemiology of abnormal glucose metabolism in a country facing its epidemic: SAUDI-DM study. *J Diabetes*. 2014 Sep 30.
- [12] Al-Daghri et al. Diabetes mellitus type 2 and other chronic non-communicable diseases in the central region, Saudi Arabia (Riyadh cohort 2): a decade of an epidemic. *BMC Medicine* 2011, 9: 76.
- [13] Cournot M, Marquié JC, et al. Relation between body mass index and cognitive function in healthy middle-aged men and women. *Neurology* 2006; 67: 1208.
- [14] Albanese E, Davis B, et al. Overweight and Obesity in Midlife and Brain Structure and Dementia 26 Years Later: The AGES-Reykjavik Study. *Am J Epidemiol* 2015; 181: 672.
- [15] Burns JM, Johnson DK, et al. Reduced lean mass in early Alzheimer disease and its association with brain atrophy. *Arch Neurol* 2010; 67: 428.
- [16] Wilson RS, Li Y, et al. Education and the course of cognitive decline in Alzheimer disease. *Neurology* 2004; 63: 1198.
- [17] Hall CB, Derby C, et al. Education delays accelerated decline on a memory test in persons who develop dementia. *Neurology* 2007; 69: 1657.
- [18] Tsuno N, Homma A. What is the association between depression and Alzheimer's disease? *Expert Rev Neurother.* 2009; 9: 1667-76.
- [19] Rubin EH, Veiel LL, et al. Clinically significant depressive symptoms and very mild to mild dementia of the Alzheimer type. *Int J Geriatr Psychiatry.* 2001; 16: 694-701.
- [20] Al-Qadhi W, Ur Rahman S. et al. Adult depression screening in Saudi primary care: prevalence, instrument and cost. *BMC Psychiatry.* 2014 Jul 3; 14: 190.
- [21] Robert S. Wilson, Ana W, et al. Bennett. Clinical-pathologic study of depressive symptoms and cognitive decline in old age. *Neurology*, August 19, 2014, 83: 702-709.
- [22] Castilla-Puentes RC, Habeych ME. Subtypes of depression among patients with Alzheimer's disease and other dementias. *Alzheimer's Dement.* 2010; 6: 63-9.
- [23] Campana Chiara, Assenza Giovanni, et al, Nonconvulsive Seizures and Dementia: A Case Report. *International Journal of Alzheimer's Disease*, Volume 2011 (2011), Article ID 690305, 4 pages.
- [24] AJ Lerner. Epileptic seizures in neurodegenerative dementia syndromes. *JOURNAL OF NEUROLOGY AND NEUROSCIENCE* (2010), Vol. 1, No. 1:3.
- [25] AF Alothman, K Mohajer, Prevalence of HIV-infection in Saudi Arabia, *BMC Proceedings* 20115 (Suppl 6): P252.
- [26] Nomenclature and research case definitions for neurologic manifestations of human immunodeficiency virus-type 1 (HIV-1) infection. Report of a Working Group of the American Academy of Neurology AIDS Task Force. *Neurology.* 1991 Jun. 41 (6): 778-85.
- [27] Constantine G. Lyketsos, Oscar Lopez, Prevalence of Neuropsychiatric Symptoms in Dementia and Mild Cognitive Impairment Results From the Cardiovascular Health Study. *JAMA.* 2002; 288 (12): 1475-1483.
- [28] Van der Linde RM et al Systematic reviews on behavioural and psychological symptoms in the older or demented population. *Alzheimers Res Ther* 2012; 4: 28.
- [29] Gorelick PB, Scuteri A, et al; Vascular contributions to cognitive impairment and dementia: a statement for healthcare professionals from the American heart association/American stroke association. *Stroke.* 2011 Sep; 42 (9): 2672-713.
- [30] Muneerah Albugami, Yasmin Al Twaijri, et al. Impact of Tube Feeding on Aspiration Pneumonia at Tertiary Care Hospital. *American Journal of internal Medicine.* 2015 2015; 3 (3): 95-102.
- [31] Elizabeth L. Sampson, et al Behavioural and psychiatric symptoms in people with dementia admitted to the acute hospital: prospective cohort study. *The British Journal of Psychiatry* Sep 2014, 205 (3) 189-196.
- [32] Elizabeth L. Sampson, Nicola White, et al. Behavioural and psychiatric symptoms in people with dementia admitted to the acute hospital: prospective cohort study. *The British Journal of Psychiatry* Sep 2014, 205 (3) 189-196.

- [33] Llinàs-Regla J, López-Pousa S, et al. Mortality after a diagnosis of dementia in a population aged 75 and over in Spain. *Neuroepidemiology*. 2008; 31 (2): 80-8.
- [34] Morrison RS, Siu AL (2000) Survival in end-stage dementia following acute illness. *JAMA* 284: 47-52.
- [35] Kua EH, Ho E, Tan HH, et al; The natural history of dementia. *Psychogeriatrics*. 2014 Sep; 14 (3): 196-201.
- [36] Villarejo A, Benito-León J. et al Dementia-associated mortality at thirteen years in the NEDICES Cohort Study. *J Alzheimers Dis*. 2011; 26 (3): 543-51.