Neonatal Seizures: Frequency, Clinical Forms, Aetiology and Outcome in Tropical Practice

Bahoura Balaka¹, Komi Assogba^{2,*}, Fidato A. Touglo¹, Damelan Kombaté², Kossivi M. Apetsè², Marie C. Ayassou-Madji¹

¹Neonatology units, paediatrics University Hospital, CHUSO, Lomé, Togo

²Neurology departments, Campus University Hospital, Lomé, Togo

Email address:

seraphinassogba@hotmail.com (K. Assogba)

To cite this article:

Bahoura Balaka, Komi Assogba, Fidato A. Touglo, Damelan Kombaté, Kossivi M. Apetsè, Marie C. Ayassou-Madji. Neonatal Seizures: Frequency, Clinical Forms, Aetiology and Outcome in Tropical Practice. *American Journal of Psychiatry and Neuroscience*. Special Issue: Clinical Neurosciences in Tropical Practice. Vol. 3, No. 5-1, 2015, pp. 1-4. doi: 10.11648/j.ajpn.s.2015030501.11

Abstract: Background: Seizures are common cause of paediatric admissions in poor resource countries and a risk factor for neurodevelopmental impairment. This study aimed to determine the frequency, clinical forms, aetiology and outcome of seizures in relation to rectal temperature in newborn. Patients and Methods: The study was located in paediatric department of our University Hospital. This prospective research had assessed consecutive admitted newborns aged 0-28 days with incident seizure during a period from January to December, 2013. Newborns with epilepsy and non-incident admission seizure were excluded. Results: Seizures were observed in 344/4836 (7.1%). The sample included 192 males and 152 females. The seizures frequency in newborn 0-21 days was 95.9% and was 52.6% in neonates < 1day. The focal seizures were reported in 68% follow by generalized tonic-clonic in 24.4%. Status epilepticus was observed in 7.6%. Seizures had occurred in a context of hypothermia in 77.9%, normal temperature, 14.2% and hyperthermia in 7.8%. The hypothermia and normal temperature seizures were more common in newborn < one week and hyperthermia was noted in older. The associated pathologies in the context of hyperthermia were neonatal infections (17.4%) and falciparum malaria (22.7%) respectively in one week newborn and older. Perinatal ischaemic/hypoxic (35.5%), falciparum malaria (13.6%) and respiratory tract infections and gastroenteritis (11.8%) were the main diseases associated with seizures in the context of hypothermia and normal temperature. The aetiologies were marked by neonatal brain injuries and perinatal asphyxia. Sixteen neonates with seizures were died giving a direct mortality rate of 4.7%. Eight (2.3%) surviving newborns had heavy neurodevelopmental deficits at discharge. Conclusion: The neonate seizures are an indicative of infection with or without fever. The causes of newborn convulsions must be prevented through a wide available public health programs.

Keywords: Convulsions, Etiologies, Incidence, Neonates, Temperature

1. Introduction

Seizures are the most common emergencies neurological conditions in the neonate.^[1,2] It occurred in 3.9% of neonates compared with 1.5% of older infants.^[2-4] The mortality rate was reported as approximately 20% to 35%.^[5,6] Neonatal seizures are an age-specific seizure syndrome and most symptomatic due to brain immature.^[7,8] These seizures differed in semiology, pathophysiology, etiology, and electroencephalographic signature and treatment compared with seizures at older ages.^[9-12] The causes were marked by hypoxic/ ischemic encephalopathy (HIE), stroke disorders, cortical malformations, infections, metabolic disturbances

and benign familial neonatal convulsions.^[9,10,13] The outcome of seizures in over 30% of survival neonates were marked by neurodevelopmental and cognitive deficits and later life epilepsy.^[3,7,14]

The objectives of the present research were to assess the frequency, a etiology and outcome of neonatal seizures in relationship with rectal temperature in neonates.

2. Subjects and Methods

2.1. Setting

The study was located in pediatric department of Lomé University Hospital. This center is the national referral hospital

for children health under 15 years and it is divided into many units. The prospective study assessed newborns admitted consecutively and covered a period from 1st January to 31st December 2013, for seizures associated to rectal temperature.

2.2. Neonates Assessment

A neonate is defined as any child admitted aged 28 days or younger. Seizures were defined as reported or observed repeated involuntary muscle contractions, abnormal tonic extensions or jerky movements of any part of the limbs, face or mouth.

At admission, the neonatologist performed emergency care procedures according to the national guidelines (correction of metabolic and electrolytes unbalance) before taking a formal admission history. The history included the number and where possible, a parental description of the seizure types and interventions already offered. The physical exam was included a formal assessment of the gestational age, delivery condition. The general aspect of neonate, determination of a possible site of infection and any injuries was done.

The collected data concerned seizures features, temperature (took with rectal thermometer), umbilical aspect (redness or pus), respiratory distress, poor Apgar score scale (<7) at 5 min, feeding problems, a bulging fontanel, and consciousness status.

A venous blood sample was drawn for a full routine analysis. Hypokalaemia was defined as plasma K⁺<3.0 mmol/L, hyperkalaemia as K⁺>5.0 mmol/L, hyponatraemia as Na⁺<125 mmol/L and impaired renal function as plasma creatinine >80 µmol/L. Hypoglycaemia was considered if a blood glucose <3 mmol/L. Where not contraindicated, a lumbar puncture was performed in all babies admitted for seizures. All received empirical antibiotic treatment in case of suspected infection. There were no facilities to identify viral causes of meningoencephalitis and for neonate EEG recording.

Premature (<37 weeks), low birth weight (<2000 grams), neonatal tetanus and any neonate presenting with other symptoms than seizures were excluded. The discharging clinician made the final diagnosis after review of the admission history, inpatient management notes and laboratory investigations. These diagnoses were checked by a supervising clinician neonatologist and pediatrics neurologist. The study was approved by the ethic institute of the university hospital. Parents, relatives or near misses were asked to sign the inform consent.

3. Results

3.1. Seizures Frequency

Seizures were observed in 344/4836(7.1%). The sample included 192 males and 152 females. The seizure frequency in newborn aged 0-21 days was 95.9% and was 52.6% in neonates < one day. The focal seizures (tonic, clonic, spasms) were reported in 68% follow by generalized tonic-clonic in 24.4%. Status epilepticus was observed in 7.6%.

3.2. Seizures and Rectal Temperature

Seizures had occurred in a context of hypothermia in 77.9%, normal temperature, 14.2% and hyperthermia in 7.8%. The hypothermia and normal temperature seizures were more common in newborn < one week and hyperthermia was noted in older.

3.3. Seizures Risk Factors and Associated Pathologies

The seizures risk factors were marked by laborious delivery (39.5%), neonate aged less than one day, (86.7%), poor Apgar score and neonatal distress (49.7%). The associated pathologies in the context of hyperthermia were dominated by neonatal infections with bacterial meningitis (17.4%), falciparum malaria (22.7%) respectively in one week newborn and older. Perinatal asphyxia/hypoxic (35.5%), falciparum malaria (13.6%) and respiratory tract infections and gastroenteritis (11.8%) were the main diseases associated with seizures in the context of hypothermia and normal temperature. In table I is reported the demographic, clinical and aetiological repartition of newborns with seizures.

3.4. Distribution of Seizures

The months of September (51, 14.8%), January (42, 12.2%) and May (40, 11.6%), and the 1st and 3rd weeks of life respectively (39.2%, 135; 30.8%, 106) have recorded more seizures in newborns. Table II and III had reported respectively the weekly and monthly distribution of neonatal seizures.

Sixteen neonates with seizures were died giving a global mortality rate of 3.3% and direct rate of 4.7%. The mortality rate in newborns younger than one week was 6.6% (12). Eight (2.3%) surviving newborns had gross neurodevelopmental and physical disabilities at discharge.

Table I. Demographic, clinical and etiologic distribution of newborns with seizures.

Age	Sex		Frequency		Clinical forms	Temperature		Etiologies	
Days	N [‡] (%§)	M [*]	F [†]	N [‡] (%§)	Seizures	N [‡] (%§)	°C	N [‡]	N [‡] (%§)
0-1	98(28.5)	79	62	180 (52.3%)	Tonic	108(31.4)	< 35°5	139	Perinatal asphyxia 122(35.5)
2-7	83(24.1)	23	20	32 (9.3%)	Tonic-clonic	84 (24.4)	35°5- 36°	84	Falciparum malaria 78 (22.7)
8-14	65(18.9)	42	37	78 (22.7%)	Clonic	72 (20.9)	36°1- 36°5	42	Bacterial Meningitis 60(17.4)
15-21	60(17.5)	34	31	40 (11.6%)	Spasms	54(15.7)	36°6- 37°	27	Neonatal brain injuries 48 (13.9)
22-28	38(11)	10	6	14 (4.1%)	Status epilepticus	26(7.5)	37°1- 37°5	20	Neonatal jaundice 20(5.8)
		192	152				37°6-38°	14	Toxoplasmosis 10 (2.9)
		55.8%	44.2%				38°1-38°5	11	Hydrocephalus 6(1.7)
	344			344		344	>38°6	7	344

Legend: M^{*}= male; F[†]=female; N[‡]= number; %§= percentage; °C^{||}= Celsius degree

Table II. Frequency of neonatal convulsions during the first four weeks of life.

	1 st week	2 nd	3 rd	4 th	Total
Number	135	62	106	41	344
Percentage (%)	39.2	18	30.8	12	100

Table III. Monthly distribution of neonatal convulsions.

	September	October	November	December	January	February	March	April	May	June	July	August	Total
Number	51	22	24	15	42	18	33	28	40	17	26	28	344
%	14.8	6.4	7	4.4	12.2	5.2	9.6	8.1	11.6	5	7.6	8.1	100

Legend: %= percentage

4. Discussion

4.1. Methodology, Strengthen and Weakness

The study concerned newborns aged 28 days with seizures in relationship to temperature. Some weaknesses were associated to this work. We were only able to document seizures that were clinically evident. It is possible that we missed subclinical seizures.^[11,12] Secondly, some children died being transferred process from home or basic clinic to hospital. We were unable to collect patients who could not afford care facilities or were more likely to attend basic hospitals or traditional healers. These have been the bias in the recruitment of patients. However, the consistency of our results with previous studies in the literature provides some validity to the data.

4.2. Study Outcome

The first and third weeks of life, and especially the first 24 hours, have emerged as the wealthiest in neonatal seizures. This hospital frequency of seizures is much lower than that reported in previous studies in the north of our country which was 26.2% fifteen years ago.^[15] This reduction may partly be the result of improved in management and especially the education of the population to change behavior towards infant's fever and convulsions. The male was prominent as reported in previous studies ranging from 55 to 56%.^[9,16,17] Convulsive seizures are more common in young neonate according to the immaturity of the brain.^[4,7,18] Some previous studies have reported that the incidence of that condition in poor resources setting is much higher than that reported from developed nations. This may reflect the high prevalence of risk factors for neonatal brain insult in these parts of the world.^[1,15, 17]

The clinical forms of neonatal seizures associated to rectal temperature can be difficult to discern as they can occur within or without fever. There are often subtle seizures given the fact that electro-clinical dissociation is more common in neonates than in other age groups.^[6,11,12] Over than 80% of EEG documented seizures were not accompanied by observable clinical seizures.^[12,20,21] These difficulties are due to immature of white matter tracts.

There is a permanent controversy regarding the dominated form of seizures in neonates. In one hand, the focal seizures are predominant in newborn and generalized tonic-clonic form in infant and child as reported in several

studies.^[2,7,13,24,25]

In other hand, for some authors, the tonic-clonic generalized seizures are more common than focal seizures.^[5,9] In fact, the divergence of these data reflects the state of brain maturity at the time of the seizures. The more the child is younger, he convulses and the process of the convulsion is unclear and willingly atypical. In children older and therefore more mature brain, tonic-clonic seizures go according to conventional neuron pathways.

The most two common diagnoses associated with neonatal seizures were infections (malaria and meningitis) and neonatal encephalopathy. Both conditions are preventable by using simple public health interventions. These measures are pregnancy surveillance, safe deliveries, appropriate neonatal resuscitation, and prevention and early treatment of neonatal infections.^[26-28] Febrile and non febrile seizures are the main condition in newborn to manifest infection. The hypothermic convulsions are mainly due to HIE by perinatal asphyxia, neonate infection and familial epilepsy. The main diseases associated with febrile seizures were severe malaria and bacterial meningitis as reported by several authors.^[5,9,30,31] In none endemic-malaria countries, febrile seizures are often due to nasopharyngeal and respiratory tract infections, and eruptive fevers or gastroenteritis.^[4,10,24] Fifteen percent of all newborns with seizures had neurological impairment at discharge.^[16,22] The long term outcome is marked by neurodevelopmental retardation, behavioral disturbance and epilepsy and is largely depended on the cause of the seizure.^[9,11,25]

5. Conclusion

The early management of fetal distress and the strengthening of neonatologist, obstetrician and wise wives capacities can help to prevent brain injuries and its related events in neonate later life. These vital measures must be disseminated through a large public health program to basic health structures.

References

- [1] Mwaniki M, Mathenge A, Gwer S, Mturi N, Bauni E, Newton CRJC, et al. Neonatal seizures in a rural Kenyan District Hospital: Aetiology, Incidence and Outcome of hospitalization. *BMC Medicine* 2010;8:16-23.

- [2] Ronen GM, Penney S, Andrews W. The epidemiology of clinical neonatal seizures in Newfoundland: a population-based study. *J. Pediatr* 1999; 134:71–5.
- [3] Jensen FE. Neonatal Seizures: An Update on Mechanisms and Management. *Clin Perinatol*. 2009; 36:881-900.
- [4] Silverstein FS, Jensen FE. Neonatal seizures. *Ann Neurol* 2007;62:112-20.
- [5] Abdou Raouf O, Tchoua RK, Ndinga JP, Guikoumbi JR, Josseaume A, Tchoua R. Paediatric emergencies at Libreville University Hospital. *Méd. d'Afrique Noire* 2002;49:475-80.
- [6] Lawn JE, Wilczynska-Ketende K, Cousens SN. Estimating the causes of 4 millions neonatal deaths in the year 2000. *Int J Epidemiol* 2006; 35:706-18.
- [7] Schmitt B, Wohlrab G, Sander T, Steinlein OK, Hajnal BL. Neonatal seizures with tonic clonic sequences and poor developmental outcome. *Epilepsy Res* 2005; 65: 161-8.
- [8] Scher MS. Neonatal seizure classification: a fetal perspective concerning childhood epilepsy. *Epilepsy Res* 2006; 70 Suppl 1:S41-57.
- [9] Tekgul H, Gauvreau K, Soul J, Murphy L, Robertson R, Stewart J, et al. The current etiologic profile and neurodevelopmental outcome of seizures in term newborn infants. *Pediatrics* 2006;117:1270-80.
- [10] Sankar R, Painter MJ. Neonatal seizures: after all these years we still love what doesn't work. *Neurology* 2005;64:776-777.
- [11] Kumar A, Gupta A, Talukdar B. Clinico-etiological and EEG profile of neonatal seizures. *Indian J Pediatr* 2007;74:33-7.
- [12] Murray DM, Ryan CA, Boylan GB, Fitzgerald AP, Connolly S. Prediction of seizures in asphyxiated neonates: correlation with continuous video electroencephalographic monitoring. *Pediatrics* 2006;118:41-6.
- [13] Pellock JM. Understanding co-morbidities affecting children with epilepsy. *Neurology* 2004 ;62 :17–23.
- [14] Idro R, Gwer S, Kahindi M, Gatakaa H, Kazungu T, Ndiritu M, et al. The incidence, aetiology and outcome of acute seizures in children admitted to a rural Kenyan district hospital. *BMC Pediatr* 2008;8:5-15.
- [15] Agbèrè AD, Tatagan K, Madikorai MG, Eklu-Avlasu E, Balaka B, Bakondé B, et al. Infant convulsive seizures in the pediatric service of Kara Regional Hospital (North-Togo). *Méd Afr Noire* 1995;42:310-4.
- [16] Gebremariam A, Gutema Y, Leuel A, Fekadu H. Early-onset neonatal seizures: types, risk factors and short-term outcome. *Ann Trop Paediatr* 2006;26:127-31.
- [17] Birbeck GL. Seizures in rural Zambia. *Epilepsia* 2000;41:277-81.
- [18] Pisani F, Cerminara C, Fusco C, Sisti L. Neonatal status epilepticus vs recurrent neonatal seizures: clinical findings and outcome. *Neurology* 2007;69:2177-85.
- [19] Saliba RM, Annegers JF, Waller DK, Tyson JE, Mizrahi EM. Incidence of neonatal seizures in Harris County, Texas, 1992–1994. *Am. J Epidemiol* 1999;150:763–9.
- [20] Toet MC, Groenendaal F, Osredkar D, van Huffelen AC, de Vries LS. Post neonatal epilepsy following amplitude-integrated EEG-detected neonatal seizures. *Pediatr Neurol* 2005;32:241-7.
- [21] Scher MS, Aso K, Beggarly ME, Hamid MY, Steppe DA, Painter MJ, et al. Electrographic seizures in preterm and full-term neonates: clinical correlates, associated brain lesions and risk for neurologic sequelae. *Pediatrics* 1993;91:128–34
- [22] Ronen GM, Buckley D, Penney S, Streiner DL. Long-term prognosis in children with neonatal seizures: a population-based study. *Neurology* 2007;69:1816-22.
- [23] Hauser WA. The prevalence and incidence of convulsive disorders in children. *Epilepsia* 1994;35 Suppl 2:S1-6.
- [24] Kriouile Y, Mdaghri AA, Hamdani S, Lamdouar BN. Neonatal convulsions: A study of 71 cases. *Rev. Maghr. Pédiatr.* 1999; 9:141-6.
- [25] Pisani F, Sisti L, Seri S. A scoring system for early prognostic assessment after neonatal seizures. *Pediatrics* 2009;124:580-7.
- [26] Rennie JM, Boylan GB. Neonatal seizures and their treatment. *Curr Opin Neurol* 2003; 16:177-81.
- [27] Hellstrom WL, Blennow G, Lindroth M, Rosen I, Svenningsen NW. Low risk of seizure recurrence after early withdrawal of antiepileptic treatment in the neonatal period. *Arch. Dis. Child.* 1995;72:F97-101.
- [28] Musoke RN, Malenga GJ. Bacterial infections in neonates at the Kenyatta Hospital Nursery. A prospective study. *East Afr Med J* 1984;61:909-16.
- [29] Lanska MJ, Lanska DJ. Neonatal seizures in the United States: results of the National Hospital Discharge Survey, 1980-1991. *Neuroepidemiology* 1996;15:117-25.
- [30] Ghedira-Besbes L, Fattaki S, Ben Meriem Ch, Chouchene S, Nouri S, Haddad S, et al. Febrile seizures in infant: A study of 511 cases. *Rev. Maghr. pédiatr* 2004;14:171-9.
- [31] Sadleir LG, Scheffer IE. Febrile seizures. *BMJ* 2007; 334:307-11.