



WBC, PLT of Women's Blood Affects by Alpha Deposition into the Body

Najeba F. Salih^{1,2,*}, Mohamad Jaafar²

¹Department of Physics, Faculty of Science and Health, Koya University, Koya, Iraq

²Department of Physics, Medical Physics, Universiti Sains Malaysia, Penang, Malaysia

Email address:

najebafarhad@yahoo.com (N. F. Salih), najeba.farhad@koyauniversity.org (N. F. Salih)

*Corresponding author

To cite this article:

Najeba F. Salih, Mohamad Jaafar. WBC, PLT of Women's Blood Affects by Alpha Deposition into the Body. *American Journal of Thoracic and Cardiovascular Surgery*. Vol. 2, No. 2, 2017, pp. 25-28. doi: 10.11648/j.ajtcs.20170202.13

Received: March 10, 2017; **Accepted:** March 23, 2017; **Published:** April 12, 2017

Abstract: This study is presented the effects of alpha deposition into the women blood samples, most of WBC, RBC and PLT counted by used automatic CBC machine before and after irradiation with radium source. The values showed statistical significance for WBC and PLT based on the mean of their differences before and after irradiation which decreased with increasing radiation. Significant differences were found in the platelet and WBC before and after treatment with radiation after 10 minutes of activities of ²²⁶Ra. Although, significant decrease in platelet and WBC was observed after 15 minutes of treatment with ²²⁶Ra. However, no significant ($p > 0.05$) change in red blood count RBC was found after 15 minutes irradiated by radium, therefore, WBC/PLT were high significant decrease ($p = 0.001$) after irradiation.

Keywords: Radiation, Blood, WBC, RBC, PLT, Alpha Deposit

1. Introduction

Radiation plays a significant role in life generally, and in the fields of life sciences and medicine. Radiation can result in several different types of biological effect, depending mainly on the part of the body irradiated [1]. The first case of radiation induced leukemia was reported in 1911. This has been demonstrated by studies on the survivors of atomic bomb explosions in Japan [2]. Radioactive radon gas is generally regarded as a health hazard, naturally because of its ability to produce several different types of biological effects [3]. Alpha particles react directly with DNA to generate double-strand breaks or interact with water to produce hydroxyl radicals, both of which affect biological materials negatively [4].

An active cell is affected negatively by the direct interaction with radiation source leading to the death or mutation of the cell. These outcomes are as a result of the cell's relative sensitivity to radiation. One-third of the assimilated decay products from radon are transported from the lungs into the bloodstream through the stomach or intestinal walls and distributed throughout the body [5]. The lymphocyte is a distinct kind of mature white blood cell that

is very radiosensitive for unknown reasons. Lymphocytes are more susceptible to the effects ionizing radiation compared other mature blood cells. White blood cell population reduces a few days after exposure, often accompanied by slow decrease in red blood cells. Red cells have longer life spans than white blood cells, so they are substituted more gradually [6]. The harmful effect of radon was elaborated as an increase in chromosome damage [7]. Aberrations in lymphocytes serve as a sensitive indicator of accidental exposures to doses greater than about 0.1 Gy [8].

2. Methodology

For this study, blood samples were collected from 11 women from Malaysia region with fertility problems. The purpose is to examine the effects of alpha particles on the fertility of women by the analysis of major blood components White blood cell (WBC), Red blood cell (RBC), and Platelet (PLT). The blood samples were collected in the clinic (Health Center) of Universiti Sains Malaysia (USM) from patients aged between 19 and 41 years at two different

times and during two different processes. An exceptional blood tube contain Ethylene Di-amine Tetra Acid (EDTA) was used. Each sample was collected from the women by used a syringe, and then transferred into a tube. The volume of the samples before division was 4ml. Then, each sample of blood was divided regularly into two (2) parts. The study used in vitro irradiation of alpha particles along with hematology Count Blood Cell (CBC) machine to examine the blood parameters (WBC, RBC, and PLT) of women blood before and after irradiation to investigate the impact of the radiation on women with fertility problems. The blood samples were irradiated at a room temperature of 24°C at the University Sciens Malaysia (USM), School of physics.

Irradiation was carried out at room temperature 24°C using ²²⁶Ra. The irradiation was repeated for 2 different time exposures (10 minutes, 15 minutes) of surface samples to the ²²⁶Ra, therefore the samples were divided into two parts and placed in the Polymerizing Vinyl Chloride (PVC) tubes (height: 12.0 cm, diameter: 3.0 cm). The distance between the source and blood surface is fixed at 10.5cm inside the PVC tube with 5 µCi activity for source ²²⁶Ra.

3. Statistical Analysis

All statistical calculations were performed using Statistical Package for the Social Sciences (SPSS) for Windows, Standard version 20.0. The blood data were analyzed for irradiation. Paired samples t test and one way ANOVAs were used to investigate the normality distribution of the data. In addition, Wilcoxon Signed Rank test and Kruskal Wallis test (non-parametric tests) were used in case of non normal distribution

4. Results and Discussion

The effects of radiation (alpha deposition) on the blood the

Table 2. Difference of blood cells between control and after irradiation.

Laboratory measurements		Mean (± SD)	Mean difference ± SD			p value
WBC	Control	6.241 ± 2.149	0.198 ± 0.408			0.009*
	Irradiation	6.043 ± 2.055				
Laboratory measurements		Ranks	Number	Mean rank	Z value	p value
		Negative ranks	22	17.23	0.070	0.133**
RBC	Irradiation - control	Positive ranks	10	17.16	-3.193	<0.001**
		Ties	1			
		Negative ranks	25	14.16		
PLT	Irradiation – control	Positive ranks	6	17.25		
		Ties	2			

* Paired t test

** Wilcoxon Signed Rank test

The WBC decreased after irradiation because clearly, the maximum level of alpha radiation based on field strength of ²²⁶Ra. WBC demonstrated high statistical significance (p = 0.009), because the mean of White Cell Count for the exposed to alpha radiation were observed to be decreased, respectively when compared to the mean White Cell Count for control.

The number of RBC showed no significant change after

women with fertility problems were determined by conducting haematology blood tests before and after irradiation by exposure to the radium source for alpha. The data were analyzed by used the statistical SPSS analysis to check and examine the presence of any relation between the effect of alpha particles on the blood samples and the 3 major blood parameters (WBC, RBC, and PLT) of the blood sample.

4.1. Participants Demographic and Laboratory Characteristics

Table 1. Participants' demographic and laboratory characteristics.

Variables (33 participants)	% (no.) / Mean (± SD)	Median
Age (years)	29.00 ± 6.356	27.00
Total WBC control	8.270 ± 3.045	7.500
Total WBC radiation	7.159 ± 2.709	6.500
Total RBC control	5.588 ± 1.918	5.06
Total RBC radiation	5.050 ± 0.990	4.95
Total PLT control	162.576 ± 64.577	162.00
Total PLT radiation	145.121 ± 51.814	143.00

There were eleven participants who agreed to be involved in current study, with mean age 29.00 ± 6.356. The participants demographic and laboratory characteristics were found in the Table 1.

4.2. Difference of Overall Blood Cells Before and After Radiation

Paired t test was used to finding the difference for the parametric variables; it used to determining the difference of WBC between control and irradiation. And Wilcoxon Signed Rank test was used to finding the difference of the non-parametric variables; also it used to find the difference of RBC and PLT before and after irradiation. Significant differences found in all blood cells, where these cells were reduced after radiation, as shown in the Table 2.

irradiation. The mean count of RBC parameter was not statistically significant (p = 0.133) as the count remained after 10 minutes of irradiation and after 15 minutes of irradiation, which is similar to the mean Red Cell Count of control, as shown in the Table 2. The results are consistent with the studies of [9, 10]. The number of PLT count has decreased after irradiation. The mean count of PLT parameter was highly significant statistically (p = 0.001) after 10 minutes

of irradiation and after 15 minutes irradiation because the mean of PLT count for the exposed to alpha radiation were observed to be decreased.

4.3. Differences in Blood Cells Before and After Irradiation with Time

Wilcoxon Signed Rank test used to find the differences of non-parametric results for blood cells before and after irradiation. No significant differences observed for blood cells before and after radiation with 10 minutes time. All blood cells got significantly lower levels after 15 minutes of radiation only WBC was significant differences in this time, as shown in the Table 3.

Table 3. Difference of blood cells before and after radiation with time.

Variable	Difference	Ranks	Number	Mean rank	p value
WBC 10 mins	Radiation – control	Negative ranks	5	6.90	0.475*
		Positive ranks	5	4.10	
		Ties	1		
RBC 10 mins	Radiation – control	Negative ranks	7	7.21	0.120*
		Positive ranks	4	3.88	
		Ties	0		
Variable	Difference	Mean (\pm SD)	Mean difference		p value
PLT 10 mins	Control – radiation	127.455 \pm 62.237	- 9.454		0.635**
WBC 15mins	Control – radiation	10.509 \pm 2.997	2.109		0.009**
RBC 15 mins	Control – radiation	6.206 \pm 1.884	0.916		0.023**
PLT 15 mins	Control – radiation	5.289 \pm 1.217	44.09		0.026**
		176.000 \pm 60.548			
		131.909 \pm 39.058			

* Wilcoxon Signed Rank test

** Paired t test

4.4. Differences Among Blood Cells With Time

Non parametric test Kruskal Wallis used to find the differences of WBC and RBC, while One Way ANOVA used to find the differences in PLT with time differences. The overall blood cells of participants were not statistically different with time $p > 0.05$, as shown in the Table 4.

Table 4. Differences among blood cells with time.

Variable	Number	Mean rank	Chi-square	p value	
Total WBC	10 mins	11	14.05	2.381	0.304*
after radiation	15 mins	11	20.36		
Total RBC	10 mins	11	15.77	0.330	0.848*
after radiation	15 mins	11	18.14		
Variable	Number	Mean (\pm SD)	F		p value
Total PLT	10 mins	11	136.909 \pm 57.341	1.479	0.244**
after radiation	15 mins	11	131.909 \pm 39.831		

* Kruskal Wallis test

** One Way ANOVA

WBC and PLT decreases after irradiation while RBC showed no considerable changes, which is in agreement with the result of [11]. No significant decrease in red blood count was found after irradiation by radium that is concert with [12]. However, the WBC reduced and susceptibility to

infection increases, because of insufficient white blood cells to fight bacteria, viruses, and other microbes that damage DNA [4] resulting in cell dysfunction (somatic effect) or mutations (genetic damage) [13]. Therefore, radiation affects the ability of women to pregnant [13, 14]. Thus, the irradiation of blood with sufficient ionizing radiation may be decrease women fertility because neither red cells nor platelets contain the genetic material for replication. Also decreased PLT after irradiation inhibits blood clotting.

5. Conclusion

WBC and PLT decreased after irradiation, while RBC increased a little after irradiation. No significant decrease in red blood count was found after irradiation by radium. Thus, the irradiation of blood with sufficient ionizing radiation may be decrease women fertility.

Abbreviations

WBC: White blood cell, RBC; Red blood cell, PLT: Platelet, EDTA: Ethylene Di-amine Tetra Acid, CBC: Count Blood Cell, USM: University Sciens Malaysia, PVC: Polymerizing Vinyl Chloride, SPSS: Statistical Package for the Social Sciences.

References

- [1] ICRP (2003) Biological Effects after Prenatal Irradiation (Embryo and Fetus). ICRP Publication 90 for the International Commission on Radiological Protection Pergamon.
- [2] Abu -Saleh RM (2005) Measurement of Radiation in Soil in the Middle of Gaza. MSc. Thesis. Islamic University of Gaza.
- [3] ICRP (1969) Radiosensitivity Distribution and Spatial of Dose. International Commission on Radiological Protection. Oxford Glasgow. Pergamon Press Ltd, No. 70-78961, London.
- [4] Jostein D, Thora JJ, Helen H, Jahn MN, Jørgen B, Anne KH Roy HL (2010) Assessment of long-term radiotoxicity after treatment with the low - dose - rate alpha - particle - emitting. European Journal of Nuclear Medicine and Molecular Imaging. Springer. 37: 93–102.
- [5] Alberigi S, Pecequilo B, Lobo H, Campos M (2004) Assessment of effective doses from radon levels for tour guides at several galleries of Santana Cave, Southern Brazil, with CR-39 detectors: preliminary results. Radiation Protection Dosimetry. Vol. 0, No. 0, pp. 0–0.
- [6] Gordon E (1998) Health and Environmental Issues Linked to the Nuclear Fuel Chain Section B: Health Effects.
- [7] Abo-Elmagd M, Manal MD, Eissa HM (2008) Cytogenetic effects of radon inhalation. Radiation Measurements. Elsevier 43, 1265–1269.
- [8] IAEA (2004) Practical radiation technical manual health effects and medical surveillance. WagramerStrasse, Vienna, Austria. 5, P. O. Box 100, A-1400.

- [9] Daniela C, Morari V, Cosma C, Neamt S, Corina C (2010) Quantification on DNA damage in human lymphocytes by comet assay, during in vitro ageing in the presence of radon. *Romanian Journal of Physics*. 20, 2, 137–148.
- [10] Hamza VZ, Mary NM (2009) Cytogenetic damage in human blood lymphocytes exposed in vitro to radon. *Fundamental and Molecular Mechanisms of Mutagenesis journal*. Mutation Research. Science Direct Elsevier 661, 1–9.
- [11] Yazdi GT (2012) Effects of ultraviolet light irradiation on hematological and morphological characteristics and potassium level of human blood effect of Thesis, Doctor of Philosophy, Universiti sains Malaysia.
- [12] Damian W, Michael FK, Hanwen Z (2011) Alpha- Versus Beta-Particle Radiopeptide Therapy in a Human Prostate Cancer Model ((²¹³Bi-DOTA-PESIN and ²¹³Bi-AMBA versus ¹⁷⁷Lu-DOTA-PESIN). *Cancer Research*. 71: 1009-1018.
- [13] Sultane AM (2004) Hematological findings in male x-ray technicians. *Saudi Medical Journal*. Original articles. 7: 852-856.
- [14] James AC, Gary FP, FACS (2007) Radiation Injury and the Surgeon. *American College of Surgeons*. Elsevier. 204, 1.