Review Article

A Review on Dendrimers and Metallo dendrimers, the Important Compounds as Catalyst in Material Chemistry

Md. Ashraful Alam^{1,*}, Aklima Jahan², Md. Wahab Khan³¹Department of Chemistry and Bioengineering, Iwate University, Morioka, Japan²Department of Chemistry, University of Chittagong, Chittagong, Bangladesh³Department of Chemistry, Bangladesh University of Engineering and Technology (BUET), Dhaka, Bangladesh**Email address:**

sdashraf84@yahoo.com (Md. A. Alam)

*Corresponding author

To cite this article:Md. Ashraful Alam, Aklima Jahan, Md. Wahab Khan. A Review on Dendrimers and Metallo dendrimers, the Important Compounds as Catalyst in Material Chemistry. *Advances in Materials*. Vol. 6, No. 5, 2017, pp. 52-56. doi: 10.11648/j.am.20170605.11**Received:** July 25, 2017; **Accepted:** August 2, 2017; **Published:** August 31, 2017

Abstract: The research and the usage of the dendrimers and metallo dendrimer compounds which are also known as supramolecules are attracted by the scientists very fast. The use of these two types of compounds are developed and spread out from the chemical to materials throughout the world. One of the important applications of these supramolecules is as catalyst in the different fields of chemistry and material sciences.

Keywords: Dendrimers, Metallo dendrimers, Synthetic, Catalyst, Materials, Supramolecules

1. Introduction

Dendrimer compounds are onion-type polymers which bear branching units or points in each shell [1-2]. Consequently, the number of branches and their molecular masses increase exponentially from the core outwards. Metallo dendrimers [3-13] are now the well-defined and well established term of research due to the efforts of both the

fields of dendrimer and inorganic chemists. They potentially provide ecologically viable and re-generable catalysts whose efficiency can be equal to those of mononuclear homogeneous catalysts and which can be easily recovered and re-used. They are also ideal hosts for supramolecular chemistry [14] and, as such, have been designed as exo-receptors for sensing and titration of biologically relevant anions [15].

Figure 1. Dendrimer and Dendron.

2. A Brief Discussion

Dendritic and metallo dendritic polymers are unique “ball shaped” polymeric substance, whose molecular architecture consists of an initial core and repeating units with branching and terminal groups (Figure 1). Each repeating units bears a branching point to which two or several new repeating units are attached. Owing to their unique properties such as solubility in water, well defined molecular architecture, and spherical shape, dendrimers and metallo dendrimers have found numerous applications in chemical, physical and biological processes [16-18].

The dendrimer and metallo dendrimer compounds can be synthesized in many ways. Convergent and divergent synthetic methodologies for the synthesis of dendrimers provide a high degree of control in terms of the molecular size, shape, and location of functional groups, leading to almost total control over the molecular architecture [19, 20]. These two polymeric compounds used in the vast range of fields in science as in synthetic organic chemistry, materials chemistry, sensors, energy transfer as well as in the molecular electronics. During the past few decades, considerable effort has been devoted to the design of dendrimers & Metallo dendrimers and to their applications in coordination chemistry. Metallic moieties can be attached at various points in dendrimers, specifically the peripheral surface, the branching unit, or multiple sites within the dendrimers, all of which can lead to a variety of applications, including catalysis [21, 22]. The dendrimers have become suitable materials to overcome the drawbacks of linear-chain polymers, and dendritic structures have been shown to act as light harvesting antennas [23-25] and to be appropriate compounds for optoelectronics applications [26, 27]. Recently, the study of directional energy transfer in dendrimers has attracted significant attention

because their unique structures can mimic the biological photosynthetic process and can be applied to optoelectronic molecular devices such as organic light-emitting diodes (OLEDs) [26, 27].

3. Applications of Dendrimers and Metallo dendrimers in Material Chemistry

In the past two decades, the design and synthesis of diverse dendrimers has received increasing attention [28] not only because of the aesthetically pleasing structures of these molecules but also as a result of their various applications in host-guest chemistry [29], material science [30], and membrane chemistry [31]. Up to now, two complementary methodologies [32, 33], the divergent and the convergent, have been employed in the preparation of dendrimers. When the dendrimer branches are sufficiently long, the redox events at the many termini of the metallo dendrimer are independent, appearing as a single wave in the cyclic voltammogram, because of very weak electrostatic effects. As a result, these metallo dendrimers have applications in the molecular recognition, sensing, and titration of anions (e.g., ATP^{2-}) and cations (e.g., transition metal complexes) (Figure 2). When the recognition properties are coupled with catalysis, the metallo dendrimers function in an enzyme-like manner. For example, Pd^{II} can be recognized and titrated using the dendrimer's terminal redox centers and internal coordinate ligands. Redox control over the number of Pd^{II} species located within a dendrimer allows us to predetermine the number of metal atoms that end up in the form of a dendrimer-encapsulated Pd nanoparticle (PdNP).

Figure 2. Dendrimer and Metallo dendrimers as the Design for Molecular Electronics.

at least two redox forms by multiple electron-transfer reactions. Early examples included the first dendrimers terminated by ferrocenyl (Scheme 1), cobaltocenyl (eq. 1) [41, 42], and [FeCp (η -arene)]⁺ (eq. 2) [36, 41] units.

Although dendrimers presenting catalysts at their branch termini can be recovered and reused readily, their inner sphere components can lead to steric inhibition of substrate approach. In contrast, star-shaped catalysts do not suffer from such steric problems, as has been demonstrated for water-soluble dendrimers bearing cationic iron-sandwich termini, which are redox catalysts of cathodic nitrate and nitrite reduction in water.

approximately 5-10 nanometers in diameter. Dendritic polymers or dendrimers provide a route to create very well-defined nanostructures suitable for drug solubilization applications, delivery of DNA and oligonucleotide, targeting drug at specific receptor site, and ability to act as carrier for the development of drug delivery system. It is assumed that within the short period, the research of the dendrimers and metallo dendrimers will be the hotcake to the scientists in different fields.

The authors are grateful to Dr. Kazuaki Shimada for his valuable suggestions to make this manuscript.

References

- [1] Zeng F, Zimmerman SC. (1997) Dendrimers in Supramolecular Chemistry: From Molecular Recognition to Self-Assembly. *Chem. Rev.* 97 (5): 1681–1712.
- [2] Schalley CA, Baytekin B, Baytekin HT, Engeser M, Thorsten F, Rang A. (2006) Mass spectrometry as a tool in dendrimer chemistry: from self-assembling dendrimers to dendrimer gas-phase host–guest chemistry. *J. Phys. Org. Chem.* (19): 479–490.
- [3] Balzani V, Campagna S, Denti G, Juris A, Serrini S, Venturi M (1998) Designing Dendrimers Based on Transition-Metal Complexes. Light-Harvesting Properties and Predetermined Redox Patterns. *Acc. Chem. Res.* 31: 26–34.

- [4] Newkome GR, Moorefield CN, Vögtle F (2001) *Dendrimers and Dendrons: Concepts, Syntheses, Applications*, Wiley-VCH, Weinheim.
- [5] Tomalia D, Fréchet JMJ (Eds.) (2002) *Dendrimers and other Dendritic Polymers*, Wiley-VCH, New York.
- [6] Newkome GR, He E, Moorefield CN (1999) Suprasupermolecules with Novel Properties: Metallo-dendrimers *Chem. Rev.* 99: 1689-1746.
- [7] Bosman AW, Jansen EW, Meijers EW, (1999) About Dendrimers: Structure, Physical Properties, and Applications *Chem. Rev.* 99: 1665-1688.
- [8] Oosterom GE, Reek JNH, Kamer PCJ, Leeuwen PWNMV (2001) Transition Metal Catalysis Using Functionalized Dendrimers. *Angew. Chem. Int. Ed. Engl.* 40: 1828-1859.
- [9] Heerbeek RV, Kamer PCJ, Leeuwen PWNMV, Reek JNH (2002) Dendrimers as Support for Recoverable Catalysts and Reagents. *Chem. Rev.* 102: 3717-3756.
- [10] (a) Knapen JWI, van der Made AW, de Wilde JC, van Leeuwen PWNM, Wijkens P, Grove DM, van Koten G, Homogeneous catalysts based on silane dendrimers functionalized with arylnickel (II) complexes. (1994) *Nature* 372: 659-663; (b) R. Kreiter, A. W. Kleij, R. J. M. Klein Gebbink, G. van Koten, in: F. Vögtle, C. A. Schalley (Eds.), *Dendrimers IV: Metal Coordination, Self Assembly, Catalysis*. (2001) *Top. Curr. Chem.* 217: 163.
- [11] Cuadrado I, Morán M, Casado CM, Alonso B, Losada J (1999) *Coord. Chem. Rev.* 395: 193-195.
- [12] Hearshaw MA, Moss JR (1999) Organometallic and related metal-containing dendrimers. *Chem. Commun.* 1-8.
- [13] (a) D. Astruc, F. Chardac, *Chem. Rev.* 101 (2001) 1991; (b) L. Gade, C. R. *Chimie* 6 (2003).
- [14] Lehn JM (1995) *Supramolecular Chemistry: Concepts and Perspectives*, VCH, Weinheim, Germany.
- [15] Astruc D, Electron and Proton Reservoir Complexes (2000) Thermodynamic Basis for C-H Activation and Applications in Redox and Dendrimer Chemistry. *Acc. Chem. Res.* 33: 287.
- [16] Jansen, J. F. G. A and Meijer, E. W., "The dendritic box: Shape selective liberation of encapsulated Guests". *J. Am. Chem. Soc.*, vol. 117, pp. 4417-4418, 1995.
- [17] Wooly, K. L., Hawker, C. J. and Frechet, M. J., "Unsymmetrical three-dimensional macromolecules: Preparation and characterization of strongly dipolar dendritic macromolecules", *J. Am. Chem. Soc.*, vol. 115, pp. 11496-11505, 1993.
- [18] Percec, V., Johansen, G., Unger, G. and Zhou, J., "Fluorophobic effect induces the self-assembly of semifluorinated tapered monodendrons containing crown ethers into supramolecular columnar dendrimers which exhibit a homeotropic hexagonal columnar liquid crystalline phase", *J. Am. Chem. Soc.*, vol. 118, pp. 9855-9866, 1996.
- [19] (a) Jiang, DL, Aida, T (2001) *Dendrimers and Other Dendritic Polymers*. (b) Fréchet, JMJ, Tomalia, DAE (Eds) (2001) John Wiley & Sons: Chichester.
- [20] (a) Newkome, GR, Moorefield, CN, Vögtle, F (2001) *Dendrimers and Dendrons*; Wiley-VCH: Weinheim (b) Grayson, SM, Fréchet, JMJ (2001) Convergent Dendrons and Dendrimers: from Synthesis to Applications. *Chem. Rev.* 101: 3819-3868. (c) Vögtle, F, Vol. Ed (2001) *Dendrimers II: Design, Dimension, Function*. *Top. Curr. Chem.* 212. (d) Vögtle, F, Gestermann, S, Hesse, R, Schwierz, H, Windisch, B (2000) Functional dendrimers. *Prog. Polym. Sci.* 25: 987-1041. (e) Vögtle, F, Vol. Ed (2000) *Dendrimers I: Architecture, Nanostructure and Supramolecular Chemistry*. *Top. Curr. Chem.* 210. (f) Bosman, AW, Janssen, HM, Meijer, EW (1999) *Chem. Rev.* 99: 1665. (g) Majoral, JP, Caminade, AM (1999) *Chem. Rev.* 99: 845.
- [21] Rehana ME and Mapolie SF (2014) Nickel metallo-dendrimers as catalyst precursors in the tandem oligomerization of ethylene and Friedel-Crafts alkylation of its olefinic products. *Inorg Chim Acta* 409: 96-105.
- [22] Tansir Ahamad and Saad MA (2014) Synthesis and characterization of first and second-generation polyamide pyridylimine nickel dihalide metallo-dendrimers and their uses as catalysts for ethylene polymerization. *Polym Int* 63: 1965-1973.
- [23] (a) Weil, T, Reuther, E, Müller, K (2002) Shape-Persistent, Fluorescent Polyphenylene Dyads and a Triad for Efficient Vectorial Transduction of Excitation Energy. *Angew. Chem., Int. Ed.* 41 (11): 1900-1904. (d) Jiang, D L, Aida, T (1997) Photoisomerization in dendrimers by harvesting of low-energy photons. *Nature* 388: 454-456. (e) Shortreed, MR, Swallen, SF, Shi, ZY, Tan, W, Xu, Z, Devadoss, C, Moore, JS, Kopelman, R (1997) Directed Energy Transfer Funnels in Dendrimeric Antenna Supermolecules. *J. Phys. Chem. B* 1997, 101 (33): 6318-6322. (f) Devadoss, C, Bharathi, P, Moore, JS (1996) Energy Transfer in Dendritic Macromolecules: Molecular Size Effects and the Role of an Energy Gradient. *J. Am. Chem. Soc.* 1996, 118 (40): 9635-9644. (g) Stewart, GM, Fox, MA (1996) Chromophore-Labeled Dendrons as Light Harvesting Antennae. *J. Am. Chem. Soc.* 118 (18): 4354-4360.
- [24] (a) Cameron, C. S.; Gorman, C. B. (2002) Effects of Site Encapsulation on Electrochemical Behavior of Redox-Active Core Dendrimers. *Adv. Funct. Mater.* 2002, 12 (1): 17-20. (b) (b) Stone, D. L.; Smith, D. K.; McGrail, P. T (2002) Ferrocene Encapsulated within Symmetric Dendrimers: A Deeper Understanding of Dendritic Effects on Redox Potential. *J. Am. Chem. Soc.* 124 (5): 856-864.
- [25] Díez-Barra E, García-Martínez JC, del Rey R, Rodríguez-Lo'pez J, Giacalone F, Segura JL, Martí'n N (2003) Synthesis and Photoluminescent Properties of 1,1'-Binaphthyl-Based Chiral Phenylenevinylene Dendrimers. *J. Org. Chem.* 68: 3178-3183.
- [26] Stewart, G. M, Fox, M. A (1996) Chromophore-Labeled Dendrons as Light Harvesting Antennae *J. Am. Chem. Soc.* 118: 4354-4360.
- [27] Kimura M, Shiba T, Muto T, Hanabusa K, Shirai H (2000) Energy transfer within ruthenium-cored rigid Metallo-dendrimers. *Tetrahedron Letters* 41: 6809-6813.
- [28] For selected reviews: (a) Grayson, S. M.; Fréchet, J. M. J. *Chem. Rev.* 2001, 101, 3819-3868. (b) Hecht, S.; Fréchet, J. M. J. *Angew. Chem., Int. Ed.* 2001, 40, 74-91. (c) Astruc, D.; Chardac, F. *Chem. Rev.* 2001, 101, 2991-3024. (d) van Heerbeek, R.; Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Reek, J. N. H. *Chem. Rev.* 2002, 102, 3717-3756. (e) Crespo, L.; Sanclimens, G.; Pons, M.; Giral, E.; Royo, M.; Albericio, F. *Chem. Rev.* 2005, 105, 1663-1682. (f) Tomalia, D. A. *Prog. Polym. Sci.* 2005, 30, 294-324.

- [29] (a) Baars, M. W. P. L.; Kleppinger, R.; Koch, M. H. J.; Yeu, S.-L.; Meijer, E. W. *Angew. Chem., Int. Ed.* 2000, 39, 1285-1288. (b) Hecht, S.; Vladimirov, N.; Fre'chet, J. M. J. *J. Am. Chem. Soc.* 2001, 123, 18-25. (c) Marsitzky, D.; Vestberg, R.; Blainey, P.; Tang, B. T.; Hawker, C. J.; Carter, K. R. *J. Am. Chem. Soc.* 2001, 123, 6965-6972. (d) Le Derf, F.; Levillain, E.; Trippe', G.; Gorgues, A.; Salle', M.; Sebastian, R.-M.; Caminade, A.-M.; Majoral, J.-P. *Angew. Chem., Int. Ed.* 2001, 40, 224-227. (e) Gong, L.-Z.; Hu, Q.-S.; Pu, L. *J. Org. Chem.* 2001, 66, 2358-2367.
- [30] (a) Freeman, A. W.; Koene, C.; Malenfant, P. R. L.; Thompson, M. E.; Fre'chet, J. M. J. *J. Am. Chem. Soc.* 2000, 122, 12385-12386. (b) Weener, J.-W.; Meijer, E. W. *Ad V. Mater.* 2000, 12, 741-746. (c) Newkome, G. R.; He, E.; Godi'nez, L. A.; Baker, G. R. *J. Am. Chem. Soc.* 2000, 122, 9993-10006. (d) Lupton, J. M.; Samuel, I. D. W.; Beavington, R.; Burn, P. L.; Ba'ssler, H. *Ad V. Mater.* 2001, 13, 258-261. (e) Maus, M.; De, R.; Lor, M.; Weil, T.; Mitra, S.; Wiesler, U.-M.; Herrmann, A.; Hofkens, J.; Vosch, T.; Mu'llen, K.; De Schryver, F. C. *J. Am. Chem. Soc.* 2001, 123, 7668-7676.
- [31] (a) Jansen, J. F. G. A.; de Brabander-van den Berg, E. M. M.; Meijer, E. W. *Science* 1994, 266, 1226-1229. (b) Jansen, J. F. G. A.; Meijer, E. W. *J. Am. Chem. Soc.* 1995, 117, 4417-4418. (c) Kovvali, A. S.; Sirkar, K. K. *Ind. Eng. Chem. Res.* 2001, 40, 2502-2511.
- [32] Grayson, S. M.; Fre'chet, J. M. J. *Chem. ReV.* 2001, 101, 3819-3868.
- [33] (a) Hawker, C. J.; Fre'chet, J. M. J. *J. Am. Chem. Soc.* 1990, 112, 7638-7647. (b) Miller, T. M.; Neenan, T. X.; Zayas, R.; Bair, H. E. *J. Am. Chem. Soc.* 1992, 114, 1018-1025. (c) Ihre, H.; Hult, A.; So'nderlind, E. *J. Am. Chem. Soc.* 1996, 118, 6388-6395. (d) Moore, J. S. *Acc. Chem. Res.* 1997, 30, 402-413. (e) Deb, S. K.; Maddux, T. M.; Yu, L. *J. Am. Chem. Soc.* 1997, 119, 9079-9080.
- [34] Moulines, F.; Astruc, D. Tentacled Iron Sandwichs. *Angew. Chem., Int. Ed. Engl.* 1988, 27, 1347-1349.
- [35] Marx, H.-W.; Moulines, F.; Wagner, T.; Astruc, D. Hexakis (but-3-ynyl) benzene. *Angew. Chem., Int. Ed. Engl.* 1996, 35, 1701-1704.
- [36] Moulines, F.; Djakovitch, L.; Boese, R.; Gloaguen, B.; Thiel, W.; Fillaut, J.-L.; Delville, M.-H.; Astruc, D. Organometallic Molecular Trees as Multi-Electron and Proton Reservoirs: CpFeInduced Nona-Allylation of Mesitylene and Phase-Transfer Catalyzed Synthesis of a Redox Active Nona-Iron Complex. *Angew. Chem., Int. Ed. Engl.* 1993, 32, 1075-1077.
- [37] Fillaut, J.-L.; Astruc, D. Tentacled Aromatics: From Central-Ring to Outer-Ring Iron Sandwich Complexes. *J. Chem. Soc., Chem. Commun.* 1993, 1320-1322.
- [38] Fillaut, J.-L.; Linares, J.; Astruc, D. Single-Step Six-Electron Transfer in a Heptanuclear Complex: Isolation of Both Redox Forms. *Angew. Chem., Int. Ed. Engl.* 1994, 33, 2460-2462.
- [39] Astruc, D.; Valerio, C.; J. -Ruiz, J.; Hamon, J. -R.; Varret, F. Electron Reservoir Sandwich Complexes: From Mono- and Bimetallic Complexes to Molecular Trees In Supramolecular Magnetism; Kahn, O., Ed.; NATO ASAI Series, Kluwer: Dordrecht, 1996; p 107.
- [40] Vale' rio, C.; Fillaut, J.-L.; Ruiz, J.; Guittard, J.; Blais, J.-C.; Astruc, D. The Dendritic Effect in Molecular Recognition: Ferrocene Dendrimers and their Use as Supramolecular Redox Sensors for the Recognition of Small Inorganic Anions. *J. Am. Chem. Soc.* 1997, 119, 2588-2589.
- [41] Alonso, E.; Vale' rio, C.; Ruiz, J.; Astruc, D. Polycationic Metallo dendrimers with Cobalticinium and FeCp (arene) termini. *New J. Chem.* 1997, 21, 1139-1141.
- [42] Vale' rio, C.; Ruiz, J.; Fillaut, J.-L.; Astruc, D. Dendritic Effect in the Recognition of Small Inorganic Anions using a Polycationic Nona-cobalticinium Dendrimer. *C. R. Acad. Sci., Paris* 1999, 2, 79-83.
- [43] Vale' rio, C.; Alonso, E.; Ruiz, J.; Blais, J.-C.; Astruc, D. A Polycationic Metallo dendrimer with 24 Organoiron Termini which Senses Chloride and Bromide Anions. *Angew. Chem., Int. Ed.* 1999, 38, 1747-1751.