Retrospective Study of Tick-Borne Pathogens and Observation of *Ehrlichia ewingii* / *Anaplasma phagocytophilum* and Hemotropic *Mycoplasma* spp. in Dogs' Blood Films

Maryam Rassouli^{1,2,*}, Ghazaleh Aghazamani³

¹Pathobiology Department, Shahmirzad School of Veterinary Medicine, Semnan University, Semnan, Iran

²Pathobiology Department, Faculty of Veterinary Medicine, Semnan University, Semnan, Iran

³Shahmirzad School of Veterinary Medicine, Semnan University, Semnan, Iran

Email address:

Mrdvmp@yahoo.com (M. Rassouli), Mrvpar@semnan.ac.ir (M. Rassouli), Ghazale.zamani3@yahoo.com (G. Aghazamani)

To cite this article:

Maryam Rassouli, Ghazaleh Aghazamani. Retrospective Study of Tick-Borne Pathogens and Observation of *Ehrlichia ewingii* / *Anaplasma phagocytophilum* and Hemotropic *Mycoplasma* spp. in Dogs' Blood Films. *Animal and Veterinary Sciences*.

Vol. 3, No. 6, 2015, pp. 171-178. doi: 10.11648/j.avs.20150306.15

Abstract: *Ehrlichia ewingii*, *Anaplasma phagocytophilum* and hemotropic *Mycoplasma* spp. are three bacteria which can infect different dog's blood cells. All of these three pathogens can be transmitted by different ticks and some reservoir hosts also play a role in their transmission. Although *E. ewingii* and *A. phagocytophilum* infect granulocytes and neutrophils of their hosts, respectively, hemotropic *Mycoplasma* spp. can infect reticulocytes. In this research, the previously taken dogs' blood films were collected randomly from different veterinary hospitals of Tehran, the capital of Iran. The blood films were reinvestigated for arthropod-borne diseases. Surprisingly, *E. ewingii*/*A. phagocytophilum* morulae and hemotropic *Mycoplasma* spp. were observed in 18% and 37.7% of samples, respectively. None of these pathogens have been reported in Iran. In addition to the new report, the changes which have been made by these pathogens, their similarities, differences and zoonotic importance are discussed.

Keywords: *Ehrlichia ewingii*, *Anaplasma phagocytophilum*, *Mycoplasma*, Dog, Iran

1. Introduction

Ehrlichiosis is caused by several bacterial species in the genus *Ehrlichia*, pronounced (err-lick-ee-uh) [1], family Anaplasmataceae, order Rickettsiales. They are small, pleomorphic Gram-negative bacteria found in circulating leukocyte or platelets of susceptible mammalian hosts. *Ehrlichia* spp. are tick-borne, obligatory intracellular pathogens. These organisms are often observed in compact colonies or phagosomal inclusions which are named morula [2]. Canine ehrlichiosis is caused by *Ehrlichia canis*. This disease is distributed worldwide and its primary vectors are brown dog tick (*Rhipicephalus sanguineus*) and *Dermacentor variabilis*. This species of *Ehrlichia* commonly infect mononuclear white blood cells [1]. In addition to *E. canis*, *E. chaffeensis*, the agent of human monocytic monocytopathic ehrlichiosis (HME) can infect dogs' mononuclear leukocytes. *E. ewingii* causes canine

granulocytic ehrlichiosis, this disease is reported in the US and its potential vector is *Amblyomma americanum*. Granulocytes are the most infected cells [1]. *Anaplasma phagocytophilum* is another tick-borne pathogen of family Anaplasmataceae which can be transmitted by different species of *Ixodes* [3]. *A. phagocytophilum* infection was first described in Human. It also can infect horses, ruminants, dogs and cats. Macaques and baboons were infected experimentally [4]. Small mammals can be reservoir of *A. phagocytophilum* [3].

Hemotropic mycoplasmosis (formerly, hemobartonellosis) is caused by wall-less Gram-negative bacterial organisms, *Mycoplasma* spp., class Mollicutes. These hemotropic *Mycoplasma* spp. attach and grow on the surface of infected reticulocytes [5]. *Mycoplasma haemocanis* (formerly, *Hemobartonella canis*) and *Candidatus Mycoplasma haematoparvum* cause hemotropic mycoplasmosis in dogs. *M. haemocanis* was experimentally transmitted by *R.*

sanguineus [6], transstadial and transovarial transmission in tick has also been described. The infection transmission through blood transfusion from clinically normal carrier dogs has also been reported [3].

Diagnosis of *E. ewingii* is usually based on tick infestation history, observation of its morula in granulocytes, hematological abnormalities and serological findings. Polymerase chain reaction (PCR) is now available [3].

A. phagocytophilum is often pleomorphic organisms are varying in size from 0.2 to 2 μm in diameter. *A. phagocytophilum* inclusions are morphologically similar to *E. ewingii* morula. More diagnostic techniques are needed for their differential diagnosis [3].

Mycoplasma spp. are easily recognized in stained blood films, *M. haemocanis* are usually occur singly with one to four cocci per red blood cell or in doublets, but ring forms and chains have not been described [7], but these bacteria should be differentiated from Basophilic stipplings and Howell-Jolly bodies [3].

Most symptoms of Ehrlichiosis among dogs were described in *E. canis* infection. This disease is divided into acute, subclinical and chronic forms; clinical disease in *E. ewingii* infection is acute rather than chronic [3]. Common clinical signs are fever, depression, anorexia, weight loss, stiffness, joint swelling [8-11], lethargy and CNS signs such as head tilt, tremors and anisocoria [10, 12]. Rarely observed signs include hemorrhage, weight loss, uveitis, pruritis, vomiting, diarrhea and organomegaly [10-13]. The most important laboratory changes can be observed as mild nonregenerative anemia and thrombocytopenia [9, 12, 14].

The symptoms of *A. phagocytophilum* infection are associated with acute phase of the disease and include: fever, lethargy, depression, anorexia, vomiting, diarrhea, musculoskeletal pain (reluctance to move, stiffness, weakness and lameness) [15, 16], polyarthritis [17], vomiting, diarrhea, respiratory signs (coughing, labored breathing) [15, 16, 18, 19]. Organomegaly is also present [15, 16, 20, 21, 22]. In clinical laboratory examinations; mild to moderate thrombocytopenia and nonregenerative normocytic normochromic anemia can be observed [15, 16, 23].

In Canine hemotropic mycoplasmosis, clinical evidences do not develop. It can cause mild to severe regenerative anemia [7]. The severe symptoms are mostly seen in splenectomized dogs, PCR detection is also available [3].

All of these three organisms can potentially be transmitted to human [24, 25], but their zoonotic importance will be discussed later.

2. Materials and Methods

61 blood films were randomly collected from dogs (33 bitches, 28 studs) which were referred to different veterinary hospitals of Tehran, the capital of Iran. Dogs' age and breed were recorded in a few cases. Previously taken blood films were observed again for hemotropic arthropod-borne pathogens. The observed abnormalities were recorded and

compared with previous diagnosis. The newly recorded diseases were analyzed among studs and bitches by chi-square test, SPSS software. The breeds and ages were not analyzed because they had not mostly been recorded in the case histories.

3. Results

Hemotropic *Mycoplasma* spp. were observed in 15 samples out of 61 (24.6%) (group A) and *Ehrlichia ewingii*/*Anaplasma phagocytophilum* morulae were observed in 3 blood films out of 61 (4.9%) (group B) and 8 samples out of 61 (13.1%) (group AB) had coinfection of *Mycoplasma* spp. and *E. ewingii*/*A. phagocytophilum* (Figs 1, 2, 3); therefore 11 samples (18%) (5 Bitches, 6 studs) were positive in *E. ewingii*/*A. phagocytophilum* and 23 samples (37.7%) (14 Bitches, 9 studs) were positive in *Mycoplasma* spp. The species of *Mycoplasma* could not be determined because there was no access to the previously referred dogs' whole blood samples and the *E. ewingii*/*A. phagocytophilum* could not be differentiate because the dogs' sera were not available in the referred veterinary hospitals. There was no significant difference between infected studs and bitches ($P > 0.05$).

In Diff Quick of group A, in 11 samples mature neutrophils were decreased but band cells increased, the other leukocytes were normal in counting. 2 cases were leukopenic. In 2 cases lymphocytosis was observed in addition to mature neutropenia and band cell increasing. In band cell increasing samples, the band cells of 3 samples was 20%-30%, 9 samples were 10-20% and 1 sample 5-10%. There were toxic changes in addition to left shift in neutrophils. Polychromatophilic macrocytes, hypochromic reticulocytes and anisocytosis were also observed in all samples (Figures 1, 3).

In group B, all of the *E. ewingii*/*A. phagocytophilum* morulae were observed in neutrophil cytoplasm. In Diff Quick, 2 samples were leukopenic with hypochromatic reticulocytes, demonstrated as discocytes. Poikilocytosis include: discocyte, codocyte, echinocyte, Elliptocyte, echinoelliptocyte and shistocyte. In 1 sample lymphopenia with monocytosis was observed. Reactive lymphocytes were also observed in this group (Figures 2, 3).

In group AB: in Diff Quick, 5 samples were leukopenic, 3 cases had mature neutropenia and band cell increasing (1 sample 30-40% and the rest 2 cases 10-20% of the counted leukocytes were band cells). In 1 case in addition to mature neutropenia and band cell increasing, Lymphopenia, monocytosis and reactive lymphocytosis could be observed. Reticulocyte changes include: hypochromasia, anisocytosis and poikilocytosis.

In all leukopenic samples, only neutrophils could be counted in low numbers.

Figure 1. a: hemotropic *Mycoplasma* sp. closed to reticulocyte membrane, black arrow. Note the discocytes, 300X, stained by Giemsa. b: hemotropic *Mycoplasma* sp. closed to reticulocyte membrane, black arrow. Note the discocytes and hyperchromatic macrocytes, 200X, stained by Giemsa. c: doublet of hemotropic *Mycoplasma* sp., black arrow, 400X, stained by Giemsa. d: hemotropic *Mycoplasma* sp. closed to two reticulocyte membranes, black arrows, 500X, stained by Giemsa.

Figure 2. a: *Ehrlichia ewingii*/*Anaplasma phagocytophilum* inclusion in a neutrophil cytoplasm, black arrow, 300X, stained by Giemsa. b: *Ehrlichia ewingii*/*Anaplasma phagocytophilum* inclusion in a neutrophil cytoplasm, black arrow, 400X, stained by Giemsa. c: Basophilic *Ehrlichia ewingii*/*Anaplasma phagocytophilum* inclusions on neutrophil nuclei, thin black arrows and hemotropic *Mycoplasma* sp. closed to reticulocyte membrane, thick black arrow. Note the discocytes and rouleaux. 400X, stained by Giemsa. d: *Ehrlichia ewingii*/*Anaplasma phagocytophilum* inclusion in a band cell cytoplasm, black arrow. 500X, stained by Giemsa. e: *Ehrlichia ewingii*/*Anaplasma phagocytophilum* inclusion in a neutrophil cytoplasm, 500X, stained by Giemsa.

Figure 3. a: poikilocytosis, note the echinoelliptocyte, black arrow, 400X, stained by Giemsa. b: hypochromasia, poikilocytosis and anisocytosis. Note the codocyte, black arrow, 200X, stained by Giemsa. c: panleukopenia and hypochromasia, 40X, stained by Giemsa.

4. Discussion

According to the research, the most important problem among the samples was history taking. Most of them were incomplete. The age and breed of dogs were rarely recorded. In spite of finding some tick-borne diseases, tick infestation history was recorded in only 6 samples. None of these observed abnormalities in the blood films were recorded as a “diagnosed disease” and most of the referred dogs diseases had been diagnosed as viral infections. The concurrent viral infections cannot be rejected but these tick-borne infections (*E. ewingii*/*A. phagocytophilum* and hemotropic *Mycoplasma* spp.) may make the infected dogs susceptible to other infectious diseases such as distemper.

Therefore, it can be concluded that blood films were not used for diagnosis and just taken as a routine procedure from referred dogs.

E. ewingii was endemic in southeastern and south-central of the United States and 26% of shelter dogs were seropositive in endemic areas of the U. S [3, 11]. *E. ewingii* DNA was detected in Cameroon and Brazil [26, 27]. *A. phagocytophilum* is distributed worldwide. In the United States, *A. phagocytophilum* is mostly in western and northern Midwestern states [15, 23, 28]. The range of the seroprevalence in infected dogs can vary widely; in the Europe, the *A. phagocytophilum* seroprevalence may vary from 5% to 70.5% [29-37] and in the United States was from 0.0% to 67.4% [23, 38-40]. In Canada, there was low seroprevalence of *A. phagocytophilum* among dogs which were tested in different provinces (0.09-0.9%) [3, 41]. *A. phagocytophilum* DNA was also detected in Europe and North Africa [3].

Canine hemotropic mycoplasmosis was reported in Europe, especially in its Mediterranean countries [42]. In France 3.3% of examined blood samples were infected with *M. haemocanis* and 9.6% with *Candidatus M. haematoparvum* [43] and high prevalence of this disease was reported among dogs in Africa [44].

None of these diseases has been reported in Iran, so; the importance of our findings is to prove that *E. ewingii*/*A. phagocytophilum* and hemotropic *Mycoplasma* spp. infections are present among dogs in Iran, therefore; further researches are needed to understand the true prevalence, the importance, species diagnosis and the potential vectors of these diseases. Seroprevalence of *E. canis* has been reported among dogs in different parts of Iran [45, 47]. There is a cross-reactivity between *E. ewingii* and *E. canis* in some serological methods [3]. Therefore some of the reported seropositive samples might be related to *E. ewingii* exposure, but there is not any cross-reactivity between *E. ewingii* and *A. phagocytophilum* [3] and serological tests can well differentiate these infections from each other. In this case, because of retrospective study, the access to referred dogs sera was impossible. In this research 23 samples (37.7%) (25.5-50%; 95% confidence interval “CI”) were positive in *Mycoplasma* spp. and 11 samples (18%) (8.4-27.6%; 95% CI) were positive in *E. ewingii*/*A. phagocytophilum* which

are relatively high. One of the results of the disease unidentification is, no treatment of infected dog will occur and consequently high prevalence of the infection among more ticks and more dogs will be observed.

Amblyomma americanum is the main vector of *E. ewingii* and white-tail deer (*Odocoileus virginianus*) serve as a reservoir for *E. ewingii* in the southern parts of the U.S [48-50]. *E. ewingii* DNA was detected in *D. variabilis* and *Rh. sanguineus* [14, 51-54]. These species and may be some other ticks are more important in other countries such as Brazil and Cameroon where *E. ewingii* has been detected [26, 27]. Once ticks are infected by *E. ewingii*, they remain infected throughout their life (transstadial transmission) [14, 55]. *A. phagocytophilum* can be transmitted by members of the *Ixodes persulcatus* complex (e.g., *I. scapularis*, *I. pacificus*, *I. ricinus* and *I. persulcatus*) [3, 56]. A minimum required feeding time for *Ixodes* spp. which can transmit *A. phagocytophilum* to susceptible mammalian hosts is 24 to 48 hours [57-59]. *A. phagocytophilum* is able to reside in the salivary glands of unfed *Ixodes* spp. ticks [60]. *R. sanguineus* serves as a vector and reservoir of *M. haemocanis* because of transstadial and transovarial transmission of the organism [3]. Unfortunately, there is no information of vectors and reservoirs of above pathogens in Iran. *A. americanum* has not been reported in Iran thus, the *E. ewingii* vectors can be the other tick genera. It is also predicted that because of the mean temperature increase in Iran, ticks can be active in more months of the year and thus tick-borne diseases will increase. Therefore, tick control is necessary for pets and farm animals.

Observable changes in blood films were recorded in dogs infected with *E. ewingii*, *A. phagocytophilum* and *M. haemocanis* and listed as below; in *E. ewingii* infection, the most important abnormality is thrombocytopenia but platelet counts are not helpful for diagnosis [9, 10, 12, 14]. Mild anemia and reactive lymphocytosis are observed [10, 11]. Biochemical abnormalities are mild and nonspecific in this disease [3]. Neutrophilic inflammation causes polyarthritis [10]. *E. ewingii* morulae can also be detected in CSF fluid, joint fluid, prostatic fluid or other body fluids [9, 10, 12]. Another problem which is caused by *E. ewingii* is the delay in neutrophil apoptosis. In normal cases neutrophils are in peripheral circulation for only 6 to 8 hours [61], but in some naturally and experimentally infected dogs this time prolonged to 5 days [62]. This phenomenon provides additional time for the pathogen replication [63, 64]. In this study, in leukopenic cases, just low numbers of neutrophils could be counted. Some of these neutrophils were in abnormal shapes; these neutrophils can be named "old neutrophils". In canine granulocytic anaplasmosis, mild to moderated nonregenerative anemia and thrombocytopenia are observed [15, 16, 23]. In Diff Quick, both neutrophilia and neutropenia can be recorded, lymphopenia, monocytosis, eosinopenia are also observed. Monocytopenia was reported in some dogs in the US [65, 66]. In hemotropic mycoplasmosis the anemia is regenerative so hematological findings include: reticulocytosis, polychromasia, anisocytosis

and presence of Howell-Jolly bodies.

Macrocytosis takes more time to develop [3]. In this case, polychromatophic macrocytes were observed, so this is another reason for chronic forms of these diseases in Iran because of the unidentification.

Neutrophilia, monocytosis and lymphopenia can occur in a response to endogenous or exogenous glucocorticoids in dogs like other animals [67-70]. Potential causes of increased endogenous releasing of glucocorticoids include fever, pain, stress, intense exercise and hyperadrenocorticism [69, 71]. Left shift is often associated with inflammatory conditions. Bacterial infections, among other infectious diseases are more important for left shift occurrence [72, 73]. Glucocorticoids increase lymphocyte sequestration and apoptosis [74-76], some secreted cytokines in bacterial and viral infections also cause lymphopenia [77-79]. Reactive lymphocytes are antigen stimulated lymphocytes which their size and their cytoplasmic basophilia were increased [73]. In this case all of the above abnormalities were observed and mentioned in details in the result section but leukopenia might occur due to chronic *E. ewingii*/*A. phagocytophilum* or other concurrent diagnosed diseases like distemper or parvovirus infection! It is also unclear if *E. ewingii*/*A. phagocytophilum* and hemotropic *Mycoplasma* spp. infections can make dogs susceptible to other infections. This question and other ambiguous aspects of these tick-borne diseases must be answered by further researches.

No breed predispositions have been reported in *E. ewingii*, *A. phagocytophilum* and *M. haemocanis* infection [15, 16, 30]. There were some reports of coinfection of *A. phagocytophilum* with *Borrelia burgdorferi* in different hosts [80]. *A. phagocytophilum* infected dogs in the US coinfecting with other *Anaplasma* spp., *Babesia canis*, *Babesia vinsonii*, *E. canis*, *Rickettsia rickettsia* [3, 81-83]. In this research only coinfection of *E. ewingii*/*A. phagocytophilum* with hemotropic *Mycoplasma* spp. was observed.

The mortality has not been reported in order to *E. ewingii* or *A. phagocytophilum* infection among dogs, thus these diseases are mild or inapparent. *E. ewingii* infection may clear spontaneously within weeks or several months [84]. Two deaths of puppies were reported due to hemotropic mycoplasmosis [3]. *M. haemocanis* could be successfully transmitted congenitally and orally [3, 85] in experimental studies. *M. haemocanis* can cause severe life-threatening anemia in splenectomized dogs [3]. According to the results, these diseases have not been diagnosed in Iran, so their mortality rate cannot be estimated.

There are some reports of *E. ewingii* in immunocompromised human [24, 86-88], but there is no report of dog-to-human transmission [3]. *A. phagocytophilum* in human is self-limiting such as dogs [3], but pet animals have an important role in epidemiological aspects of *A. phagocytophilum* in an area [25, 89]. *E. ewingii* and *A. phagocytophilum* infections are also important in sheepdogs because both can also infect small ruminants [90, 91]. *Mycoplasma haemofelis* was detected in an immunocompromised man from Brazil [92]. This case and

some similar cases in human [93, 94] suggest vector, bite, or scratch transmission, or handling blood from animals may transmit the infection into human, especially in immunocompromised individuals [3].

E. ewingii can be treated rapidly by suitable antimicrobial therapy. Doxycycline is a choice drug but other tetracycline can also be used [95]. Supportive treatment is necessary in polyarthritis cases [3]. *A. phagocytophilum* responds to doxycycline, rifampin and levofloxacin. Other tetracycline derivatives are also useful [96]. Canine hemotropic mycoplasmosis is never treated completely and after treatment latent infection is still in infected dogs. This disease can be limited by orally administered tetracyclines. Blood transfusion is necessary when the anemia is severe [3].

5. Conclusion

E. ewingii and *A. phagocytophilum* infection make some similar symptoms in infected dogs and their morulae cannot be differentiated in the blood films [3, 21], but fortunately both bacteria are susceptible to tetracyclines [95]. It is recommended that if the morula was observed in the infected dog's blood film, treatment must start immediately either the infectious agent is *E. ewingii* or *A. phagocytophilum* but the laboratory should record the result as *E. ewingii/A. phagocytophilum*. Afterward, if it is necessary, the serum is tested by serological techniques. Greene (2012) also mentioned that differential diagnosis between *E. ewingii* and *A. phagocytophilum* is usually done for zoonotic concern (for the owner or academic importance) [3]. In developing countries such as Iran because all of the veterinary laboratories have not been equipped for serological and molecular tests, the differential diagnosis between *E. ewingii* and *A. phagocytophilum* may get in trouble. In this research the accurate diagnosis of the reported infections didn't occur due to retrospective study limitations which were mentioned above but further researches to complete our findings are necessary.

Acknowledgments

Special thanks to Dr. Fathipour for his help and there is no competing of financial interests exists.

References

- [1] Parker, J. N. and Parker, P. M. (2002). The official patient's sourcebook on Ehrlichiosis. ICON Health Publication, San Diego, CA, USA, pp. 11-14.
- [2] Williams, J. C. and Kakoma, I. (1990). Ehrlichiosis, a vector-borne disease of animals and humans. Kluwer Academic Publishers. Biston, USA, pp. 1-5.
- [3] Greene, C. E. (2012). Infectious diseases of the dog and cat, forth ed. Elsevier Saunders, Saint Louis, Missouri, USA, pp. 227-258, 310-319.
- [4] Raoult, D. and Parola, P. (2007). Rickettsial diseases. Informa healthcare, New York, USA.
- [5] Peters, I. R., Helps, C. R., McAuliffe, L., Neimark, H., Lappin, M. R., Gruffydd-Jones, T. J., Day, M. J., Hoelzle, K., Willi, B., Meli, M. L., Hofmann-Lehmann, R. and Tasker, S. (2008). RNase P RNA gene (rnpB) phylogeny of hemoplasmas and other *Mycoplasma* species. J. Clin. Microbiol. 46: 1873-1977.
- [6] Seneviratna, P., Weerasinghe, N. and Ariyadasa, S. (1973). Transmission of *Haemobartonella canis* by the dog tick *Rhipicephalus sanguineus*. Res. Vet. Sci. 14: 112-114.
- [7] Sykes, J. E., Ball, L. M., Bailiff, N. L. and Fry, M. M. (2005). *Candidatus Mycoplasma haematoparvum*, a novel small haemotropic mycoplasma from a dog. Int. J. Syst. Evol. Bact. 55: 27-30.
- [8] Bellah, J. R., Shull, R. M., Shull Selcer, E. V. (1986). *Ehrlichia canis*-related polyarthritis in a dog. J. Am. Vet. Med. Assoc. 189: 922-923.
- [9] Stockham, S. L., Schmidt, D. A., Curtis, K. S., Schauf, B. G., Tyler, J. W. and Simpson, S. T. (1992). Evaluation of granulocytic ehrlichiosis in dogs of Missouri, including serologic status to *Ehrlichia canis*, *Ehrlichia equi*, and *Borrelia burgdorferi*. Am. J. Vet. Res. 53: 63-68.
- [10] Goodman, R. A., Hawkins, E. C., Olby, N. J., Grindem, C. B., Hegarty, B. and Breitschwerdt, E. B. (2003). Molecular identification of *Ehrlichia ewingii* infection in dogs: 15cases (1997-2001). J. Am. Vet. Med. Assoc. 222: 1102-1107.
- [11] Liddell, A. M., Stockham, S. L., Scott, M. A., Sumner, J. W., Paddock, C. D., Gaudreault-Keener, M., Arens, M. Q. and Storch, G. A. (2003). Predominance of *Ehrlichia ewingii* in Missouri dogs. J. Clin. Microbiol. 41: 4617-4622.
- [12] Gieg, J., Rikihisa, Y. and Wellman, M. (2009). Diagnosis of *Ehrlichia ewingii* infection by PCR in a puppy from Ohio. Vet. Clin. Pathol. 38: 406-410.
- [13] Goldman, E. E., Breitschwerdt, E. B., Grindem, C. B., Hegarty, B. C., Walls, J. J. and Dumler, J. S. (1998). Granulocytic ehrlichiosis in dogs from North Carolina and Virginia. J. Vet. Intern. Med. 12: 61-70.
- [14] Anziani, O. S., Ewing, S. A. and Barker, R. W. (1990). Experimental transmission of a granulocytic form of the tribe Ehrlichieae by *Dermacentor variabilis* and *Amblyomma americanum* to dogs. Am. J. Vet. Res. 51: 929-931.
- [15] Greig, B., Asanovich, K. M., Armstrong, P. J. and Dumler, J. S. (1996). Geographic, clinical, serologic, and molecular evidence of granulocytic ehrlichiosis, a likely zoonotic disease, in Minnesota and Wisconsin dogs. J. Clin. Microbiol. 34: 44-48.
- [16] Egenvall, A. E., Hedhammar, A. A. and Björnsdörff, A. I. (1997). Clinical features and serology of 14 dogs affected by granulocytic ehrlichiosis in Sweden. Vet. Rec. 140: 222-226.
- [17] Foley, J. E., Drazenovich, N., Leutenegger, C. M. and Chomel, B. B. (2007a). Association between polyarthritis and thrombocytopenia and increased prevalence of vectorborne pathogens in Californian dogs. Vet. Rec. 160: 159-162.
- [18] Pusterla, N., Huder, J., Wolfensberger, C., Litschi, B., Parvis, A. and Lutz, H. (1997). Granulocytic ehrlichiosis in two dogs in Switzerland. J. Clin. Microbiol. 35, 2307-2309.

- [19] Carrade, D., Foley, J., Borjesson, D. and Sykes, J. E. (2009). Canine granulocytic anaplasmosis: a review. *J. Vet. Intern. Med.* 23: 1129-1141.
- [20] Maretzki, C. H., Fisher, D. J. and Greene, C. E. (1994). Granulocytic ehrlichiosis and meningitis in a dog. *J. Am. Vet. Med. Assoc.* 205: 1554-1556.
- [21] Kohn, B., Galke, D., Beelitz, P. and Pfister, K. (2008). Clinical features of canine granulocytic anaplasmosis in 18 naturally infected dogs. *J. Vet. Intern. Med.* 22: 1289-1295.
- [22] Granick, J. L., Armstrong, P. J. and Bender, J. B. (2009). *Anaplasma phagocytophilum* infection in dogs: 34 cases (2000-2007). *J. Am. Vet. Med. Assoc.* 234: 1559-1565.
- [23] Beall, M. J., Chandrashekar, R., Eberts, M. D., Cyr, K. E., Diniz, P. P., Mainville, C., Hegarty, B. C., Crawford, J. M. and Breitschwerdt, E. B. (2008). Serological and molecular prevalence of *Borrelia burgdorferi*, *Anaplasma phagocytophilum*, and *Ehrlichia* species in dogs from Minnesota. *Vector. Borne. Zoonotic. Dis.* 8: 455-464.
- [24] Buller, R. S., Arens, M., Hmeil, S. P., Sumner, J. W., Rikhisa, Y., Unver, A., Gaudreault-Keener, M., Manian, F. A., Liddell, A.M., Schmulewitz, N. and Storch, G.A. (1999). *Ehrlichia ewingii*, a newly recognized agent of human ehrlichiosis. *N. Engl. J. Med.* 341: 148-155.
- [25] Cleaveland, S., Meslin, F. X. and Breiman, R. (2006). Dogs can play useful role as sentinel hosts for disease. *Nature.* 440: 605.
- [26] Ndip, L. M., Ndip, R. N., Esemu, S. N., Dickmu, V. L., Fokam, E. B., Walker, D. H. and McBride, J. W. (2005). Ehrlichial infection in Cameroonian canines by *Ehrlichia canis* and *Ehrlichia ewingii*. *Vet. Microbiol.* 111: 59-66.
- [27] Oliveira, L. S., Oliveira, K. A., Mourao, L. C., Pescatore, A. M., Almeida, M. R., Conceição, L. G., Galvão, M. A. and Mafra, C. (2009). First report of *Ehrlichia ewingii* detected by molecular investigation in dogs from Brazil. *Clin. Microbiol. Infect.* 15 (suppl 2): 55-56.
- [28] Arsenault, W. G. and Messick, J. B. (2005). Acute granulocytic ehrlichiosis in a Rottweiler. *Am. Anim. Hosp. Assoc.* 41: 323-326.
- [29] Pusterla, N., Pusterla, J. B. and Deplazes, P. (1998). Seroprevalence of *Ehrlichia canis* and of canine granulocytic *Ehrlichia* infection in dogs in Switzerland. *J. Clin. Microbiol.* 36: 3460-3462.
- [30] Egenvall, A., Bonnett, B. N., Gunnarsson, A., Hedhammar, A., Shoukri, M., Bornstein, S. and Artursson, K. (2000). Seroprevalence of granulocytic *Ehrlichia* spp. and *Borrelia burgdorferi* sensu lato in Swedish dogs 1991-94. *Scand. J. Infect. Dis.* 32: 19-25.
- [31] Barutzki, D., De Nicola, A., Zeziola, M. and Reule, M. (2006). Seroprevalence of *Anaplasma phagocytophilum* infection in dogs in Germany. *Berl. Munch. Tierarztl. Wochenschr.* 119: 342-347.
- [32] Solano-Gallego, L., Lluall, J., Osso, M., Hegarty, B. and Breitschwerdt, E.B. (2006). A serological study of exposure to arthropod-borne pathogens in dogs from northeastern Spain. *Vet. Res.* 37: 231-244.
- [33] Torina, A. and Caracappa, S. (2006). Dog tick-borne diseases in Sicily. *Parasitol.* 48: 145-147.
- [34] Amusategui, I., Tesouro, M. A., Kakoma, I. and Sainz, A. (2008). Serological reactivity to *Ehrlichia canis*, *Anaplasma phagocytophilum*, *Neorickettsia risticii*, *Borrelia burgdorferi* and *Rickettsia conorii* in dogs from Northwestern Spain. *Vector Borne. Zoonotic. Dis.* 8: 797-803.
- [35] Ebani, V., Cerri, D., Fratini, F., Ampola, M. and Andreani, E. (2008). Seroprevalence of *Anaplasma phagocytophilum* in domestic and wild animals from central Italy. *New. Microbiol.* 31: 371-375.
- [36] Ravník, U., Tozon, N., Strasek, K. and Avsic Zupanc, T. (2009) Clinical and haematological features in *Anaplasma phagocytophilum* seropositive dogs. *Clin. Microbiol. Infect.* 15(suppl 2): 39-40.
- [37] Santos, A. S., Alexandre, N., Sousa, R., Nuncio, M. S., Bacellar, F., and Dumler, J. S. (2009). Serological and molecular survey of *Anaplasma* species infection in dogs with suspected tickborne disease in Portugal. *Vet. Rec.* 164: 168-171.
- [38] Suksawat, J., Hegarty, B. C. and Breitschwerdt, E. B. (2000). Seroprevalence of *Ehrlichia canis*, *Ehrlichia equi* and *Ehrlichia risticii* in sick dogs from North Carolina and Virginia. *J. Vet. Intern. Med.* 14: 50-55.
- [39] Foley, J. E., Foley, P. and Madigan, J. E. (2001). Spatial distribution of seropositivity to the causative agent of granulocytic ehrlichiosis in dogs in California. *Am. J. Vet. Res.* 62: 1599-1605.
- [40] Foley, J. E., Brown, R. N., Gabriel, M. W., Henn, J., Drazenovich, N., Kasten, R., Green, S.L. and Chomel, B. B. (2007b). Spatial analysis of the exposure of dogs in rural north-coastal California to vectorborne pathogens. *Vet. Rec.* 161: 653-657.
- [41] Gary, A. T., Webb, J. A., Hegarty, B. C. and Breitschwerdt, E.B. (2006). The low seroprevalence of tick-transmitted agents of disease in dogs from southern Ontario and Quebec. *Can. Vet. J.* 47: 1194-1200.
- [42] Willi, B., Novacco, M., Meli, M., Wolf-Jackel, G. A., Boretti, F. S. and Wengi, N. (2010). Haemotropic mycoplasmas of cats and dogs: transmission, diagnosis, prevalence and importance in Europe. *Schweiz. Arch. Tierheilkd.* 152: 237-244.
- [43] Kenny, M. J., Shaw, S. E., Beugnet, F. and Tasker, S. (2004). Demonstration of two distinct hemotropic *Mycoplasmas* in French dogs. *J. Clin. Microbiol.* 42 (11): 5397-5399.
- [44] Barker, E. N., Tasker, S., Day, M. J., Warman, S. M., Woolley, K., Birtles, R., Georges, K. C., Ezeokoli, C. D., Newaj-Fyzul, A., Campbell, M. D., Sparagano, O. A. E., Cleaveland, S. and Helps, C. R. (2010). Development and use of real-time PCR to detect and quantify *Mycoplasma haemocanis* and "*Candidatus* *Mycoplasma haematoparvum*" in dogs. *Vet. Microbiol.* 140: 167-170.
- [45] Akhtardanesh, B., Ghanbarpour, R., Blourizadeh, H., 2010. Serological evidence of canine monocytic ehrlichiosis in Iran. *Comp. Clin. Path.* 19: 469-474.
- [46] Avize, R., Mosallanejad, B., Razi Jalali, M. H. and Alborzi, A.R. (2010). Seroprevalence of *Ehrlichia canis* in dogs referred to Veterinary Hospital of Shahid Chamran University of Ahvaz, Iran. *Arch. Razi. Inst.* 65: 1-5.
- [47] Ansari-Mood, M., Khoshnegah, J., Mohri, M. and Rajaei, S.M. (2014). Seroprevalence and risk factors of *Ehrlichia canis* infection among companion dogs of Mashhad, North East of Iran, 2009-2010. *J. Arthropod. Borne. Dis.* 9(2): 184-194.

- [48] Beaufile, J. P. (1997). Ehrlichiosis: clinical aspects in dogs and cats. *Compend. Cont. Educ. Pract. Vet.* 19: 57-61.
- [49] Yabsley, M. J., Varela, A. S., Tate, C. M., Dugan, V. G., Stallknecht D. E., Little, S. E. and Davidson, W. R. (2002). *Ehrlichia ewingii* infection in white-tail deer (*Odocoileus virginianus*). *Emerg. Infect. Dis.* 8: 668-671.
- [50] Murphy, G. L., Ewing, S. A., Whitworth, L. C., Fox, J. C. and Kocan, A. A. (1998). A molecular and serologic survey of *Ehrlichia canis*, *E. chaffeensis*, and *E. ewingii* in dogs and ticks from Oklahoma. *Vet. Parasitol.* 79: 325-339.
- [51] Varela-Stokes, A. S. (2007). Transmission of bacterial agents from lone star ticks to white-tailed deer. *J. Med. Entomol.* 44: 478-483.
- [52] Steiert, J. G. and Gilfoy, F. (2002). Infection rates of *Amblyomma americanum* and *Dermacentor variabilis* by *Ehrlichia chaffeensis* and *Ehrlichia ewingii* in southwest Missouri. *Vector. Borne. Zoonotic. Dis.* 2: 53-60.
- [53] Childs, J. E. and Paddock, C. D. (2003). The ascendancy of *Amblyomma americanum* as a vector of pathogens affecting humans in the United States. *Annu. Rev. Entomol.* 48: 307-337.
- [54] De Shields, A., Borman-Shoap, E., Peters, J. E., Gaudreault-Keener, M., Arens, M. Q. and Storch, G. A. (2004). Detection of pathogenic *Ehrlichia* in ticks collected at acquisition sites of human ehrlichiosis in Missouri. *Mo. Med.* 101: 132-136.
- [55] Varela, A. S., Moore, V. A. and Little, S. E. (2004). Disease agents in *Amblyomma americanum* from northeastern Georgia. *J. Med. Entomol.* 41: 753-759.
- [56] Woldehiwet, Z. (2010). The natural history of *Anaplasma phagocytophilum*. *Vet. Parasitol.* 167: 108-122.
- [57] Hodzic, E., Fish, D., Maretzki, C. M., De Silva, A. M., Feng, S. and Barthold, S. W. (1998). Acquisition and transmission of the agent of human granulocytic ehrlichiosis by *Ixodes scapularis* ticks. *J. Clin. Microbiol.* 36: 3574-3578.
- [58] Katavolos, P., Armstrong, P. M., Dawson, J. E. and Telford, S. R. (1998). Duration of tick attachment required for transmission of granulocytic ehrlichiosis. *J. Infect. Dis.* 177: 1422-1425.
- [59] Des Vignes, F., Piesman, J., Heffernan, R., Schulze, T. L., Stafford, K. C., Fish, D. and (2001). Effect of tick removal on transmission of *Borrelia burgdorferi* and *Ehrlichia phagocytophila* by *Ixodes scapularis* nymphs. *J. Infect. Dis.* 183: 773-778.
- [60] Kidd, L. and Breitschwerdt, E. B. (2003). Transmission times and prevention of tick-borne diseases in dogs. *Compend. Cont. Educ. Pract. Vet.* 25: 742-750.
- [61] Christopher, M. J. and Link, D. C. (2007). Regulation of neutrophil homeostasis. *Curr. Opin. Hematol.* 14: 3-8.
- [62] Xiong, Q., Bao, W., Ge, Y. and Rikihisa, Y. (2008). *Ehrlichia ewingii* infection delays spontaneous neutrophil apoptosis through stabilization of mitochondria. *J. Infect. Dis.* 197: 1110-1118.
- [63] Yoshiie, K., Kim, H. Y., Mott, J. and Rikihisa, Y. (2000). Intracellular infection by the human granulocytic ehrlichiosis agent inhibits human neutrophil apoptosis. *Infect. Immun.* 68: 1125-1133.
- [64] Scaife, H., Woldehiwet, Z., Hart, C. A. and Dwards, S. W. (2003). *Anaplasma phagocytophilum* reduces neutrophil apoptosis in vivo. *Infect. Immun.* 71: 1995-2001.
- [65] Johansson, K. E., Pettersson, B., Uhlen, M., Gunnarsson, A., Malmqvist, M., and Olsson, E. (1995). Identification of the causative agent of granulocytic ehrlichiosis in Swedish dogs and horses by direct solid phase sequencing of the PCR products from the 16S rRNA gene. *Res. Vet. Sci.* 58: 109-112.
- [66] Daniels, T. J., Falco, R. C., Schwartz, I., Varde, S. and Robbins, R. G. (1997). Deer ticks (*Ixodes scapularis*) and the agents of Lyme disease and human granulocytic ehrlichiosis in a New York City park. *Emerg. Infect. Dis.* 3: 353-355.
- [67] Osbaldiston, G. W. and Greve, T. (1978). Estimating adrenal cortical function in dogs with ACTH. *Cornell. Vet.* 68: 308-309.
- [68] Kaufman, J. (1984). Diseases of the adrenal cortex of dogs and cats. *Mod. Vet. Pract.* 65: 429-434.
- [69] Davis, J. M., Albert, J. D., Tracy, K. J., Calvano, S. E., Lowry, S. F., Shires, G. T., and Yurt, R. W. (1991). Increased neutrophil mobilization and decreased chemotaxis during cortisol and epinephrine infusions. *J. Trauma.* 31: 725-731.
- [70] Yamada, R., Tsuchida, S., Hara, Y., Tagawa, M. and Ogawa, R. (2002). Apoptotic lymphocytes induced by surgical trauma in dogs. *J. Anesth.* 16: 131-137.
- [71] Brenner, I., Shek, P. N., Zamecnik, J. and Shephard, R. J. (1998). Stress hormones and the immunological responses to heat and exercise. *Int. J. Sports. Med.* 19: 130-143.
- [72] Kogan, D. A., Johnson, L. R., Jandrey, K. E. and Polard, R. E. (2008). Clinical, clinicopathologic, and radiographic findings in dogs with aspiration pneumonia: cases (2004-2006). *J. Am. Vet. Med. Assoc.* 233: 1742-1747.
- [73] Stevens, A., Lowe, J. S. and Scott, I. (2011). *Veterinary Hematology. A Diagnostic guide and color atlas.* Elsevier Saunders, Saint Louis, Missouri, USA, pp. 122-174.
- [74] Bloemena, E., Weinreich, S., Schellekens, P. T. A. (1990). The influence of prednisolone on the recirculation of peripheral blood lymphocytes in vivo. *Clin. Exp. Immunol.* 80: 460-466.
- [75] Toft, P., Lillevang, S. T., Tonnesen, E., Svendsen, P. and Höhndorf, K. (1993). Redistribution of lymphocytes following *E.coli* sepsis. *Scand. J. Immunol.* 38: 541-545.
- [76] Ammersbach, M. A., Kruth, S. A., Sears, W. and Bienzle, D. (2006). The effect of glucocorticoids on canine lymphocyte marker expression and apoptosis. *J. Vet. Intern. Med.* 20: 1166-1171.
- [77] Ulich, T. R., del Castillo, J., Ni, R. X. and Bikhazi, N. (1989). Hematologic interactions of endotoxin, tumornecrosis factor alpha (TNF alpha), interleukin 1, and adrenal hormones and the hematologic effects of TNF alpha in *Corynebacterium parvum* primed rats. *J. Leukoc. Biol.* 45: 546-557.
- [78] Van Miert, A. S., van Duin, C. T. and Wensing, T. (1992). Fever and acute phase response induced in dwarf goats by endotoxin and bovine and human recombinant tumour necrosis factor alpha. *J. Vet. Pharmacol. Ther.* 15: 332-342.
- [79] Sheridan, W. P., Hunt, P., Simonet, S. and Ulrich, T. R. (1997). Hematologic effects of cytokines. In: Remick DG, Friedland JS, eds. *Cytokines in Health and Disease*, second ed. Marcel Dekker Inc., New York, USA, pp. 487-505.

- [80] Nieto, N. C. and Foley, J. E. (2009). Meta-analysis of coinfection and coexposure with *Borrelia burgdorferi* and *Anaplasma phagocytophilum* in humans, domestic animals, wildlife, and *Ixodes ricinus*-complex ticks. Vector. Borne. Zoonotic. Dis. 9: 93-102.
- [81] Du Plessis, J. L., Fourie, N., Nel, P. W. and Evezard, D. N. (1990). Concurrent babesiosis and ehrlichiosis in the dog: blood smear examination supplemented by the indirect fluorescent antibody test, using *Cowdria ruminantium* as antigen. Onderstepoort. J. Vet. Res. 57: 151-155.
- [82] Dykstra, E. A., Slater, M. R. and Teel, P. D. (1997). Perceptions of veterinary clinics and pest control companies regarding tick-related problems in dogs residing in Texas cities. J. Am. Vet. Med. Assoc. 210: 360-365.
- [83] Kordick, S. K., Breitschwerdt, E. B., Hegarty, B. C., Southwick, K. L., Colitz, C. M., Hancock, S. I., Bradley, J. M., Rumbough, R., Mcpherson, J. T. and MacCormack, J. N. (1999). Coinfection with multiple tick-borne pathogens in a Walker hound kennel in North Carolina. J. Clin. Microbiol. 37: 2631-2638.
- [84] Stockham, S. L., Tyler, J. W., Schmidt, D. A. and Curtis, K. S. (1990). Experimental transmission of granulocytic ehrlichial organisms in dogs. Vet. Clin. Pathol. 19: 99-104.
- [85] Krakowka, S. (1977). Transplacentally acquired microbial and parasitic diseases of dogs. J. Am. Vet. Med. Assoc. 171: 750-753.
- [86] Paddock, C. D., Folk, S. M., Shore, G. M., Machado, L. J., Huycke, M.M., Slater, L. N., Liddell, A. M., Buller, R. S., Storch, G. A., Monson, T. P., Rimland, D., Sumner, J. W., Singleton, J., Bloch, K. C., Tang, Y. W., Standaert, S. M. and Childs, J. E. (2001). Infections with *Ehrlichia chaffeensis* and *Ehrlichia ewingii* in persons coinfecting with human immunodeficiency virus. Clin. Infect. Dis. 33: 1586-1594.
- [87] Thomas, L. D., Hongo, I., Bloch, K. C., Tang, Y. W. and Dummer, S. (2007). Human ehrlichiosis in transplant recipients. Am. J. Transplant. 7: 1641-1647.
- [88] Ganguly, S. and Mukhopadhyay, S. K. (2008). Tick-borne ehrlichiosis infection in human beings. J. Vector. Borne. Dis. 45: 273-280.
- [89] Duncan, A. W., Correa, M. T., Levine, J. F. and Breitschwerdt, E. B. (2005). The dog as a sentinel for human infection: prevalence of *Borrelia burgdorferi* C6 antibodies in dogs from southeastern and mid-Atlantic States. Vector. Borne. Zoonotic. Dis. 5: 101-109.
- [90] Dumler, J. S., Barbet, A. F., Bekker, C. P. J., Dasch, G. A., Palmer, G. H., Ray, S. C., Rikihisa, Y. and Rurangirwa, F. R. (2001). Reorganization of genera in the families Rickettsiaceae and Anaplasmataceae in the order Rickettsiales: unification of some species of *Ehrlichia* with *Anaplasma*, *Cowdria* and *Ehrlichia*, and *Ehrlichia* with *Neorickettsia*, descriptions of six new species combinations and designation of *Ehrlichia equi*, and "HGE agent" as subjective synonyms of *Ehrlichia phagocytophila*. Int. J. Syst. Evol. Microbiol. 51: 2145-2165.
- [91] Loftis, A. D., Levin, M. L. and Spurlock, J. P. (2008). Two USA *Ehrlichia* spp. cause febrile illness in goats. Vet. Microbiol. 130: 398-402.
- [92] Dos Santos, A. P., Dos Santos, R. P., Biondo, A. W., Dora, J. M., Goldani, L. Z., de Oliveira, S. T., de Sá Guimarães, A. M., Timenetsky, J., de Moraes, H. A., González, F. H. D. and Messick, J. B. (2008). Hemoplasma infection in HIV-positive patient Brazil. Emerg. Infect. Dis. 14: 1922-1924.
- [93] Yuan, C. L., Liang, A. B., Yao, C. B., Yang, Z. B., Zhu, J. G., Cui, L., Yu, F., Zhu, N. Y., Yang, X. W. and Hua, X. G. (2009). Prevalence of *Mycoplasma suis* (*Eperythrozoon suis*) infection in swine-farm workers in Shanghai, China. Am. J. Vet. Res. 70: 890-894.
- [94] Sykes, J. E., Lindsay, L. L., Maggi, R. G. and Breitschwerdt, E. B., (2010). Human coinfection with *Bartonella henselae* and two hemotropic *Mycoplasma* variants resembling *Mycoplasma ovis*. J. Clin. Microbiol. 48: 3782-3785.
- [95] Quinn, P. J., Markey, B. K., Leonard, F. C., Fitz Patrick, E. S., Fanning, S. and Hartiga, P. J. (2011). Veterinary microbiology and microbial disease, second ed. Wiley-Blackwell, London, UK, pp. 1048-1076.
- [96] Maurin, M., Bakken, J. S. and Dumler, J. S. (2003). Antibiotic susceptibilities of *Anaplasma* (*Ehrlichia*) *phagocytophilum* strains from various geographic areas in the United States. Antimicrob. Agents. Chemother. 47: 413-415.