
Versatility of RNA-Binding Proteins in Living Cells

Riki Kurokawa^{1,*}, Akihide Takeuchi², Nobuyuki Shiina^{3,4,5}, Masato Katahira^{6,7}, Takefumi Yamashita⁸, Yoko Matsuno⁹, Keisuke Hitachi¹⁰, Shinsuke Ishigaki^{11,12,13}, Nesreen Haamad^{6,7}, Ryoma Yoneda¹, Naomi Ueda¹, Kei Iida², Motoyasu Hosokawa², Masatoshi Hagiwara², Mamiko Iida^{6,7}, Tsukasa Mashima^{6,7}, Yudai Yamaoki⁶, Masatomo So¹⁴, Takashi Nagata^{6,7}, Gen Sobue^{11,12}, Keiko Kondo⁶, Hiroki Watanabe¹⁵, Takayuki Uchihashi¹⁵

¹Division of Gene Structure and Function, Research Center for Genomic Medicine, Saitama Medical University, Hidaka, Japan

²Department of Anatomy and Developmental Biology, Graduate School of Medicine, Kyoto University, Kyoto, Japan

³Laboratory of Neuronal Cell Biology, National Institute for Basic Biology, Okazaki, Japan

⁴Exploratory Research Center on Life and Living Systems, Okazaki, Japan

⁵Department of Basic Biology, The Graduate University for Advanced Studies (SOKENDAI), Okazaki, Japan

⁶Institute of Advanced Energy, Kyoto University, Kyoto, Japan

⁷Graduate School of Energy Science, Kyoto University, Kyoto, Japan

⁸Laboratory for Systems Biology and Medicine, Research Center for Advanced Science and Technology, the University of Tokyo, Tokyo, Japan

⁹Division of Clinical Preventive Medicine, Niigata University, Niigata, Japan

¹⁰Division for Therapies Against Intractable Diseases, Institute for Comprehensive Medical Science, Fujita Health University, Toyoake, Japan

¹¹Research Division of Dementia and Neurodegenerative Disease, Nagoya University Graduate School of Medicine, Nagoya, Japan

¹²Brain and Mind Research Center, Nagoya University, Nagoya, Japan

¹³Department of Neurology, Nagoya University Graduate School of Medicine, Nagoya, Japan

¹⁴Institute for Protein Research, Osaka University, Osaka, Japan

¹⁵Department of Physics, Nagoya University, Nagoya, Japan

Email address:

rkurokaw@saitama-med.ac.jp (R. Kurokawa)

*Corresponding author

To cite this article:

Riki Kurokawa, Akihide Takeuchi, Nobuyuki Shiina, Masato Katahira, Takefumi Yamashita, Yoko Matsuno, Keisuke Hitachi, Shinsuke Ishigaki, Nesreen Haamad, Ryoma Yoneda, Naomi Ueda, Kei Iida, Motoyasu Hosokawa, Masatoshi Hagiwara, Mamiko Iida, Tsukasa Mashima, Yudai Yamaoki, Masatomo So, Takashi Nagata, Gen Sobue, Keiko Kondo, Hiroki Watanabe, Takayuki Uchihashi. Versatility of RNA-Binding Proteins in Living Cells. *Biomedical Sciences*. Vol. 5, No. 1, 2019, pp. 7-13. doi: 10.11648/j.bs.20190501.12

Received: July 23, 2019; **Accepted:** August 14, 2019; **Published:** August 29, 2019

Abstract: This conference paper aims at demonstration of complexity or multiplicity of RNA-binding protein (RBP) and its related RNA. The data in the manuscript are based upon the third Annual Meeting of the RNA study group, held at the Tokyo office of the Kyoto University on May 23 in 2019. The purpose of the meeting is to present recent progress of the study group and have discussion regarding RNA and related molecules, especially focusing on RBP in this year. We had fourteen sessions and documented ten of them in the manuscript. There were exciting and fruitful debates. Actual presentations are shown as follows. Specific RBP Sfpq and Qk are involved in neuronal development in mouse system. Also in mouse, liquid- and solid-like RNA granule formation plays pivotal roles in neuronal functions. Biochemical experiments demonstrate that affinity profiles categorize RNA-Binding proteins into distinctive groups. This presents basis for biological divergence of RBPs. In the context of pathology, RNA aptamers against prion and Alzheimer's diseases may present therapeutic outcome. Computational analysis is another utility for approaching riddle of RBPs. The study of X-chromosome inactivation is pioneering field of long noncoding RNAs. Redundant triplex interactions between lncRNA and LINE-1s are supposed to be crucial in X-chromosome inactivation. Myogenesis-related lncRNA, *Myoparr* promotes skeletal muscle atrophy caused by denervation through the regulation of

GDF5/BMP14 expression. In neurodegenerative disorders functional loss of TLS/FUS induces onset of frontotemporal lobar degeneration. Conformational change of TLS is induced with binding of various nucleic acid molecules. TLS is also involved in phase separation into aggregation to pathogen generation of neurodegenerative diseases. The meeting has been successfully prorogued. We utilize the data to boost the activity of the field of RNA and RBP. The achievements of the meeting have confirmed pivotal roles of RBPs in living cells. It should be in complexity. We will be able to analyze intricate phenomena of RBP biological actions and contribute to uncover the veiled rules in divergent biological programs.

Keywords: RNA-Binding Protein, Long Noncoding RNA, Amyotrophic Lateral Sclerosis, TLS FUS, X Chromosome Inactivation, Phase Separation, Neuron, Computer Simulation

1. Introduction

The divergent biological activity of RNA-binding protein (RBP) has been focused owing to its contribution to biological programs including homeostasis, development, and heredity. The intention of this conference paper is to announce activity of the 3rd Annual Meeting of the RNA Study group related to divergent property of RNA-binding protein (RBP). Divergent researchers also contribute to this article. This implies that divergent readers should be attracted to the article.

The Research activity of RBPs has been radically in progress. Actually, it has been shown that more than 1500 proteins function as RBP in the human genome [1]. These RBPs contain established or canonical RNA binding domains, which include the RNA recognition motif, the heterogeneous ribonucleoprotein K homology domain, and C3H1 zinc-finger domain [1]. Some of the proteins have previously unidentified or non-canonical RBPs [2]. RBPs are involved in all aspect of RNA metabolism. Recently, it has shown that many RNA processing steps are well coupled with transcription [3]. RBPs are prevalently localized on active chromatin regions. The promoters of active genes are major hot spots for interaction with RBPs. Multiple RBPs are directly involved in transcriptional regulation [3]. It has been reported a novel algorithm called mCross to accurately define RBP specificity by registering protein-RNA crosslink sites utilizing CLIP data [4]. This technique was applied to more than one hundred RBPs and identified a non-canonical binding motif SRSF1, which denotes the protein in regulating phase separation [4]. Now, phase separation is one of the most exciting topics in the RBP field [5-7]. At the conference, we have been exploring these kinds of attractive enigmas of basic sciences.

On Thursday, May 23 in 2019, the third Annual Meeting of the RNA study group was held at the Tokyo office of the Kyoto University, located at the Tokyo station. This gave us a lot of beneficial discussion. We really appreciate every attendee there. We have obtained a plan of the meeting at the chat of the dinner after the session at the 39th annual meeting of the Molecular Biology Society of Japan in Yokohama in December, 2016. Katahira, Yamashita, and Kurokawa voluntarily have started to take charge of the meeting. We decided to have a first meeting at the Research Center for Advanced Science and Technology (RCAST) on Thursday, May 18 in 2017, and also had second meeting on Thursday,

May 24 in 2018 there [8]. We did not have any funding on it. This means that it should be a small meeting with less luxurious setting, but have the freedom to take time for discussion. We have prepared for the meeting with groping for a way to make better one. Fortunately, we have attained a great success on it. We have it as a closed format for secured exchange of unpublished data between attendees. Therefore, limited numbers of people were permitted to attend. Then, we had a plenty of time to have productive discussion at the sessions. Official meeting of the big academic society has stringent regulation of the program because of time limitation. Then, we have appreciated the liberty of discussion there.

We have determined to have the date of the fourth meeting on Thursday, May 21 in 2020 somewhere in Japan. This will lead another successful history of the Meeting.

2. Multiplicity in RBPs in Living Cells

Akihide Takeuchi *et al.* present the data regarding RBP regulation in neuronal development. To decipher physiological roles of mRNA regulation especially in nervous system development, we are analyzing the function of RBPs with multidisciplinary approaches from *in vitro* to *in vivo*. Using transcriptome studies, we identified Sfpq and Qk as highly and selectively expressed RBPs in embryonic mouse brains. Sfpq is expressed in differentiating neurons which are generated from neural stem cells. Using gene targeted mice and genome wide RNA biology approaches, we found that Sfpq critically facilitates transcriptional elongation of long neural genes >100 kbp in length and comprehensively regulates developmentally essential genes as functional clusters called "regulon [9, 10]. Qk is specifically expressed in neural stem cells in embryonic mouse brains and functions as the significant regulator in neurogenesis-to-gliogenesis switching. Qk regulates mRNAs of the certain pathway that is essential for the fate determination of neural stem cells. From series of these studies, we propose the significance of mRNA regulation in neural functions and developmental processes of brain.

Nobuyuki Shiina has shown that RNA granules are membrane-less assemblies of RNA-protein complexes formed by liquid-liquid phase separation. Recently, RNA granules have been shown to contain solid-like cores rather than just liquid droplets [11]. However, it remained unclear how these different substructures are formed and whether different RNA granule scaffold proteins induce the formation

of different substructures. We found that RNA granule scaffold proteins, when expressed separately in cultured fibroblasts, induce the formation of either liquid-type or solid-type granules. When co-expressed in the same cells, the liquid-type and solid-type scaffold proteins were combined into the same granules and formed liquid-type shells and solid-type cores therein, respectively. Being combined with liquid-type scaffold proteins, solid-type scaffold proteins increased their mobility and reduced their dose-dependent translation inhibition in the granules [12]. RNG105 (also known as caprin1) was a liquid-type scaffold protein with remarkably strong activity to change the dynamics and translation inhibition ability of solid-type scaffold proteins. This activity of RNG105 suggested that in RNG105 knockout neurons, RNA granule cores could be very immobile and translation activity in the granules could be decreased. These changes in RNA granule dynamics and translation activity might underlie reduced dendritic localization of mRNAs in RNG105 knockout neurons and severe defects in long-term memory formation in RNG105 conditional knockout mice [13].

Riki Kurokawa have established a novel assay to detect interaction of RBPs with RNA, using biotinylated RNA oligos to capture RBPs with Western blot of specific antibodies against RBPs. The assay detects RNA binding more confidently than the traditional gel shift assay. Starting with completely randomized RNA oligos from 5mer through 12mer length, their binding was examined with HeLa cell nuclear extract (NE). Coomassie brilliant blue-based (CBB) staining did not detect any strong signal. Western blot analysis of typical six RBP antibodies showed four RBPs bound with the random RNA oligos. hnRNPUL2 bound to all from 5mer to 12mer of RNA oligos, while no TLS and hnRNPH1 signal was detected in the random RNA oligo samples. Next, base specificity was examined using sets of oligos of RNA fixed at G, A, U, and C (GAUC RNA oligos). The RNA oligos fixed at "G" (G RNA oligos) have the most prominent protein bands. A, U, and C of RNA oligos were shown to bind less numbers of protein. Western blot indicated that hnRNPUL2 and hnRNPU bind all four oligos of GAUC at the 10mer length. Contrarily, TLS and hnRNPH1 have no binding with these oligos of GAUC. Then, poly G, A, U and C of RNA at the length of 100mer were tested to see binding profile of RBPs. The CBB staining of the fractions bound with these four polymers of RNA showed that more bands were bound than GAUC RNA oligos. hnRNPU bound well to poly G, A, and, U, but slightly less to poly C. Intriguingly, TLS and hnRNPH1 have binding only to poly G, and also to their common specific sites consisting of GGUG motifs. These data demonstrate that RNA binding is regulated with three factors, length, base composition, and sequence. Furthermore, hnRNPU and hnRNPUL2 have low specificity binding to RNAs, while TLS and hnRNPH1 exert high specific binding. These different propensities in bindings of RBPs are supposed to support specific biological roles in living cells [14].

Mamiko Iida and Masato Katahira identified an RNA

aptamer against a prion protein, r (GGAGGAGGAGGA) (R12). We showed that R12 forms a unique quadruplex structure with a tetrad and a hexad by NMR and reduces a level of the abnormal prion protein (PrP), PrP^{Sc}, in the mouse neuronal cells, implying its therapeutic potential as to prion diseases [15, 16]. This time, we developed an RNA aptamer that exhibits much higher anti-prion activity than R12 through the logical design of the RNA sequence on the basis of the R12 sequence. The structure of this aptamer was determined by NMR. The origin of the much higher anti-prion activity was rationally interpreted by this structure. This aptamer has therapeutic potential as to the prion disease. The neurodegenerative disorder of Alzheimer's disease (AD) is associated with the accumulation of misfolded proteins. It was revealed that the amyloid β (A β) oligomers bind to prion protein, which acts as a receptor on the cell membrane, resulting in AD [17]. Thus, it is thought that compounds that can disrupt the formation of the prion-A β oligomer complex may prevent AD. Fluorescence anisotropy revealed that R12 is capable of binding to PrP, resulting in dissociation of PrP with A β . This is the first demonstration of disruption of the PrP-A β complex by a nucleic acid [18]. This ability of R12 implied its therapeutic potential as to AD.

Takefumi Yamashita presents a theoretical approach using molecular dynamics (MD) simulations of proteins and RNAs. MD simulations have been applied to many phenomena of biological polymers (e.g. protein-protein interaction) [19-24] as well as chemical reactions [25-27]. In these studies, detection of structural change is essential. Recently, we have proposed that the mathematics-based approach and machine learning-based approach are useful to detect the characteristic structural change efficiently [28, 29]. For example, the persistent homology analysis method, one of the computational mathematics methods, characterizes point clouds. By applying this method to the water molecules of the antigen-antibody complex solution system, we quickly detected water molecules located in the antigen-antibody interface [29], which should largely influence the binding entropy [22]. The persistent homology analysis does not require any thermodynamic knowledge, but can automatically detect characteristic water distributions, which may influence thermodynamic properties.

Yoko Matsuno presents data of redundant triplex interactions between lncRNA and LINE-1s in X-chromosome inactivation [30]. Associations between X-inactive transcript (Xist) -long noncoding RNA (lncRNA) and chromatin are crucial intermolecular interactions in the process of X-chromosome inactivation (XCI). Even with high-resolution analyses of the Xist RNA-binding sites, specific interaction sequences are yet to be identified. Using a combination of bioinformatics, computational simulations, and quartz crystal microbalance measurements *in vitro*, we characterized redundant UC/TC (r-UC/TC) or AG (r-AG) motifs (≥ 5 nucleotides in length) that are abundantly localized in Xist/XIST/RNA-on-the-silent X (R_{sx}) RNAs and long interspersed nuclear element 1 (LINE-1) in mouse, human, and opossum genomes. These multiple short motifs in lncRNA and

DNA molecules could be brought together to form Hoogsteen or reverse Hoogsteen triplexes, which co-operatively enhance their associations. In addition, we showed the conservation of r-AG motifs in LINE-1s of any length (100-7,000 bp) or base composition, possibly through convergence in LINE-1 nucleotide variations, which provided a similar motif-length distribution across the three species (Kolmogorov–Smirnov distance <0.100). On the basis of this conserved motif-length distribution pattern, the triplex interactions would preferentially occur via shorter motifs (5-7 nucleotide in length) with their high occupancy rate, whereas longer motifs (≥ 8 nucleotides in length), although fewer in number, would provide more chances for multiple shorter target motifs to bind. This mode of interaction may impart redundant properties to the association between lncRNAs and LINE-1s in the XCI process.

Keisuke Hitachi *et al.* have shown that *Myoparr* promotes skeletal muscle atrophy caused by denervation through the regulation of *GDF5/BMP14* expression. Hundreds of long non-coding RNAs (lncRNAs) are expressed from the cis-regulatory regions including promoter and enhancer. The promoter region of the *myogenin* gene, coding for a pivotal regulator of myogenesis, produces a lncRNA *Myoparr*.

Myoparr regulates myogenesis through the activation of neighboring *myogenin* gene expression and skeletal muscle miRNA expressions (*miR-133b/206* and *miR-675*) [31]. Intriguingly, *Myoparr* also promotes skeletal muscle atrophy caused by surgical denervation. However, the downstream genes of *Myoparr* during muscle atrophy remain unclear. We found that *Myoparr* activates the expression of *myogenin* gene, which also promotes neurogenic muscle atrophy by inducing the expression of E3 ubiquitin ligases. In addition, RNA-seq analysis revealed that *Myoparr* decreases the expression of *GDF5/BMP14* gene to repress the activity of BMP signaling [32]. BMP signaling increases the anabolic pathway and antagonizes neurogenic muscle atrophy. Thus, our findings indicate that *Myoparr* promotes neurogenic muscle atrophy by increasing *myogenin* expression and decreasing *GDF5/BMP14* expression. Moreover, our comparative transcriptome analysis between mouse myoblast cell line C2C12 and mouse tibialis anterior muscles showed that the downstream genes of *Myoparr* largely differ between myogenic differentiation and muscle atrophy condition, indicating that the cell status and context affect the downstream genes of lncRNAs, possibly depending on their binding proteins.




Figure 1. Summary of the Talks.

Shinsuke Ishigaki and Gen Sobue show Importance of Functional Loss of FUS in in Neurodegenerative Disorders. Fused in sarcoma (FUS) is an RBP that regulates RNA metabolism including alternative splicing, transcription, and RNA transportation. FUS is genetically and pathologically involved in amyotrophic lateral sclerosis (ALS) and frontotemporal lobar degeneration (FTLD). FUS-related ALS/FTLD pathology is characterized by mislocalization of FUS to the cytoplasm with a reduction in nuclear expression in affected neurons, implying that functional loss of FUS

could lead to neuronal dysfunction and/or neuronal cell death. We have reported that FUS and SFPQ regulate alternative splicing of *Mapt* gene at exon10 which generates two isoforms of neural microtubule-associated protein tau (Tau) protein. Silencing of FUS or SFPQ in the hippocampus of mice resulted in the increased ratio of 4-repeat tau (4R-tau) / 3-repeat tau (3R-tau) followed by FTLD-like abnormal behaviors, accumulation of phosphorylated tau, and neuronal loss [33]. Additionally, loss of FUS function can affect dendritic spine maturations by destabilizing mRNAs such as

Glutamate receptor 1 (GluA1) [34], and Synaptic Ras GTPase-activating protein 1 (SynGAP1) [35]. Since abnormalities in post-synapse are often observed in neurodegenerative diseases, a link among loss of FUS function, tau isoform alteration, aberrant post-synaptic function, and phenotypic expression, might be a possible sequential biological cascade leading to FTLD.

Nesreen Haamad and Masato Katahira have shown the conformational change of translocated in liposarcoma, TLS, upon binding to various non-coding nucleic acids. Translocated in liposarcoma (TLS), also known as fused in sarcoma (FUS), is a multifunctional RNA/DNA binding protein. TLS mutations are associated with numerous neurodegenerative disorders, such as amyotrophic lateral sclerosis (ALS). TLS plays a series of critical roles in RNA processing, including transcription, splicing, transport, translation, and decay. DNA damage triggers the transcription of a long non-coding RNA (lncRNA) from the 5' upstream region of a cyclin D1 gene (*CCND1*), named promoter-associated non-coding RNA (pncRNA) [36, 37]. Subsequently, TLS is recruited to the damage site. The C-terminal region of TLS interacts with pncRNA, which allows N-terminus TLS to bind CREB-binding protein (CBP)/E1A-binding protein P300 (p300) and to regulate the histone acetyltransferase HAT activity, resulting in transcription repression of *CCND1*. Although it was proposed that the interaction between TLS and pncRNA can convert TLS from closed to open conformation, this structural change has not been experimentally confirmed, so far. This time, we have spectroscopically detected the structural change of TLS upon binding pncRNA. Then, we have further confirmed the structural change on the basis of images of TLS.

Ryoma Yoneda shows phase separation and fibrillization of TLS/FUS-effect of mutation, RNase, and drugs. TLS/FUS is a DNA/RNA binding protein whose mutations cause familial amyotrophic lateral sclerosis (ALS). Severe ALS-related mutations occur in nuclear localization signal at the C-terminal end of TLS/FUS. These mutations cause TLS/FUS aggregation in the cytoplasm, and leads to cytotoxicity. In order to reproduce the TLS/FUS aggregation *in vitro*, we constructed GFP-fusion TLS/FUS with two kinds of ALS-related mutations (H517D and R521G). We first examined the phase separation of wild type (wt) and mutated TLS/FUS in different NaCl concentration, and observed promoted phase separation in low NaCl concentration, but no specific difference between wt and mutated TLS/FUS. RNase treatment also facilitated phase separation of all three TLS/FUS. We finally tested the effect of bisox, a reagent induces TLS aggregation [38], and similar dipeptide named bLH on TLS/FUS fibrillization. As expected, bisox induced fibrillization of all three TLS/FUS examined, and interestingly, bLH inhibited the TLS/FUS aggregation formed by bisox. Further analysis is required to reveal the inhibitory effect of bLH on bisox independent TLS/FUS fibrillization.

3. Conclusion




Figure 2. Conclusion of the Conference RNA-binding protein plays a central role in RNA function in living cells.

The presentations of the meeting cover variety of biomedical sciences, indicating the significance of RBPs and cognate RNA molecules in each field (Figure 1). The achievements of the meeting are to make sure a point of view of versatility of RBPs in living cells. It should be in complexity or chaos. We will focus on analyzing such intricate phenomena of RBP actions and unraveling the hidden rules. We will have more fruitful activity in the next year's meeting.

RNAs with divergent sequences are transcribed from the human genome. The RNAs have their own intrinsic property, but not so many RNAs are functional themselves. Most of RNAs needs to capture their binding partner for exerting their function. It is RBPs which mediate or integrate divergent sequence information of the RNAs into biological actions in living cells (Figure 2). Therefore, RBPs function as mediator or integrator for divergent RNA sequences. The possible model is presented there (Figure 2). Then, we will extensively examine the model to demonstrate exact molecular mechanism of RBPs in living organisms.

Funding

Takeuchi: This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (MEXT, JSPS KAKENHI 19500269, 25500288, 21249013, 15H05721), Innovative Cell Biology by Innovating Technology (Cell Innovation), a Core Research for Evolutional Science and Technology (CREST) grant from the Japan Science and Technology Agency (JST), a grant from the Japan Agency for Medical Research and Development (AMED), the Asian CORE Program of JSPS, iCeMS Cross-Disciplinary Research Promotion Project of Kyoto University, and the Fujiwara Memorial Foundation; Shiina: This research was supported by the Takeda Science Foundation, the Astellas Foundation for Research on Metabolic Disorders, and the Orion Project of Okazaki Institute for Integrative Bioscience; Kurokawa: This work

was supported in part by a Grant-in-aid for “Support Project of Strategic Research Center in Private Universities” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) to Saitama Medical University Research Center for Genomic Medicine, and by KAKENHI from JSPS to MEXT of Japan (KibanB nos22390057; 25293073, KibanC 18K06939); Katahira: This work was supported by the Ministry of Education, Culture, Sports, Science and Technology of Japan (18H04550 and 18K19397); Yamashita: This research used computational resources of the TSUBAME through the HPCI System Research Project (hp180044, hp190117), and of the Research Center for Computational Science, Okazaki, Japan. T. Y. acknowledges support from the SIP program (Structural Materials for Innovation), Scientific Research (JSPS KAKENHI (S) 15H05752, (C) 15KT0103, (C) 18K05025), and Innovative Drug Discovery Infrastructure through Functional Control of Bio-molecular Systems (hp180191, hp190171); Hitachi: This work was supported by JSPS KAKENHI (17K08646 and 19H03427); Yoneda: This work was supported in part by KAKENHI from JSPS to MEXT of Japan (17K18065).

Acknowledgements

Kunihiro Tsuchida, at the Division for Therapies against Intractable Diseases, Institute for Comprehensive Medical Science, Fujita Health University, has a significant experimental contribution to the project.

References

- [1] Gerstberger S, Hafner M & Tuschl T (2014) A census of human RNA-binding proteins. *Nature Reviews Genetics* 15, 829, doi: 10.1038/nrg3813
<https://www.nature.com/articles/nrg3813#supplementary-information>.
- [2] Bao X, Guo X, Yin M, Tariq M, Lai Y, Kanwal S, Zhou J, Li N, Lv Y, Pulido-Quetglas C, et al. (2018) Capturing the interactome of newly transcribed RNA. *Nature Methods* 15, 213, doi: 10.1038/nmeth.4595
<https://www.nature.com/articles/nmeth.4595#supplementary-information>.
- [3] Xiao R, Chen J-Y, Liang Z, Luo D, Chen G, Lu ZJ, Chen Y, Zhou B, Li H, Du X, et al. (2019) Pervasive Chromatin-RNA Binding Protein Interactions Enable RNA-Based Regulation of Transcription. *Cell* 178, 107-121. e118, doi: 10.1016/j.cell.2019.06.001.
- [4] Feng H, Bao S, Rahman MA, Weyn-Vanhentenryck SM, Khan A, Wong J, Shah A, Flynn ED, Krainer AR & Zhang C (2019) Modeling RNA-Binding Protein Specificity In Vivo by Precisely Registering Protein-RNA Crosslink Sites. *Molecular Cell* 74, 1189-1204. e1186, doi: 10.1016/j.molcel.2019.02.002.
- [5] Hnisz D, Shrinivas K, Young RA, Chakraborty AK & Sharp PA (2017) A Phase Separation Model for Transcriptional Control. *Cell* 169, 13-23, doi: 10.1016/j.cell.2017.02.007.
- [6] Boija A, Klein IA, Sabari BR, Dall’Agnese A, Coffey EL, Zamudio AV, Li CH, Shrinivas K, Manteiga JC, Hannett NM, et al. (2018) Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell* 175, 1842-1855. e1816, doi: 10.1016/j.cell.2018.10.042.
- [7] Ries RJ, Zaccara S, Klein P, Olarerin-George A, Namkoong S, Pickering BF, Patil DP, Kwak H, Lee JH & Jaffrey SR (2019) m6A enhances the phase separation potential of mRNA. *Nature* 571, 424-428, doi: 10.1038/s41586-019-1374-1.
- [8] Kurokawa R, Komiya R, Oyoshi T, Matsuno Y, Tani H, Katahira M, Hitachi K, Iwashita Y, Yamashita T, Kondo K, et al. (2018) Multiplicity in Long Noncoding RNA in Living Cells. *Biomedical Sciences* 4, 18-23, doi: 10.11648/j.bs.20180402.11.
- [9] Takeuchi A, Iida K, Tsubota T, Hosokawa M, Denawa M, Brown JB, Ninomiya K, Ito M, Kimura H, Abe T, et al. (2018) Loss of Sfpq Causes Long-Genes Transcriptopathy in the Brain. *Cell Reports* 23, 1326-1341, doi: 10.1016/j.celrep.2018.03.141.
- [10] Hosokawa M, Takeuchi A, Tanihata J, Iida K, Takeda S & Hagiwara M (2019) Loss of RNA-Binding Protein Sfpq Causes Long-Genes Transcriptopathy in Skeletal Muscle and Severe Muscle Mass Reduction with Metabolic Myopathy. *iScience* 13, 229-242, doi: 10.1016/j.isci.2019.02.023.
- [11] Jain S, Wheeler JR, Walters RW, Agrawal A, Barsic A & Parker R (2016) ATPase-Modulated Stress Granules Contain a Diverse Proteome and Substructure. *Cell* 164, 487-498, doi: 10.1016/j.cell.2015.12.038.
- [12] Shiina N (2019) Liquid- and solid-like RNA granules form through specific scaffold proteins and combine into biphasic granules. *The Journal of biological chemistry* 294, 3532-3548, doi: 10.1074/jbc.RA118.005423.
- [13] Nakayama K, Ohashi R, Shinoda Y, Yamazaki M, Abe M, Fujikawa A, Shigenobu S, Futatsugi A, Noda M, Mikoshiba K, et al. (2017) RNG105/caprin1, an RNA granule protein for dendritic mRNA localization, is essential for long-term memory formation. *eLife* 6, doi: 10.7554/eLife.29677.
- [14] Ueda N & Kurokawa R (2018) Affinity Profiles Categorize RNA-Binding Proteins into Distinctive Groups. *Biomedical Sciences* 4, 24-31, doi: 10.11648/j.bs.20180403.11.
- [15] Mashima T, Nishikawa F, Kamatari YO, Fujiwara H, Saimura M, Nagata T, Kodaki T, Nishikawa S, Kuwata K & Katahira M (2013) Anti-prion activity of an RNA aptamer and its structural basis. *Nucleic Acids Research* 41, 1355-1362, doi: 10.1093/nar/gks1132.
- [16] Hayashi T, Oshima H, Mashima T, Nagata T, Katahira M & Kinoshita M (2014) Binding of an RNA aptamer and a partial peptide of a prion protein: crucial importance of water entropy in molecular recognition. *Nucleic Acids Research* 42, 6861-6875, doi: 10.1093/nar/gku382.
- [17] Lauren J, Gimbel DA, Nygaard HB, Gilbert JW & Strittmatter SM (2009) Cellular prion protein mediates impairment of synaptic plasticity by amyloid-beta oligomers. *Nature* 457, 1128-1132, doi: 10.1038/nature07761.
- [18] Iida M, Mashima T, Yamaoki Y, So M, Nagata T & Katahira M (2019) The anti-prion RNA aptamer R12 disrupts the Alzheimer’s disease-related complex between prion and amyloid beta. *The FEBS journal* 286, 2355-2365, doi: 10.1111/febs.14819.

- [19] Yamashita T (2018). Toward rational antibody design: recent advancements in molecular dynamics simulations. *International Immunology* 30, 133-140, doi: 10.1093/intimm/dxx077.
- [20] Yamashita T (2016) Towards Physical Understanding of Molecular Recognition in the Cell: Recent Evolution of Molecular Dynamics Techniques and Free Energy Theories. *Biomedical Sciences* 2, 34-47, doi: 10.11648/j.bs.20160205.11.
- [21] Koyama T, Nakamoto M, Morishima K, Yamashita R, Yamashita T, Sasaki K, Kuruma Y, Mizuno N, Suzuki M, Okada Y, et al. (2019) A SNP in a Steroidogenic Enzyme Is Associated with Phenotypic Sex in *Seriola* Fishes. *Current Biology* 29, 1901-1909 e1908, doi: 10.1016/j.cub.2019.04.069.
- [22] Yamashita T, Mizohata E, Nagatoishi S, Watanabe T, Nakakido M, Iwanari H, Mochizuki Y, Nakayama T, Kado Y, Yokota Y, et al. (2019) Affinity Improvement of a Cancer-Targeted Antibody through Alanine-Induced Adjustment of Antigen-Antibody Interface. *Structure* 27, 519-527 e515, doi: 10.1016/j.str.2018.11.002.
- [23] Miyanabe K, Yamashita T, Abe Y, Akiba H, Takamatsu Y, Nakakido M, Hamakubo T, Ueda T, Caaveiro JMM & Tsumoto K (2018) Tyrosine Sulfation Restricts the Conformational Ensemble of a Flexible Peptide, Strengthening the Binding Affinity for an Antibody. *Biochemistry* 57, 4177-4185, doi: 10.1021/acs.biochem.8b00592.
- [24] Yamashita T & Takamatsu Y (2017) An ensemble docking calculation of lysozyme and HyHEL-10: Insight into the binding mechanism. *AIP Conference Proceedings* 1906, 030022, doi: 10.1063/1.5012301.
- [25] Yamashita T (2014) Properties of a Hydrated Excess Proton Near the Cholesterol-Containing Phospholipid Bilayer. *JPS Conference Proceedings* 1, 013086, doi: 10.7566/JPSCP.1.013086.
- [26] Xu J, Yamashita T, Agmon N & Voth GA (2013) On the origin of proton mobility suppression in aqueous solutions of amphiphiles. *The Journal of Physical Chemistry B* 117, 15426-15435, doi: 10.1021/jp4051726.
- [27] Yamashita T (2015) Improvement in Empirical Potential Functions for Increasing the Utility of Molecular Dynamics Simulations. *JPS Conference Proceedings* 5, 010003, doi: 10.7566/JPSCP.5.010003.
- [28] Yamashita T, Okajima R & Shoji N (2018) Efficiency strategy for peptide design: A comparative study on all-atom, coarse-grained, and machine learning approaches. *AIP Conference Proceedings* 2040, 020014, doi: 10.1063/1.5079056.
- [29] Sasaki K, Okajima R & Yamashita T (2018) Liquid structures characterized by a combination of the persistent homology analysis and molecular dynamics simulation. *AIP Conference Proceedings* 2040, 020015, doi: 10.1063/1.5079057.
- [30] Matsuno Y, Yamashita T, Wagatsuma M & Yamakage H (2019) Convergence in LINE-1 nucleotide variations can benefit redundantly forming triplexes with lncRNA in mammalian X-chromosome inactivation. *Mobile DNA* 10, 33, doi: 10.1186/s13100-019-0173-4.
- [31] Hitachi K, Nakatani M, Takasaki A, Ouchi Y, Uezumi A, Ageta H, Inagaki H, Kurahashi H & Tsuchida K (2019) Myogenin promoter-associated lncRNA Myoparr is essential for myogenic differentiation. *EMBO Reports* 20, e47468, doi: 10.15252/embr.201847468.
- [32] Hitachi K, Nakatani M & Tsuchida K (2019) Long Non-Coding RNA Myoparr Regulates GDF5 Expression in Denervated Mouse Skeletal Muscle. *Non-coding RNA* 5, E33, doi: 10.3390/ncrna5020033.
- [33] Ishigaki S, Fujioka Y, Okada Y, Riku Y, Udagawa T, Honda D, Yokoi S, Endo K, Ikenaka K, Takagi S, et al. (2017) Altered Tau Isoform Ratio Caused by Loss of FUS and SFPQ Function Leads to FTL-like Phenotypes. *Cell Reports* 18, 1118-1131, doi: 10.1016/j.celrep.2017.01.013.
- [34] Udagawa T, Fujioka Y, Tanaka M, Honda D, Yokoi S, Riku Y, Ibi D, Nagai T, Yamada K, Watanabe H, et al. (2015) FUS regulates AMPA receptor function and FTL/ALS-associated behaviour via GluA1 mRNA stabilization. *Nature communications* 6, 7098, doi: 10.1038/ncomms8098.
- [35] Yokoi S, Udagawa T, Fujioka Y, Honda D, Okado H, Watanabe H, Katsuno M, Ishigaki S & Sobue G (2017) 3'UTR Length-Dependent Control of SynGAP Isoform alpha2 mRNA by FUS and ELAV-like Proteins Promotes Dendritic Spine Maturation and Cognitive Function. *Cell Reports* 20, 3071-3084, doi: 10.1016/j.celrep.2017.08.100.
- [36] Wang X, Arai S, Song X, Reichart D, Du K, Pascual G, Tempst P, Rosenfeld MG, Glass CK & Kurokawa R (2008) Induced ncRNAs allosterically modify RNA-binding proteins in cis to inhibit transcription. *Nature* 454, 126-130, doi: nature06992[pii]10.1038/nature06992.
- [37] Song X, Wang X, Arai S & Kurokawa R (2012) Promoter-associated noncoding RNA from the CCND1 promoter. *Methods in molecular biology* 809, 609-622, doi: 10.1007/978-1-61779-376-9_39.
- [38] Ueda N, Kashiwazaki G, Bando T & Kurokawa R (2017) Biotin-Lys-His Blocks Aggregation of RNA-binding Protein TLS, a Cause of Amyotrophic Lateral Sclerosis. *Biomedical Sciences* 3, 67-77, doi: 10.11648/j.bs.20170304.11.