



---

# Application of Lemongrass and Cinnamon Pesticides on Rice Ear Bug (*Leptocorisa oratorius* Fabricius) in Padi Plant

**Christina Salaki, Vivi Montong**

Study Program of Plant Protection, Faculty of Agriculture, Sam Ratulangi University, North Sulawesi, Indonesia

**Email address:**

christinasalaki@gmail.com (C. Salaki)

**To cite this article:**

Christina Salaki, Vivi Montong. Application of Lemongrass and Cinnamon Pesticides on Rice Ear Bug (*Leptocorisa oratorius* Fabricius) in Padi Plant. *International Journal of Applied Agricultural Sciences*. Vol. 5, No. 5, 2019, pp. 105-108. doi: 10.11648/j.ijaas.20190505.11

**Received:** July 24, 2019; **Accepted:** August 13, 2019; **Published:** September 4, 2019

---

**Abstract:** Lemongrass and cinnamon plant are known holding active compounds that can be used as raw materials of plant pesticides. It is related with their ability to kill, extrude, and prevent the pest insects from feeding, including rice ear bug *Leptocorisa oratorius*, one of the dangerous and uncontrollable padi plant pests. This study was aimed at finding the best formula of lemongrass and cinnamon-based plant pesticide formulation to control rice ear bug *L. oratorius*, to know the killing ability, and to do the pathogenicity test. It was carried out in a greenhouse of Pest and Disease Department, Faculty of Agriculture, UNSRAT, Manado, in March to September 2018. The treatments consisted of mixed lemongrass and cinnamon at concentrations of 2 cc/L, 4 cc/L, 6 cc/L, 8 cc/L, 10 cc/L, and control. Parameters observed were nymph mortalities at 12, 24, 48, and 72 hours after application. The treatments were administered by spraying the insect, the plant, and the insect on the plant. Results showed that spraying the insect with rice ear bug pesticide could result in 36.7 – 86.7% of mortality and with cinnamon pesticide could kill 56.7 – 93.3%, spraying the plant with rice ear bug and cinnamon pesticides could cause 3.3 – 36.7% and 6.7 – 40% of mortality, respectively, while spraying the insect on the plant could make 60 – 100% of mortality using rice ear bug pesticide and 70 – 100% of mortality using cinnamon pesticide at 12 and 72 hours after the application.

**Keywords:** Formulation, Pathogenicity, Insect, Mortality, Spraying Method

---

## 1. Introduction

Paddy is an important agricultural commodity in Indonesian people's life and has become priority in agricultural program development. Therefore, padi planting has still been the backbone of rural economy up to now [1]. One of the constraints in growing padi in Indonesia is invasion of rice ear bug *L. oratorius* (Hemiptera; Alydidae). This insect has become important pest for padi plant. It is easily recognized from the form with body length of 2 cm, grey-brown colour, and has proboscis to suck the plant liquid. *L. oratorius* sucks the plant liquid of panicle (paniculae) and padi fruit at maturing phase and makes the plant lack of nutrient and turn yellow (chlorosis), and gradually weaken. The name of the insect has exhibited its defensive mechanism by releasing stinky unpleasant aroma [2].

*L. oratorius* is cause of production loss. Invasion of one

individual per night per week can cause 27% production loss. Maintaining padi productivity in Indonesia from decline needs the way to prevent the invasion of this insect pest, one of which is the use of essential oil active compound-based plant pesticide [3, 4]. Essential oil is one of the plant materials derived from leaf, flower, stem, seeds, and flower bud. One of the promising essential oil as plant insecticide is lemongrass and cinnamon oil [5-7].

This plant contains about 32-45% citronella, 10-12% geraniol, 11-15% citronellol, 3-8% geranyl acetate, 2-4% citronellal acetate, and some sesquiterpenes. Lemongrass can control flour beetles (*Tribolium* sp), weevils (*Sitophilus* sp), cowpea weevils (*Callosobruchus* sp), nematode (*Melodogyne* sp), and fungi [8-9]. It is also potentially used as plant pesticide because lemongrass contains methyl heptanone that is repellent to insects [10-12] insecticidal to flies [13, 14], and can be used to control the mosquito bugs *Helopeltis antonii* [15-17].

Citronellol content of lemongrass oil [18] is toxic and reduces the reproductive ability of insects. Citronellal compound is a contact toxin and causes dehydration, so that the insect will dehydrate and die [19], besides lemongrass oil aroma has efficacy to exclude mosquitoes [20, 21]. Lemongrass oil at concentration of 2.5% is efficient to reduce number of flies resting and their live larvae on the dead carp [22]. Laboratory experiment [23] shows that at the concentration of 0.4% lemongrass oil can reduce 55-66% egg placement of *Helicoverpa armigera*.

Cinnamon (*Cinnamomum burmanii*) belongs to potential plant to be developed as source of raw material of plant pesticide industry since it contains numerous toxic components to insects as anti-feedant (affecting feeding) and antihormonal (disturbing hormone formation). Major content of cinnamon is cinnamaldehyde [24].

Essential oil of cinnamon wastes is evident to be able to prevent plant pathogenic fungi, such as *Fusarium oxysporum*, *F. sp. vanillae* as cause of rotten trunk disease of vanilla, *F. oxysporum* f. sp. *Zingiberi* as cause of rotten rhizome of ginger, and *Phytophthora capsici* as cause of rotten trunk of pepper plant. In relation with the potential of plant-based pesticides to control plant pests and diseases [17, 25, 26]. This study was conducted in order to obtain the best plant insecticide formula of lemongrass and cinnamon oil that effectively control the attack of *L. oratorius* on padi plant. This study was aimed to know the lethal concentration of lemongrass oil and cinnamon oil based plant pesticide against *L. oratorius* on padi plant.

## 2. Research Methodology

The experiment was carried out in the greenhouse of Plant Pest and Disease Department, Faculty of Agriculture UNSRAT Manado from March to September 2018. The study used big wire cage, small wire cage, polybag, 30 days old-padi plant, *L. oratorius* nymphs, lemongrass and cinnamon pesticides, hand sprayer, and other materials.

The test insects *L. oratorius* were collected from padi plant production centre. Eggs and nymphs were reared and multiplied on padi plant placed in the wire cage. The nymphs were taken from the third instar nymphs of the second imago generation.

The study used with 6 treatments and 4 replications. Treatments employed 2 types of pesticides, cinnamon and lemongrass at the concentration of 2 cc/L, 4 cc/L, 6 cc/L, 8 cc/L, and 10 cc/L, and water as control.

Pesticide application employed 3 techniques: 1) Insect spraying. Ten test insects were placed in the flask and sprayed with sufficient amount of pesticide, then invested on the padi plant in the cage; 2) Plant spraying. Entire padi plants were evenly sprayed with pesticide solution and placed in the cage. After wind-dried, 10 test insects were placed on each individual plant; 3) Spraying insects on plant. Thirty insects were invested into the cage of padi plant and sprayed evenly with pesticide. Parameter observed was nymph mortality at 12, 24, 48, and 72 hours after

administration. Mortality was calculated following Abbot [27] as follows:

$$M = \frac{A}{B} \times 100 \tag{1}$$

Where M = mortality, A= number of insects die, and B= number of test insects

## 3. Results and Discussion

### A. Lemongrass application to *L. oratorius*

Mortality level of *L. oratorius* seems to be higher in direct spray on the insect than on the plant (Table 1). The pesticide works as contact toxin and nerve toxin. It enters through body pores or mouth together with food. This compound then goes into the digestive organ and absorbed by the intestinal wall and translocated to the nerve centre so that the nerve system is disturbed and affects the equilibrium of nerve system ions and kills the insect [28].

**Table 1.** Mortality of rice ear bug (*L. oratorius*) treated with lemongrass-based pesticide.

Method	Concentration (cc/L)	Mortality (%)			
		12 h	24 h	48 h	72 h
Insect spraying	2 cc/L	0	13.3	20	36.7
	4 cc/L	0	20	33.3	40
	6 cc/L	10	36.7	50	63.3
	8 cc/L	13.3	53.3	60	73.3
	10 cc/L	26.7	60	73.3	86.7
	Control	0	0	0	0
Plant spraying	2 cc/L	0	0	0	3.3
	4 cc/L	0	0	0	6.7
	6 cc/L	0	0	3.3	20
	8 cc/L	3.3	3.3	6.7	26.7
	10 cc/L	3.3	6.7	26.7	36.7
	Control	0	0	0	0
Insect spraying on plant	2 cc/L	3.3	20	53.3	60
	4 cc/L	3.3	26.7	66.7	76.7
	6 cc/L	10	50	73.3	83.3
	8 cc/L	13.3	60	80	93.3
	10 cc/L	16.7	70	93.3	100
	Control	0	0	0	0

Other factor affecting the mortality of *L. oratorius* beside spraying method application is active compound. The presence of essential oil in lemongrass, such as citronella and limonoid, influences the movement of the insect *L. oratorius* [29]. The higher the concentration of essential oil sprayed is, the faster the pesticide kill the insects.

Observations showed the pesticide spray made the insect's limb be stationary and stiff. It could result from that the essential oil of the lemongrass enters through the cuticle of *L. oratorius*. As intestinal toxin, limonoid could enter the digestive tract through digestive tract. The pesticide will go into the digestive organ and disturb the metabolism of the insect so that the insect will lack of energy for living activities and kill the insect. The essential oil could directly penetrate the integument, trachea, or other sensor glands [29].

Lemongrass leaf contains saponin, flavonoid, alkaloid, polyphenol, and citronella that were potential to kill insects.

Besides these compounds are toxic, they could also inhibit the appetite. Similar findings are also reported in previous studies that the active compounds in the essential oil function as insect killer and reduce the insect reproductive ability [30-32]. The concentration of lemongrass-based pesticide at 1-7% can kill 81.15-98.06% of aphid insects [30].

Citronella content in the lemongrass oil is repellent at low concentration and insecticidal at high concentration. Concentration of 4.000 ppm could kill the mosquito bug *Helopeltis* spp., cacao fruit sucking pest insects [15]. Besides being insect repellent, the citronella works as contact toxin. Contact toxin of citronella works to inhibit acetylcholinesterase enzyme so that phosphorylation of amino acid serine occurs. Poisoning symptoms appears due to acetylcholine hoarding that causes the central nerve system disturbance, seizures, respiration failure, and mortality [15].

#### B. Cinnamon pesticide application on *L. oratorius*

Cinnamon pesticide could cause > 70% mortality at the concentration of 4 cc/L, 6 cc/L, 8 cc/L, and 10 cc/L at 72 hours, and it is much higher than that in control treatment.

**Table 2.** Mortality of rice ear bug (*L. oratorius*) treated with cinnamon-based pesticide.

Method	Concentration (cc/L)	Mortality			
		12 h	24 h	48 h	72 h
Insect spraying	2 cc/L	0	20	50	56.7
	4 cc/L	3.3	43.3	56.7	70
	6 cc/L	10	53.3	70	86.7
	8 cc/L	10	63.3	90	93.3
	10 cc/L	13.3	66.7	93.3	93.3
Plant spraying	Control	0	0	0	0
	2 cc/L	0	0	0	3.3
	4 cc/L	0	0	0	6.7
	6 cc/L	0	0	3.3	10
	8 cc/L	3.3	3.3	16.7	26.7
Spraying insect on plant	10 cc/L	3.3	6.7	30	40
	Control	0	0	0	0
	2 cc/L	3.3	40	60	70
	4 cc/L	6.7	46.7	66.7	80
	6 cc/L	6.7	53.3	70	86.7
Control	8 cc/L	13.3	66.7	80	93.3
	10 cc/L	26.7	76.7	90	100
	Control	0	0	0	0

Results showed that the mortality of *L. oratorius* varied at different concentration with the highest at 10 cc/L in insect on plant spraying method and the lowest in control treatment (Table 2). This study also found that higher concentration level increased the toxin content of the cinnamon extract indicated with higher mortality of *L. oratorius*. Previous study has found that cinnamon contains anti-feedant compound that reduces the appetite and the reproductive ability [33]. The use of 1% (b/v) cinnamon-based pesticide can kill 95.7% pest insects [34, 35].

Cinnamon oil-based pesticide at the concentration of 0.8% is effective to control *Helopeltis theivora* with killing ability of 30% [16, 36]. Cinnamon oil has also insecticidal effect on the biological aspect of *Aspidomorpha milliaris* [24].

The examination of cinnamon extract on *L. oratorius* could influence the physiology of the insect growth indicated by the behaviour and the feeding activity. The insect growth inhibition feeding on the cinnamon extract-containing plant could result from energy allocation for toxic compound detoxification. Inhibition of protein synthesis also occurs due to the disturbance of protein breakdown of the food that causes increased mortality during the molting process.

## 4. Conclusion

The use rice ear bug and cinnamon plant pesticides could kill the pest *L. oratorius*. The experiment showed that 10% concentration resulted the highest mortality of the pest, while the most effective method used 2 cc/L at 72 hours and could give  $\geq 50\%$  mortality. However, the use of these types of pesticides needs to be tested in the field.

## References

- [1] P. M. Sirrapa (2011). Study on padi culture technology improvement through the use of superior seeds and *Legowo*-planting system to increase padi productivity for food self-sufficiency. *J. Budidaya Pertanian*, 7 (2): 79-86. [in Indonesian]
- [2] Z. F. Rozi, Y. Febrianti, Y. Telaumbanua (2018). Potential of *Diascurea hispida* L juice as bio-insecticide of rice ear bug *Leptocorisa oratorius* on padi plants *Oryza sativa* L. *J. Biogenesis*. Vol. 6, No. 1: 18-22. [in Indonesian]
- [3] A. F. Sitompul, O. Syahrial, Y. Pangestiniingsih (2014). The effectivity test of plant insecticide on the mortality of *Leptocorisa acuta* T (Hemiptera; Alydidae P) on padi plants (*Oryza sativa* L.) in the greenhouse. Program Studi Agroekoteknologi, Fakultas Pertanian. Universitas Sumatera Utara. Medan. *J. Agroekoteknologi* ISSN No. 2337-6597. Vol. 2 (3): 1075-1080. [in Indonesian]
- [4] Rianiyati (2013) The influence of plant pesticide combination application on the production of padi plant (*Oryza sativa* L.) Varietas Inpara-3 of The System of Rice Intensification (SRI). *Jurnal Bioplantae*. Vol. 2 No. 4.: 169-174. ISSN: 2302-6472. [in Indonesian]
- [5] S. B. Moraes, W. B. Moraes, G. Camara, K. S. Maciel, P. A. Lima, A. Ferreira, R. S. Alexandre and J. C. Lopes (2018). Cinnamon and citronella essential oil in the in vitro control of the fungi *Aspergillus* sp. and *Sclerotinia sclerotiorum*. *African Journal of Agricultural Research*. 13 (35): 1811-1815. August, 2018.
- [6] R. Samarasekera, K. S. Kalhari, and L. S. Weerasinghe (2006). Insecticidal activity of essential oils of *Ceylon Cinomomum* and *Cymbopogon* species against *Musca domestica*. *Journal of Essential Oil Research*. 18 (3): 352-354.
- [7] H. Idris and Nurmansyah (2017). Plant pesticide of cinnamon and lemongrass for pest *Nilam Pachyzancla stultalis* control. *Buletin Penelitian Tanaman Rempah dan Obat*. Vol. 28 No. 2: 163-170. [in Indonesian]
- [8] M. Isnaini, E. R. Pane, S. Wiridianti (2015). Testing of several plant insecticides on the rice bug (*Sitophilus oryzae* L). *Jurnal Biota*. Vol. 1: 1-8. [in Indonesian]

- [9] M. A. Ootani, D. R. Brito, G. P. S. Maciel, L. A. Lopes, and R. W. S. Aguiar (2016). Effect of essential oil and Citronellal compound on bean seeds stored microflora. *Revista Verde de Agroecologia e Desenvolvimento Sustentavel*. 11 (1): 49-56.
- [10] R. W. Geetha and A. Roy (2014). Essential oil repellents. A short review. *International Journal Drug Development and Research* 6 (2), 20-27.
- [11] C. S. Rante, D. T. Sembel, E. R. M. Meray, M. M. Ratulangi, M. F. Dien, and D. S. Kandowanko (2013). The use of plant insecticide to control pests on tomato plant. *J. Eugenia* 19 (2): 97-102. [in Indonesian]
- [12] Supriadi (2013). Optimization of various pesticides utilization to control plant pests and diseases. *J. Penelitian dan Pengembangan Pertanian* 32 (1): 1-9. [in Indonesian]
- [13] Ch. Salaki and J. Pelealu (2012). The use of *Barringtonia asiatica* and *Annona muricata* on disease vector insects on chili plant. *J. Eugenia Vol* 18 (1): 22-28. [in Indonesian]
- [14] L. Tigaw and Ch. Salaki (2015). Effectivity test of garlic and tobacco extract as bio-insecticide on green peach aphid *Myzus persicae* on ada tanaman cabai. *J. Eugenia Vol*. 21 (3): 135-141 [in Indonesian]
- [15] Nurmansyah (2011). The effectivity of lemongrass on sucking pest of cocco plant *Helopeltis antonii*. *Bul. Littro*. 22 (2): 205-213. [in Indonesian]
- [16] M. Darwis and W. R. Atmadja (2010). [The use of 10 medicinal and aromatic plants for pest *Helopeltis theivora* Watch prevention.] In; *Prosiding Seminar Nasional VI. Peranan Entomology dalam Mendukung Pengembangan Pertanian Ramah Lingkungan dan Kesehatan Masyarakat*. Bogor, Perhimpunan Entomologi Indonesia. p. 328-336. [in Indonesian]
- [17] Z. Jiang, H. Jiang and P. Xie (2013). Antifungal activities against *Sclerotinia sclerotiorum* by *Cinnamomum cassia* oil and its main components. *J. essential oil Research* 25 (6): 444-451.
- [18] Rohimatum and I. W. Laba (2013). The effectivity of lemongrass oil and clove oil insecticides on pepper pests. *Bul. Littro*. 24 (1): 26-34.
- [19] F. M. Howlett (1912). The effect of oil of citronella on two species of *Dacus*. *Transactions of the Royal Entomological Society of London*. 60 (2): 412-418.
- [20] C. Kongkaew, I. Sakunrag, N. Chaiyakunapruk and A. Tawatsin (2011). Effectiveness of citronella preparations in preventing mosquito bites; Systematic Review of Controlled Laboratory Experimental Studies. *Tropical Medicine & International Health*. 16 (7): 802-810.
- [21] C. Kongkaew, L. Sakunrag, N. Chaiyakunapruk, and A. Tawatsin (2011). Effectiveness of citronellain preventing mosquito bites. *Tropical Medicine and International Health* 16 (7): 802-810.
- [22] L. R. Lindsay, G. A. Surgeoner, J. D. Heal and G. J. Gallivan (1996). Evaluation on the efficacy of 3% citronella candles and 5% Citronella incense for protection against field populations of *Aedes* mosquitoes. *J. the American Mosquito Control Association*. 12 (2 Pt 1): 293-294.
- [23] W. R. Atmadja (2010). The use of shrub, clove, lemongrass-based insecticides for worm *Spodoptera litura* prevention. *Prosiding Seminar Nasional VI Peranan Entomologi dalam Mendukung Pengembangan Pertanian Ramah Lingkungan dan Kesehatan Masyarakat*. Indonesian Entomology Association, Bogor Branch. p. 191-200. [in Indonesian]
- [24] H. Idris (2014). Effect of cinnamon-based bio-insecticide on the biological aspects of *Aspidomorpha milliaris* F. *Buletin Ilmiah Eka Sakti* 26 (1): 68-77. [in Indonesian]
- [25] Nurmansyah (2006). The effect of plant and solvents administration on the capability of cinnamon waste oil-based pesticide antifungal. *Jurnal Dinamika Pertanian*. 21 (2), 115-120 [in Indonesian]
- [26] M. S. Khan and I. Ahmad (2011). In vitro antifungal, anti-elastase and anti-keratinase activity of essential oil of *Cinnamomum*, *Syzygium* and *Cymbopogon*-species against *Aspergillus fumigatus* and *Trichophyton rubrum*, *Phytomedicine* 19 (1): 48-55.
- [27] W. S. Abbott (1925). A Method of Computing the Effectiveness of an Insecticide. *J. Am. Mosq. Control. Assoc.* 1987. 3; 302-303.
- [28] Nakahara, Kazuhiko, N. S. Alzoreky, T. Yoshihashi, H. T. Nguyen and G. Trakoon-tivakorn (2013). Chemical composition and antifungal activity of essential oil from *Cymbopogon nardus* (Citronella Grass). *Japan Agricultural Research Quarterly*. 37 (4): 249-252.
- [29] R. O. B. Wijsekera (1973). The chemical composition and analysis of Citronella oil. *J. the Natural Science Council of Sri Lanka*. 1: 67-81.
- [30] C. L. Abramson, P. A. Wanderley, M. J. A. Wanderley, A. J. S. Mina, O. B. D. Souza, and C. Edgar (2006). Effective essential oil from citronella and alfazema on Fennel Aphids *Hyadaphis foeniculli passerine* (Hemiptera: Aphididae) and its predator *Cycloneda sanguinea* (coleopteran; Coccinelidae). *American Journal of Enviromental Science* 3 (1): 9-10.
- [31] A. Ahmad, M. Kaleem, Z. Ahmed, and H. Shafiq (2015). Therapeutic potential of flavonoids and their mechanism of action against microbial and viral infection –A review. *Food Research International* 77 (2): 221-235.
- [32] M. A. Ootani, R. W. Aguiar, A. C. Ramos, D. R. Brito, J. B. D. Silva, and J. P. Cajazeira (2013). Use of essential oil in agriculture. *J. Bioteknologi and Biodiversity* 4 (2): 162-175.
- [33] D. Ganjewala (2009). Cymbopogon essential oils; chemical compositions and bioactivities. *International journal of essential oil therapeutics*. 3 (2-3): 56-65.
- [34] H. Idries (2014). The effect of cinnamon bio-insecticide on the biological aspects of *Aspidomorpha milliaris* F. *Buletin Ilmiah Eka Sakti*. 26 (1): 68-77. [in Indonesian]
- [35] Clemente, M. Aparecido, C. M. Monteiro, M. G. coralik, F. T. Gomes, M. C. Prata, and E. Daemon (2010). Acaricidal activity of the essential oils from *Eucalyptus citriodora* and *Cymbopogon nardus* on larvae of *Amblyomma cajennense* (Acari; Ixodidae) and *Anocentor nitens* (Acari; Ixodidae). *Parasitology Research* 107 (4): 987-992.
- [36] H. Idris (2014). Plant insecticide formula for pest *Pachyzancla stultalis* control in patchouli plant. *Bul. Littro*. 25 (1): 69-76. [in Indonesian]