Prognostic Ability of Simplified Pulmonary Embolism Severity Index (PESI) Score in Real World: A Brief Report

Luca Masotti^{1, 7, 25, *}, Grazia Panigada², Giancarlo Landini¹, Filippo Pieralli³, Francesco Corradi³, Salvatore Lenti⁴, Rino Migliacci⁵, Carlo Nozzoli³, Maddalena Grazzini³, Lucia Ciucciarelli³, Alessandro Morettini³, Sara Bucherelli³, Alessandra Petrioli³, Carlotta Casati³, Mario Felici⁴, Luciano Ralli⁴, Stefano Arrigucci⁴, Laila Teghini², Giovanni Antonio Porciello⁵, Stefano Spolveri⁶, Daniele Baldoni⁶, Anna Frullini⁶, Barbara Cimolato¹, Gianni Lorenzini⁷, Alessandro Pampana⁷, Guidantonio Rinaldi⁸, Maria Chiara Bertieri⁸, Raffaele Laureano⁹, Stefano Tatini⁹, Alberto Fortini¹⁰, Chiara Angotti¹⁰, Valerio Verdiani¹¹, Anna Maria Romagnoli¹¹, Irene Cascinelli¹¹, Alberto Camaiti¹², Nicola Mumoli¹², Marco Cei¹², Stefano Giuntoli¹², Massimo Alessandri¹³, Alessandro De Palma¹³, Maurizio Manini¹⁴, Veronica De Crescenzo¹⁴, Michele Piacentini¹⁵, Carlo Passaglia¹⁶, Giancarlo Tintori¹⁶, Carlo Palermo¹⁷, Alba Dainelli¹⁷, Roberto Andreini¹⁸, Giuseppa Levantino¹⁸, Plinio Fabiani¹⁹, Lucia Raimondi¹⁹, Massimo Di Natale²⁰, Filippo Risaliti²⁰, Rossella Nassi²¹, Roberta Mastriforti²¹, Roberto Cappelli²², Michele Voglino²², Paola Lambelet²³, Stefano Fascetti²³, Adriano Cioppi²³, Valentina Carli²³, Alessandro Tafi²⁴, Simone Meini²⁴, Emilio Santoro²⁶, Claudia Rosi²⁶

¹Internal Medicine, Santa Maria Nuova Hospital, Florence, Italy

²Internal Medicine, Pescia Hospital, Pistoia, Italy

³Internal Medicine, Careggi Hospital, Florence, Italy

⁴Internal Medicine, Arezzo Hospital, Arezzo, Italy

⁵Internal Medicine, Cortona Hospital, Arezzo, Italy

⁶Internal Medicine, Borgo San Lorenzo Hospital, Florence, Italy

⁷Internal Medicine, Cecina Hospital, Livorno, Italy

⁸Internal Medicine, Barga Hospital, Lucca, Italy

⁹Internal Medicine, Santa Maria Annunziata Hospital, Florence, Italy

¹⁰Internal Medicine, San Giovanni di Dio Hospital, Florence, Italy

¹¹Internal Medicine, Grosseto Hospital, Grosseto, Italy

¹²Internal Medicine, Livorno Hospital, Livorno, Italy

¹³Internal Medicine, Massa Marittima Hospital, Grosseto, Italy

¹⁴Internal Medicine, Orbetello Hospital, Grosseto, Italy

¹⁵Internal Medicine, Piombino Hospital, Livorno, Italy

¹⁶Internal Medicine, Cisanello Hospital, Pisa, Italy

¹⁷Internal Medicine, Campostaggia Hospital, Siena, Italy

¹⁸Internal Medicine, Pontedera Hospital, Florence, Italy

¹⁹Internal Medicine, Portoferraio Hospital, Livorno, Italy

²⁰Internal Medicine, Prato Hospital, Prato, Italy

²¹Internal Medicine, San Sepolcro Hospital, Arezzo, Italy

²²Internal Medicine, Le Scotte Hospital, Siena, Italy

²³Internal Medicine, Viareggio Hospital, Viareggio, Italy

²⁴Internal Medicine, Volterra Hospital, Volterra, Italy

²⁵Internal Medicine, Santa Maria Nuova Hospital, Florence, Italy

²⁶Internal Medicine, Bibbiena Hospital, Bibbiena, Italy

Email address:

luca.masotti@tin.it (L. Masotti), g.panigada@usl3.toscana.it (G. Panigada), filpiero@tin.it (F. Pieralli), lentisa@libero.it (S. Lenti)

To cite this article:

Luca Masotti, Grazia Panigada, Giancarlo Landini, Filippo Pieralli, Francesco Corradi, Salvatore Lenti, Rino Migliacci, Carlo Nozzoli, Maddalena Grazzini, Lucia Ciucciarelli, Alessandro Morettini, Sara Bucherelli, Alessandra Petrioli, Carlotta Casati, Mario Felici, Luciano

Ralli, Stefano Arrigucci, Laila Teghini, Giovanni Antonio Porciello, Stefano Spolveri, Daniele Baldoni, Anna Frullini, Barbara Cimolato, Gianni Lorenzini, Alessandro Pampana, Guidantonio Rinaldi, Maria Chiara Bertieri, Raffaele Laureano, Stefano Tatini, Alberto Fortini, Chiara Angotti, Valerio Verdiani, Anna Maria Romagnoli, Irene Cascinelli, Alberto Camaiti, Nicola Mumoli, Marco Cej, Stefano Giuntoli, Massimo Alessandri, Alessandro De Palma, Maurizio Manini, Veronica De Crescenzo, Michele Piacentini, Carlo Passaglia, Giancarlo Tintori, Carlo Palermo, Alba Dainelli, Roberto Andreini, Giuseppa Levantino, Plinio Fabiani, Lucia Raimondi, Massimo Di Natale, Filippo Risaliti, Rossella Nassi, Roberta Mastriforti, Roberto Cappelli, Michele Voglino, Paola Lambelet, Stefano Fascetti, Adriano Cioppi, Valentina Carli, Alessandro Tafi, Simone Meini Emilio Santoro, Claudia Rosi. Prognostic Ability of Simplified Pulmonary Embolism Severity Index (PESI) Score in Real World: A Brief Report. *International Journal of Clinical and Experimental Medical Sciences*. Vol. 1, No. 2, 2015, pp. 22-26. doi: 10.11648/j.jcems.20150102.14

Abstract: Background and aim: Prognostic stratification of acute pulmonary embolism (PE) represents the cornerstone of modern management of this potentially life-threatening disease. In the latest years, a lot of clinical prognostic models have been validated. However, these are yet underused in clinical practice, especially in real world populations. The aim of our study was to test the prognostic ability of the Simplified Pulmonary Embolism Severity Index (sPESI) score in a real world population. Methods: Data records of 452 patients that were discharged for acute PE from 28 Internal Medicine wards of Tuscany (Italy) were retrospectively analysed. sPESI was calculated in the identical manner as the original study. Prognostic ability of sPESI score for predicting in-hospital all-cause and PE-related mortality was tested by using Areas under Receiver Operating Characteristics (ROC) curve (AUC). Results: 15.2% of patients were classified as sPESI score 0, whereas 84.8% were classified as sPESI ≥ 1 . All causes of in-hospital mortality were 10.95% (5.75% PE-related) in patients with sPESI score ≥ 1 and 0% (0% PE-related) in sPESI score 0. AUC for all causes of mortality was 0.694 (95% CI: 0.650-0.736), whereas it was 0.702 (95% CI: 0.657-0.743) for PE-related mortality. Conclusion: In a real world population, sPESI is a good prognosticator for all causes of in-hospital and PE-related mortality and its use should be encouraged.

Keywords: Pulmonary Embolism, Prognosis, PESI Score

1. Introduction

Acute pulmonary embolism (PE) remains one of the leading causes of mortality and morbidity in cardiovascular setting, especially when is associated with hemodynamic instability (1).

Prognostic stratification of acute PE is of utmost importance for the choice of appropriate treatment and setting of care. It can be assured by using clinical, instrumental and laboratory assessment (2, 3).

In the latest years, many prognostic models for mortality risk stratification in acute PE have been proposed. Of them, simplified PESI score (sPESI) showed to be a good prognosticator for 30-day mortality (4). The sPESI score considers the presence or absence of age over 80 years, history of cancer, heart failure or chronic obstructive pulmonary disease (COPD), heart rate (HR) ≥ 110 beats for minute, systolic blood pressure (SBP) ≤ 100 mmHg, oxygen arterial saturation $\leq 90\%$.

The contemporary absence of all variables (sPESI score 0) identifies low risk patients (30-day mortality $\leq 1\%$) whereas the presence of at least one of variables (sPESI score ≥ 1) identifies patients with high risk (30-day mortality around 11%) (4). Despite, sPESI score has been used as prognostic model for identifying low risk patients candidate for home treatment in clinical trials (5-7) and much recently European Society of Cardiology (ESC) has proposed a prognostic model that is based on sPESI score as first prognostic tool for subdividing low risk patients from high risk patients needing for further prognostic assessment (8), to now sPESI score is

still underused in clinical practice, especially in real world population.

Therefore, the aim of the present study was to test the prognostic ability of sPESI score as prognosticator of all causes of in-hospital and PE-related mortality.

2. Materials and Methods

We performed a multicenter, observational, retrospective, cohort study aimed to evaluate characteristics and clinical management of acute PE none selected patients that were admitted in internal medicine wards of Tuscany (Italy). Demographic, clinical and prognostic data of at least ten patients consecutively were discharged for acute PE between 2012 and 2013 years from each center were retrospectively provided. sPESI was calculated in the identical manner of the study validation (4).

For testing the prognostic ability of sPESI, we calculated the Areas under Receiver Operating Characteristic (ROC) curves (AUCs) for each study endpoint.

3. Results

Table 1 shows the general characteristics of patients.

Table 1. General characteristics of patients.

Number	452
Males/Females	39.8%/60.2%
Mean age (years) $\pm$ SD	76.01 $\pm$ 12.34
Age ≥ 80 years	45.3%

Number	452
Cancer	30.2%
Cardiopulmonary diseases	22.4%
Heart rate ≥ 110 beats for minute	19.8%
Systolic blood pressure ≤ 100 mmHg	7.3%
Oxygen arterial saturation $\leq 90\%$	27.4%
All cause in-hospital mortality	9,5%
PE related in-hospital mortality	5.7%
sPESI score distribution	
0Low risk	15.2%
≥ 1 High risk	84.8%
1	39.2%
2	27.7%
3	12.6%
≥ 4	5.3%

Legend= PE: Pulmonary embolism, sPESI: Pulmonary embolism severity

index score

452 patients were enclosed in the study. 15.2% of patients were classified as sPESI score 0, whereas 84.8% were classified as sPESI ≥ 1 , 66.9% of patients being classified as sPESI 1 or 2.

All causes of in-hospital mortality was 10.95% in patients with sPESI score ≥ 1 and 0% in sPESI score 0 ($p < 0.001$). PE-related mortality was 5.75% in patients with sPESI score ≥ 1 and 0% in sPESI score 0 ($p < 0.001$). All causes of in-hospital mortality increased from 0% in patients with sPESI score 0 to 29.1% in patients with sPESI score ≥ 4 , whereas PE-related in-hospital mortality increased from 0% in patients with sPESI score 0 to 20.8% in patients with sPESI score ≥ 4 (Figure 1).

Figure 1. sPESI score and in-hospital mortality.

AUC for all causes of mortality was 0.694 (95% Confidence Interval (CI): 0.650-0.736), whereas it was 0.702 (95% Confidence Interval (CI): 0.657-0.743) for PE-related mortality. Table 2 summarizes the statistical analysis.

Table 2. Summary of statistical analysis.

Area under the ROC curve (AUC) for all cause in-hospital mortality	0,694
Standard Error	0,0384
95% Confidence interval	0,650 to 0,736
z statistic	5,055
Significance level P (Area=0.5)	<0,0001
Youden index J	0,2674
Associated criterion	≤ 1
Sensitivity	56,97
Specificity	69,77

Criterion values and coordinates of the ROC curve [Show]

Area under the ROC curve (AUC) for PE-related in-hospital mortality	0,702
Standard Error	0,0507
95% Confidence interval	0,657 to 0,743
z statistic	3,975
Significance level P (Area=0.5)	0,0001
Youden index J	0,2996
Associated criterion	≤ 2
Sensitivity	83,80
Specificity	46,15

ROC: Receiver Operating Characteristic

Figure 2 shows the ROC curves.

(1)

(2)

Figure 2. ROC curves for sPESI as prognosticator of in-hospital all-cause mortality (1) and PE-related mortality (2).

4. Discussion

Prognostic stratification of acute pulmonary embolism (PE) represents the cornerstone of modern management of this disease (9). sPESI score is now considered as a safe prognostic model that is able to classify patients in 30-day low mortality risk (sPESI 0) and high mortality risk (sPESI ≥ 1). A meta-analysis showed that the odds ratio for all causes of mortality in low risk sPESI versus high risk (3 events on 770 patients with low risk sPESI versus 72 events on 1404 patients with high risk sPESI) was 0.09 (95% CI: 0.03-0.26), whereas it was 0.08 (95% CI: 0.04-0.16) for PE-related mortality (8 events on 1003 patients with low-risk sPESI versus 205 events on 1981 patients with high risk sPESI) (10).

Prognostic stratification by using sPESI score is of utmost importance for making decision on appropriate treatment and setting of care. New ESC recommendations on PE management suggest that patients with low risk sPESI score should not receive further investigations such as right heart dysfunction and myocardial damage assessment and could be early discharged from hospital and/or treated at home after diagnostic and prognostic assessment are performed in emergency department (8). Much recently, a meta-analysis of

studies on home treatment of low risk PE, a lot of them classifying low risk patients by using sPESI score, has demonstrated that 14-day and 90-day all causes of mortality were 0.41% and 1.58, respectively (11).

From a true real world perspective, our study confirms that mortality may be extremely low if not completely absent in low risk sPESI patients.

5. Conclusion

Prognostic stratification is a key point in the modern management of acute PE. Despite it was limited by the retrospective methodology, our study was performed in more than 80% of internal medicine wards of one of the most populated regions of Italy, demonstrates that sPESI score, a simple and easily performable score, seems to be a good prognosticator for stratifying the in-hospital prognosis. Therefore, its wide use should be strongly encouraged.

Appendix

The Simplified Pulmonary Embolism Severity Index (sPESI)

Predictors	Scoring
• Age over 80 years	1
• History of cancer	1
• History of cardiopulmonary diseases	1
• Heart rate ≥ 110 bpm	1
• Systolic blood pressure < 100 mmHg	1
• Arterial oxygen saturation $< 90\%$	1
Total _____	

0 Low risk
30-day mortality 1%

≥ 1 High risk
30-day mortality 10.9%

References

- [1] Wood KE. Major pulmonary embolism. Review of a pathophysiologic approach to the golden hour of hemodynamically significant pulmonary embolism. *Chest* 2002; 121: 877-905.
- [2] Masotti L, Righini M, Vuilleumier N, Antonelli F, Landini G, Cappelli R, Ray P. Prognostic stratification of acute pulmonary embolism: focus on clinical aspects, imaging, and biomarkers. *Vasc Health Risk Manag* 2009; 5: 567-75.
- [3] Becattini C, Agnelli G. Acute pulmonary embolism: risk stratification in the emergency department. *Intern Emerg Med* 2007; 2: 119-29.
- [4] Jiménez D, Aujesky D, Moores L et al.; RIETE Investigators. Simplification of the pulmonary embolism severity index for prognostication in patients with acute symptomatic pulmonary embolism. *Arch Intern Med* 2010; 170: 1383-9.
- [5] Lankeit M, Konstantinides S. Is it time for home treatment of pulmonary embolism? *Eur Respir J* 2012; 40(3): 742-9.
- [6] Squizzato A, Galli M, Dentali F, Ageno W. Outpatient treatment and early discharge of symptomatic pulmonary embolism: a systematic review. *Eur Respir J* 2009; 33: 1148-55.

- [7] Aujesky D, Roy PM, Verschuren F, Righini M, Osterwalder J, Egloff M, Renaud B, Verhamme P, Stone RA, Legall C, Sanchez O, Pugh NA, N'gako A, Cornuz J, Hugli O, Beer HJ, Perrier A, Fine MJ, Yealy DM. Outpatient versus inpatient treatment for patients with acute pulmonary embolism: an international, open-label, randomized, non-inferiority trial. *Lancet* 2011; 2; 378(9785): 41-8.
- [8] Authors/Task Force Members, Konstantinides S, Torbicki A, Agnelli G *etv al.*; Authors/Task Force Members. 2014 ESC Guidelines on the diagnosis and management of acute pulmonary embolism: The Task Force for the Diagnosis and Management of Acute Pulmonary Embolism of the European Society of Cardiology (ESC) Endorsed by the European Respiratory Society (ERS). *Eur Heart J* 2014; 35: 3033-69.
- [9] Masotti L, Mannucci A, Antonelli F, Maurini V, Testa R, Marchetti S, Landini G, Cappelli R. The Risk-based Treatment of Acute Pulmonary Embolism. *J Clin Med Res* 2009; 1: 1-7.
- [10] Zhou XY, Ben SQ, Chen HL, Ni SS. The prognostic value of pulmonary embolism severity index in acute pulmonary embolism: a meta-analysis. *Respir Res* 2012; 13: 111
- [11] Piran S, Le Gal G, Wells PS, Gandara E, Righini M, Rodger MA, Carrier M. Outpatient treatment of symptomatic pulmonary embolism: a systematic review and meta-analysis. *Thromb Res* 2013; 132: 515-9.