Review Article

Interleukin 4 (IL-4) in Urinary Schistosomiasis as a Diagnostic Biomarker During Its Reinfection- A Review

Aigbogun Edwin Ighodaro¹, Mohammed Sani Sambo Datsugwai¹, Umar Nabeela Bello¹, Afangide Christopher Stephen²

¹Department of Microbiology, Faculty of Science, Kaduna State University, Kaduna, Nigeria

²Zamani College, Kaduna, Nigeria

Email address:

aigbogunighodaro@gmail.com (A. E. Ighodaro)

To cite this article:

Aigbogun Edwin Ighodaro, Mohammed Sani Sambo Datsugwai, Umar Nabeela Bello, Afangide Christopher Stephen. Interleukin 4 (IL-4) in Urinary Schistosomiasis as a Diagnostic Biomarker During Its Reinfection- A Review. *International Journal of Clinical and Experimental Medical Sciences*. Vol. 3, No. 5, 2017, pp. 57-60. doi: 10.11648/j.ijcems.20170305.11

Received: August 16, 2017; **Accepted:** September 5, 2017; **Published:** October 10, 2017

Abstract: Urinary schistosomiasis is caused by *Schistosoma haematobium* worm and it is most common source of haematuria (blood in urine) worldwide. This disease has long been recognized as serious public health to man especially in countries like Nigeria, Egypt, with currently more than 790 million people at risk. Since IL-4 stimulates IgE production, the hypothesis that Helper T cell 2 (Th2) associated with cell immunity participate in protection to reinfection in an anti-inflammatory dichotomy manner. IL-4 confers proliferation of T cells, induction of class switching immunoglobulins and also serve as a biomarker in co association with IL-5 for (Th2). The aim of this review paper is to simplify the immunological mechanism that happens when Schistosome worm elicit strong humoral and cellular response in definite host.

Keywords: Helper T Cell, Schistosomiasis, Interleukin 4, Immunoglobulin

1. Introduction

Schistosomiasis or bilharzia is a disease condition caused by a trematode schistosome, which was discovered by Theodor Bilharz in 1852. Species of the worm that prevalently infect man are *Schistosoma haematobium*, *S. mansoni* and *S. japonicum*. This condition is characterized by inflammation mainly in the bladder, intestine and liver [1, 2]. Schistosomiasis can be acute which is characterized by feverish syndrome (Katayama fever), systemic hypersensitivity reaction within few weeks to months after primary infection [3] and chronic; abdominal pain, inflamed liver, blood in the stool or blood in the urine, and problems passing urine [4].

Epidemiologically, about 200-300 million people are infected worldwide and more than 790 million people are at risk [4, 5]. Immunopathologically, schistosome eggs trapped in the tissue for years leading to obstruction of the urinary bladder, urogenital inflammation and scarring by *S. haematobium* or intestine or liver leading to inflammation and

fibrosis by *S. mansoni* and *S. japonicum* respectively [3] and the chronic stage of this infection is characterized by the release of IL-4 [4].

IL-4 was discovered by Maureen Howard and William Paul and Ellen Vitella and her research group in 1982. They are cytokines produced by leukocyte in the blood. IL-4 has compact, globular fold, stabilized by three disulphide bonds.

Figure 1. Interleukin 4 cytokine protein [6].

diagnostic biomarker [10], elicit differentiation and proliferation of Helper T cell, cell metastasis, induces differentiation of naive helper T cells Th0 cells to Th2 cells. Upon activation by IL-4, Th2 cells subsequently produce additional IL-4 in a positive feedback loop [11] and has recently be found to initiate a regeneration cascade through phosphorylation of its intracellular effector STAT6 in an experimental Alzheimer's disease model in adult zebra fish brain [12].

The latter delivers cytokine that further activates the APC [13]. The IL-4 released by the Th0 in higher concentration further orchestrate the proliferation and differentiation of Th 2 cells and inhibit Th1 cells [14].

KEY:
Tho- Naïve T cell
Th- Helper T cell
Th1-Helper T cell 1
Th2-Helper T cell 2
DC – Dendritic cell
APC –Antigen presenting cell

2.1. Stimulation of Activated B Cells

IL-4 induces the stimulation of activated B cells against the worm antigen. The differentiated Th2 cells further releases IL-4 towards B cells which causes cytochange of B cells, thereby differentiating it to plasma cells which further produces immunoglobulin (Ig) and memory B cell against

further reinfection of the exogenous antigen of the worm [13]. One condition for this to happen is that, MHC of Th2 must recognize B cell receptor, adhere, synthesize IL-4 and CD4 ligand which orchestrates cytoskeletal changes in activated B cells [15].

Figure 4. Diagram indicating adherence of B cell to Th2. [16].

The production of antibodies against the antigen of the worm lead to protective outcome such as precipitation, agglutination and complement system activation [17].

It is important to understand that infected individuals of all ages have varying immunoglobulin class and subclass. IgE for example, is associated with resistance to reinfection after treatment with praziquantel, whereas, IgM, IgG 4 and IgG2 are associated with susceptibility to reinfection after treatment while IgA has been implicated with eosinophil mediated killing of the parasite, this is to confirm that IL-4 induce class switching that makes other immunoglobulin to have some specificity but different biological characteristics [18]. In 2007, [19] showed relationship between age and intensity of infection as connected to immunoglobulin profile in cohort group. The result showed that IgM and IgE are extremely high in serum of infected population compared to uninfected population. It synoptically means that without the help of or release of IL-4, there will be no protection to fighting parasitic reinfection, and 2016 result of [20] arrived at the same conclusion with Nmorsi. Hence, measurement of immunoglobulin concentration can be used as indicators of heavy and acute infection of *S. haematobium* in our endemic localities. Reports exists where IgE has been indicated as diagnostic biomarker of infections as it correlates positively with the intensity of infection. Serologically, these levels of Ig can be used as markers of chronic schistosomiasis in clinical and laboratory practice, especially where clinical presentations of the infection are covert and sub clinical [21].

2.2. Allergies or Hypersensitivity

Allergy or hypersensitivity is allergen which is a substance

that causes allergic reaction in susceptible individuals [22].

Researches had been done and facts have been concluded that IgE have protective outcome against parasitic worm and also cause immediate (type 1) hypersensitivity characterized by a reaction in a sensitized individual within minutes of exposure to antigen [23].

Upon exposure, the allergen readily combines with the cell fixed IgE. At least two cell bound IgE molecules for reaction to occur. Within milliseconds, the IgE-antigen attachment and cross linkage of IgE in the cell membrane cause the mast cell or eosinophil to release histamine, leukotrienes, prostaglandins and cytokines in the process called degradation. These mediators are direct causes of hives, hay, fever, asthma and anaphylactic shock [24].

2.3. Diagnostic Biomarker

Parasitological techniques (stool examination and/or urine filtration/microscopy) are still the diagnosis of choice in national schistosomiasis control programmes around the globe. However it comes with its cost of labour and time wallowing procedures. There is an undying need for a more sensitive approach especially in endemic area and persons with prolong cases of Schistosomiasis [10]. The presence of IL-4 can be measured by the amount or concentration of CD4 using flow cytometry or ELISPOT kit [25].

3. Conclusion

Minute amount of IL-4 against urinary schistosomiasis does not produce significant reactions in the living system except in increased concentration or co association with IL-5. However,

it is a key player in in dealing with parasitic infection caused by *S. haematobium* because of its multi facet dynamics in function as part of the immune system to actualizing its primary aim to protect infected individuals against reinfection and regulate proliferation, differentiation, and apoptosis in multiple cell types of haematopoietic and non-haematopoietic source.

References

- [1] Nmorsi, O. P. G. (2009). Principles of Parasitology. PON Publication, 100-107.
- [2] Abdulkadir, A., Ahmed, M., Abubakar, B. M., Suleiman, I. E., Yusuf, I., Imam, I. M., and Musa, B. M. (2017). Prevalence of Urinary schistosomiasis in Nigeria Systematic Review and Meta-analysis. *African Journal of Urology*, 1994–2015.
- [3] Gryseels, B., Polman, K., Clerinx, J. and Kestens, L. (2006). Human schistosomiasis. *The Lancet*, 368(9541), 1106-1118.
- [4] Colley, D. G., Bustinduy, A. L., Secor, W. E. and King, C. H. (2014). Human schistosomiasis. *The Lancet*, 383(9936), 2253-2264.
- [5] Steinmann, P., Keiser, J., Bos, R., Tanner, M. and Utzinger, J. (2006). Schistosomiasis and Water Resources Development: Systematic Review, Meta-analysis, and Estimates of People at Risk. *The Lancet infectious diseases*, 6(7), 411-425.
- [6] Mueller, T. D., Zhang, J. L., Sebal, W., and Duschl, A. (2002). Structure, Binding, and Antagonists in the IL-4/IL-13 Receptor System. *Biochimica et Biophysica Acta (BBA)Molecular Cell Research*, 1592(3), 237-250.
- [7] Paul, W. E. (2015). History of interleukin-4. *Cytokine*, 75(1), 3-7.
- [8] Guo, L., Urban, J. F., Zhu, J. and Paul, W. E. (2008). Elevating Calcium in Th2 Cells Activates Multiple Pathways to Induce IL-4 Transcription and mRNA Stabilization. *Journal of Immunology*.181:3984–3993.
- [9] Price, A. E., Liang, H. E., Sullivan, B. M., Reinhardt, R. L., Eisley, C. J., Erle, D. J., and Locksley, R. M. (2010). Systemically Dispersed Innate IL-13-expressing Cells in Type 2 Immunity. *Proceedings of the National Academy of Sciences*,107:11489–11494.
- [10] Zhu, Y. C. (2005). Immunodiagnosis and its Role in Schistosomiasis Control in China: a Review. *Actatropica*, 96(2), 130-136.
- [11] Hosoyama, T., Aslam, M. I., Abraham, J., Prajapati, S. I., Nishijo, K., Michalek, J. E. and Keller, C. (2011). IL-4R drives dedifferentiation, mitogenesis, and metastasis in rhabdomyosarcoma. *Clinical Cancer Research*, 17(9), 2757-2766.
- [12] Bhattarai, P., Thomas, A. K., Cosacak, M. I., Papadimitriou, C., Mashkaryan, V., Froc, C. and Kizil, C. (2016). IL4/STAT6 signaling activates neural stem cell proliferation and neurogenesis upon Amyloid- β 42 aggregation in adult zebrafish brain. *Cell reports*, 17(4), 941-948.
- [13] Nester, A., Roberts, P., Pearsall, N. N. and Anderson, D. G. (2009). *Microbiology: A Human Perspective*, eight edition, Mc Graw Hill Education.
- [14] Arinola, O. G. (2005). Immunological Aspects of Urinary Schistosomiasis in Ibadan, Southwestern Nigeria. *Journal of the Association of Resident Doctors*, 3(1), 5.
- [15] de Moraes, C. G. V., Castro Lima, A. K., Terra, R., dos Santos, R. F., Da-Silva, S. A. G. and Dutra, P. M. L. (2015). The dialogue of the host-parasite relationship: *Leishmania* spp. and *Trypanosoma cruzi* infection. *Bio Medical research international*, 2015.
- [16] Janeway, C. A., Travers, P., Walport, M. and Shlomchik, M. (2001). *Immunology*. New York: Garland Science.
- [17] Colley, D. G., and Secor, W. E. (2014). Immunology of Human schistosomiasis. *Parasite Immunology*, 36(8), 347-357.
- [18] Jung, Y. and Rothenberg, M. E. (2014). Roles and Regulation of Gastrointestinal Eosinophils in Immunity and Disease. *The Journal of Immunology*, 193(3), 999-1005.
- [19] Nmorsi, O. P. G., Ukw, N. C. D., Isaac, C., Egwunyenga, A. O. and Olague, N. H. (2007). Immunoglobulin Profile of some Nigerians with *Schistosoma haematobiu* Infection. *African Journal of Microbiology Research*, 1(7), 113-116.
- [20] Elfaki, T. E. M., Arndts, K., Wiszniewsky, A., Ritter, M., Goreish, I. A., Misk El Yemen, A. and Hoerauf, A. (2016). Multivariable Regression Analysis in *Schistosomamansoni*-Infected Individuals in the Sudan Reveals Unique Immunoepidemiological Profiles in Uninfected, Egg+ and non-egg+ infected Individuals. *PLOS Neglected Tropical Disease*, 10(5).
- [21] Kardoush, M. I. (2016). Identification of Candidate Serum Biomarkers for Schistosomiasis Infection using Mass Spectrometric Approaches (Doctoral dissertation, McGill University) 20-23.
- [22] Murphy, K. and Weaver, C. (2016). *Janeway's immunobiology*. Garland Science. 48-64.
- [23] Wagner, B. (2016). Immunoglobulin E and allergy. *Equine veterinary journal*, 48(1), 13-14.
- [24] Fitzsimmons, C. M., Falcone, F. H., and Dunne, D. W. (2014). Helminth Allergens, Parasite-specific IgE and its Protective Role in Human Immunity. *Frontiers in Immunology*, 47-65.
- [25] Douglas, S. D. (2016). Introduction. In *Manual of Molecular and Clinical Laboratory Immunology*, Eighth Edition. 259-262.