

**Review Article**

A Review of Impacts of Gas Flaring on Vegetation and Water Resources in the Niger Delta Region of Nigeria

Enetimi Idah Seiyaboh, Sylvester Chibueze Izah *

Department of Biological Sciences, Niger Delta University, Wilberforce Island, Nigeria

Email address:

chivestizah@gmail.com (S. C. Izah)

*Corresponding author

To cite this article:Enetimi Idah Seiyaboh, Sylvester Chibueze Izah. A Review of Impacts of Gas Flaring on Vegetation and Water Resources in the Niger Delta Region of Nigeria. *International Journal of Economy, Energy and Environment*. Vol. 2, No. 4, 2017, pp. 48-55.

doi: 10.11648/j.ijeee.20170204.11

Received: May 12, 2017; **Accepted:** June 6, 2017; **Published:** July 14, 2017

Abstract: Nigeria is an oil and gas producing nation. The Niger Delta is the Nigerian oil and gas province. About 10 – 40% of produced associated gas is underutilized and is flared into the environment. This is the usual trend in the Nigerian oil and gas industry. Gas flaring has adverse impacts on the environmental components and its associated biota. This paper reviews the impacts of gas flaring on the vegetation and water quality resources in the Niger Delta region of Nigeria. The study found that gas flaring alters water ions (especially sulphate, carbonate, nitrate), pH, conductivity heavy metals (such as lead and iron) concentration especially in rainwater. It also affects vegetation leading to decrease in growth and productivity probably due to changes in soil quality parameters. The paper concludes by suggesting promulgation/implementation of gas flaring laws in Nigeria.

Keywords: Gas Flaring, Impacts, Vegetation Structure, Water Quality

1. Introduction

Following the discovery of crude oil in commercial quantity in Nigeria, the mainstay of the nation economy shifted from agriculture to crude oil and natural gas. Till date, crude oil and natural gas accounts for significant source of revenue and foreign earning to the economy [1]. According to Ohimain [2, 3], Izah and Ohimain [4], about 85% and 90% of Nigerian earning and export respectively are provided by petroleum. Furthermore over 80% of money used in financing national budget is from oil rich region of Nigeria [5]. Nigeria also have other resources including mineral resources such as natural gas, tin, iron ore, coal, lead, zinc limestone, niobium and arable land for agricultural purposes [6]. The country has several other renewable energy resources such as biomass, solar, wind, hydropower etc [4].

Crude oil and natural gas account for about 50% of global energy resources [7]. On global perspective, Nigeria is ranked 7th and 12th largest exporters and producers crude oil respectively [8]. It has been reported that Nigeria crude oil and natural gas resources include 35 – 36.22 billion barrel of crude

oil, 187 trillion standard cubic feet barrel of natural gas and 31 billion barrel of oil equivalent of tar sand [2, 3, 9, 10].

Nigeria have been reported to produce about 6 billion standard cubic feet of gas per daily [10] and 2.2 – 2.7 million of crude oil daily [3, 10, 11]. But the production rate varies due to several factors such as the activities of militants in the oil rich region of Nigeria, pipeline vandalism, sabotages, oil spills, among other factors. Of these, the activities of militia in the region have significantly affected the production rate in Nigeria. For instance, crude oil production significantly decreased in first half of 2016 due to activities of militancy and resource control.

The Nigeria crude oil and natural gas is domicile in the Niger Delta including Ondo, Edo, Delta, Bayelsa, Rivers, Abia, Imo, Akwa Ibom and Cross Rivers states. The crude oil and natural gas is located in both offshore and onshore in the region. Two major products are produced include crude oil and natural gas. Both have adverse effect on the environment. Crude oil could spill into the environment and change the characteristics of the receiving environment including soil [12 – 14] and water quality. In some cases, the volatile

components of the crude oil are released into the air. The natural gas are basically utilized while the excess is flared into the atmosphere through combustion processes.

Gas flaring is common in the Niger Delta region of Nigeria. During flaring several pollutants gases are released into the environment including nitrogen dioxides, sulphur dioxide, volatile organic compounds like benzene, toluene, xylene, polyaromatic hydrocarbons, hydrogen sulfide, benzopyrene and dioxins [15, 16], and particulates.

As such, several impacts (including human and environmental) are associated to gas flaring. Several studies have indicated the impacts of gas flaring in the Niger Delta including vegetation and physical infrastructure such as roofing sheet, buildings/structures, artifacts, monuments, paints [15, 17 – 23], pathological and psychological impacts on human health [24]. As such, several diseases have been reported to be associated with gas flaring including excessive heat and discomfort [18], gastrointestinal problems, skin diseases, cancer, neurological, reproductive and developmental effects, haematological and respiratory ailments [15], heart (cardiovascular) related illness including atherosclerosis, hypertension and ischaemic heart disease [25, 26], renal and related diseases [27], bronchitis, asthma, cancers and several other diseases [20]. Most of the impacts are due to indirect effects resulting from acid rain. These could also affect other biodiversity resources including humans. For instance, acid rain which could result from gas flaring has the tendency to cause lung related diseases, and affect aquatic organisms such as fishes and other wildlife and natural forest resources such as vegetation. Furthermore, Noise emanating from the flare could also affect humans residing close/ and or working close to the vicinity.

As such the exploration of oil and gas and flaring of natural gas has several environmental impacts. Therefore, this study reviews the impacts gas flaring on groundwater resources and vegetation structure and cover in the Niger

Delta region of Nigeria. The paper is organized into 5 sections. Section 1 is the introduction providing information on oil and gas resources in Nigeria and overview of the impacts of gas flaring in the Niger Delta. Section 2 discussed gas flaring in Nigeria. Section 3 discussed the effect of season and distance of disposal of emission. Section 4 discussed the impact of gas flaring to vegetation and water resources. Section 5 is conclusion and the way forward.

2. Gas Flaring: A Nigerian Scenario

Globally, high amount of gas is flared in to the environment by oil and gas producing countries. Nigeria being among the world producing nations, flare a significant amount of natural gases into the environment through vertical and horizontal flaring stack (Figure 1). Globally about 110 billion cubic meters of associated gas is flared per annum [28]. Ogbé [29] opined that Nigeria account for about 12.5% global flared gases per annum [29]. The flaring of gases is a global issue for some decades now [3]. Emam [31] described gas flaring as the used combustion device (flare stack) to remove unwanted gases and liquids during operation in many industrial processes, such as oil-gas extraction, refineries, chemical plants, coal industry and landfills to prevent unplanned over-pressuring. Soltanieh *et al.* [32] also noted that gas is flared in producing nations due to a number of reasons such as inadequate infrastructure to collect, treat, transport and utilize the associated gases; location of the production site is remote from the market demand (such as offshore sites); small volume of the gas and its fluctuation, which make the design of facilities more uncertain and therefore uneconomical investment; Impurities in the gas that require hard and expensive treatment methods (such as highly acidic gases); safety and operational reasons. Gas flaring leads to release of three major components including noxious gases, heat and noise.




Figure 1. Gas flaring in a location in the Niger Delta.

Natural gas can be converted into different form for downstream applications including electricity generation and cooking gas. But due to inadequate resources for its conversation and utilization, oil and gas companies prefers to flare the gases and pay compensation. According to Donwa *et al.* [15], wastage of gases through flaring is carried out due to problems associated to processing, storing and transporting it in Nigeria setting.

Nigeria flare significant amount of natural gas into the environment leading to loss of substantial amount of money per annum. For instance, World Bank reported that 150 to 170 billion m³ of gases are flared annually, worth up to about \$ 30.6 billion, the price equivalent of one-quarter of the United States' gas consumption or 30% of the European Union's yearly gas consumption [31]. Lower amount of gas equivalent have been reported to loss due to gas flaring to the tune of \$2.0 billion per annum [29], \$2.5 billion [15, 33].

Despite the incentives to capture the associated gas and

bring it to market, the volume of gas flared is still high. About 70million /m³ of natural gas are flared per day [34]. On yearly basis, Donwa *et al.* [15], Ishisone [35] reported that Nigeria flare about 17.2 billion m³ of natural gas in the Niger Delta. Between 2006 to 2014 several oil wells were explored in Nigeria. Table 1 presents information the details of crude oil and condensate production, total gas and quantity utilized and flared between 2006 to 2014. Within the period the production, utilization and quantity flared varies. Also, the amount targeted could not be met. This is typically attributed to several reasons including delays in upgrade of facilities to floody terrain, inadequate line, limited facilities, obsolete equipment, community disturbance, activities of militia etc. The demand for gas increase as a result of new opportunities for gas micro power, combined cycle turbines, independent power plant, gas to liquids and expansion in liquefied natural gas trade also contributed to the utilization.

Table 1. Crude oil and gas production and utilization in Nigeria between 2004 to 2014.

Years	crude oil and condensate production, barrels		Total gas, Billion Standard Cubic Feet	Quantity of gas utilized, BSCF	Quantity flared, BSCF	References
	Total	Daily average, mmb/pd				
2014	798,541,589	2.19	2,524.27	2,233.49	289.60	[36]
2013	800,488,102	2.19	2,325.14	1,916.53	409.31	[37]
2012	852,776,653	2.27	2,580.17	1,991.50	588.67	[38]
2011	866,245,232	2.37	2,400.40	1,781.37	619.03	[39]
2010	896,043,406	2.45	2,392.84	1,811.27	581.57	[40]
2009	780,347,940	2.14	1,837.28	1,327.93	509.35	[41]
2008	768,745,932	2.10	2,282.44	1,664.97	617.62	[42]
2007	803,000,708	2.20	2,415.65	1,626.10	789.55	[43]
2006	869,196,506	2.38	2,182.43	1,382.43	799.99	[44]

Furthermore, Nigeria flares about 11 – 42.54% of total natural gas produced. These make Nigeria one of the largest gas flaring nations in the World. Several oil wells are built with the period of 2004 to 2014, with an average of 155 per annum. According to Oniemola and Sanusi [45], Nigeria has about 160 oil fields having about 1500 oil wells that produces 2.2 to 2.7 million barrels/day. The authors reported that of these, 17 billion m³ of associated gas are flared leading to the releasing 2,700, 160, 5400, 12million and 3.5 million tons of particulate, sulphur oxides, carbon monoxide, carbon dioxide and methane respectively. The number of wells has far increased due to continual exploration.

3. Effect of Season and Distance of Disposal of Emission

Nigeria has two predominant season including wet season (April to October) and dry season (November to March of the following year). The rainfall pattern is usually optimum in June, July and September. The effect of meteorology especially temperature, rainfall, relative humidity, wind speed have been reported in the Niger Delta specifically in Bayelsa state [46].

Season affects the rate of dispersal of pollutants and noise. Dispersal of pollutants resulting from emissions is affected by

season. For example, Anomohanran [18] reported pollution from thermal plant within a distance of 2.15 km and 2.06km in wet and dry season respectively. Several metrological parameters affects dispersal rate of pollutant gases. Some of the notable once include wind speed, wind direction, closeness to ocean.

Distance in another major factor that could affect the emission of pollutants. Emission rate typically decreases as distance from the source of pollution is increased. Ojeh [47] reported that in a gas flaring station, the concentration of noxious gases reduces as the distance increased in gas flaring site. But the effects of gas flared could be felt within 450m radius of the flare stack which could be depend on the volume of gas flared, wind speed, temperature, temperature, velocity of discharged and height of the stack [47]. In non-gas flaring site, specifically on oil palm processing mill, emissions also decreases as the distance from the sources increased [48, 49]. Though, instances of fluctuation in wind speed and direction during measurement could lead to shorter distance from emission source having lower values instead of higher values compared to long distance.

4. Impacts of Gas Flaring on Water Quality and Vegetation Resources

Gas flaring typically has two valves, the high and low gas

pressure valves. Hence, the concentration of emissions, noise, released is a function of the valve used in flaring the gas as well as the physical composition of the gas prior to flaring. The flared gases affect several environmental components as well as its biotic composition. The section of the paper affects the impact of gas flaring on water quality and vegetation resources.

4.1. Changes in Water Quality

Water resources are typically needed for the sustenance of life, growth and development [50 – 62]. Water serve as habitant to several biodiversity especially fisheries (shelled and finfish), aquatic reptiles, mammals, birds and breeding ground for several other diversity such as frog, parasite of medical importance such as Schistome, and vectors transmitting diseases such as mosquitoes. As such water has a unique place for life to thrive.

Water is also used for domestic purposes such as washing, cooking, bathing [63]. Most water resources are from surface water, groundwater and rainwater [52]. Izah et al. [50] is with the opinion that groundwater is mostly consumed in the Niger Delta and surface water and rainwater also serve as potable water sources in regions that groundwater/ borehole water is unavailable.

Water is typically impacted by gas flaring activities. Authors have variously reported that some water quality parameters of surface water, ground water and rain water are affected in gas flaring locations in the Niger Delta. For instance, Dami et al. [64] studied the impacts gas flaring and oil spillage on rainwater quality for domestic use in Okpai and Beneku areas of Delta State and reported that temperature, taste, color, conductivity, total dissolved, salts and alkalinity were altered when compared to the permissible limits specified by National Agency for Food and Drug Administration and Control, United State Environmental Protection Agency and World Health Organization for drinking water. Dami et al. [65] studied the impacts of gas flaring and oil spillage on groundwater quality for domestic use in Okpai and Beneku areas of Delta State and reported that color is majorly impacted and to lesser extent conductivity were affected in some locations due to accumulation of dissolved salts and other organic materials when compared to World Health Organization standard. Nwankwo and Ogagarue [66] studied the effect of gas flaring in surface and groundwater quality in Delta state and reported that water from gas flaring area contain higher concentrations of metals such as barium, cyanide, selenium, cadmium, chromium, iron, manganese and copper, conductivity, color and taste when compared to non-flaring location. Emumejaye [67] studied the effects of gas flaring on surface and ground water in Irri town and environs and reported that high content of iron and lead in the water which the author attributed to gas flaring in the area. Ezenwaji et al. [68] studied the effects of gas flaring on rainwater quality in Bayelsa State and reported that all the parameters studied including temperature, lead, conductivity, total dissolved solid, nitrate, carbonate, sulphate and pH had values above World Health Organization permissible limits.

The authors further reported that nitrate contributing the highest of 38.44% to poor quality of rain water in the area. Amadi [17] reported that concentration of the major anions (bicarbonate, sulphate and nitrate), temperature, conductivity, heavy metal content and total hydrocarbon content increase significantly in the vicinity of the gas flaring location and steadily declined in surface water, groundwater and rainwater away from the flaring location. Efe [69] studied the effect of acid rain and reported that pH values (which is usually used to measure the acidity of the water) were lower (4.96) in the Niger Delta compared to coastal region of the Northern Nigeria with a mean pH of 5.36 which indicate the impact of acid rain. Other effects of acid rain on water quality have been comprehensively documented in literature by Ogunkoya and Efi [70], Ubani and Onyejekwe [23], Olobaniyi et al. [71], Efe and Mogborukor [72].

Among the water quality parameters commonly affected by gas flaring is nitrate, carbonate, sulphate, nitrate, lead etc. Amadi [17] reported that gas flaring reduces pH (pH tending toward acidity), moisture content and bacteria density/population. The intensity of the flare with regards to microbial density in water is usually higher in horizontal flare compared to the vertical flare stack. The heat from the flare could enhance the temperature of the environment including water. High occurrence of sulphate and nitrate ions may be connected to the emission of carbon, sulphur and nitrogen oxide during flaring activities. The occurrence of ions in the water could lead to high conductivity level in water close to gas flaring location.

4.2. Impact on the Vegetation

Plants are a major source of active nutrients required by the body. Plants are source of carbohydrate, protein, lipid and other minerals and vitamins depending on species. Vegetation is also source of food for livestock especially omnivorous animals such as goat, cow, grass cutter, rabbits etc. Vegetation cover is also a source of habitat to several wildlife species especially bushmeat that do not burrow. Some notable vegetation is found in close to farmland and residential area such as cassava and oil palm. These are two predominant vegetation cover used as food found close to gas flaring location in the Niger Delta. Gas flaring has the tendency to affect several plant species [73] especially productivity and growth [74]. For instance, Lawanson [75] reported that gas flaring decreases the length and weight of cassava and increase its amino acid and total sugar contents as the distance from the flares decreases. The authors furthered reported that such decreases were also correlated with decreases in the content of starch and ascorbic acid (vitamin C) in the tubers. Based on survey study in the Niger Delta region, 77% of the resident is with the opinion that gas flaring affect vegetation and agricultural activities [76]. In a similar study the resident of Ebedei community in Delta state 94.6, 90, 98.75, 50.4 and 5% is with the opinion that gas flaring affect food such as yam, cassava, okra, plantain and potatoes [74].

Gas flaring can also cause deforestation and acid rain [74]. According to Ezenwaji et al. [68], exploitation of crude oil and

associated gas flaring is a major contributor of acid rain in the Niger Delta region of Nigeria has had a fairly long history [68]. Acid rain could lead to loss of vegetation [17] and several symptoms in plants that could lead to their death. Some of the notable symptoms include chlorosis, abscission and yellowing of leaves, wilting of the leaf tips and accelerated senescence, root and shoot of plants are also destroyed and microbial community that aid in decompositions processes [69]. The impacts of acid rain on vegetation structures and cover is most severe close to gas flaring stack [69].

Acid rain result in the decline in productivity and growth of some major food crops such as cassava, sweet potatoes, maize, melon, plantain, and cash crop like rubber [69]. The impacts in the growth and productivity of crops could also be an indication that the soil fertility have been impacted upon. This may lead to loss of vital soil nutrients that encourage the growth of plants. Other authors have comprehensively reported the effect acid rain on vegetation have been documented by Jacobson [77], Neufeld et al. [78], Efe [79].

Plants are known to pharmacological and bioactive composition [80, 81]. The composition of the bioactive constituents play a significant role in determining their medicinal properties. The effect of gas flaring on the nutritional and bioactive component of vegetation established. For instance, Ifemeje [82] reported gas flaring could change the anti-nutrient compositions (alkaloid, phytate, oxalate, Saponin, tannin and cyanogenic glycosides) in some common vegetables used for food purposes such as scent leaf, bitter leaf, water leaf and fluted pumpkin leaf. Ujowundu *et al.* [83] also reported impacts in phytochemical (alkaloid, tannin, cyanogenic glycoside, phytate), proximate composition (moisture, ash, protein and carbohydrate), micronutrients (calcium, sodium, magnesium, potassium and phosphorus) and vitamins (riboflavin, vitamin E and C) in African breadfruit and Bambara groundnuts planted close to gas flaring stack. Anacleto *et al.* [84] also reported that phytochemical (alkaloids, flavonoids, saponins and tannins) and trace metal (iron, lead, cadmium and zinc) constituents of fluted pumpkin could be affected by gas flaring.

Vegetation plays several ecological roles. For instance Izah et al. [85] reported that vegetation prevents soil erosion. Gas flaring is known to cause physical damage to plant close to the flare stack [17]. This could lead to other downstream impacts. In addition, gas flaring could alter soil quality parameters [86] including physiochemical and microbial characteristics. Some notable soil quality parameters such as pH, temperature, soil moisture, soil microbial population are commonly impacted by gas flaring [23]. Okeke and Okpala [86] reported that soil quality parameters from flaring sites such as temperature and bulk density decreased with distance from the flare point while other such as CEC, organic matter, moisture content etc) increases with distance. The authors further asserted that soil nutrient were lower in gas flaring environment compared to the control in Eket and Izombe area of the Niger Delta. Variation in soil characteristics especially the nutrient related characteristics may affect the crop productivity indirectly. Microbes being unique, changes in the soil properties could

alter the microbial diversity and density. Typically microbes play essential role in nutrient and biogeochemical cycling.

In a survey study, the resident of Okpai, Ndokwa East Local Government Area, Delta State is with the opinion that gas flaring is having impact on the soil including its fertility and productivity for food crops such as cassava, plantain, yam [87]. Also maize is also affected by gas flaring and cassava [88, 89].

5. Conclusion and the Way Forward

The Niger Delta region is one of the most productive and fragile ecosystem in Nigeria. The area is rich in vegetation with several pharmacological properties and water resources. These resources are constantly being destroyed by the activities of man in quest for industrialization and urbanization. Several oil and gas installations are found in the region. Gas is used by gas-turbine for electricity generation. Despite the huge source of gas available for electricity, the gas is constantly flared into the environment. While electricity supply remain epileptic in the region. Nearly 10 – 40% of natural gas produced is under-utilized and therefore flared into the environment. Gas flaring are known to have impact on air quality, physical infrastructure, biodiversity composition including plants and animals especially insects, impacts on human health over a prolong period of time and water resources especially rainwater. Acid rain has been widely attributed to impact of gas flaring especially in the Niger Delta region of Nigeria. On water quality, gas flaring alters ions especially sulphate, carbonate, nitrate, pH, conductivity, lead and iron concentration especially in rainwater. On vegetation perspective, it could lead to loss of vegetation cover, reduced growth and productivity/yield probably due to changes in soil quality parameters.

Based on the review, the attendant impacts associated with gas flaring on vegetation and water quality could be reduced through:

- a. utilization of the gas and generation of revenue from it, promulgation of associated gas re-injection and amendment of flaring policy to more reasonable amount [67].
- b. enforcement of laws aimed at minimizing the amount of gas flared into the atmosphere [68].

References

- [1] Mogborukor, J. O. A. (2014). The Impact of Oil Exploration and Exploitation on Water Quality and Vegetal Resources in a Rain Forest Ecosystem of Nigeria. *Mediterranean Journal of Social Sciences*, 5 (27): 1678 – 1685.
- [2] Ohimain, E. I. (2013). Can the Nigerian biofuel policy and incentives (2007) transform Nigeria into a biofuel economy? *Energy Policy*, 54: 352 – 359.
- [3] Ohimain, E. I. (2013). The challenges of liquid transportation fuels in Nigeria and the emergence of the Nigerian automotive biofuel programme. *Research Journal of Applied Sciences, Engineering and Technology*, 5 (16): 4058 – 4065.

- [4] Izah, S. C. and Ohimain, E. I. (2015). Bioethanol production from cassava mill effluents supplemented with solid agricultural residues using bakers' yeast [*Saccharomyces cerevisiae*]. *Journal of Environmental Treatment Techniques*, 3 (1): 47 – 54.
- [5] Ede, P. N. and Edokpa, D. O. (2015). Regional Air Quality of the Nigeria's Niger Delta. *Open Journal of Air Pollution*, 4: 7-15.
- [6] Ohimain, E. I., Emeti, C. I., Izah, S. C. (2014). Employment and socioeconomic effects of semi-mechanized palm oil mill in Bayelsa state, Nigeria. *Asian Journal of Agricultural Extension and Sociology*, 3 (3): 206-216.
- [7] Ohimain, E. I. (2010). Petroleum Geomicrobiology. In: Jain, S. K., Khan, A. A., Rain, M. K., (Editors). *Geomicrobiology: Biodiversity and Biotechnology*. CRC Press/Taylor and Francis, Boca Raton, Florida, USA. Pp. 139 – 174.
- [8] Ohimain, E. I. (2013). Environmental impacts of smallholder ethanol production from cassava feedstock for the replacement of kerosene household cooking fuel in Nigeria. *Energy Sources*, Part A. 35: 1560 – 1565.
- [9] Ohimain, E. I. (2013). A review of the Nigeria biofuel policy and incentives (2007). *Renewable and Sustainable Energy Reviews*, 22: 246 – 256.
- [10] Sambo, A. S. (2008). The role of energy in achieving millennium development goal (MDGs). Keynote address at the national engineering technology conference (NETec 2008). Ahmadu Bello University, Zaria held on 1st of April, 2008.
- [11] Fadare, D. A., Bamiro, O. A. and Oni, A. O. (2009). Energy analysis for production of powdered and pelletised organic fertilizer in Nigeria. *ARPJ Journal of Engineering and Applied Sciences*, 4 (4): 75 – 82.
- [12] Aigberua, A. O., Ekubo, A. T., Inengite, A. K. and Izah, S. C. (2017). Assessment of some selected heavy metals and their pollution indices in an oil spill contaminated soil in the Niger Delta: a case of Rumuolukwu community. *Biotechnological Research*, 3 (1): 11- 19.
- [13] Aigberua, A. O., Ekubo, A. T., Inengite, A. K. and Izah, S. C. (2016). Evaluation of Total Hydrocarbon Content and Polycyclic Aromatic Hydrocarbon in an Oil Spill Contaminated Soil in Rumuolukwu Community in Niger Delta. *Journal of Environmental Treatment Techniques*, 4 (4): 130 – 142.
- [14] Aigberua, A. O., Ekubo, A. T., Inengite, A. K. and Izah, S. C. (2016b). Seasonal variation of nutrient composition in an oil spill contaminated soil: a case of Rumuolukwu, Eneka, Port Harcourt, Nigeria. *Biotechnological Research*, 2 (4): 179-186.
- [15] Donwa, P. A., Mgbame, C. O., Utomwen, O. A. (2015). Gas flaring in the oil and gas sector in Nigeria. *International Journal of Commerce and Management Research*, 1 (1): 28-39.
- [16] Adamu, H. and Umar, B. A. (2013). Occurrence and Chemistry of Co-contamination of Nitrate and Hydrocarbon Pollutants in Gas-Flared Areas of Niger-Delta, Nigeria. *International Journal of Environmental Monitoring and Analysis*, 1 (4): 139-146.
- [17] Amadi, A. N. (2014). Impact of Gas-Flaring on the Quality of Rain Water, Groundwater and Surface Water in Parts of Eastern Niger Delta, Nigeria. *Journal of Geosciences and Geomatics*, 2 (3): 114-119.
- [18] Anomohanran, O. (2012). Thermal Effect of Gas Flaring at Ebedei Area of Delta State, Nigeria. *The Pacific Journal of Science and Technology*, 13 (2): 555 – 560.
- [19] Iyorakpo, J. and Odibikuma, P. W. (2015). Impact of gas flaring on the built Environment: the case of Ogba/Egbema/Ndoni Local Govt Area, Rivers State, Nigeria. *European Scientific Journal*, 11 (26): 83 – 95.
- [20] Olukoya, O. A. P. (2015). Negative Effects of Gas Flaring On Buildings and Public Health in Oil Producing Communities: The OgbiaCommunity, Bayelsa State Case. *International Journal of Environmental Monitoring and Protection*, 2 (5): 52-61.
- [21] Nkwocha, E. E. and Pat-Mbano, E. C. (2010). Effect of Gas Flaring on Buildings in the Oil Producing Rural Communities of River State, Nigeria. *African Research Review*, 4 (2): 90-102.
- [22] Abua, M. A. and Ashua, S. W. (2015). The Impact of Gas Flaring on Plant Diversity in Ibeno Local Government Area. *Journal of Agriculture and Ecology Research International*, 4 (1): 10-17.
- [23] Ubani, E. C. and Onyejekwe, I. M. (2013). Environmental impact analyses of gas flaring in the Niger delta region of Nigeria. *American Journal of Scientific and Industrial Research*, 4 (2): 246-252.
- [24] Nriagu, J., Udofia, E. A., Ekong, I. and Ebuk, G. (2016). Health Risks Associated with Oil Pollution in the Niger Delta, Nigeria. *International Journal of Environmental Research and Public Health*, 13, 346; doi: 10.3390/ijerph13030346.
- [25] Egwurugwu, J. N., Nwafor A, Chinko BC, Oluronfemi OJ, Iwuji SC, Nwankpa, P. (2013). Effects of prolonged exposure to gas flares on the lipid profile of humans in the Niger Delta region, Nigeria. *American Journal of Research Communication*, 1 (5): 115-145.
- [26] Egwurugwu, J. N. and Nwafor, A. (2013). Prolonged Exposure to Oil and Gas Flares Ups the Risks for Hypertension. *American Journal of Health Research* 1 (3): 65-72.
- [27] Egwurugwu, J. N., Nwafor, A., Oluronfemi, O. J., Iwuji, S. C. and Alagwu, E. A. (2013). Impact of Prolonged Exposure to Oil and Gas Flares on Human Renal Functions. *International Research Journal of Medical Sciences*, 1 (11): 9-16.
- [28] Ismail, O. S. and Umukoro, G. E. (2012). Global Impact of Gas Flaring. *Energy and Power Engineering*, 4: 290-302.
- [29] Ogbe, E. (2010). Optimization of strategies for natural gas utilization: case study of the Niger Delta. A Thesis presented to the department of petroleum engineering, African University of Science and Technology.
- [30] Otio, D. (2013). Gas Flaring Regulation in the Oil and Gas Industry: A Comparative Analysis of Nigeria and Texas Regulations.
- [31] Emam, E. A. (2015). Gas Flaring In Industry: An Overview. *Petroleum and Coal*, 57 (5): 532-555.
- [32] Soltanieh, M., Zohrabian, A., Gholipour, M. J. and Kalnay, E. (2016). A review of global gas flaring and venting and impact on the environment: Case study of Iran. *International Journal of Greenhouse Gas Control*, 49: 488–509.
- [33] Ajugwo, A. O. (2013). Negative Effects of Gas Flaring: The Nigerian Experience. *Journal of Environment Pollution and Human Health*, 1 (1): 6-8.

- [34] Emoyan, O. O., Akporborie, I. A. and Akporhonor, E. E. (2008). The Oil and Gas Industry and the Niger Delta: implications for the Environment. *J. Appl. Sci. Environ. Manage.*, 12 (3): 29 – 37.
- [35] Ishisone, M. (Undated). Gas Flaring in the Niger Delta: the Potential Benefits of its Reduction on the Local Economy and Environment. <http://nature.berkeley.edu/classes/es196/projects/2004final/Ishone.pdf>, Assessed June 4th, 2016.
- [36] Nigerian National Petroleum Corporation (NNPC) (2014). Annual statistical bulletin. 1st Edition. Corporate planning and development division (CPDD).
- [37] Nigerian National Petroleum Corporation (NNPC) (2013). Annual statistical bulletin. 1st Edition. Corporate planning and development division (CPDD).
- [38] Nigerian National Petroleum Corporation (NNPC) (2012). Annual statistical bulletin. 1st Edition. Corporate planning and development division (CPDD).
- [39] Nigerian National Petroleum Corporation (NNPC) (2011). Annual statistical bulletin. 1st Edition. Corporate planning and development division (CPDD).
- [40] Nigerian National Petroleum Corporation (NNPC) (2010). Annual statistical bulletin. 1st Edition. Corporate planning and development division (CPDD).
- [41] Nigerian National Petroleum Corporation (NNPC) (2009). Annual statistical bulletin. Draft Edition. (Web Version). Corporate planning and development division (CPDD).
- [42] Nigerian National Petroleum Corporation (NNPC) (2008). Annual statistical bulletin. 1st Edition. (Web Version). Corporate planning and development division (CPDD).
- [43] Nigerian National Petroleum Corporation (NNPC) (2007). Annual statistical bulletin. 2nd Edition. (Web Version). Corporate planning and development division (CPDD).
- [44] Nigerian National Petroleum Corporation (NNPC) (2006). Annual statistical bulletin. 1st Edition. (Web Version). Corporate planning and development division (CPDD).
- [45] Oniemola, P. K. and Sanusi, G. (2009). The Nigerian biofuel policy and incentives (2007); a need to follow the Brazilian pathway. International Association for Energy Economics, 4th Quarter, pp: 135-139.
- [46] Izah, S. C., Angaye, T. C. N. and Ohimain, E. I. (2015). Climate change: some meteorological indicators and perception of farmers in Yenagoa metropolis, Bayelsa state, Nigeria. *International Journal of Geology, Agriculture and Environmental Sciences*, 3 (1): 56 – 60.
- [47] Ojeh, V. N. (2012). Sustainable Development and Gas Flaring Activities: a Case Study of Ebedei Area of Ukwuani LGA, Delta State, Nigeria. *Resources and Environment*, 2 (4): 169-174.
- [48] Ohimain, E. I. and Izah, S. C. (2013). Gaseous emissions from a semi-mechanized oil palm processing mill in Bayelsa state, Nigeria. *Continental Journal of Water, Air and Soil Pollution*, 4 (1): 15 – 25.
- [49] Ohimain, E. I., Izah, S. C. and Abah, S. O. (2013). Air quality impacts of smallholder oil palm processing in Nigeria. *Journal of Environmental Protection*, 4: 83-98.
- [50] Izah, S. C., Chakrabarty, N. and Srivastav, A. L. (2016a). A Review on Heavy Metal Concentration in Potable Water Sources in Nigeria: Human Health Effects and Mitigating Measures. *Exposure and Health*, 8: 285–304.
- [51] Izah, S. C. and Ineyougha, E. R. (2015). A review of the microbial quality of potable water sources in Nigeria. *Journal of Advances in Biological and Basic Research*, 1 (1): 12 – 19.
- [52] Izah, S. C. and Srivastav, A. L. (2015). Level of arsenic in potable water sources in Nigeria and their potential health impacts: A review. *Journal of Environmental Treatment Techniques*, 3 (1): 15 – 24.
- [53] Agedah, E. C., Ineyougha, E. R., Izah, S. C., and Orutugu, L. A. (2015). Enumeration of total heterotrophic bacteria and some physico-chemical characteristics of surface water used for drinking sources in Wilberforce Island, Nigeria. *Journal of Environmental Treatment Techniques*, 3 (1): 28 – 34.
- [54] Ogamba, E. N., Izah, S. C. and Oribu, T. (2015). Water quality and proximate analysis of Eichhornia crassipes from River Nun, Amassoma Axis, Nigeria. *Research Journal of Phytomedicine*, 1 (1): 43 – 48.
- [55] Ogamba, E. N., Izah, S. C. and Toikumo, B. P. (2015). Water quality and levels of lead and mercury in Eichhornia crassipes from a tidal creek receiving abattoir effluent, in the Niger Delta, Nigeria. *Continental Journal of Environmental Science*, 9 (1): 13 – 25.
- [56] Ogamba, E. N., Seiyaboh, E. I., Izah, S. C., Ogbugo, I. and Demedongha, F. K. (2015c). Water quality, phytochemistry and proximate constituents of Eichhornia crassipes from Kolo creek, Niger Delta, Nigeria. *International Journal of Applied Research and Technology*, 4 (9): 77 – 84.
- [57] Ogamba, E. N., Ebere, N. and Ekuma, C. G. (2017). Physicochemistry and Ichthyofauna of Ikoli Creek, Niger Delta, Nigeria. *Biotechnol Res* 3 (2): 43-49.
- [58] Seiyaboh, E. I., Izah, S. C. and Oweibi, S. (2017). Physico-chemical Characteristics of Sediment from Sagbama Creek, Nigeria. *Biotechnological Research*, 3 (1): 25-28.
- [59] Seiyaboh, E. I., Izah, S. C. and Oweibi, S. (2017). Assessment of Water quality from Sagbama Creek, Niger Delta, Nigeria. *Biotechnological Research*, 3 (1): 20-24.
- [60] Seiyaboh, E. I., Harry, G. A. and Izah, S. C. (2016). Length-Weight Relationship and Condition Factor of Five Fish Species from River Brass, Niger Delta. *Biotechnological Research*, 2 (4): 187-192.
- [61] Seiyaboh, E. I., Inyang, I. R. and Izah, S. C. (2016). Spatial Variation in Physico-chemical Characteristics of Sediment from Epie Creek, Bayelsa State, Nigeria. *Greener Journal of Environment Management and Public Safety*, 5 (5): 100 – 105.
- [62] Seiyaboh, E. I., Inyang, I. R. and Izah, S. C. (2016). Seasonal Variation of Physico-Chemical Quality of Sediment from Ikoli Creek, Niger Delta. *International Journal of Innovative Environmental Studies Research*, 4 (4): 29-34.
- [63] Oyoroko, E. and Ogamba, E. N. (2017). Effects of detergent containing linear alkyl benzene sulphonate on behavioural response of Heterobranchus bidorsalis, Clarias gariepinus and Heteroclaris. *Biotechnol. Res.* 3 (3): 55-58.

- [64] Dami, A., Ayuba, H. K. and Amukali, O. (2012). Effects of Gas Flaring and Oil Spillage on Rainwater Collected for Drinking in Okpai and Beneku, Delta State, Nigeria. *Global Journal of Human Social Science Geography & Environmental GeoScience*, 12 (13): 25- 29.
- [65] Dami, A., Ayuba, H. K. and Amukali, O. (2013). Ground water pollution in Okpai and Beneku, Ndokwa east local government area, delta state, Nigeria. *E3 Journal of Environmental Research and Management*, 4 (1): 0171-0179.
- [66] Nwankwo, C. N. and Ogagarue, D. O. (2011). Effects of gas flaring on surface and ground waters in Delta State Nigeria. *Journal of Geology and Mining Research*, 3 (5): 131-136.
- [67] Emumejaye, K. (2012). Effects of Gas Flaring On Surface and Ground Water in Irri Town and Environs, Niger-Delta, Nigeria. *IOSR Journal Of Environmental Science, Toxicology And Food Technology*, 1 (5): 29-33.
- [68] Ezenwaji, E. E., Okoye, A. C. and Otti, V. I. (2013). Effects of gas flaring on rainwater quality in Bayelsa State, Eastern Niger-Delta region, Nigeria. *Journal of Toxicology and Environmental Health Sciences*, 5 (6): 97-105.
- [69] Efe, S. I. (2011). Spatial Variation of Acid Rain and its Ecological Effect in Nigeria. Proceedings of the Environmental Management Conference, Federal University of Agriculture, Abeokuta, Nigeria, pp 381 – 396.
- [70] Ogunkoya, O. O. and Efi, E. J. (2003). Rainfall quality and sources of rainwater acidity in Warri area of the Niger Delta, Nigeria. *Journal of Mining and Geology*, 39 (2): 125-130.
- [71] Olobaniyi, S. B.; Ogban, F. E.; Ejechi, B. O. and Ugbe, F. C. (2007), Quality of Groundwater in Delta State, Nigeria. *Journal of environmental Hydrology*, 15: 1-9.
- [72] Efe, S. I. and Mogborukor J. O. A. (2008). Acid rain in the Niger Delta region: Implication on Water Resource Crises, Proceedings, International Conference on the Nigerian State, Oil industry and the Niger Delta, 11-13th March, 2008, Glory land Cultural Centre, Yenagoa, Bayelsa State, Nigeria.
- [73] Achi, C. (2003). Hydrocarbon exploitation, environmental degradation and poverty: the Niger Delta experience. Diffuse Pollution Conference, Dublin 2003 2B Policy / Socio-Economics.
- [74] Ozabor, F. and Obisesan, A. (2015). Gas Flaring: Impacts on Temperature, Agriculture and the People of Ebedei in Delta State Nigeria. *Journal of Sustainable Society*, 4 (2): 5-12.
- [75] Lawanson, A. O., Imevbore, A. M. A. and Fanimokun, V. O. (Undated). The Effects of Waste-Gas Flares on the Surrounding Cassava Plantations in the Niger Delta Regions of Nigeria. http://www.istrc.org/images/Documents/Symposiums/Sixth/6th_symposium_proceedings_0041_section_3_239.pdf. Accessed September 21st, 2016.
- [76] Adewale, O. O. and Mustapha, U. (2015). The Impact of Gas Flaring in Nigeria. *International Journal of Science, Technology and Society*, 3 (2): 40-50.
- [77] Jacobson, J. S. (1984). Effect of Acidic Aerosol, fog, Mist and Rain on Crops and Trees, *Phill. Trans. Roy. Soc. Lond. B* 49: 327-338.
- [78] Neufeld, H. S., Jernstedt, J. A. and Haines, B. L. (1985). Direct Foliar Effects of Simulated Acid Rain, I. Damage, Growth and Gas Exchange *New Phytologist*, 99 (3): 389-405.
- [79] Efe, S. I. (2010). Spatial variation in acid and some heavy metal composition of rainwater harvesting in the oil producing region of Nigeria. *Natural Hazard*, DOI 10.1007/s11069-010-9526-2.
- [80] Epidi, J. O, Izah, S. C. and Ohimain, E. I., Epidi, T. T, (2016). Phytochemical, antibacterial and synergistic potency of tissues of *Vitex grandifolia*. *Biotechnological Research*, 2 (2): 69-76.
- [81] Epidi, J. O., Izah, S. C. and Ohimain, E. I. (2016). Antibacterial and Synergistic Efficacy of Extracts of *Alstonia boonei* Tissues. *British Journal of Applied Research*, 1 (1): 0021-0026.
- [82] Ifemeje, J. C. (2015). Effect of Gas Flaring on the Anti-nutritional Composition of Four Green Leafy Vegetables from Eleme in Rivers State, Nigeria. *Columbia International Publishing International Journal of Environmental Pollution and Solutions*, 3 (1): 31-37.
- [83] Ujowundu, C. O., Nwaogu, L. A., Ujowundu, F. N. and Belonwu, D. C. (2013). Effect of Gas Flaring on the Phytochemical and Nutritional Composition of *Treculia Africana* and *Vigna subterranean*, *British Biotechnology Journal*, 3 (3): 293-304.
- [84] Anacletus, F., Adisa, O. I. and Uwakwe, A. (2014). Effect of Gas Flaring on Some Phytochemicals and Trace Metals of Fluted Pumpkin (*Telferia occidentalis*). *Journal of Environment and Earth Science*, 4 (16): 1 – 5.
- [85] Izah, S. C., Angaye, T. C. N. and Ohimain, E. I. (2016). Environmental Impacts of Oil palm processing in Nigeria. *Biotechnological Research*, 2 (3): 132-141.
- [86] Okeke, P. N. and Okpala C. Q. (2014). Effects of gas flaring on selected arable soil quality indicators in the Niger Delta, Nigeria. *Sky Journal of Soil Science and Environmental Management*, 3 (1): 001 – 005.
- [87] Olisemauche, O. O. and Avwersuoghene, O. P. (2015). The effect of gas flaring on Agricultural production of Okpai, Ndokwa East Local Government Area, Delta State, Nigeria. *Standard Scientific Research and Essays*, 3 (9): 266-272.
- [88] Odjugo, P. A. O. and Osemwenkhae, E. J. (2009). Natural gas flaring affects microclimate and reduces maize (*Zea mays*) yield. *Int. J. Agri. Biol.*, 11: 408–412.
- [89] Odjugo, P. A. O. (2007). Some Effects of Gas Flaring on the Microclimate of Yam and Cassava Production in Erhorike and Environs. Delta State, Nigeria. *Nigerian Geographical J.* 5: 43-54.