

Characterization on PM_{2.5}/PM₁₀ During Winter Period in Guangzhou, China

Tian Lixue

School of Geography Sciences, South China Normal University, Guangzhou, China

Email address:

li-xue.tian@m.scnu.edu.cn

To cite this article:

Tian Lixue. Characterization on PM_{2.5}/PM₁₀ During Winter Period in Guangzhou, China. *International Journal of Environmental Monitoring and Analysis*. Vol. 3, No. 5, 2015, pp. 331-333. doi: 10.11648/j.ijema.20150305.23

Abstract: With the rapid economic development and urbanization over the past several decades, air pollution and degradation of visibility are common in Guangzhou, especially in winter. Our objective is to generalize the variation characteristic of PM_{2.5} and PM₁₀ during winter period in Guangzhou and provide some theoretical evidence for the government and the relevant administration to defend PM_{2.5} and PM₁₀. This article collects the concentration data of PM_{2.5} and PM₁₀ in the area of Guangzhou from 2014.11-2015.1 (the winter period), and the concentration of PM_{2.5} and PM₁₀ were $(55.01 \pm 78.19) \mu\text{g}/\text{m}^3$ and $(78.34 \pm 101.76) \mu\text{g}/\text{m}^3$, respectively. The mean ratio of PM_{2.5} to PM₁₀ was 70.59% and the PM_{2.5}/PM₁₀ value ranges from 54.61% to 94.37%, indicating that fine particle (PM_{2.5}) occupy high proportion in the PM₁₀. In addition, PM_{2.5} has a good linear relationship with PM₁₀. All evidence provides us conclusions that fine particle has a higher proportion in the atmospheric aerosol, and make a major contribution in the air environmental pollution.

Keywords: PM_{2.5}, PM₁₀, PM_{2.5}/PM₁₀, Variation Characteristic, Guangzhou

1. Introduction

With the rapid economic growth and urbanization during the last several decades, air pollution has become a pressing environmental problem in China [1]. PM_{2.5}, the particulate matter with an aerodynamic diameter of less than 2.5 μm and PM₁₀, the particulate matter with an aerodynamic diameter of more than 2.5 μm and less than 10 μm , were increasingly emphasized [2]. The epidemiological studies have demonstrated that the PM exposure is associated with the occurrence of acute respiratory infections, lung cancer and chronic respiratory and cardiovascular diseases [3, 4, 5].

Compared to the foreign, the study of PM_{2.5}, PM₁₀ in China started relatively later, and the research is focused on a few big cities, such as Beijing [6], Shanghai [7], Hong Kong [8], Guangzhou [9]. Guangzhou (112°57'E - 114°03'E, 22°26'N - 23°56'N), the capital city of Guangdong Province in China, has a population of more than 10 million and is located in the northern Pearl River Delta (PRD). With the rapid economic development and urbanization over the past several decades, air pollution and degradation of visibility are common in the PRD region [10-11].

Big cities have more seriously haze phenomenon, even Guangzhou, the China's most southern coastal city has also

suffered. However, the research of large cities' haze phenomenon especially in winter is rare. This paper focuses on the study of Guangzhou in winter and combines the PM_{2.5} with PM₁₀. More importantly, this paper has a higher reference value owing to the use of the latest data.

2. Material and Methods

Table 1. Ambient air quality standards (GB 3095-2012).

	Average time	Concentration limit		Unit
		primary	secondary	
PM _{2.5}	Annual average	15	35	ug/m ³
	24hour average	35	75	
PM ₁₀	Annual average	40	70	
	24hour average	50	150	

Our data are collected from the Guangdong Environmental Monitoring Center, which starts on Mar 8, 2012, meanwhile the PRD region become the first city agglomeration that publish monitoring indicators and evaluate air quality after implementing the new «Ambient air quality standards» (GB3095-2012) (Table.1). There are thirteen monitoring stations in Guangzhou, and data used in this paper are from the average 24 hours concentration values of thirteen monitoring stations. PM_{2.5} and PM₁₀ are measured by the Particle Analyzer

based on the Tapered Element Oscillating Microbalance (TEOM) working principle.

3. Results and Discussion

3.1. $PM_{2.5}$ and PM_{10}

There are ninety-two samples during the winter period (from Nov 1, 2014 to Jan 31, 2015). The variations of the $PM_{2.5}$ mass concentration are shown in Figure 1. The mass concentration values of $PM_{2.5}$ range from 9.0 ug/m^3 to 133.2 ug/m^3 (Table 2). And the mean mass concentration of $PM_{2.5}$ was 55.01 ug/m^3 . The highest concentration of $PM_{2.5}$ was

133.2 ug/m^3 , in excess of 60 ug/m^3 than standard (75 ug/m^3) approximately, and the least values were 9.0 ug/m^3 , which was favorable air environment for living. Among the total samples, 10 samples' $PM_{2.5}$ mass concentration exceed standard (75 ug/m^3) (Table 2), occupying 10.87% of total sample number.

The variations of the PM_{10} mass concentration are also shown in Figure 1, ranging from 14.6 ug/m^3 to 180.1 ug/m^3 , and the average values was 78.34 ug/m^3 . The peak of PM_{10} mass concentration (180.1 ug/m^3) just a little more than the standard 150 ug/m^3 , and this phenomenon is better than $PM_{2.5}$. There are 4 samples exceed standard, and they have a proportion of 4.35% in total samples.

Table 2. Summary of $PM_{2.5}$ and PM_{10} , $PM_{2.5}/PM_{10}$ from Nov 2014 to Jan 2015.

	Mean/(ug/m^3)	Max./(ug/m^3)	Min./(ug/m^3)	Total sample number/d	Exceed standard number/d	The percentage of exceed standard number to sample number/%
$PM_{2.5}$	55.01	133.2	9.0	92	10	10.87
PM_{10}	78.34	180.1	14.6	92	4	4.35
$PM_{2.5}/PM_{10}$	70.59%	94.37%	54.61%	92		

Seen from Figure 1, there are several peaks, such as Dec 15, 2014, Dec 24-26, 2014, Dec 30-31, 2014, Jan3, 2015, Jan 20-21, 2015. By viewing historical weather data, during this peak period, the weather was dry and without or little wind

and rain, which inhibits the dispersion of pollutants. Therefore, the $PM_{2.5}$ and PM_{10} mass concentration were closely associated with weather conditions.




Figure 1. Variation characteristic of $PM_{2.5}$, PM_{10} , and $PM_{2.5}/PM_{10}$.

Throughout the values of $PM_{2.5}$ and PM_{10} , the author has come to the proportion of pollution levels in the winter period, as shown in Figure 2. The benign environmental condition has the largest proportion, that is 73.91%. The optimal and mild pollution have the same proportion, and the rest is 2.17%, proportion of moderate pollution. The percentage of exceed standard is 14.13% and the overall is good air quality.

3.2. $PM_{2.5}/PM_{10}$

The ratio of $PM_{2.5}/PM_{10}$ indicates the percentage composition of $PM_{2.5}$ fine particle in the PM_{10} . Shown from Table 2 and Figure 1, among the total ninety-two samples, the

mean ratio of $PM_{2.5}$ to PM_{10} is 70.59%. The $PM_{2.5}/PM_{10}$ value ranges from 54.61% to 94.37%. Compared with other domestic cities, such as Wuhan (61%), Lanzhou (52%), Chongqing [12] (65%) and Hong Kong [13] (82%), the ratio of $PM_{2.5}$ to PM_{10} in Guangzhou during the sampling period is higher than some inner cities and the data of Guangzhou is closed to that of Hong Kong as probably they are both in the Pearl River Delta of China. This evidence provide us a conclusion that fine particle make a major contribution in the air environmental pollution. For example, in recent years, the increased number of hazy may have a close relationship with that in the region.




Figure 2. The proportion of pollution levels.

3.3. The Relationship Between PM_{2.5} and PM₁₀




Figure 3. Correlation of PM_{2.5} and PM₁₀.

Figure 3 shows plots of PM_{2.5} and PM₁₀ concentrations and the linear relationships between them. PM_{2.5} has a good linear relationship with PM₁₀, with correlation coefficients (R^2) exceeding 0.9, and the regression equation is $PM_{2.5} = 0.66 \cdot PM_{10} + 3.1$ (Figure 3). In the study, PM_{2.5} and PM₁₀ are closely correlated and there is an interaction relationship between PM_{2.5} and PM₁₀. This evidence also indicates that higher percentage composition of PM_{2.5} fine particle in the PM₁₀ and both of them make a combined contribution in the air environmental pollution.

4. Conclusion

The concentration of PM_{2.5} and PM₁₀ in Guangzhou during the winter period was $(55.01 \pm 78.19) \mu\text{g}/\text{m}^3$ and $(78.34 \pm 101.76) \mu\text{g}/\text{m}^3$, respectively. The mean ratio of PM_{2.5} to PM₁₀ was 70.59% and the PM_{2.5}/PM₁₀ value ranges from 54.61% to 94.37%, indicating that fine particle (PM_{2.5}) occupy high proportion in the PM₁₀. Compared with PM₁₀, PM_{2.5} has higher percentage of exceeding standard number. In addition, PM_{2.5} has a good linear relationship with PM₁₀. From the above, all evidence provides us conclusions that fine particle has a higher proportion in the atmospheric aerosol, and make a major contribution in the air environmental pollution.

Therefore, government should aim to control fine particles for priority in the particulate matter control strategy.

Acknowledgments

The study was financially supported by the National Natural Science Foundation of China (41571187) for which the author is grateful. Thanks are giving to the Guangdong Environmental Monitoring Center for the sample data.

References

- [1] Feng, J., et al., Characteristics and seasonal variation of organic matter in PM_{2.5} at a regional background site of the Yangtze River Delta region, China, *Atmos, Environ*, 2015, 1-10.
- [2] Xu, L.L., et al., Seasonal variations and chemical compositions of PM_{2.5} aerosol in the urban area of Fuzhuo, China, *Atmos Res*. 2012, 104-105: 264-272.
- [3] Pope III, C.A., et al., Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *J. Am. Med. Assoc.* 2002, 287: 1132-1141.
- [4] Kok, T.M.C.M., et al., Toxicological assessment of ambient and traffic-related particulate matter: a review of recent studies, *Mutat. Res.* 2006, 613: 103-122.
- [5] Taus, N., et al., Respiratory exposure to air pollutants, *J. Environ. Prot and Ecol.* 2008, 9: 15-25.
- [6] Zhao, X.J., et al., Seasonal and diurnal variations of ambient PM_{2.5} concentration in urban and rural environments in Beijing. *Atmos. Environ.* 2009, 43: 2893-2900.
- [7] Huang K, et al., How to improve the air quality over mega-cities in China? Pollution characterization and source analysis in Shanghai before, during and after the 2010 World Expo. *Atmos. Chem. Phys.* 2013, 13: 3379-3418.
- [8] Shi W Z, et al., Analysis of airborne particulate matter (PM_{2.5}) over Hong Kong using remote sensing and GIS. *Sensor.* 2012, 12: 6825-6836.
- [9] Garland R M, et al., Aerosol optical properties in a rural environment near the mega-city Guangzhou, China: implication for regional air pollution radiates forcing and remote sensing. *Atmos. Chem. Phys.* 2008, 8: 5161-5186.
- [10] Wu, D., et al., An extremely low visibility event over the Guangzhou region: A case study, *Atmos. Environ.* 2005, 39: 6568-6577.
- [11] Deng, X.J., et al., Long-term trend of visibility and its characterizations in the Pearl River Delta (PRD) region, China, *Atmos. Environ*, 2008, 42:1424-1435.
- [12] Wei, F., et al., Ambient Concentrations and Elemental Compositions of PM₁₀ and PM_{2.5} in Four Chinese Cities, *Environ. Sci. Technol.* 1999, 33: 4188-4193.
- [13] Chan. L.Y., Spatial Variation of Mass Concentration of Roadside Suspended Particulate Matter in Metropolitan Hong Kong. *Atmos. Environ*, 2001, 35: 3167-3176.