

A Quasi-Unitary Pre-Quantum Theory of Particles and Fields and Some Theoretical Implications

Marius Arghirescu

State Office for Inventions and Trademarks, Bucharest, Romania

Email address:

arghirescu.marius@osim.ro

To cite this article:

Marius Arghirescu. A Quasi-Unitary Pre-Quantum Theory of Particles and Fields and Some Theoretical Implications. *International Journal of High Energy Physics*. Special Issue: Symmetries in Relativity, Quantum Theory, and Unified Theorie. Vol. 2, No. 4-1, 2015, pp. 80-103.

doi: 10.11648/j.ijhep.s.2015020401.17

Abstract: Through a phenomenological approach using the concept of sub-quantum fluid, the theory argues the possibility of a cold genesis of elementary particles and of fields, explaining the electro-magnetic and the gravitic fields by equations of ideal fluids applied to the subquantum and the quantum “primordial dark energy”. The possibility to explain the cold genesis of “dark” photons and of “dark” elementary particles is obtained by a CF -chiral soliton model of lepton, resulted as vortex of „primordial dark energy”, respective- as Bose-Einstein condensate of gammonic (e^+e^-)-pairs confined in a very strong magnetic field, in the Protouniverse’s period of time. This possibility results by a model of primordial ‘gravistar’ with a self-growing property given by the confining of “primordial dark energy” into “dark photons” and into “dark particles” by a “vortex cascade” mechanism induced by its magnetic field and gravitationally sustained. The supposed primordial “big bang” of the Universe results as a period of gravistars transforming into magnetars, supernovae and into (micro) quasars. The resulted model of expanding Universe gives a semi-sinusoidal variation of the expansion speed. The approach, even if does not propose an enough unitary equation of the known basic fields, it explains naturally the fundamental interactions, by the same basic concept.

Keywords: Unitary Theory, Gravistar, Chiral Soliton, Dark Energy, Dark Matter, CF-Particles, Bosons Confining, Expanding Universe

1. Introduction

In a recent book: “The Cold Genesis of Matter and Fields”, of the author [1], was argued the possibility to explain the fundamental fields of particles by the classic mechanics of the ideal fluids in a galileian relativity, by a simple or composite chiral soliton model of fermion considered as formed “at cold”, according to a fractalic „vortices cascade” process and the next structure of the subquantum and quantum medium:

(A_c) – sub-quantum medium; ($m_s \ll m_h = h/c^2$;
 $S_s^* \cong 0$; maximal speed: $v_s \approx \sqrt{2}c$):

-gravitons; (g-etherons): $m_g = (10^{-68} \div 10^{-72})$ kg, acting as gravitic field quanta and having contribution as etheronic wind to the genesis of gravitomagnetic quantum-vortices;

-sinergons; (s-etherons): $m_s = (10^{-59} \div 10^{-61})$ kg, acting as sinergonic quanta of vortices of gravitomagnetic chiral solitons but also as gravitonic quanta ;

(B_c) – quantum medium, $m_b \geq m_h = h/c^2$:

-quantons: $m_h = h \cdot 1/c^2 = 7.37 \times 10^{-51}$ Kg; $S_h^* \ll \frac{1}{2}\hbar$, acting as quanta of the B- magnetic field and forming the μ_p

-magnetic moment of fermion; similarly, the pseudomagnetic moment of quanton: μ_h , results as a sinergonic vortex formed around a quantonic superdense centrol having the mass: $m_{hc} = m_h$, the quanton being -in the theory, the smallest hard-core (pseudo)fermion;

-vectors (vectorial photons): $m_v = 3 \times 10^{10} m_h = 2.2 \times 10^{-40}$ kg; $S_v = S_v^* = \frac{1}{2}\hbar$; acting as electrostatic field quanta, resulted as hard-core semiphotons of the cosmic 3K-background radiation;

-vexons; $m_w \geq 10m_v$; $S_w = S_w^* = \frac{1}{2}\hbar$; structured as vectonic chiral soliton assembly, acting as constituents of quantum volume of the elementary particles and of luxons;

-pseudoscalar photons, (particularly-luxons): $m_f = n \cdot v \cdot m_h = 2n \cdot m_w$, $S_f = \hbar$; acting as electromagnetic radiation pseudoscalar quanta, formed by ‘n’ pairs of vectorial photons: $m_f = n \cdot (m_w - m_w)$.

The reason of the previous theoretical considerations was based on the fact that the masses of the stable/quasistable free photons or etherons are in the relation:

$$m_s^k \approx (K^v)^{-1} \cdot m_s^{k+1} ; \text{with: } K^v \in (10^{-9} \div 10^{-11}); \quad k \geq 1 \quad (1)$$

and these (quasi)stable free photons or etherons can be field quanta or pseudo-quanta or/and constituent quanta of the elementary particles with bigger mass.

According to the model, the electron results as a CF chiral soliton with the inertial me-mass with exponential density variation, $\rho_e = \rho_e^0 \cdot e^{-r/\lambda}$, given by the confining of paired and un-paired vortons around a electronic hard control, in a quantum volume of radius: $r_e = a = 1.41\text{fm}$, by the electron's magnetic moment vortex: $\Gamma_e = \Gamma_A + \Gamma_\mu$, of sinergons and quantons, the Γ_A -vortex generating the magnetic potential, A , and the Γ_μ -vortex generating the magnetic induction, $B = \text{rot}A$, by the impulse density gradient of the Γ_A -vortex.

The exponential form of the nuclear potential is theoretically re-found through a cold genesis nucleon model, formed as B-E condensate of paired quasi-electrons (degenerate electrons) and by an Eulerian expression, as being generated by the vortexial dynamic pressure created inside the nucleonic quantum volume.

According to the proposed Cold Genesis Theory (CGT), the vectorial photon and the electron results as vortexes of „primordial dark energy”, with confined inertial mass and an evanescent part, the elementary mesons and baryons resulting as Bose-Einstein condensate of gammonic (e^+e^-)-pairs, formed by vortexes of “primordial dark energy” in a very strong magnetic field, in the Protouniverse's period of time.

This possibility results by a model of primordial ‘gravistar’ with a stong vortex of “primordial dark energy”- considered as being composed by etherons and quantons, and with a self-growing property of its kernel formed “at cold”, from the vortexially formed leptons, which may be transformed into a mini-“black hole”, this gravistar model being similar to the hypothetical “gravastar” proposed by E. Mottola and P.O. Mazur [3], but relative different.

According to this fractalic mechanism of the particles genesis, it results that in the Protouniverse's period existed Majorana neutrins which -through theirs vortexial

confinement, generated massive neutrins (postulated as components of the Protouniverse also by the Dark matter Universe model) and micro- and mini-black-holes with growing mass and magnetic field.

The possibility of “dark particles” forming by the confinement of “primordial dark energy”, as “dark chiral solitons”, is sustained in concordance with some other theories [2].

For an unifying model of particles, is argued in CGT the conclusion of a vortexial sub-structure with a radius: $a \approx 1.41\text{fm}$ of the inertial mass volume, for all elementary particles excepting the photons and the electronic neutrino.

The considered proto-“dark energy” structure gives the possibility to explain the gravitational and the gravito-magnetic force and field by an unitary theory, based on the similitude between the gravitational force and the electric force, with a similar charge model.

For the weak and the strong interactions, even if the resulted models not results from a more general (unitary) equation of fields, as particular case, these models are based to the same fundamental model of composite fermion resulted in CGT.

2. An Unitary Theory of the Electric and Gravitic Fields

In accordance also with the charge model of quantum mechanics, according to a classical etheronic theory of fields, [1], the charge $Q(M)$ of a particle, results as being given by a spheric-symmetric distribution of charge's quanta: $n \cdot m_c(r_c)$ around the particle having the electron radius: $r_a = a$, (fig.1), i.e.: $\rho_a \cdot r^2 = \rho_a^0 \cdot a^2$, which generates a pressure force over the semi-surface $S^x = S^0/2 = 2\pi r_0^2$ of a pseudocharge $q_s(m_0)$ approximated as equal with its scattering cross section: $\sigma = \pi(r_0+r_c)^2 \approx S^x$, ($r_c \approx 0.41r_0$; $r_0 = a$), given by the impulse density variation : $\Delta p_c = p_c(r) - p_c(-r) = 2n \cdot m_c \cdot v_c$:

$$F(r) = S^x \cdot \frac{\Delta(p_c)_r}{\Delta t} = S^x \cdot \frac{\Delta(n \cdot m_c \cdot v_c)_r}{\Delta t} = S^0 \cdot \rho_v(r) \cdot v_c^2 = q_s \cdot E(r); \quad n \cdot m_c = n_0 \Delta r \cdot m_c = \rho_v \Delta r \quad (2)$$

By a gauge constant: $k_1 = S^0/e = 4\pi a^2/e$ gauged by the electron with elastic interaction of m_c , the electric E-field has the form:

$$E_c = k_1 \rho_c(r) \cdot v_c^2 = k_1 \rho_a^0 \frac{a^2}{r^2} \cdot v_c^2; \quad \rho_a^0 = \rho_c(a); \quad v_c = c; \quad k_1 = \frac{4\pi a^2}{e}; \quad \rho_c(r) = \rho_a^0 \frac{a^2}{r^2} \quad (3)$$

The magnetic B-field is generated when the (pseudo)charge: $q_s = S^0/k_1$ has a perpendicular $v_0 = v_p \cdot \cos\theta$ - speed relative to the E-field, (fig.2), according to the impulse density theorem

for ideal fluids derived from a Gauss- Ostrogranski relation, which gives the relation for the total electrodynamic force (including the Lorentz force), in the form [1]:

$$F_i = m_p a_i = \frac{S^0}{k_1} (k_1 \rho_c v_c^2 + k_1 \rho_c v_c v_0) n_i = q_s (E_i^0 + B_j \cdot v_0) = F_i^0 + F_i^l; \quad B_j = k_1 \rho_c v_c; \quad v_c \equiv c \quad (4)$$

the eq. (4) resulting by the impulse density: $p_i = \rho_c v_c$ included in the tensor Π_{ik} , that is:

$$F_i = m_p \cdot a_i = - \frac{d}{dt} \int_s \rho_c \cdot v_c \cdot d\tau = \int \Pi_{ik} \cdot dS_k \quad (5)$$

with: k_1 given by the equality: $\frac{1}{2}\epsilon_0 E^2(a) = \epsilon_{v_c} = \frac{1}{2}\rho_a^0 \cdot c^2 = \frac{1}{2}(k_1^{-1})E(a)$, ($\epsilon_{v(a)}$ -the quanta energy) and: Π_{ik} -the impulse density tensor:

$$\begin{aligned} \Pi_{ik} = P_c \cdot \delta_{ik} + \rho_c (v_i \cdot v_k) \quad ; \quad \text{with: } \delta_{ik} = (n_i n_k) = n_j; \quad |n_i| = |n_k| = 1; \quad dS_k = n_k dS \\ (n_i, n_k \text{-unit vectors}) \quad ; \quad P_c = \rho_c \cdot v_c^2 \quad ; \quad v_i = v_c \cdot n_i; \quad v_k = v_0 \cdot n_k \quad ; \end{aligned} \quad (6)$$

For $\Pi_{ik} = \text{constant}$ and $\int dS_k = S^0 \cdot n_k$, with: $S^0 = 4\pi a^2$, for an elastic interaction of the field quanta with the e-charge and with: $a = e^2/8\pi\epsilon_0 m_e c^2 \approx 1.41 \text{ fm}$, (i.e. the e-charge in surface), it results that: $k_1 = 1.57 \times 10^{-10} [\text{m}^2/\text{C}]_{\text{si}}$.

For the elementary electric charge 'e' of the electron, the charge sign depends on its intrinsic chirality: ζ_e . Also, a vortex $\Gamma_e = \Gamma_A + \Gamma_B$ of the A- potential and of the magnetic

induction $B = \text{rot} A$, generates the field lines of the induction B by the gradient of the impulse density: $\nabla_r p_A = dp_A/dr$, which induces ξ_B -vortex-tubes of the B -induction around the vectons of the electric E- field.

The Maxwell's electromagnetic field equations results in CGT according to eqs. (3)-(4), in a general vectorial form, of a vectorial E- or H- field intensity reciprocal generation :

$$\vec{B} = \frac{1}{c^2} \nabla_E \times \vec{E} \quad \vec{E}' = \nabla_0 \times \vec{B} = -\nabla_B \times \vec{B} \quad (7)$$

another specific field equation resulting also in a general way from the continuity equation:

$$\frac{\partial \rho_c}{\partial t} = -\nabla(\rho_c \cdot v_E); \quad \frac{1}{c^2} \cdot \frac{\partial(k_1 \rho_c c^2)}{\partial t} = -\nabla(k_1 \rho_c v_E); \Rightarrow \frac{1}{c^2} \frac{\partial E}{\partial t} = -\nabla \cdot B = -\text{div} B \quad (8)$$

For the electron, according to eq. (4), for $r \gg r_\mu = 3.86 \text{ fm}$ representing its Compton radius, the spinning of quantons in the Γ_B -vortex around the e-charge, is realized in conditions of quantum non-equilibrium, according to the vortexial kinetic

moment conservation law:

$\Gamma_B = 2\pi r \cdot v_{ct} = 2\pi r_\mu c = ct$, and $B(r)$ has the form found by the classic magnetism:

$$B(r) = k_1 \rho_v v_{ct}^r \cong k_1 \rho_a^0 \frac{a^2}{r^2} \cdot \frac{r_\mu c}{r} = k_1 \rho_B c = \frac{\mu_0 \cdot \mu_e}{2\pi r^3}; \quad \rho_a^0 = \frac{\mu_0}{k_1^2}; \quad \rho_B = \frac{v_v^r}{c} \rho_v; \quad r > r_\mu; \quad (9)$$

(ρ_B –the density of ξ_B -vortex-tubes), the magnetic potential resulting in the form:

$$A_k(r) = \frac{B_j(r) \cdot r}{2} = \frac{k_1 r_\mu c}{2} \rho_a^0 \frac{a^2}{r^2} = \frac{k_1 \cdot \Gamma_A(r_\mu)}{4\pi} \rho_s(r); \quad r \geq r_\mu; \quad \rho_s(r) = \rho_a^0 \frac{a^2}{r^2}; \quad \Gamma_A(r_\mu) = 2\pi r_\mu c \quad (10)$$

Also, the Lorentz force results of Magnus type - according also to other theories [4], considering a pseudo-cylinder (barrel like) form of the electron with the high $l_e = 2a$ and a

relative impulse density of the E-field vectons: $p_v = \rho_e v_v^r$, generating the B-field according to eq. (4):

$$F_L = 2a \cdot \Gamma_a^* \cdot \rho_B \cdot v_e = q \cdot B \cdot v_e = e \zeta_e \cdot k_1 (\rho_e v_v^r)_r \cdot v_e; \quad \Gamma_a^* = 2\pi \cdot a \cdot c \cdot \zeta_e; \quad \rho_B = \rho_e(r) \cdot [v_v^r / c] \quad (11)$$

The m_p -particle being formed by n_p quantons having the m_h -mass, the eq. (4) is generalisable for the gravito-dynamic force and field, by the relation: $S_g^0 = n_p \cdot \frac{1}{2} S_h = (m_q/m_h) \cdot \frac{1}{2} S_h$, ($S_h = 4\pi r_h^2$; r_h -the quanton radius), resulted from its penetrability to the g- and s-etherons action and an elastic

interaction of the etherons with the S_g^0 -surface.

For the attracted m_p -mass and for the gravitic field of an attractive mass M of a particle or of a body, it may be assigned an "electrogravitic" pseudo-charge, q_G , respective- by eq. (4), -also an "electrogravitic" field, $E_G(r, Q_G)$, i.e.:

$$\begin{aligned} F_i^g = m_p a_{Gi} = q_G \cdot E_G(r, Q_G) = -k_h \cdot m_p (\rho_g v_g^2 + \rho_g \langle v_g \cdot v_0 \rangle) \cdot n_i; \quad k_h = S_h / 2m_h \\ \text{with: } q_G = S_g^0 / k_1; \quad E_G = \pm k_1 \rho_g c^2; \quad S_g^0 = k_h m_p \end{aligned} \quad (12a, b)$$

In the expression (12b) of the electrogravitic field intensity, E_G , the meaning of the sign: \pm is that the electrogravitic Q_G -charge generating the E_G -field is given by an uniform spheric distribution of an etheronic flux with a non-compensated component, i.e. -by the difference between the received etheronic flux and the etheronic flux reflected by the super-dense centrols of the inertial M-mass structure, in the case of an attractive, gravitic M-charge.

Therefore, considering this non-compensated etheronic component as a gravitonic field flux, having the impulse density $p_g(r) \uparrow \downarrow r$, the generation of the gravitation force, F_N ,

complies with the Lesage's hypothesis [5] which presumes the screening of the m_p -mass by the M-mass in report with the cosmic etheronic winds that comes radial-symmetrically towards the M-mass, (fig.2).

The etheronic flux formed by a M-mass with disturbed sinergonic vortex which emits s-etherons, gives an antigravitic pseudocharge, generating a positive, i.e. repulsive E_G -field.

For the variation of $\rho_g(r)$ -density of the gravitonic wind, in compliance with eq. (12) of the electrogravitic $q_G(M)$ -charge of the M-mass having the radius r_0 and for $v_g = c$, the gravitic force results from eq. (12) as having the form:

$$F_i^g = -k_h m_p \rho_g c^2 \left(1 + \frac{v_0}{c}\right) n_i = -G \frac{m_p M}{r^2} \left(1 + \frac{v_0}{c}\right) n_i; \quad \rho_g(r) = \rho_g^0 \frac{r_0^2}{r^2} \approx \frac{M}{m_h} \rho_g^h \frac{r_h^2}{r^2} \quad (13)$$

where: ρ_g^0 and ρ_g^h are the density of the gravitonic flux (i.e.-of the uncompensated etheronic wind) at the $M(r_0)$ -mass surface and- respectively- at the $m_h(r_h)$ -quanton surface.

If the m_p -mass represents a photon having the speed $v_0 = c$, the value of the F_i^g -force, acting as a gravitic type force, results from the equation (13) as being: $F^g(r,c) = 2 F^g(r,0)$, -of a double value comparing to Newtonian static gravitational force, in accordance with the Einstein's theory of relativity and the astrophysical observations.

A form with lorentzian type term of the total gravitation

$$G = \frac{k_h \rho_g^0 r_0^2 c^2}{M} = \frac{k_h \rho_g^h r_h^2 c^2}{m_h} = \frac{2\pi \rho_g^h r_h^4 c^2}{m_h^2} = 6,67 \times 10^{-11} \frac{N \cdot m^2}{kg^2} \quad (14)$$

The value of the density ρ_g^0 of the uncompensated etheronic wind on a black-hole surface -for example, characterizes only the local (not also the intergalactic) etheronic density: ρ_e^0 , because that it results by the speed's statistic distribution of the etherons emitted by the solitonic quantum-vortices of the elementary particles, i.e. it is proportional with the mass

force F_i^g , is obtained also in the tensorial theory of gravitation for a weak gravitational field, giving as solutions the gravitational analogs to Maxwell's equations for electromagnetism, (Lano, Fedosin, M.Agop, N.I.Pallas, [6], [7]), the increasing of F_i^g with the v-speed, being equivalent with an transversal relativistic effect of the gravitational mass growth: $F_v = g_g \cdot m_p(1+\beta) = g_g \cdot m_p^v$, ($\beta = v_0/c$).

The eq. (13) gives for the G-gravitation constant, the expression:

$$F_i^g = q_G \cdot E_G(r, Q_G) = -\frac{m_p}{m_e} e \cdot k_1 \rho_g c^2 = -k_h m_p \rho_g c^2 \Leftrightarrow \rho_g^0 \approx k_p \rho_a^0; \quad q_G = \frac{m_p}{m_e} e; \Rightarrow k_h = \frac{4\pi a^2}{m_e} \quad (15)$$

In this case, for an unitary form of the electric and electrogravitic fields, we may obtain a plausible gauge value of k_h and of ρ_g^h , considering for the electron case the gauge condition: $q_{Ge} \approx e$, which complies with the expression obtained by M. Agop [7], starting from the acceleration of an electron in the field of another:

$$a_i^e = \frac{F_N^e}{m_e} + \frac{F_g^e}{m_e} = a_{Gi}^e + \left(\frac{e}{m_e}\right) \cdot \frac{e}{4\pi \epsilon \cdot r^2} = \left(\frac{e}{m_e}\right) \cdot (E_G^e(r) + E_e^e(r)); \quad E_G^e = \left(\frac{m_e}{e}\right) \cdot a_{Gi}^e \quad (16)$$

which gives by (15) the gauge constants: $k_h = 27.4 [m^2/kg]$, $r_h = 1.79 \times 10^{-25} m$; $\rho_h = \rho_c^M = 3 \times 10^{23} kg/m^3$, $\rho_g^h = 6.2 \times 10^{-30} kg/m^3$ and respectively, by eq.(14): $\rho_g^0 = 1.24 \times 10^{-29} kg/m^3$. Also, by eq.(12), it results that: $Q_G = 4\pi \epsilon_0 GM \cdot (m_e/e)$.

If the g- and s- etheron has the same ρ_c^M density as the quanton, it results also the size order of the graviton's and of the sinergon's radius: $r_g \approx 10^{-31} m$; $r_s \approx 10^{-28} m$ and the ratio: $r_s/r_g \approx r_h/r_s \approx 10^3$.




Fig. 1. Static type Charge.




Fig. 2. The Gravitational Interaction.

By the galileian relativity, it is possible also to avoids the paradoxes of the physical interpretation of Einstein's special relativity equations, such as the so-called "the twins paradox",

by the concept of cosmic ether, by a physical re-interpretation of the Einstein's relativistic equations such as the equation of speed- dependent mass variation, considering the

sub-quantum medium as an ideal fluid with a ρ_s mean density, in which- according to a specific equation for ideal fluids, the acceleration a_p of the m_0 -particle “falling” is dependent on the

“falling” v_p - speed because the resistance force of the sub-quantum fluid: $F(r,v) = \frac{1}{2}S_g^0 \rho_s v^2$, ($S_g^0 = k_h m_p$), in the form:

$$a_{ps} = a_0 \left(1 - \frac{v_p^2}{w^2}\right); \quad a_p = \frac{F_{(r,v_p)}}{m_p}; \quad a_0 = \frac{F_{(r,0)}}{m_p}; \quad F_{(r,0)} = S_g^0 \rho_s c^2; \quad w = \sqrt{2} \cdot c \quad (17)$$

If we consider formally a constancy of the accelerating force $F(r,v_p)$, the eq. (17) is mathematically equivalent to a longitudinal relativist effect of the inertial mass variation:

$$m_p^*(v_p) = m_p^0 / [1 - v_p^2/w^2] = m_0/\beta'; \quad \text{with: } w = \sqrt{2} \cdot c \quad (18)$$

According to (18), the etherons may be also tachyonic quanta, with a maximal speed: $w = \sqrt{2}c$.

The expression (15) for the gravito-electric field corresponds also to the Schiff-Barnhill effect which states that in the presence of a gravitational field and in stationary conditions, there is a small electric field generated in a conductor or superconductor: $E = -(m/e) \cdot a_G$.

Combined with the result of eq. (13), the relativist expression of the gravitation force results similar to those of I. Somacescu's theory, [4]:

$$F_i^g(r;v) = F_i^g(r;0) \cdot \frac{1 + v_0/c}{1 - v_0^2/2c^2}; \quad v_0 = v_p \cos \alpha \perp v_r$$

3. The Elementary Particles as Chiral Solitons

3.1. The Photon

For explain the wave-particle dualism of the light photon, H. A. Múnera considered a semi-classic model of vortical particle-antiparticle pair, which explains the frequency and the repose mass of the photon, [8], by preonic vortexes of quasi-constant density, formed in the quantum vacuum.

Geoffrey Hunter and L. P. Wadlinger [9] proposed a solitonic model of photon corresponding to the Einstein's concept, considered as a confined electromagnetic wave in a volume of an ellipsoid with the length along the propagation axe- equal to the wave- length, λ , and with the photon diameter: $d_f = \lambda/\pi$.

It may be observed that if the Múnera's model of photons is dimensioned like the Hunter- Wadlinger model, considering the simple photon as a doublet of two vectorial photons with mutually anti-parallel spins $S = \hbar/2$ and a mean diameter: $d_w = \lambda/\pi$ and considering the hard-gamma quanta as a doublet: negatron-positron, $\gamma_c = (e^+e^-)$, with opposed spins and an energy: $\epsilon_\gamma = h\nu = 2m_e c^2$, it results that the electron may be assimilated with a vectorial (semi) photon, m_w^e , with a Compton radius: $r_\lambda = \lambda/2\pi$, ($\lambda = h/m_e c$).

Because that according to CGT, the magnetic field B is given by a vortex of sinergons and of quantons which has the rotation speed: $\omega \cdot r = c$ inside the pseudo-cylindrical volume of radius

$r_\lambda : v_\lambda \approx 2\pi r_\lambda^3 = \lambda^3/4\pi^2$, from the equality between the vortical energy: $w_\mu^f = \frac{1}{2}\mu_0 H^2 \sim \frac{1}{2}m_s(\omega_h \cdot r)^2$ and $w_e^f = \frac{1}{2}\epsilon_0 E^2 \sim \frac{1}{2}m_s c^2$ - given by the translation energy of a spinorial Γ_s vortex of quantons which do not contribute to the inertial mass $m_{v(w)}$ of the photon core, it results for the proposed revised model of photon, the equation:

$$E_E^c = E_\mu^c \Rightarrow \frac{\sum m_h \cdot c^2}{2} = \frac{m'_s (\omega \cdot r)^2}{2} = \frac{h\nu}{4} = \frac{m_w c^2}{2}; \quad \omega \cdot r = c \quad (19)$$

In this case, it results that:

$$E_v = 2(E_E + E_\mu) = 2m_s c^2 = h\nu = 2m_w c^2,$$

by the conclusion that the spinorial (vortical) m_s - mass of the vorton is equal with its inertial mass: m_w (considered in CGT as cluster of vectons with super-dense kernel, contained in a

volume $v_f(r_f \leq a)$) which explains its gravitational charge, the photoelectric effect and the atomic electrons transitions.

This equality is argued in CGT by the fact that- for a vectorial photon, by a ρ_w density variation with r^{-1} -for a pseudo-cylindrical v_λ -volume or with r^{-2} -for a spherical v_λ -volume, the vectorial photon spin results of value:

$$S_v^* = \int_{r_a}^{r_v} r \cdot c \cdot dm \cong 4\pi r_a^2 \rho(r_a) \cdot c \cdot \frac{r_v^2}{2} \cong m_s \cdot c \cdot \frac{r_v}{2} = m_s c \cdot \frac{\lambda}{4\pi} = m_s \cdot c \cdot \frac{h}{4\pi \cdot m_v c} = \frac{1}{2} \frac{h}{2\pi} \quad (20)$$

- equal with the value considered for fermions by the quantum mechanics.

The explaining of the equality: $m_s = m_v$, may be given by the chiral sub-solitons forming condition [10] which specifies that the energy $E_\Gamma = m_\Gamma c^2 = 2E_\mu$ of the mass-generating chiral soliton field, given- in this case by the sinergono-quantonic vortex $\Gamma_\mu = \Gamma_\Lambda + \Gamma_H = 2\pi \cdot c$, should be double, at least,

comparing to the mass energy : $E_w = m_w c^2$ of the generated sub-solitons; ($E = 2E_w$).

This condition imply also the conclusion that the particles mass may increase with the speed only by a $\Gamma_\mu(v)$ -vortex increasing but cannot exceed the value $2m_0$, (with m_0 -the particle rest mass).

The fact that the photon length is -normally: $l_v = \lambda > 2d_w =$

$2\lambda/\pi$, may be explained by the conclusion that the vectorial photon volume v_λ is a little expanded over the value d_w to at least one direction.

Because that some experiments [11] confirmed the value: λ/π for the diameter of the photon section, we take into account the vexon's volume expansion along the axe $x // v_p = c$; this possibility suppose that the vortexial energy E_μ of vectorial photon is contained by an ellipsoidic volume with the length: $l_w = \lambda/2$, i.e.: $v_\lambda^c \approx \pi(\lambda/\pi)(\lambda/2\pi)(\lambda/4) \approx \lambda^3/8\pi$, the inertial m_w –mass of vexon being closer to the focal point of the ellipsoid which is the first in contact with the pseudo-stationary (brownian) etherons and quantons of the quantum vacuum, of total density: $\rho_r = \rho_{rc} + \rho_{rs}$, which acts as a relativist wind, deforming the vexonic quantum volume along $x // v_p$, by the sinergonic component: $P_s = \rho_{rs} v_p^2$, until a value: $l_w = \lambda$, (fig.3), i.e.:

$$p_r = \rho_r v_p \approx \text{const.} \Rightarrow l_w = (2r_\lambda + \Delta l_w) \sim v_p, \quad (21a)$$

relation which results in the hypothesis of a relativist etherono-quantonic impulse density: $p_r = \rho_r v_p$ of quasi-constant value.

By eq. (21a), the revised model of photon may explain also the photon wavelength variation with the photon's speed in an optical medium with a specific index of refraction, n , i.e.: $v_w = c/n \Rightarrow \lambda = \lambda_0/n$; ($v = 2m_w/m_h = \text{constant}$), in the form:

$$\Delta v = (c - v_p) \sim (\rho_r - \rho_r^0); l_w / l_{wc} = v_p / c; \quad (21b)$$

$$(l_{wc} = \lambda_0(c); l_w = \lambda(v_p); v_p \approx v_w)$$

in which ρ_r^0 is the initial density of the etherono-quantonic medium in the quantum vacuum.

Also, because that- in the case of light propagation in an optical medium in the presence of an intense electric E-field

$$v_p \approx \frac{K_n h}{2kaK_s v_r c} \cdot \frac{1}{\rho_r \lambda_0} \approx K_p \frac{1}{\rho_r \lambda_0}; \quad n_1 = \frac{c}{v_1} = \frac{c \cdot \rho_{r1} \lambda_0}{K_p}; \quad n_2 = \frac{c}{v_2} = \frac{c \cdot \rho_{r2} \lambda_0}{K_p} = \frac{c \cdot (\rho_{r1} + \Delta \rho_r) \lambda_0}{K_p} = n_1 + \Delta n \quad (23)$$

If: $\Delta \rho_r \approx k_E \cdot \epsilon_0 E^2 / c^2$ and: $\Delta \rho_r \approx k_H \cdot \mu_0 H^2 / c^2$, ($k_E \sim \epsilon_r$; $k_H \sim \mu_r$ –proportionality constants depending on the optical medium and on the relative permittivity/ permeability), it results that:

$$\Delta n_E \approx \frac{k_E \epsilon_0 E^2}{c \cdot K_p} \lambda_0 = \lambda_0 K_E E^2; \quad (K_E = \frac{k_E \epsilon_0}{c \cdot K_p}); \quad \Delta n_H \approx \frac{k_H \mu_0 H^2}{c \cdot K_p} \lambda_0 = \lambda_0 K_H H^2; \quad (K_H = \frac{k_H \mu_0}{c \cdot K_p}) \quad (24)$$

where K_E and K_H are the Kerr and the Cotton-Mouton constants of the optical medium. So, we obtain a microphysical explanation for the Kerr and the Cotton-Mouton effects.

The decomposing of a monochromatic polarised light beam into two waves: ordinary and extraordinary, in a strong magnetic field, may be also explained by the revised model of photon, considering also the difference of the magnetic interaction with the external B-field of the two vectorial photons which gives the light photon.

Also, in the inverse Cotton-Mouton effect, when an oscillating field passes through a dielectric medium in such a way that it's oscillations are similar to the polarization induced by the C-M effect, then a static magnetic field is created

or magnetic B-field the etherono-quantonic density ρ_r of the quantum vacuum is supplemented with a value: $\Delta \rho_r \sim \epsilon_0 E^2$ or




Fig.3. Revised Model of Multi-photon, ($l_p = 1,5 \lambda$).

$\Delta \rho_r \sim \mu_0 H^2$ –according to CGT, it is possible to explain by the model also the Kerr effect [12] and the Cotton-Mouton effect [13], of the refractive index changement in an applied E-field or H- field or by the intensity I of the light beam, (the AC Kerr effect), by the hypothesis that the photon speed: $v_p \leq c$ results by a quantum Stokes force: $F_s = K_s \cdot \eta \cdot d \cdot v_p$, ($\eta = \rho_r v_r$; v_r –kinematic viscosity of quantum vacuum; ρ_r – the etherono-quantonic density), acting over the m_p inertial mass of photon and generating a quasi-constant mean deceleration- according to the eqn. (21a), i.e.:

$$a_e = k_n P_e = \frac{F_s}{m_p} = \frac{K_s \rho_r v_r d \cdot v_p}{h} \cdot c \lambda_0 \approx K_n \quad (22)$$

Because that in CGT is considered a radius: $r_w = k \cdot a$ with $k \approx (0.4 \div 1)$ for the inertial mass of photons, ($k \approx 0.7$ for light photons), it results that: $d = 2k \cdot a$, ($a = 1.41 \text{ fm}$), resulting for v_p and n that:

perpendicular to the direction of propagation of light. This effect is also explained by the fact that the $\Gamma\mu$ -vortexes of light quantum vexons induces parallelly oriented vortexes of a H-field in the quantum medium of the dielectric body.

By the model, the photon may be considered un-deformed until a speed:

$$v_p^c = 2d_p c / \lambda_0 = 2\lambda_0 \cdot c / \pi \lambda_0 = (2/\pi)c \approx 0.64c .$$

The refraction angle may be also explained by a model of double multiphoton.

If the dielectric body has a speed $v_D // v_p^*$ with $v_p^* \neq v_p$, if $K_n^*(v_p^*) \approx K_n(v_p)$, it results by eq. (22) that: $v_p^r = v_D + v_p^* = v_p$, (i.e.- the invariance of v_p^r - speed of light).

The possibility: $K_n^*(v_p^*) \approx K_n(v_p)$ corresponds to the conclusion that the deceleration a_e is equilibrated –inside the dielectric body, by an etheronic wind pressure P_e given by atomic vortexes, which may explain the fact that- when the photon get out from the optical body, it regains its normal speed, c .

3.2. The Electron

According also to CGT, the electron may be considered –in a degenerate form, a semiphoton of a hard-gamma quantum, considered as a pair: $e^{+*}e^{-*}$, (degenerate negatron- degenerate positron), which may be splitted in an electron-positron pair in the field of a nucleus.

Because that- according to some other opinions, the charge variation inside the quantum volume of the electron's inertial mass, m_e , corresponds to the density variation, in a electron model of confined vexons distributed inside the electron according to a Boltzmann type statistic distribution: $\rho_e(r) = \rho_e^0 \cdot |\psi(r)|^2$ (that characterizes also the mixtures of bosons and fermions), by similitude with the structure of proton, in CGT it results the following substructure of the electron, (figure 4, [1]) :

-an “impenetrable” supersaturated quantum volume having the radius $a_i = 0.5 \div 0.6 \text{ fm}$, composed of vexonic layers-in even number for positrons and odd number for negatrons, with paired and magnetically coupled vexons to the radial and the meridian direction;

The q_w^* -pseudo-charge of vexons of the last layer of “impenetrable” quantum volume, attracts light vexons with oppsed q_w^* pseudo-charge.

-a charge's volume which is also of strong interaction quantum volume with the thickness: $\Delta a = a - a_i$, ($a=1.41 \text{ fm}$), formed by un-paired light vexons: m_w^* , attracted by the last layer of the “impenetrable” quantum volume and polarized with the pseudo-magnetic moments: μ_w on the meridian direction by the magnetic lines: ξ_B of the electron magnetic moment, μ_e , (ξ_B vortex-tubes).

The pseudo-charge q_w^* of the polarised vexons from the strong interaction quantum volume of the electron gives the electron's charge: $e = \Sigma(q_w^*)$.

- According to the model, the attractive or repulsive interaction is carried through vectorial quanta (“vectons”) of the electron e-charge, obtained from the bosonic pairs of the

background radiation (from the quantum vacuum), attracted by the Γ_e vortex of the μ_e -magnetic moment and “splitted” by the m_w^* -vexons of the e-charge, the m_v -vectons having the same q^* -pseudo-charge as the m_w^* -vexons of the electron's charge being rejected with an oriented spin, forming the E-field, the remained antivectons being absorbed by the m_w^* -vexons and used for the regeneration of the Γ_e -vortex .




Fig. 4. Model of Chiral Soliton Electron.

According to the model, the polarization rate of m_w^* -vexons giving the electron e-charge, and implicitly- the value of the vectonic flux: $\Phi_v(E)$, are proportional to the impulse density of the Γ_e -vortex in the strong interaction quantum volume of the electron, by the dependence:

$$e \sim \mu_e(\Gamma_e) \sim \rho_\mu(r) \cdot c^2 ; (r_i < r \leq a), \quad (25)$$

In accordance with the experiments [14], it results also the existence of a super-dense electronic centroid (control) having the density $\rho^m \geq 10^{19} \text{ kg/m}^3$ and the radius: $r_0 = 10^{-18} \text{ m}$, so –which is a very penetrant particle, being in the model a half of electronic neutrino, with the mass: $m_0 \approx 2 \times 10^{-4} m_e = 1.82 \times 10^{-34} \text{ kg}$ - in accordance with an experimental result [15] for the superior limit of the neutrino rest mass.

In the model, the electron's control is characterized also by an intrinsic chirality: $\zeta_e = \pm 1$, corresponding to a hypothetical helical form which determines the sense of the induced soliton vortex, Γ_e .

The electron spin results according to eq.(20) by a spinorial mass: $m_s = m_e$ given by a vortex $\Gamma_s(v_\lambda)$ of light photons.

The mass of the electron's quantum volume: $v_e(a)$, results according to the eqn:

$$m_e^v = \int_0^a 4\pi r^2 \rho_e(r) \cdot dr = 9,109 \times 10^{-31} \text{ kg} ; \quad \rho_e(r) = \rho_e^0 \cdot e^{-\frac{r}{\eta_e}} = \rho_e^0 \cdot |\Psi_e|^2 ; \quad \rho_e(a) = \mu_0 / k_1^2 = 5.17 \times 10^{13} \text{ kg / m}^3 \quad (26)$$

resulting a value: $\eta_e \equiv 0.965 \times 10^{-15} \text{ m}$, that is relative close to the value: $\eta_p = 0.895 \text{ fm}$ of the root-mean-square radius of proton charge distribution, experimentally deduced by Ingo Sick [16] and to the isoscalar magnetic mean radius: $r_m = 0.92 \text{ fm}$, given with the Skyrmion soliton model of proton, [17].

From eq. (26) it results also:

$$\rho_e^0 = 22,24 \times 10^{13} \text{ kg/m}^3.$$

In conditions of quantum equilibrium: $\varepsilon_h(r) = \gamma(k_B / \hbar) \cdot S_h(r)$, ($S_h(r) = 2\pi r \cdot m_h c$),

(ε_h – the negentropy per quanton; $\gamma = 64$, [1]), the sub-solitons forming condition [10] applied to the electron's volume, v_e , is respected for an identical variation of the densities: $\rho_e(e)$, $\rho_\mu(\Gamma_\mu)$ and $\rho_s(\Gamma_\lambda)$ -of the inertial mass and of the magnetic moment's quantono-sinergonic vortex, i.e.:

$$\rho_s(r) = \rho_\mu(r) = \rho_e(r) = \rho_r(r)/2, \quad (27)$$

$$(\rho_r(r) = \rho(\Gamma_r^c) = \rho_s(r) + \rho_\mu(r)) .$$

In accordance with other soliton models of electron [19], the stability equation of the Γ_e soliton vortex may be expressed by the Schrödinger nonlinear equation (NLS) with soliton-like solutions, identifying in this equation the term: $k_n \cdot |\Psi|^2$, (k_n -the nonlinearity constant), with the strong self-potential: $V_p(r)$ of

$$i\hbar \frac{\partial \Psi}{\partial t} = \hat{H} \Psi = E_{cf} + V_p = -\frac{\hbar^2}{2\delta m_e} \frac{\partial^2 \Psi}{\partial x^2} - k_n \cdot |\Psi|^2 \Psi = 0 ; \quad \Psi = R \cdot e^{i\frac{S_0}{\hbar}} ; \quad -\frac{1}{2} \delta v_e \cdot \rho_\mu(r) \cdot c^2 = V_p(r) \quad (28)$$

with: $k_n = -V_p^0(o)$; $S_\mu = (\delta m_e)_r \cdot c \cdot x$.

The form (28) of the fermion strong self-potential corresponds to an eulerian attractive force of quantum dynamic pressure gradient: $f_p = \nabla_r V_p = -\delta v_e \cdot \nabla_r P_d$, generated by a pseudostationary quantonic medium accumulated by the Γ_A sinergonic vortex, having the same density variation and the relativistic c -speed in report with $(\delta m_e)_x$.

The same force f_p is generated also by the Γ_μ -vortex acting upon a (pseudo)stationary mass having a volume: δv_e .

Because the solitonic nature of vexons, by the eq. (27) it results that the quantum intrinsic energy of electron, which is liberate at electron-positron annihilation, is given by the intrinsic vortexial energy of vexons-induced by Γ_e -vortex and the kinetic energy of the electron's magnetic moment:

$$E_w = \frac{1}{2} \sum_e m_w c^2 + \frac{1}{2} \sum_\mu m_c (\omega r)^2 = m_e c^2 \quad (29)$$

in accordance with the quantum mechanics.

By CGT, the previous electron model results as an unifying model of fermion, the mesons and the elementary baryons

$$F_G = \nabla Q_G = (-1/\eta) \cdot Q_G = (1/\eta) \cdot h/2 = m_\mu c^2 / r = h/r_i ; \Rightarrow r_i = 2\eta \approx 1.92 \text{ fm} > a \quad (31)$$

For the electron cold genesis, the eq. (27) resulted from the chiral sub-solitons forming condition [10], impose- by the relation: $2\pi a^3 \cdot \rho_a^0 = m_e$, (specific also to a proto-electron), the condition: $\rho_s^* \rightarrow \rho_a^0 = 5.17 \times 10^{13} \text{ kg/m}^3$, resulting: $\mu_c \rightarrow 1.36 \times 10^{-46} \text{ A} \cdot \text{m}^2$; $B_S \rightarrow 2 \times 10^{12} \text{ T}$.

A consequence of the chiral soliton model of charged particle is the conclusion that- at the fermions vibration under energy shocks, the vexons from the particle's quantum volume are easier destroyed and their vortexial structure is disturbed, decreasing the elastic character given by the photon interaction with vexons of the e -charge's surface; in this case,

$$n \cdot \varepsilon_v + m_p c^2 \rightarrow (E_v) \rightarrow m_p^* c^2 + \varepsilon_w ; n \cdot \varepsilon_v \equiv \varepsilon_w ; E_v \geq E_v^0 = \varepsilon_w / K_v \quad (32)$$

where ε_v ; ε_w -represents the energy of the captured photons and, respectively, of the emitted scalar quanta and K_v is a constant which can be also of over-unity value, in accordance with the energy conservation law, the eq. (32) explaining in this case the results of some experiments, such as the kinetobaric effect [20], the experiment of photons-electron interaction, (Stanford, 1997, [21]), of γ -rays emission by interaction with green laser pulse and the observations of γ -rays emission generated by thunderstorm, (italian group, 2000, [22]).

Another consequence of the chiral soliton model results by

the particle, generated by its Γ_μ -vortex of quantum volume, by the condition: $i\hbar \cdot (\partial \psi / \partial t) = 0$ (without contraction or expansion), which express the equality between $V_p(r)$ and the centrifugal potential: $E_{cf}(r)$ of an infinitesimal vortex volume: $\delta v_e = (\delta m_e / \rho_\mu)_r$, i.e:

resulting as composed by cold formed quarks, resulted as clusters of degenerate electrons, i.e: of quasi-electrons, with degenerate mass, charge and magnetic moment.

It results that -in the fermion's genesis process, in a very strong magnetic field, (as those of a magnetar), at quantum equilibrium, when: $\Gamma_c = 2\pi m_e c$, the genesis Q_G - quantum potential resulted by the formation of the sinergonic Γ_A -vortex and acting over a quanton moved to the Γ_μ -vortex line: $l_r = 2\pi r$ ($r \leq a$), compensates the quanton's centrifugal potential: $Q_G = -E_{cf} = -p_c^2 / 2m_e$, the nature of the Q_G potential being a magneto-gravitic genesis field, acting by a pseudomagnetic (sinergonic) B_S - induction in report with the μ_c -pseudomagnetic moment of the quanton. According also to the eq. (4) of the B-induction, it results that:

$$Q_G = -\mu_c \cdot B_S(r) = -\mu_c \cdot k_l \cdot \rho_s^* \cdot c = -p_c^2 / 2m_e = -h/2 = -E_{cf} \quad (30)$$

The dynamic equilibrium of forces, for $\rho_s = \rho_s^0 e^{-r/\eta}$; $\eta \approx 0.96 \text{ fm}$ -in accordance with eq. (26), gives:

photons which in the unperturbed state are reflected, can penetrate the quantum volume.

Through the Γ_μ quantum vortex of the particle, these photons are converted, inside the impenetrable quantum volume, into vexons with bigger mass which may be emitted as stable-bounded vexon-antivexon bosonic double pairs: $2(m_w - \bar{m}_w)$, having a null prequantum spin, (i.e- a scalar radiation quantum), under the action of the soliton quantum-vortex of the μ magnetic moment, according to equation:

the fact that the dependence: $\varepsilon_h(r) \sim S_h(r) \sim r$, for the electron, shows that the vortexiality gives negentropy, i.e.- an energy which may be converted as in the case of some functional „free energy” devices using long life magnets, (magnetic motors, etc.).

3.3. The Proton Model and the Nuclear Force

In CGT, the proton is a (N^p)-quasi-electrons protonic cluster, with quantonic Γ_μ^* -vortices of paired degenerate electrons, induced by the sinergonic $\Gamma_A^* = \Gamma_A^e$ vortices around each electronic control, with the degenerated density variation of

the Γ_{μ}^* quantonic vortex of μ_c^* -magnetic moment, corresponding by eqs.(25)-(27), to a degenerate charge: $e^* = {}^2/3e$, (characterizing the quark's charge), i.e., to:

$\rho_{\mu}(a) = {}^2/3\rho_e(a) = 3.44 \times 10^{13} \text{ kg/m}^3$, which corresponds by eq.(26), to: $\eta^* = 0.755 \text{ fm}$, the degenerate quasidelectron mass resulting by eq. (27): $m_c^* = 0.8722 \cdot m_e = f_d \cdot m_e$, (f_d –mass degeneration coefficient).

The proton's e^+ -charge results in the model by an attached positron, placed axially (at $r^+ > \eta^*$) in the strong interaction quantum volume.

In this case, the neutral proton cluster is formed by : $N^p = 1835.1/f_d \cong 2104$ paired quasidelectrons, according to the model, i.e. by degenerate gammonic pairs ($e^+ - e^*$), confined „at cold”, in a very strong magnetic field and forming a Bose-Einstein condensate of gammons, according to CGT.

$$\mu_p = k_p \frac{m_e}{m_p} \mu_e = k_p \frac{\bar{\rho}_e}{\bar{\rho}_n} \mu_e = k_p \frac{1}{f_d \cdot N^p} \mu_{Bp} = \frac{e \cdot c \cdot r_{\mu}^p}{2}; \quad (33a)$$

$$k_p = \frac{g_p}{g_e} = 2.79 = \frac{\rho_n(r^+)}{\rho_n^0} = e^{-\frac{r^+}{\eta_d}} \quad (33b)$$

in which: k_p -the gyromagnetic ratio; $\bar{\rho}_e$; $\bar{\rho}_n$ – the mean density of electron and nucleon;

r^+ -the position of protonic positron control in report with the proton centre.

The interpretation given by eqs. (33) of the particle mass-depending magnetic moment variation, explains also the fact that- when the proton is transformed into neutron, the emitted positron regains the μ_e magnetic moment value of free state, by the negentropy of the quantum and the subquantum medium, given by quantonic and etheronic winds.

The virtual radius of the proton magnetic moment: $r_{\mu}^n = 0.59 \text{ fm}$ - resulted from eq. (33a), may be considered approximately equal to the radius of the impenetrable nucleon volume, of value: $r_{\mu}^n \cong r_i(m_{ni}) \cong 0.6 \text{ fm}$ - used in the Jastrow expression of the nuclear potential, [23], by the conclusion that the impenetrable nucleonic volume being supersaturated with quantons, limitates the radius decreasing of the $\Gamma_{\mu}^p = 2\pi r_{\mu}^p$ -quantonic vortex, at the value: $r_{\mu}^0 = r_i$.

-The value $\mu_N = \mu_e/1836$ for the nuclear magneton gives -by eq. (33), a magnetic moment radius: $r_i^0 = 0.21 \text{ fm}$, that represents the Compton radius of the proton, close to the experimentally deduced proton core radius r_m , (0.3fm-[24]) and to the experimentally deduced proton quark radius, [24].

The relation (33b) also gives: $r_e^+ = 0.96 \text{ fm}$ for the protonic positron axial position.

Because that—for the electron CF-model case, the vexons of electron surface has a degenerate Compton radius approximate equal with the electron Compton radius: $r_w^e \cong r_{\mu}^e$,

$$N(n) = Q(n)/e = (\sigma_e \cdot 2\pi r_e l_{\sigma})/e = 2n^2; Q(1) = 2e; r_0 = e/(\sigma_e \cdot \pi \cdot l_{\sigma}); r_e = n^2 \cdot r_0 \quad (34)$$

According to the model, the transition on under-fundamental level ($n = 1/2$) is particular to the hydrogen atom, by the condition $Q(1/2) = e$, (H-atom having a single

The loosed part of electron energy: $\Delta \epsilon_e = (m_e - m_e^*) \cong 0.13 m_e c^2 = 65.3 \text{ keV}$, in the bound degenerate gammon formation process, has the signifiante of a binding energy, as in the case of the deuteron's forming.

A small repulsive static type charge of quasidelectron control, given by a sinergonic static pressure, impede the confinement of the proton's kernel until a density: $\rho_k = \rho^m$.

The virtual radius: r_{μ}^n , of the proton μ_p -magnetic moment, compared to the electron, decreases when the protonic positron is included in the N^p cluster volume, from the value: $r_{\mu}^e = 3.86 \times 10^{-13} \text{ m}$, to the value: $r_{\mu}^p = r_i = 0.59 \text{ fm}$, as consequence of the increasing of the impenetrable quantum volume mean density in which is included the protonic positron control: m_0 , from the value: ρ_e , to the value: $\bar{\rho}_n \cong f_d \cdot N^p \cdot \rho_e$, conform to:

it results by eq. (33), for a vexon which is in the proton's surface, ($r_w \cong 1.4 \text{ fm}$), that: $r_w^n \cong (r_{\mu}^e/1836) \cdot e^{1.4/0.93} = 0.946 \text{ fm}$, so: $r_{\mu}^a = a + r_w^n = 2.35 \text{ fm}$, r_{μ}^a being the virtual radius of μ_p , for which: $\Gamma_{\mu}^p = 2\pi r_{\mu}^a$.

The resulted value $r_{\mu}^a \rightarrow 2a$, is conformed also to a pre-quantum soliton model of atom, (which degenerates in the Bohr-Sommerfeld's model at $T > 0 \text{ K}$), for which:

$$v_e(r)/c = \sqrt{2a/r}; v_e(r_0)/c = \sqrt{2a/r_0} = 1/137;$$

$$(r_0 = 0.53 \text{ \AA}; v_e(r) = v_e(r_0)/n; r_n = n^2 r_0),$$

by the conclusion that the sinergonic Γ_{Λ} -vortex of the protonic positron gives the $v_e(r)$ - speed of the atomic electrons by the action of a tangent force: $F_{\Lambda}(r)$, given by the impulse density: $p_s(r) = \rho_s(r) \cdot w$, ($w = \sqrt{2}c$) of the Γ_{Λ} vortex, in a dynamic equilibrium with the advancing resistance force: $F_R(r)$, given by a density: ρ_R of an equivalent pseudo-stationary sinergonic medium, without photons emission.

This solitonic model of atom allows also the explaining of the electron transition on under-fundamental level ($n = 1/2$) in the hydrogen atom, (for hydrino atom), observed in some experiments of cold nuclear fusion, [25], by the hypothesis that the quantification of the electron number of an energy level: $N(n)$, corresponds to a superficial charge density σ_e of constant value for an energy layer considered of quasi-cylinder (barrel- like form), with the height: l_{σ} and quantified radius, $r_e = n^2 r_0$, i.e.:

electron), condition which gives a radius for the under-fundamental level orbital: $r_0^* = e/(\sigma_e \cdot 2\pi \cdot l_{\sigma}) = r_0/2$.

For other atoms, with bigger mass, the transition on

under-fundamental level: $(n=1) \rightarrow (n'=1/2)$ results as possible by stimulated electronic transition, by which may be obtained also „mascons”, i.e- metals with concentrated atomic mass, (bigger density than the normal density), according to the model.

At the proton's level, the superposition of the (N^p+1) quantonic vortices: Γ_μ^* of the quasidelectrons magnetic moments, generates-inside the volume with the radius:

$$\nabla V_s^n(r) = -v_i \cdot \nabla P_d(r); \quad V_s^n = -V_s^0 e^{-\frac{r}{\eta^*}} = \frac{1}{2} \rho_n v_i c^2 = v_i P_d; \quad P_d = P_d^0 e^{-\frac{r}{\eta^*}}; \quad r \leq r_\mu^a = 2.35 \text{ fm} \quad (35)$$

the proton density in its centre resulting: $\rho_n^0 = (N^p+1) \cdot \rho_e^0 = 2105 \cdot \rho_e^0 = 4.68 \times 10^{17} \text{ kg/m}^3$, ($\rho_e^0 = 22.24 \times 10^{13} \text{ kg/m}^3$) and with: $\eta^* = 0.755 \text{ fm}$, the potential well resulting: $V_s^0 = -118.4 \text{ MeV}$. Thus, by eq. (35) is theoretically re-found the form of the exponential nuclear potential, with specific constants.

At the distance $d \cong 2 \text{ fm}$ between deuteron nucleons (considered as the dimension of the nuclear potential well), it results from the relation (35) that the scalar nucleonic potential $V_s^n(r)$ has the value: $V_s^n(d) = -8.37 \text{ MeV}$ - value which corresponds to the known mean binding energy inside the stable nuclei: $-7.5 \dots -8.5 \text{ MeV}$.

The mesonic theory of the nuclear force results -in consequence, by our interpretation of the eq. (33), as formal.

According to eq. (35) it results also that the deuteron self-resonance decreases the value of scalar nuclear potential,

$$V_{Mg}(r) = -\frac{v_e}{2} \rho_s(r) \cdot w_t^2 = -\frac{m_p}{2\rho^M} \rho_a^0 \frac{a^2}{r^2} \cdot c^2 = V_{Mg}^0 \left(\frac{a}{r}\right)^2; \quad V_{Mg}^0 = -\frac{m_p}{\rho^M} \rho_a^0 c^2 \quad (36)$$

3.4. The Neutron Model and the Weak Force

Complying with the CF proton soliton model, the neutron results in the theory conforming to a Lenard-Radulescu dynamid model, (Dan Radulescu, 1922, [26]) , as being composed by a proton center and a negatron revolving around it with the speed $v_e^* < c$, at a distance $r_e^* \leq a$ at which-

$$\mu_n - \mu_p = (\mu_e^L + \mu_e^S) = (-1,91 - 2,79)\mu_N = -4,7\mu_N; \quad \text{with: } \mu_e^L = \frac{e \cdot v_e^* \cdot r_e^*}{2} \quad (37)$$

The neutronic negatron orbital rotation takes place under the action of the dynamic pressure: $\frac{1}{2} \cdot \rho_\mu(r_e) c^2$ of the Γ_μ^n vortex quantons forming the proton magnetic moment μ_p and inside

$$\rho_\mu(r_e^*) c^2 = \rho_n(r_e^*) v_e^{*2}; \Rightarrow \rho_\mu^0 c^2 \approx f_d \cdot \rho_n^0 v_e^{*2}, \quad f_d = 0.8722 \quad (38)$$

With: $\rho_\mu^0 = \rho_e^0 = 22.24 \times 10^{13} \text{ kg/m}^3$; $\rho_n^0 = 4.68 \times 10^{17} \text{ kg/m}^3$, it results that: $v_e^* = 0.0233 \cdot c = 7 \times 10^6 \text{ m/s}$.

Also, by eq. (33) applied to the electron magnetic moment's

$$\mu_e^S = \mu_N \cdot \frac{\rho_n^0}{\rho_n(r_e^*)}; \quad \rho_n(r_e^*) = \rho_n^0 e^{-\frac{r_e^*}{\eta_d}}; \quad \eta_d = 0,93 \text{ fm}; \quad (39)$$

Thus, from (37), (38) and (39) it results:

$$r_e^* = 1.41 \text{ fm}; \quad \mu_e^L \cong -0.1563\mu_N; \quad \mu_e^S \cong -4.554\mu_N,$$

$r_\mu^a = 2.35 \text{ fm}$, according to the superposition principle, a total dynamic pressure: $P_n = (1/2)\rho_n(r) \cdot c^2$ having a variation according to eqs. (26) and (28) and acting over the impenetrable nucleonic volume $v_i(r_i; m_{ni}) \approx 0.9 \text{ fm}^3$ of another nucleon, (the rest part being penetrable to the quantons action).

This dynamic pressure P_n , generates a nuclear potential and force, according to eqn.:

[1].

The sinergonic dynamic pressure: $P_d^s(r)$ of the Γ_μ^n vortices of (N^p+1) -protonic cluster, generates a scalar gravito-magnetic potential $V_s^n(r)$ similar to the nuclear potential but acting upon a volume: $v_c^n \cong m_p/\rho^M = 1.67 \times 10^{-27}/8.8 \times 10^{23} \cong 1.9 \times 10^{-51} \text{ m}^3$ given by the quantonic and electronic super-dense centrols of the proton mass, so- of ~ 100 times smaller than the nuclear potential, but contributing to the stability of the proton's chiral soliton.

However, the magneto-gravitic potential is of an important value for "black holes" and for "magnetars". For $r > r_\mu^a$, by eq. (35) it results that the magneto-gravitic potential generated by an elementary particle over another particle having the mass m_p , has the expression:

according to eq. (33), it has a degenerate μ_e^S magnetic moment and S_e^n -spin, (figure 5).

The revolving of the neutron negatron, generates a negative orbital magnetic moment: μ_e^L , the neutron's magnetic moment resulting by:

the quantum volume having the density $\rho_n(r)$, so we can consider the equilibrium relation of the dynamic pressures, acting over a revolved degenerate negatron area: $S' \cong \pi a^2$, i.e.:

degeneration at the incorporation of the neutronic electron in the protonic quantum volume, it results that:

which leads to the conclusion that the neutronic negatron has the impenetrable quantum volume v_i^c with the centrol positioned in the surface of protonic quantum volume: v_n , (figure 5), comparative with the positronic proton for which

the eq. (33) in which:

$$\mu_e^+ = \mu_p = 2.79\mu_N, \text{ gives: } r_e^+ = 0.96 \text{ fm.}$$

$$v_e = v_e/2\pi r_e = 0.79 \times 10^{21} \text{ Hz}; S_e^n = \mu_e^S \cdot (m_e/e) = 0.0025 \hbar, (\hbar = h/2\pi), \quad (40)$$

-in accordance with the equality between the spin of proton and those of the neutron, ($S_n = S_p = 1/2\hbar$), resulted in the quantum mechanics.

So, by eq. (33), our model of CGT solve the classical problem of the nucleon's spin and of the magnetic moment values, problem which determined the abandonment of the classical nucleon models presuming incorporate nucleonic electron(s). For the anti-neutron, the model imply the rotation of a positron around of an antiproton.

The continuous energy spectrum of the β radiation at neutron's transformation, corresponding to a speed v_e of the β -electron up to $0.7 \div 0.92c$, is explained- in accordance with eq. (34), (35), through the acceleration given to the β -electron by the vortex Γ_μ of the protonic μ_p^- magnetic moment, the given energy depending on the angle of the electron's initial impulse, θ (p_β, r_p).

The fact that- according to the neutron "dynamide" model, the protonic positron coexists with the neutronic negatron inside its quantum volume until the neutron's transformation, may be explained by the model through the hypothesis that the difference of approximate $2.53 m_e$ between the neutron mass and the proton mass is given by a binding γ^* -gammon, called

$$(M_n^* + \gamma^0 + \sigma) \rightarrow (M_n^* + e^+) + e^- + \bar{\nu}_e + \epsilon_\sigma (889 \text{ keV}); (M_n^* + e^+) = {}^1p_r \quad (42b)$$

as given by a gammonic dynamid: $\gamma^0 = e^- + e^+$, splitted by the transformation of the σ -gluol.

According to the model, the electronic pair: negatron-positron of the neutron, represents a gammonic metastable state: $\gamma^0 = e^- + e^+$, attached to the particle cluster M^* with rotating negatron.

Also, the central couple having the mass: $2m_0$, of the disintegrated σ -gluol, is emitted under the form of an electronic antineutrino, having the approximate superior limit of repose mass: $m_\nu(v_e) = 2m_0 \cong 4 \times 10^{-4} m_e = 3.6 \times 10^{-34} \text{ kg}$, [15], by the action of the local quantonic and sinergonic pressure of Γ_μ^n and of Γ_Λ^n vortexes, with the speed $v \rightarrow c$.

$$e^+ + e^- \rightarrow \gamma \rightarrow \nu_e + \Delta \epsilon_\gamma (\sim 1 \text{ MeV}); \pi^0 + \gamma \rightarrow \pi^0 + \nu_e + \Delta \epsilon_\gamma \quad (43)$$

that may occur at high energies of matter compression or by collision with relativist mesons or baryons, with the releasing of the quantonic but also of the sinergonic energy of γ -quantum:

$$\Delta \epsilon_\gamma = (2m_e - m_\nu)c^2, (m_\nu \text{ -the neutrino rest mass}).$$

Also, it results that similarly, the protonic degenerate electron that gives the proton charge and has an axial position at $r_e^+ = 0.96 \text{ fm}$, is linked by another σ -gluol, which explains similarly the β^+ decay:

$$\text{Energy} + p^+ \rightarrow n^0 + e^+ + \nu_e$$

The neutron's spin and the revolving frequency of the neutronic negatron around the proton centre results by the relations:

" σ -gluol" in our model, which has the intrinsic energy:

$$\epsilon_\sigma = 2m_e^*c^2 \cong 1.74m_e c^2 \cong 0.889 \text{ MeV} \quad (41)$$




Fig.5. The Neutron Model.

The reaction of neutron transforming, [15]:

$${}^0n_e \rightarrow {}^1p_r + {}^{-1}\beta + \bar{\nu}_e + Q_k (780 \text{ keV}) \quad (42a)$$

results from the proposed model, in the form:

Also, because that the Γ_Λ^n -vortex generated by the protonic positron, contains also tachyonic sinergons, with $w \rightarrow \sqrt{2}c$ (according to CGT), it is possible also neutrino speed: $v_\nu \rightarrow 1.1c$, as those observed in the OPERA experiment [27], according to the model, [1].

The neutrino model of the theory is complying with the Majorana model, which considers the neutrino as a superposing of two Majorana fields having equal masses and opposed CP parities and with the fact that the neutrino is a very penetrating particle.

According to the model, there are possible also the transformations:

of proton's transforming.

The escape of β -electron from the nuclear field results in conditions of the neutron self-resonance with a intrinsic vibration energy, E_ν^c , of the neutronic electron, induced by a vibration energy of the neutron $E_\nu^n(d)$, which satisfy the conditions:

$$E_\nu^n(d) + \epsilon_\sigma \rightarrow E_\nu^0(d) = 2.226 \text{ MeV}; \quad (44a)$$

$$E_\nu^c \rightarrow m_e c^2 = 0.511 \text{ MeV} \quad (44b)$$

In this case, the natural equivalent of a W^\pm -boson mediating the weak interaction of β -disintegration, used in the quantum

mechanics standard model, is the „weson”: $w^- = (\sigma + e^-)$, which generates the beta disintegration in the form:

$$w^- \rightarrow e^- + \bar{\nu}_e + \epsilon_\sigma \text{ when: } \sigma \rightarrow \nu_e + \epsilon_\sigma, \quad (45)$$

The reciprocally opposed quantum helicities of the negatron and positron, remarked in the β^- and β^+ disintegration (Wolfenstein [28]), are explained in the theory by the spin dependence of the ζ_e -intrinsic chirality of the m_0 -electronic control which- by its considered helix form, passing through a quantum and sub-quantum medium, determines the orientation of the electron spin-parallel or antiparallel with the impulse direction. by electron capture: ${}^1p_r + e^- \rightarrow {}^0n_e + \nu_e$, may be explained similarly by the conclusion that the captured negatron and the protonic positron forms a metastable gammonic state: $\gamma^0 = e^- + e^+$, which is transformed into an ν_e -electronic neutrino, (eq. (43)).

Also, according to the model, at high temperatures as those of a supernova, by the perturbation of the nucleonic structure by the vibration of its kernel, the e^+ gammonic positron may be not retained by the neutronic cluster M_n^* and the neutron may be transformed with a temperature-dependent probability, by gammonic emission:

$$(M_n^* + \gamma^0 + \sigma) \rightarrow M_n^* + \gamma^0 + \bar{\nu}_e + \epsilon_\sigma (889 \text{ keV}) \quad (46)$$

$$\tau_k = \frac{\tau^0}{k_v \cdot 10^{2n}}; \quad \text{with: } \tau^0 \cong 10^{-13} \text{ sec.}; \quad k_v = \frac{\Delta m_p}{m_p} = \frac{n \cdot \epsilon_v}{\epsilon_v^0} = \frac{n \cdot T}{T_d}; \quad T_d \cong 10^{12} \text{ K} \quad (47)$$

in which: $\epsilon_c^0 = k_B T_d \cong h\nu_c^0$ represent the critical phononic energy of particle vibration which determines the quark deconfination, (at: $T_d \approx T_D = 3 \times 10^{12} \text{ K}$, according to the experiments [30]).

As a consequence of eq. (47), when a particle pass with the v - speed through a quantum medium of the space, the dynamic

$$\epsilon_v(v) = h \cdot \nu_i = k_p \cdot k_B T_q = k_p \frac{P_c(v) \cdot m_h c^2}{P_c^0}; \quad P_c(v) = P_c^0 - \frac{1}{2} \rho_c^0 v^2; \quad P_c^0 = \rho_c^0 c^2 \quad (48)$$

which is equivalent with a relation for the intrinsic quantum temperature variation, of the form:

$$T_q(v) = T_q(0) \cdot (1 - v^2/2c^2) = T_q(0) \cdot \beta^2; \quad k_B \cdot T_q(0) = m_h c^2 \quad (49)$$

similar to the Einstein's relation: $T = T_0 \cdot \beta$, but with β^2 in the classic form (18). Also, by (48) we have:

$$\frac{\epsilon_v(v)}{\epsilon_v(0)} = \frac{P_c(v)}{P_c^0} = \left(1 - \frac{v^2}{2c^2}\right) = \frac{\tau_k(0)}{\tau_k(v)}; \quad (50a)$$

$$\tau_k(v) = \tau_k(0) \cdot \left[1 - \frac{v^2}{2c^2}\right]^{-1} \quad (50b)$$

The eqns. (49), (50) explains also the lifetime increasing for relativistic μ^\pm - mesons or other relativistic particles with: $v \rightarrow c$, the eq. (92b) being mathematically quasi-equivalent to the einsteinian relativistic relation used by Rossi and Hall, [31], but naturally obtained.

This theoretical conclusion may explain the cosmic pulses of gamma radiation detected as coming from the direction of Oort cosmic cloud [29] and resulting by collision of nuclear components-phenomenon not enough understood.

By eq. (46) the phenomenon may be produced also by pulsatile contraction of the volume of a supernova or of a neutronic star.

The particle deconfination may be explained also by the CF particle model as a result of the vibration energy: ϵ_v , of n current mass quarks, structured as cluster of quasidelectrons [1].

Considering the lepton μ^+ , having a lifetime:

$\tau_\mu = 2.2 \times 10^{-6} \text{ sec.}$ [15], as a single-particle cluster, and taking into account that the majority of baryons (considered with three quarks in the cluster M^* sub-structure) have a lifetime value: $\tau \cong 10^{-10} \text{ sec.}$ and the majority of mesons have a lifetime of approximately 10^{-8} sec. at the ordinary temperature: $T \cong 300 \text{ K}$ of the medium which contains the particle, the lifetime of the elementary particles results -by the dependence on the destroyed mass: $\tau_k \sim 1/\Delta m_p(T)$, induced by the total intrinsic ϵ_v -vibration energy of the M^* -cluster, according to an semi-empiric relation of approximation:

quantum pressure generated in a relativistic way by the quanta and subquanta of this medium, has a cooling effect for the M^* -particle cluster, (which explains also the existence of polarized quantum vacuum bosons as metastable particles), according to the equation:

3.5. The Deuteron's Self-resonance and the Quasicrystallin Nuclear Model

In the deuteron's case, the experiments evidenced a binding energy: $\Delta E(d) = -2.226 \text{ MeV}$ for the real deuteron having parallel nucleonic spins and of about -0.07 MeV for the virtual deuteron, having anti-parallel nucleonic spins, [15].

Comparative to the binding energy value:

$V_n(d) = -8.4 \text{ MeV}$, ($d = 2 \text{ fm}$), of the unperturbed deuteron state from stable multi-nucleonic nuclei, the value $\Delta E(d) = -2.226 \text{ MeV}$ indicates, by eq. (35), a decrease of the quantonic dynamic pressure: $P_d(r) = 1/2 \rho_c(r) \cdot v_{ct}^2$ in the composite chiral soliton of the N^p protonic cluster.

This decrease is generated by the decrease of the r_μ^a - radius of the exponential part of quasidelectron chiral soliton, for the

N^p -cluster, at a value: $r_\mu^c < r_\mu^a = 2.35 \text{ fm}$, as consequence of the perturbations caused by an intrinsic vibration inside the deuteron nucleons, with an energy E_v which decreases also the value of nuclear potential well: V_s^0 , in accordance with eq. (35), by the energy: $\varepsilon_v = \Delta m_v c^2$ of destroyed vexons, to a value:

$$V_s^{0*} = k_v^* \cdot V_s^0 \sim E_v(d, l_v); (k_v^* \leq 1),$$

with: $E_v(d, l_v)$ the mean vibration energy permitted by the vibration liberty: $l_v = A_v$, i.e.:

$$V_s^{0*} = k_v^* \cdot V_s^0; k_v^* \equiv \left(1 - \frac{\varepsilon_v(d, l_v)}{\varepsilon_v^0}\right) \equiv \left(1 - \frac{E_v(d, l_v)}{E_v^0(d, l_v^0)}\right) \quad (51)$$

in which ε_v^0 ; $E_v^0(d, l_v^0)$ represents the critical values of ε_v and $E_v(d, l_v)$ which cancel the attractive potential: $V_s^*(d)$.

Because that the mass defect: $\Delta m_D = (m_p + m_n - m_D) \equiv 2.23 \text{ MeV}/c^2$ results at the deuteron's formation as destroyed vexons mass: ε_v^0/c^2 , which corresponds to the binding energy:

$$V_s^*(d, E_v) = V_s(d + l_v^*) = V_s(d) \cdot e^{-\frac{l_v^*}{\eta^*}} = V_s(d) \cdot k_v^* \left(\frac{r_\mu^c}{d}\right)^2 \equiv V_s(d) \cdot \left(\frac{r_\mu^c}{r_\mu^a}\right)^2 = \Delta E_D; \quad \eta^* = 0,755 \text{ fm}; \quad (52)$$

For: $r_\mu^a = 2.35 \text{ fm}$, $k_v^* = 0.72$, $l_v^* = 1 \text{ fm}$, it results from eq.(52): $r_\mu^c \cdot k_v^* \equiv 1 \text{ fm}$; $r_\mu^c \equiv 1.2 \text{ fm}$ and: $E_v(l_v^*) = 0.66 \text{ MeV}$.

This theoretical result complies with the conclusion of quantum mechanic's deuteron model, which shows that- on average, the deuteron nucleons are found outside the limits of the potential well having the length: $d_d = 2 \text{ fm}$, the probabilistic deuteron radius being: $R_D = 4.32 \text{ fm}$, [15].

-In the virtual deuteron case, the nucleons having anti-parallel spins, the neutronic negatron revolves as in its free state around the proton center of the neutron, passing periodically with the frequency: $\nu_e = 0.79 \times 10^{21} \text{ Hz}$ between the two deuteron's protonic centers, (figure 6).

Each time when the neutronic negatron passes between the two deuteron protonic centers with a magnetic moment parallel with the proton's magnetic moment, it intervenes

$$V_s^n(r) = -V_s^0 \cdot e^{-\frac{r}{\eta^*}} \cdot e^{-\frac{l_v^*}{\eta^*}} [\text{MeV}]; \quad l_v^* = l_v^0 \cdot \left(\frac{3}{2} - \frac{1}{2} \vec{\tau}_p \cdot \vec{\tau}_n\right); \quad \vec{\tau} = \frac{\vec{s}_n}{s_n}; \quad (53)$$

with: $V_s^0 = 118.4 \text{ MeV}$; $l_v \equiv A_v$; $l_v^0(E_v) \equiv 1 \text{ fm}$ -for the deuteron; ($l_v(E_v=0) = 0$).

The deuteron model of quantum mechanics considers also a self-resonance vibration mechanism for deuteron, but in a different way, explaining the decreasing of the deuteron's binding energy to the value: $E_D = -2.226 \text{ MeV}$ by considering a reciprocal vibration: $E_v \equiv 20 \text{ MeV}$, [15], of the deuteron nucleons - value which is in a relative discrepancy with the value E_D of the binding energy.

Comparative with the plastic interaction of nucleons, with $A_v \rightarrow 0$, when the vexon's energy: $\Delta \varepsilon_v(\Delta \rho_n^0)$ of the nucleon superficial destruction is emitted, ($\Delta \varepsilon_v = \Delta m_n c^2$), in the deuteron's case, this energy is a nucleon's re-separation energy, which-by partial absorption, is re-used for the regeneration of

ΔE_D , it results that:

$$E_v^0(d, l_v^0) = \frac{1}{2} m_p v_p^2(d) = \varepsilon_v^0 = \Delta E_D = 2.226 \text{ MeV}.$$

This conclusion is in concordance with the Onsager's observations regarding the circulation value decrease for a super-fluid perturbed over a critical value. From an energetic point of view, the effect of the vibration energy: $E_v \leq \Delta E_D$ may be explained by the contribution of the nuclear potential $V_s(d)$ to the deuteron self-resonance state, through an alternately „destruction-regeneration” mechanism of the deuteron state.

According to the model, (fig. 6), simplifying, we may approximate also that the initial value: $V(r_\mu^a)$ of the potential well is recovered by the negentropy of the etheronic winds at the distance-limit between proton and neutron: $r_d = d + A_v$ for which the nuclear potential, given by eq. (35), has the approximate value: $V_s(r_d) = \Delta E_D = -2.22 \text{ MeV}$, i.e: $r_d \equiv 3 \text{ fm}$ and $A_v = l_v^* = 1 \text{ fm}$, according to:

against the proton with a repulsive magnetic potential: $V_\mu^n(d_d/2) \equiv 0.3 \text{ MeV}$.

The deuteron's protonic centers, as a consequence of the induced deuteron's self-resonance, are thus re-separated to a distance: $r_d^* = d + A_v^*$ with $A_v^* > 2r_i$, which determines -in accordance with eq. (52), a maximum decrease of the degenerate value: r_μ^c at the value: $r_\mu^p = 0,6 \text{ fm}$, which corresponds at: $l_v^* = A_v^* \equiv 2 \text{ fm}$, the scalar nuclear potential decreasing at a minimal value: $V_s^*(d; l_v^*) \equiv -0.6 \text{ MeV}$ which is canceled by the remained nucleon's vibration energy, explaining the fact that the deuteron with anti-parallel nucleon spins is a virtual state.

In a conventional very simplified form, the spin-dependent nuclear potential may be expressed- in accordance with the resulted model, in the form:

the nucleon's mass and vorticity, by the Γ_A^* -vortices.

Conforming to the solitonic CF nucleon model and to the observations regarding the nuclear stability that shows a maximum stability for the even-even nuclei, the pre-quantum nuclear model, of $T \rightarrow 0 \text{ K}$, results as a quasi-crystalline cluster having nucleons coupled in deuteronic pairs, and corresponding also to the α -particle cluster model, to the “nuclear molecule” model and to the extreme-uniparticle type model, [32], the stable nuclei, with a “magic” number of protons or/and neutrons: 2; 8; 20; 28; (40); 50; 126, being- in the model, symmetrical quasi-crystal forms, resulted from the superposition of square root forms with an integer n^2 -number of α particles, having $2n^2$ protons [1]:

$$Z = \Sigma(2n^2), (n = 1.2 \dots 7, \text{ figure 6}), \text{ and with tendency to a}$$

minimum deformability: $2; 2 \times 2^2 = 8; (2 \times 3^2 = 18); 18 + 2 = 20; 20 + 8 = 28; (2 \times 4^2 = 32); 2 \times 5^2 = 2 \times 3^2 + 2 \times 4^2 = 50; 50 + 32 = 82$, or of quasi-stable triangular forms (^{10}Ne) or hexagonal forms (^{19}K) completed with additional neutrons, for $Z > 20$.

The $^{82}\text{Pb}^{208}$ nucleus corresponds to the initial form: $^{104}\text{N}^{208}$ ($Z=2(4^2+6^2)$) in which 22 protons was transformed into neutrons by β^- -emission giving $Z = 82$, according to the model. Similarly it may be formed a nucleus with $A = 4(5^2+7^2) = 296$, with $Z=112+114$, (close to the predicted stable form 114/298).

The weakly bound excedentary nucleons or alpha-particles formed from the valence nucleons, are revolved around a nuclear quasicrystal, (Lonnroth, [33]), as in the extreme-uniparticle (Schmidt, [32]) model, by the action of quantonic Γ_μ^N -vortex of the nuclear magnetic moment, which explains also the nuclear centrifugal potential.

A quasi-crystalline nuclear structure was evidenced by experiments of α -particles scattering on heavy nuclei, (W.Bauer, [34]). The nuclear fission is explained through the deuteronic self-resonance mechanism of weakly bound nucleons, which decreases the nuclear potential in regions with incompleteness of the quasi-crystal network or with exceeding nucleons, according to eq. (53).

Through the same relations (46), by the deuteron's self-resonance mechanism and without the hypothesis of exciting energy concentration on a single nucleon, it is also possible to explain the following:

- the compound nucleus transformation mechanism by excitation with particles having low energy, up to 2MeV, as in the case of Be9 which can be transformed with a gamma quantum of only 1.78MeV, even if the binding energy, given by its nucleons, is 58 MeV;

- some reactions with thermal neutrons (having some tens of eV) as in the reaction: $-\text{Li}7+\text{H}1 \rightarrow \text{Be}8+2\text{He}4+\gamma$, generated with only 125eV proton energy, or generated by thermal neutrons in typical reactions ($n;\alpha$), such as the reaction: $\text{B}10+n \rightarrow \text{Li}7+\alpha$, even if normally there are necessary neutrons with 0.5...10MeV; [15];

- nuclear transformation with particles having only 1÷2MeV: $\text{Ca}(p, n)\text{Sc}; \text{Al}(p, \alpha)\text{Mg}$;

- the super-asymmetrical nuclear fission, [35].




Fig. 6. The deuteron and the Quasicrystal nucleus.

3.6. The “Dark Matter” as Bosons of the Quantum Vacuum

According to CGT, in the strong interaction of particles, the interaction energy generates real (q- \bar{q})-pairs of quarks, considered in a quasidelectrons cluster model, with the charge given by an un-paired quasidelectron ($+2/3e$) or by a quasi-positron and a degenerate electron, ($-1/3e$), these

(q- \bar{q})-pairs resulting from the polarized quantum vacuum bosons, (named “zerons” in CGT), considered as particle-antiparticle pairs with self-resonance realised according to the incertitude relation: $\Delta E_v \cdot \Delta \tau_v \cong \hbar$; ($\Delta \tau_v$; ΔE_v -the vibration period and energy), the “darkness” of the vacuum M_b -boson being given by a very low self-vibration, $\Delta x_v \ll A_v$, with $A_v = \hbar/m_z c$, (by $T = E_v/k_B \rightarrow 0\text{K}$), as in the eqn:

$$v_\mu + p_r + Q_i \rightarrow v_\mu + p_r + 2(m_1 + \bar{m}_1^-) + (m_2 + \bar{m}_2^-) \rightarrow v_\mu + p_r + \pi^+ + \pi^- + \pi^0; (m_2 + \bar{m}_2^-) = m_{z2}$$

So, the hypothesis of neutral Z^0 boson or Higgs boson of Q.M. is not strictly necessary for explain the particles cold forming and theirs interactions, according to CGT, the generating of particles with bigger mass than those of particles that enters in reaction being explained- in our theory, by the decomposing of quantum vacuum “zerons” of m_z -mass and $x_r = a$ radius, in real (q- \bar{q})-pairs, by the Q_i - interaction energy, considered also in quantum mechanics, when: $O_i \approx E_q = m_z c^2$.

These “zerons” of ,quantum vacuum’ are- in our theory, a classic equivalent of the bosonic background of “dark matter” and may be considered as bosonic m_z -particles with self-resonance, (oscillons), with a phononic intrinsic vibration energy of paired quarks, given by: $E_v \cong (\Delta p \cdot \Delta x_v / \Delta \tau) < E_q$, ($E_q = m_z c^2$; $\Delta x_v \leq 2a$; $\Delta \tau$; $p \Delta x_v$ -the self-resonance period and amplitude), which explains the existence of pseudo-virtual paired quarks and fermions in the “quantum vacuum”.

The cold formed quarks, have- in CGT, a current mass which gives the particle’s rest mass according to the same rule, [1].

It results also by CGT the possibility of exotic particles cold forming as hexaquarks or nine-q clusters and the quark \rightarrow fermion transforming, ($q^{+2/3} \rightarrow p^{+1}$), at $T \gg 0$, by the relative detaching and moving in the quark interaction quantum volume (Δa), of the un-paired quasidelectron e^* which gives its charge $e^* = \pm 2/3e$ and which is auto-transformed in this case in degenerate electron with e-charge (and degenerate magnetic moment and spin), i.e.: $q^{+2/3} \rightarrow p^{+1}$; ($e^{*+2/3} \rightarrow e^{+1}$, by the quantum medium negentropy).

4. Implications of the Theory in Cosmology

4.1. A phenomenological Model of Expanding Universe

According to the CF-model of the theory, it results also that the fermions entropization at high temperatures with partial destruction, generates a temperature-dependent mass decreasing and a pseudo-antigravitic field of a Q_a -pseudocharge (eq. (43)), having the form (10) and a value proportional with the particle vibration energy: $\epsilon_v = k_B T$.

This theoretical conclusion may explain the observed temperature-dependent gravitational mass decreasing for which Shaw and Davy [36] obtained a relation of temperature-dependent gravitational force with the form:

$$F_G(T) = F_0(1 - \alpha T); F_0 = -G \cdot (M \cdot m) / r^2 \quad (54)$$

$$\alpha = 1/T_G = 2.0 \times 10^{-6} [\text{K}^{-1}], (T_G = 5 \times 10^5 \text{K}).$$

Because that the quantity of destroyed intrinsic vexons is

$$m_p(T) = m_p^0 - \Delta m_p(T) = m_p \cdot \left(1 - \frac{T}{T_c}\right); \Delta m_p(T) = m_p^0 \cdot \frac{T}{T_c}; T_c = T_N \approx 3 \times 10^{12} \div 10^{13} \text{K} \quad (55)$$

The value $T_N \approx 10^{13} \text{K}$ results from the energy necessary to nucleonic kernel: $N^p m_0$, for penetrate the proton impenetrable quantum volume, at speed $v_0 \rightarrow c$, in a classic expression permitted by eq. (18), which gives an approximate value: $E_0 = \frac{1}{2} N^p m_0 c^2 \approx 0.11 \text{MeV}$, that is obtained by the proton's vibration with an energy:

$$\epsilon_p^0 = \frac{1}{2} m_p c^2 = 0.47 \text{GeV}$$

and a critical frequency of its destruction:

$$v_c^0 = 1/\tau_c = c/a = 2 \times 10^{23} \text{Hz}.$$

The energy which must be given to the proton for its destruction is obtained by the relativist expression of mass: $m_p^r = m_p/\beta^r$, given by eq. (18) with $v^0 \rightarrow c$, and corresponds to a proton energy: $\epsilon_p^r = \frac{1}{2} m_p^r c^2 = 2\epsilon_p^0 = m_p c^2 = 0.94 \text{GeV}$ - equal with its intrinsic energy, which explains the proton destruction mechanism in concordance with the inferior limit of the proton's destruction energy obtained by the quantum mechanics.

The quasar's energy generated by nucleon's mass \rightarrow energy transformation, by a nuclear temperature: $T_N = \epsilon_p^r/k_B \approx 10^{13} \text{K}$, is more plausible than those imposed by the Big-bang model of Universe, (10^{14}K).

This phenomenon indicates as plausible an open-type phenomenological model of expanding Universe, according to CGT.

For a model of the Universe evolution, the Hubble's law of the cosmic expansion: $v_R = H \cdot R$, which is confirmed for the case of our cosmic time: t_L and our location from the Universe centre: R_L , results in CGT as a particular case, by the existence of many repulsive antigravitic charges of ultra-hot stellary sources, with a mean density: $\rho_a(R, t_E)$, the total mean gravitic

proportional with the vibration energy: $\Delta m_p c^2 \approx k_s \cdot \epsilon_v = k_s \cdot k_B T$, ($k_s < 1$ constant), it is logical to consider a temperature-dependent decreasing for the inertial mass for all particles, in the form:

charge density: $\rho_{G^r} = (\rho_M + \rho_a)_R$, being given by a gravitic and an anti-gravitic charge density of the matter.

Considering also a Macronucleus of Universe with a R^0 radius having a macro-black-hole with a Macro-vortex around it, which has the density variation proportional with the mean matter density: $\rho_m(R) \sim R^{-1}$, we may consider also a variation of the etheronic pressure: $P_c(R) \sim [R^{-1} \div R^2]$ with the R-distance from the supposed Macronucleus, specific to a magneto-gravitic pseudo-vortex with similar density variation as the matter density, the gravity G- constant depending on the quantum pressure: $P_c(R)$ by the etheronic density, ρ_G^0 , according to eq. (26).

Thus, close to the limit $R = R_u$ - considered as the structured Universe radius, the gravity force and the quantum vortices intensity becomes too weak for forming or conserving vortical structures. In this case, we may consider that the zone: $\Delta R_u \leq (3R_u/4 \div R_u)$ represents a zone of "stellar cemetery" (S.C.) in which the stellary structures disintegrates at the distance: $R_d \geq 3R_u/4$ and that the protons and the neutrons disintegrates at the distance close to $R = R_u$ - as a consequence of the decreasing of the nucleonic strong interaction potential, according to the CGT's pre-quantum chiral soliton model of particle.

This conclusion corresponds- partially, to the Universe bubble model, (i.e.-limited Universe).

The expanding of an ellipsoidal quasi-flat Universe with a mass: $M_{R_u} \sim 2R_u^0 \cdot \pi R_u^2 \cdot \rho_u$ for which the local mean matter density is: $\rho_m(R) \sim R^{-1}$, may be approximated according to the Poisson's equation as being equivalent with a deformed spheric Universe with $\rho_m'(R) \sim R^{-2}$ having the same mass with it for each R- radius, (fig. 7), by the equations:

$$M_{R^0} \equiv \int 2R^0 \cdot 2\pi R \cdot \rho_m(R) dR \equiv \int 4\pi R^2 \cdot \rho_m'(R) dR = M_{sR} \Leftrightarrow$$

$$\Leftrightarrow \rho_m(R) = \rho_m^0 \cdot (R^0/R); \rho_m'(R) = \rho_m^0 \cdot (R^0/R)^2 \quad (56)$$

$$a_u(R) = \ddot{R} = -G \frac{4\pi R^3 (\rho_m + \rho_R + \rho_a)}{3R^2} = (H^2 + \dot{H}) \cdot R; \rho_R = \frac{3p_R}{c^2} \quad (57)$$

with: ρ_R ; p_R - the space radiation density and pressure, (mainly- of 3K), ρ_m - the mean matter density; ρ_a - the mean antigravitic charge density and: $v(R) = dR/dt = H \cdot R$.

The eq. (57a) is classically equivalent to eq.

Einstein-Friedmann for the flat Universe ($k=0$) with negligible matter pressure, p_m , by: $|\rho_a| = 2\rho_\Lambda^*$, (ρ_Λ^* - the "dark energy" density, $\rho_a(R) \sim T_u(R)$, (T_u - the mean Universe's temperature- eq. (55)), i.e. -with the equation:

$$3 \frac{\ddot{a}}{a} = \Lambda c^2 - 4\pi G \left(\rho_m + \frac{3p_m}{c^2} \right) = -4\pi G \left(\rho_m + \frac{3p_m}{c^2} - 2\rho_\Lambda \right); \rho_\Lambda = \frac{\Lambda c^2}{8\pi G} \quad (58)$$

According to eqs. (57)+(58), the Universe expansion is

obtained by the antigravitic charge of the total matter, of

density ρ_m , but given by the part of the observed matter, for which:

$\Omega_M = \rho_M/\rho_c \cong 0.05$, (the rest being “dark matter”) and with $\Omega_\Lambda = \rho_\Lambda/\rho_c \cong 0.75$, ($\Omega_m = \rho_m/\rho_c \cong 0.25$; $\rho_c = 3H^2/8\pi G$), by: $|\rho_a| = 2\rho_\Lambda^*$ and by a mean temperature T_M of the visible matter, ρ_M , i.e.:

$$|\rho_a^e| = 2\rho_\Lambda^* \cong \frac{T_u}{T_G} \cdot \rho_m^e = \frac{T_M}{T_G} \cdot \rho_M^e \approx 6 \cdot \rho_m^e \cong 30 \cdot \rho_M^e$$

In the field of the Macronucleus, the disintegration of nucleons occurs also because an ultra-high nuclear temperature close to the critical value: $T_N \cong 10^{13}$ K, generated periodically by a big black hole -according to CGT.

For a position with $R > R_u/2$ of the cosmic body, the winds coming from “stellar cemetery” (S.C.) zone generates a pressure in the sense of slowing the Universe expansion, so we may approximate the Universe’s expansion law by the eqs:(59a) $v_e = \partial_t R = v_M \cdot \sin(\pi R/R_u)$; $v_M \cong k_e \cdot c$; $R \leq R_L = (1/6)R_u \Leftrightarrow \sin(\pi R/R_u) \cong (\pi R/R_u)$; (59b) in which the maximum value, $v_M \cong k_e \cdot c < c$, was considered as the maximum expansion speed, (a value: $k_e \approx 0.5$ corresponding to the redshift of the

quasar 3C295: $v_q = 0.46c$).




Fig. 7. The Expanding Universe model.

According to the model, the Hubble law is valid in the zone of the local galaxy super cluster (Virgo) and its surroundings, and it may be regained from eq. (59a) by the condition (59b), i.e.:

$$\frac{v_e}{v_M} \approx \frac{H \cdot R}{k_e \cdot c}; \Rightarrow H = \frac{k_e \pi \cdot c}{R_u}; R \leq R_u / 6; \quad (60a)$$

$$F_e = F_a - F_d = M_s^* \frac{dv_e}{dt} = \frac{\pi \cdot k_e^2 M_s^* \cdot c^2}{2R_u} \sin \frac{2\pi R}{R_u}; R < \frac{4}{5} R_u; M_s^* = M_s^0 / \left(1 - \frac{v^2}{2c^2}\right) \quad (60b)$$

With the mean value: $H_a = 75 \text{ Km/s-Mps}$, deduced by A. Sandage in 1958 [37] and by a plausible value: $k_e \approx 0.5$, it results from eq. (60a) that: $R_u = 6.28 \times 10^3 \text{ Mps}$, ($27.3 \times 10^9 \text{ l.y.}$) - of two times bigger than those deduced by the Big-Bang cosmological model-corresponding to an Universe filled with stars.

For a cosmic body M_s , the expansion force has the form (60b) in which F_a represent the accelerating force - given by the pressure of the stellar winds (mainly-etheronic winds) coming with the radial mean intensity I_a from the zone of the expansion center and F_d represent the decelerating force, given by the pressure of stellar winds coming with the radial mean intensity I_d from the C.S.-zone and by the resistance force to advancing, given by the density of the sub-quantum and quantum medium of the cosmic “vacuum”, (giving an apparent mass variation).

The mass: M_s^* represents the virtual mass given by the relativistic relation (18), the eq. (60b) being in accordance with the linearized form of the Einstein-de Sitter equation:

$$\rho_s^* > 10^2 \cdot \Delta \rho_g^M = 5.47 \times 10^{-27} \text{ kg/m}^3 > \rho_m \cong \Omega_m \cdot \rho_c \cong 4 \times 10^{-27} \text{ kg/m}^3, \quad (63)$$

so- corresponding to the mean “dark energy” density value deduced in accordance with cosmological observations [38]:

$$\rho_s^* = \rho_\Lambda \cong 1.2 \times 10^{-26} \text{ kg/m}^3.$$

For a pair of quantons, in our galaxy ($R=R_l$), because the very small quanton radius, the gravitonic component: $\rho_g^h(m_h)$ which gives the G-value by eqs. (25)-(26) may be considered

approximate equal to the value $\Delta \rho_g(R_l; G)$, i.e.:

$\Delta \rho_g(R_l; G) = \Delta \rho_g^M \cdot \sin(2\pi R_l/R_u) \approx \rho_g^h(m_h) = 6.2 \times 10^{-30} \text{ kg/m}^3$, so G_M corresponds to: $G_M(R_u/4) = 8.8G(R_l)$ and to: $R_l = 1.8 \times 10^{-2} R_u$ - for the position of our galaxy, Milky Way.

The increasing of the expansion force F_e until the maximum value F_e^M is explained in the model by the increasing of the R-depending number of dark energy sources which generates

$$R_{ik} - \frac{1}{2} g_{ik} \cdot R + \Lambda \cdot g_{ik} = T_{ik} = 0 \quad (61)$$

We may consider also that the intensities I_a and I_d of the stellar winds generating the expansion force are given mostly by etheronic winds, acting upon the mass M_s , so the expansion force F_e results conform to eq. (16) of the gravitation force, the maximum value of F_e -force being given for $R = R_u/4$, by the eqn.:

$$a_e^M = \frac{F_e^M}{M_s^*} = \frac{\pi \cdot k_e^2 \cdot c^2}{2R_u} = \frac{S_h}{m_h} (I_a - I_d) \frac{\pi}{4} \cong k_h \cdot \Delta \rho_g^M \cdot c^2 \quad (62)$$

With the gauge value: $k_h \cong 27.4 \text{ [m}^2/\text{kg]}$ resulted from the theory, considering that: $k_e \approx 0.5$, ($v_M \approx 0.5c$), it results from eq. (62) a value:

$\Delta \rho_g^M \cong 5.47 \times 10^{-29} \text{ kg/m}^3$, and because that the mean etheronic density, ρ_s^* , which ensures the gravitational stability of the material structures in the intergalactic space must be with at least two size order bigger, it results that :

the $I_a(R)$ intensity, (i.e. with pulsatory antigravitic charge), contained by the $S_U(4\pi R^2)$ sphere of Universe, (eq. (57)).

The recently observed distribution of quasars in the Universe sustains the previous explanation looking the “dark

energy” provenience. The estimated value for ρ_Λ^* gives an important effect of “radiation aging” which may explain the Olbers paradoxe and which contributes to the total redshift effect, according to the equations:

$$\Delta E_v = h \cdot v_i - h \cdot v_f = F_f \Delta R = \frac{1}{2} k_h \cdot m_f \rho_s \cdot c^2 \cdot \Delta R = \frac{1}{2} k_h \cdot \rho_s \cdot h \cdot v_i \Delta R \quad (64a)$$

$$v_f = v_i \cdot (1 - \frac{1}{2} k_h \cdot \rho_s \cdot \Delta R); z_a = \Delta v / v_f = \frac{1}{2} k_h \cdot \rho_s \cdot \Delta R / (1 - \frac{1}{2} k_h \cdot \rho_s \cdot \Delta R); \quad (64b)$$

Considering the position of the local supercluster of galaxies (Virgo) at $R_v = R_1 \approx 1.8 \times 10^{-2} R_U$, it results from eq. (64b) the condition to receive photonic radiation from the

margin of the stellary Universe, considered at: $R_M = \frac{3}{4} R_U$, according to the model:

$$\Delta v / v_i < 1 \Rightarrow \rho_s^c < 2 / k_h \cdot \Delta R = 3.77 \times 10^{-28} \text{ Kg/m}^3; (\Delta R = R_M - R_1 \approx R_M; k_h = 27.4); \quad (65)$$

From eq. (65) it results the conclusion that- because that we have: $\rho_s^* > \rho_M \approx 6 \times 10^{-28} \text{ kg/m}^3$, we cannot receive photonic radiation from the margin of the stellary Universe, with: $\rho_s^* \approx$

$\rho_\Lambda \approx 1.2 \times 10^{-26} \text{ kg/m}^3$ resulting that the maximal distance ΔR_c from which we can receive photonic radiation is given by:

$$\Delta v / v_i = 1 \Rightarrow \Delta R_c = 2 / k_h \cdot \rho_s^* = 6.08 \times 10^{24} \text{ m} = 6.4 \times 10^8 \text{ l.y.} = 2.36 \times 10^{-2} R_U. \quad (66)$$

Comparing the z_a redshift with the redshift given by relativistic Doppler effect:

$$z_r = [\sqrt{(1+v/c)/\sqrt{(1-v/c)}}] - 1 \approx H \cdot \Delta R / c, \quad (67)$$

with: $\rho_s \approx \rho_\Lambda \approx 1.2 \times 10^{-26} \text{ kg/m}^3$, for: $\Delta R = 10^{-2} R_U$, we have: $z_a \approx 0.424 / (1 - 0.424) = 0.73$ and $z_r \approx 0.016$, so it results that the redshift z_a given by the “tired light” effect is much greater than the redshift z_r given by the Universe inflation- in accordance with the conclusions of “sub-quantum kinetics” theory of Paul LaViolette which showed (1987) that the tired-light model fits observational data better than the “expanding Universe” model, as showed also Tolman, (1985). So, the red-shift of 1.4 or higher, observed to many visible galaxies may be explained by eqs. (64) as “aging radiation” effect, explaining similarly also the high value of the red- shift observed to distant supernovae (of Ia type).

It results also that the high value of some quasars redshift: z

= 4, ($v_m = 0.92c - 1986$) and $z = 6.3$, ($v_m = 0.92c - 2001$) is given by an intense “tired light” effect generated partially by the density of quantum and sub-quantum medium, increased by the strong magnetic field of a rotational (Kerr - type) “black hole” and by its gravitational attraction .

Also, the proposed inflation scenario of CGT based on the pulsatory antigravitic charge model, eliminates the hypothesis of “inflaton”, (quantum-particle of the inflation field).

Because that the density of the un-compensated etheronic winds, $\Delta \rho_g$, acts as a gravitic flux: $\Delta \phi = \frac{1}{2} \Delta \rho_g c^2$, generated by a total mean gravitic charge density: $\rho_{Gt} = (\rho_m + \rho_a)_R$ of the Universe mass, $M_U(R)$, by eqs. (56) and (60), neglecting the value of space radiation density ρ_R , it results also that:

$$a_u(R) = \ddot{R} = \frac{k_e c \cdot H}{2} \sin \frac{2\pi R}{R_u} = -\frac{4\pi G}{3} (\rho_m + \rho_a)_R \cdot R; \quad R < \frac{3R_u}{4} \quad (68)$$

The variation of the mean total gravitic charge density of the Universe mass, $M_U(R)$, given by the Universe expansion, results from eq. (68), in the form:

$$\rho_{Gt}(R) = (\rho_m + \rho_a)_R = -\frac{3k_e c H}{8\pi G} \cdot \frac{1}{R} \sin \frac{2\pi R}{R_u}; \quad \rho_a = -\frac{T_u}{T_G} \cdot \rho_m; \quad R < \frac{3R_u}{4}; \quad (69)$$

Also, because that for $v(R) \approx H \cdot R$, ($R \leq R_u/6$), we have: $|\rho_a| = 2\rho_\Lambda^* \approx 6\rho_m$, it results that: $R_1 \leq R_u/6 \Rightarrow T_u \approx 6 \cdot T_{G(R_u/2)}$. For: $H = H_a = 75 \text{ Km/s.Mps}$, ([37]), and the eq. (68) becomes:

$$a_u(R) = \ddot{R} \approx H_a^2 \cdot R = -\frac{4\pi G}{3} (\rho_m + \rho_a)_R \cdot R; \quad \Rightarrow \quad \rho_m \approx \frac{3H_a^2}{20\pi G} \approx 4.23 \times 10^{-27} \text{ kg}; \quad (\rho_a \approx -6\rho_m) \quad (70)$$

so: $\rho_\Lambda^* \approx 3\rho_m \approx 1.27 \times 10^{-26} \text{ kg}$ - in accordance with the known determined value: $\rho_\Lambda \approx \Omega_\Lambda \cdot \rho_c = 1.2 \times 10^{-26} \text{ kg}$.

The equality: $\rho_m(R_u/2) = -\rho_a(R_u/2)$ resulted from eq. (70) is explained with the eq. (54):

$$\rho_m(R) \leq -\rho_a(R) \quad \Leftrightarrow \quad T_u \geq T_G(R_u/2); \quad R \leq R_u/2 \quad (71)$$

The value $\rho_a \equiv 0$ corresponds in the model to the “thermal death” of the Universe’s stars.

4.2. Exotic stars

4.2.1. The Gravistar as Genesis Structure

Relative to the Protouniverse structure, it results in CGT, by

the similitude principle, an anisotropic model of “gravistar”- considered as a hard-core rotation ellipsoid of “primordial dark energy” containing vortexially generated “dark photons” and “dark particles”- formed as Bose-Einstein condensates at distinct levels of density. This possibility is argued also by the known model of “gravastar” with very cold core formed by a “dark energy” fluid, which may create Bose-Einstein condensate in the outer core, [39], but which suppose an already existent central “black hole”.

In the proposed model of hard-core gravistar, [1], the “gravitational vacuum” region specific to a “gravastar”, not exists, the quasi-stability of the hard-core deformed ball of “dark” energy, forming a relativist vortex of quantons: $\Gamma_\mu = 2\pi \cdot v_c$, ($v_c \rightarrow c$), being given- in the model, similarly to the electron case, by a quantum potential, $V_T(r)$ which- in the volume with exponential variation of the proto-dark energy, satisfy the stability condition in agreement with a NLS equation of (28)- form, in which: $i \hbar \cdot (\partial\psi/\partial t) = 0$, (null variation with time of $\rho_c(r)$ by expansion or contraction).

If $p_c(r) = (\rho_c \cdot v_c)_r$ is the impulse density of the relativist quantonic component of the “dark energy”, forming the gravistar vortex: $\Gamma_G = \Gamma_\mu + \Gamma_s$ of quantons and sinergons, a

$$\rho_g(R^*) = \rho_g^0(a) \cdot (M/m_n) \cdot (a/R^*)^2 = 1837 \rho_g^c \cdot (R^*/a) \approx 1.61 \times 10^{-11} \cdot R^* [\text{kg/m}^3], \quad (72)$$

is given- for a gravitation constant $G^* \approx G$ and $\rho_g^c(a) \approx 1.24 \times 10^{-29} \text{ kg/m}^3$, (CGT), according to eq.(13):

$$F_{gs} = -2 \cdot m_c (k_h \cdot \rho_g c^2) = -2G^* (m_c M_0 / R_e^2) = -m_c c^2 / R_e; \rho_g = \frac{1}{2k_h R_e} \quad (73)$$

With $m_c = m_s$ and the values: $r_s \approx 10^{-28} \text{ m}$ and $r_h/r_s \approx 10^3$, obtained in CGT, it results that $\rho_g^0(R^*)$ necessary by eq. (73) for maintain vortexed sinergons to l_v - vortex line, is smaller than $\rho_g^{0'}(R^*)$ necessary for maintain quantons at the M_0 - core surface, for which the eq. (73), gives:

$$\rho_g^{0'} = 1/(2k_h R^*) \approx 1.8 \cdot 10^{-2}/R^*, (\rho_g^{0'} \approx (r_h/r_s^2) \rho_g^0; k_h = 27.4) \quad (74)$$

So, the M_0 hard-core is formed gradually, by quantons confining and thereafter by “dark” photons confining, the vortex Γ_c of quantons being formed after the pseudovortex Γ_s of sinergons, with the contribution of the Q_G -potential.

(14) to specific values of ratio: $(M_0/R^2) = \rho_g^0 \cdot (k_h c^2/G^*)$.

Inside a zone $\Delta R = R_0 + R_G$ of quantum equilibrium, i.e. having the entropy per quanton:

$$\varepsilon_h(r) = \gamma \cdot (k_B/\hbar) \cdot S_h(r), (S_h(r) = 2\pi r \cdot m_c c; \gamma \sim 1/M_0, [1]) \quad (75)$$

the variation of the “proto-dark energy” impulse density results- in the model, by the similitude principle, as in the electron’s case, i.e.- with exponential variation of the quantons energy (forming dark photons in the gravitic and the pseudomagnetic field of the gravistar -eq. (26)), with: $\rho_c \sim e^{-(r \cdot R^*)/\eta}$ in the zone with formed dark photons of the gravistar- considered as having the effective R_G radius, and with: $\rho_c' \sim r^{-2}$ in the outer zone, (in the evanescent part).

The dark photons were formed vortexially, according to the model, by the ξ_B vortex-tubes of the hard-core magnetic field having the induction: $B_\mu(r) \sim k_i \nabla \rho_s c$, with: $\rho_s(r) \approx \rho_c(r)$, initially being formed the vectorial photons.

These ξ_B vortex-tubes favored the negatrons forming, which are vortexially more stable than their antiparticles, explaining the spontaneous symmetry breaking in the particles genesis process and their magnetic moment anomaly:




Fig. 8. The Gravistar model.

It results that the M_0 hard-core growing increased the density of vortexed sinergons and quantons at its surface until values of “frozen” photons and of “frozen” electrons genesis: $\rho_{Av} \approx 3.7 \times 10^4 \text{ Kg/m}^3$, $\rho_{Ac} \approx 5 \times 10^{13} \text{ Kg/m}^3$, which corresponds by eq.

δm_p - mass of vortexially formed “dark” photons or of “dark” particles, is attracted until a tangential v_{pt} -speed for which the δm_p - particle remains at the same r - distance from the gravistar’s center.

This Γ_G -vortex is resulted initially as a small perturbation which may generates electronic neutrinos by quantons confinement and thereafter- massive neutrinos with own magnetic moment, given by the Γ_G - vortex, at values of the dark energy density: $\rho_c \geq \rho_a^0 = 5.17 \times 10^{13} \text{ kg/m}^3$, (equal to those of a proto-electron, i.e. -of a cold barrel-like electron).

According to CGT, it results that the force which ensures the gravistar forming is generated as in the electron’s genesis case, i.e. - by the quantum pseudomagnetic potential of the pseudo-vortex: $\Gamma_s = 2\pi r \cdot w$, ($\sqrt{2}c \geq w > c$), being given by the $Q_G(r)$ potential, (eq. (30)):

$Q_G = -\mu_c \cdot B_s(r) = -\mu_c \cdot k_1 \cdot \rho_s \cdot c = -h/2$, which maintains also quantons with $v_{ct} \approx c$ to a vortex-line $l_v = 2\pi r$, but supplemented by the gravitic force, F_{gs} .

The gravitation force F_{gs} necessary for maintain quantons with $v_c = c$ to a given vortex-line, l_v , in particular -at the surface of the star’s hard-core- considered as compact cluster of neutrons (of m_n -mass), for which:

$$(\mu_m - \mu_m^-) \sim m$$

It results also that the formed electrons gives a negative electric charge to the gravistar's kernel.

The dynamic equilibrium between the pseudomagnetic and

$$Q_G = Q_G^0 \cdot e^{-\frac{r-R^*}{\eta}} = -Q_{CF}; \Leftrightarrow \mu_v \cdot B_c(r) = \mu_v \cdot k_1 \rho_c^* c = \frac{1}{2} \cdot m_v v_f^2; \rho_c^* = \rho_c^0 \cdot e^{-\frac{r-R^*}{\eta}}; Q_G^0 = -\frac{1}{2} \cdot m_v c^2 \quad (76a)$$

with: $v_f^2 = v_0^2 \cdot e^{-(r-R^*)/\eta}$. For: $\rho_c^* = \rho_c^0(R^*) \rightarrow \rho_a^0(a) = 5.17 \times 10^{13} \text{ kg/m}^3$, it results: $B_s \rightarrow 2 \times 10^{12} \text{ T}$; $\mu_v = 3 \cdot 10^{10} \mu_h = 4 \times 10^{-36} \text{ A/m}^2$. Generally: $(v_f^2)_e = 2\mu_v k_1 \rho_c^* c / m_v$, the value of r^* resulting by eq.:

$$F_{GM} = \nabla Q_G(r^*) = -\eta^{-1} \cdot Q_G(r^*) = m_v v_f^2 / r^* = -2Q_G(r^*) / r^*; \Leftrightarrow r^* = 2\eta \quad (76b)$$

r^* being a pseudomagnetic equivalent of the gravitational Schwarzschild radius.

The values: η , ρ_c^* and r^* increases gradually, proportional with R^* , according to the sub-solitons forming condition, [10].

$$m_v v_f^2 / r_e = (m_w / m_h) \cdot (F_g(r_e) + F_{GM}(r_e)), (F_{GM}(r_e) = (-\eta^{-1} \cdot Q_G(r_e, m_h) = -(h/2\eta) e^{-(r-R^*)/\eta}) \quad (76c)$$

($F_g(r_e)$ and $F_{GM}(r_e)$ acting over a single quanton).

The equations (76) explains the gravistar hard-core gradual growing:

-in the field of the gravistar's magnetic moment: $\Gamma_\mu(r) = 2\pi r v_c$, the pseudoscalar "cold" photons and the vectorial photons with lower speed, were attracted with oriented μ_c to the M_0 - hard-core surface, where generated- at a specific ρ_A -density, cold electrons and thereafter- ultracold nucleons, formed as Bose-Einstein condensate of photons, respective- of quasi-electrons, the M_0 - hard-core becoming in time a magnetar-type star, a supernova or a rotational "black hole" with (super)dense neutronic shell, but with a density:

$$\rho_f(r) \cdot v_f^2 = \rho_s^*(r) \cdot (w - v_f)^2; \text{ with: } \rho_s^*(r) = \rho_s^0 \cdot e^{-(r-R^*)/\eta}; v_f^2 = v_0^2 \cdot e^{-(r-R^*)/\eta} \quad (77)$$

For $r \gg R^*$, eq. (77) imply: $\rho_f(r) \approx 2\rho_s^0$, i.e- a condition which may not be satisfied, according to the sub-solitons forming condition [22] which imply the necessity to ensures a negentropy value specific to the quantum equilibrium (75).

It results that the conditions (74); (77) are realised only at M_0 hard-core surface, when: $\rho_f(R^*) = \rho_s^0 (w/v_0 - 1)^2$, resulting also the condition of M_0 -hard-core growing:

$$\rho_s^0(R^*) \geq \rho_f(R^*) > \rho_s^0 (w/c - 1)^2 \approx 0.17 \cdot \rho_s^0(R^*) \quad (78)$$

The transformation of the gravistar into a "black hole" results conform to eq. (76c).

So, according to the model, if $G^* \geq G$, the gravistar hard-core which is transformed into black hole is initially a neutronic rotational star grewed initially from a superdense "seed-core" with density $\rho_n^* \rightarrow \rho_h \approx 3 \times 10^{23} \text{ kg/m}^3$.

The conclusion of the electron/proton genesis as B-E condensate of 3K-photons is sustained also by the fact that the confining temperature for the electron forming, results by B-E equation of value: $T_c \cong 3.31 \cdot \hbar^2 n^{2/3} / (m \cdot k_B) \approx 6 \times 10^{-10} \text{ K}$, for $n \approx \rho_e / m_e$, i.e. - bigger than the quantons temperature: $T_h \approx 5 \times 10^{-11} \text{ K}$.

It results that the cold genesis of "dark" photons and of "cold" elementary particles was possible in the Protouniverse's period by gravistars forming which- in this

the centrifugal potential, is realised for vortexially formed vectorial photons, m_v , with $\mu_c \uparrow \uparrow B_s$ and a tangential speed v_f according to the equations:

Adding the gravitic force, (eq.(73)), for a formed pseudoscalar photon: $m_f = (m_v - m_e)$, the radius r_e of "photon sphere" of the gravistar results by the dynamic equilibrium equation:

$$\rho_n \ll \rho_c^M = 3 \times 10^{23} \text{ kg/m}^3.$$

Thereafter, by the gravitostatic F_{gs} force, a formed "black hole" may generate in the neutronic shell, nucleons destruction with emission of γ -rays and neutrins, at: $\rho_c = \rho_s^* \gg \rho_n^0 = 4.6 \times 10^{17} \text{ kg/m}^3$, transforming the gravistar into a GRB star of magnetar type or into a quasar- by the antigravitic pseudocharge, conform to eqs. (12), (43).

The maintaining of the formed photons inside the gravistar's volume is conditioned also by a dynamic equilibrium equation on the tangential direction, similar to eq. (22), with $w = \sqrt{2c}$, i.e:

case, may explains also the supposed "big-bang" scenario of the Universe genesis by a fractalic process of multi-gravistars forming and by theirs transformation into supernovae and into (micro) quasars containing a "black hole" of "magnetar" type, in the first stage.

In a similiary way it may be explained also the Multi-universe with a structure of expansionary pseudo-bubbles, for example, in accordance with the conclusions of the Fractal cosmology, (L. Pietronero, 1987, [40]).

So, according to CGT, the Protouniverse's period had some Eras specific to:

1. the gravistars forming from gravistarc "seeds", (v_μ, v_τ - heavy neutrino with Γ_G -vortex; electronic control clusters);
2. the dark photons confining and the formation of "dark electrons";
3. the "dark particles" forming and confining; -the "atonium" states forming;
4. the forming of "black holes" and of micro/mini-quasars.

The hypothesis of an Universe's Macronucleus forming, having a macro-vortex of "dark energy", may be also sustained by the conclusion that- locally, the biggest gravistar determined the attraction of other locally formed gravistars in its magnetic field, forming a super-magnetar with super- black

hole, by the transformation of the gravistars into magnetars.

4.2.2. The Magnetar as Source of Particles Genesis and/or Acceleration

The existence of magnetars as neutronic stars converting rotational energy into magnetic energy to more than 10^{11} tesla B-induction [41] and of microquasars: sources of high energy with only 10^3 km diameter [42], sustains indirectly the previous conclusions regarding the particles cold genesis in

$$m_Y = m_e \cdot K^V \approx 5x(10^{14} \div 10^{16}) \text{ eV and } m_Z = m_p \cdot K^V \approx 9.4x(10^{17} \div 10^{19}) \text{ eV} \quad (79)$$

This theoretical result, for $m_Z \approx 9x10^{19}$ eV, explains the zetta-particles, (10^{20} - 10^{21} MeV), detected by AGASA ("Akeno Giant Air Shower Array", [43]).

The forming of supermassive particles, ($m_p > 10^{10}$ GeV/c²), in the primordial Universe is deduced also by unified gauge theories of elementary particles [44], but as formed "at hot".

According to CGT, after the gravistar's transforming into a black hole, it may obtain also- in particular conditions, a pulsatory antigravitic charge, by matter \Rightarrow energy conversion ($\sim 10\%M_0$ - for supernovae), with pulsatory emission of light, of gamma quanta and of neutrins, which may generate- in particular cases, also laser emission, as a known Eta Carinae supernova.

It is know -in this sense, that the known magnetars, such as the SGR 1806-20 magnetar, are typically soft gamma repeaters, which suggest- in CGT, also the existence of a periodically matter destruction, generating pulsatory antigravitic (pseudo)charge- according to the theory.

If the formed black hole is of rotational (Kerr) type, it is possible to exists also a very strong magnetic field, characteristic to a magnetar type star, which may be given- according to CGT, by an electric charge of the magnetar surface or by a sub-quantum and quantum macro-vortex:

$\Gamma_{BH} = \Gamma_A + \Gamma_B$ of sinergons, generating the magnetic

$$\delta p_e = m_e(v_f - v_i) = m_e a \cdot \delta t = e \cdot E_q \cdot \delta t = e \cdot \delta A; E_q = k_1 \rho_s c^2 \approx \delta A / \delta t; \delta A \approx A \quad (80)$$

with:

$$A = \frac{1}{2} B \cdot r = \frac{1}{2} k_1 \rho_B(r) \cdot c \cdot r = \frac{1}{2} k_1 \rho_c(r) \cdot v_c(r) \cdot r = \frac{1}{2} k_1 \rho_s(r) \cdot r_{\mu} c \quad (81)$$

E_q acting as a gravito-electric field over a gravito - electric e-charge and giving a final impulse:

$$p_f = m_e v_i + \delta p_e = m_e v_i \pm e \cdot A; \delta p_e = e \cdot \delta A \approx \pm e \cdot A \quad (82)$$

An identic effect is obtained also in the magnetic field of a star, but also for the electrogravitic charge, q_G , of a neutral particle- according to CGT and to the Schiff-Barnhill effect, by the force: $F_q = q_G \cdot E_q$, phenomenon that may explain the possibility of the particles accelerating in the field of a pulsar or of a „black hole” type star- considered in the actual explicative model as a particularly gravitational effect generated near the event horizon, [45].

4.2.3. Stellary Pulsating Sources

According to CGT, a strong enough „black hole” having an source of matter in its proximity, such as a matter accretion disk or a ordinary star, will generate- at the black hole's

the Protouniverse's period by gravistars forming, which indicates that the electric charge of the magnetars is a negative charge, given by negatrons.

Also, extrapolating the eq. (1) of the theory for bigger m-mass of stable/quasistable particles, it results two supermassive quasistable particles, formed in a very strong magnetic field as clusters of $\frac{1}{2}K^V$ pairs of degenerate quasi-electrons:

A-potential (eq.(10)) and the magnetic induction, B, (eq.(9), with a density variation conformed to those of a gravistar, (eq. (77)).

Relating to this case, in CGT are explained microphysically [1] also some magneto-electric and magneto-mechanic effects, such as: Einstein – De Haas, Barnett, Joffe-Kapitza, Aharonov-Böhm, Hooper-Monstein effects and the London's equations of superconductivity, by the hypothesis of the gravito-magnetic nature of the magnetic potential, A, in particular- by the conclusion that the modification of the canonic impulse of a moving electron: $\delta p_e = \pm e \cdot A$ is given as a real impulse modifying by the dynamic sinergonic pressure of the Γ_A -vortex :

$$P_s = \frac{1}{2} \rho_s c^2, [1].$$

For example, by CGT, we may conclude that the Aharonov-Böhm effect, consisting in the electron phase chagement at passing through a magnetic field of null B-induction but non-null magnetic A-potential, with an additive phases: $\delta \lambda_e = h / \delta p_e$ with $\delta p_e = \pm eA$, is done by a speed/impulse modification by a (quasi)electric field: $E_q = \delta A / \delta t$ considered as being generated by the electron entering in the field: $\delta A = \pm A$ in a time δt in which this field $E_q \sim \rho_s c^2$ acting over the electron, determines an impulse variation:

surface, a high pressure/density and matter destruction, at $T \geq 1011$ K, (close to $T_N \approx 1012 \div 1013$ K), releasing the particles' energy: mc^2 , in the form of quantonic winds and gamma and neutronic fluxes (given as pairs of controls of the released degenerate gammons which -in normal state, constitutes the N_p neutral cluster of nucleons).

Also, according to CGT, the sinergonic (etheronic) winds generated to the gammon \rightarrow neutrino conversion, (eq. (43) –when also the sinergonic vortexes of the paired electrons are reciprocally annihilated), contributes to the mean value ρ_A of the „dark energy” as fluxes of quanta of an antigravitic (pseudo)charge of the „black hole”.

Writing the electric field energy of an electron in the form:
 $\epsilon_E = \frac{1}{2} a \cdot F_e(a) = m_e c^2$, for: $F_{ea} = -e^2/4\pi\epsilon_0 a^2$ and $F_{eN} = -G \cdot m_e^2/a^2$,

$$\epsilon_G = \frac{1}{2} a \cdot F_{eN}(a) = m_e^2 G/2a; \Rightarrow \epsilon_E/\epsilon_G = \rho_a^0/\rho_g^0 = 2ac^2/m_e G = 4 \times 10^{42}, \quad (83)$$

i.e.- the gravitic field energy of the m_e -gravitic charge is of $\sim 10^{42}$ times smaller than the etheronic energy contained by the sinergonic Γ_A -vortex of the particle's magnetic moment: $\epsilon_s = m_e c^2/2$.

If the mean flux of relativist particles which are destroyed is a little higher than the critical value Φ_0^a which cancel the M_G -gravitic charge of the "black hole", the generated antigravitic charge of BH: $M_a = -k_a M_G$ with $k_a > 1$, will determine the rejection of the matter and will cancel its cause, decreasing the M_a value at $k_a < 1$, and because that the induced acceleration and deceleration of particles, corresponding to $k_a < 1$ or $k_a > 1$, is realized quickly but gradually, it results that the antigravitic charge M_a of the BH is a pulsatory, oscillating charge, which not impede its growing and not affect its stability. So, the hypothesis of a BH antigravitic charge is non-contradictory.

Also, if the central black hole is of rotational (Kerr) star of magnetar type, with a strong magneto-gravitic field: $V_{MG} \sim r^{-3}$ (eq. 10), this field can exceed the resulted antigravitic field: $V_{at} \sim r^{-1}$, under a critical limit, r_1 , continuing in this way the cause of the M_a antigravitic (pseudo)charge generation:

$$r < r_1 \Rightarrow V_{MG}(r) = V_{MG}^0(r^0/r)^3 > V_{at}^0(r^0/r)^2 \quad (84)$$

It results in this case, by CGT, that the quasars, the galactic centers, some supernovae or/and hypernovae or also some magnetars with BH, may have a pulsatory antigravitic charge-conclusion which is in concordance with some astrophysical observations, such as:

a) In the case of Quasars:

It is considered that the power of quasar results from the accretion disk of a central supermassive black holes, that are believed to exist at the core of all galaxies and which can convert on the order of 10% of the mass of an object into energy- compared to 0.7% for the p-p chain nuclear fusion. But because that even the light cannot escape from the black hole's field, it results the conclusion that the escaping energy is actually generated outside the event horizon, by gravitational stresses and immense friction on the incoming material, [46].

But this hypothesis is not enough fitted with the fact that- for create a luminosity of 10^{40} watts (the typical brightness of a quasar), a super-massive black hole must consume the material equivalent of 10 stars per year.

So, the hypothesis of CGT regarding the pulsatory antigravitic (pseudo)charge, may explain better the quasar's energy.

It is deductible also that the period τ of the radiation emission is inversely proportional wispecific to a neutronic star, th the mass of the central BH, (a bigger mass generating faster the critical pressure, p_c).

The previous explanation is in concordance also with the next astrophysical observations:

- the OVV blazars (compact quasars with optical violent

it results for the electron's (electro)gravitic energy, that:

variability) such as the quasar 3C279 has luminosity variations with 50% even in a day ;

-was observed also a periodically varying luminous quasar : PSO J334.2028 with a period of light variation of 542 days, caused by a specific mass accretion rate, [47].

-was observed quasars in pairs but also in triplets and even a quartet of quasars , [48] ;

-the X-rays emission from Cygnus X1, considered as being a "black hole", has repetitive variations, (quasi-periodic oscillations) and γ -rays emission in the very high energy band, $E > 100$ GeV, observed in the same time with the hard-X-rays, suggesting a causal link;

-there were some suggestions that quasars were made of some hitherto unknown form of stable antimatter which may explain their brightness.

b) In the case of a Hyper- or Super- nova:

Generally, it is considered that the releasing of gravitational energy by matter falling towards a massive black hole is the only process known that can produce high power continuously, in the case of quasars, but stellar explosions—supernovas and gamma-ray bursts - can do likewise, for a few weeks.

- In the case of a collapsar, i.e.-a hypernova which produces a BH after the explosion, it is considered that if the star is rotating quickly enough, then the fallback to the black hole will produce relativistic jets whose energy, transferred into the ejected shell and will render the visible outburst substantially more luminous than those of a standard supernova, the jets containing high energy particles and gamma rays and generating x-ray or gamma-ray bursts of several seconds or longer, corresponding to long-duration gamma-ray bursts; but the model do not explain the short-duration gamma-ray bursts.

-Also, in 1966 Colgate and White [49] calculated that neutrinos carry away most of the gravitational energy released by the collapse of massive stars, (the case of Type Ib and Ic and Type II supernovae) because the proton- electron combining, with the forming of a neutron and an electronic neutrino at the stars collapse, at densities specific to a neutronic star, (10^{17} kg/m³), but resulting that a second and more important neutrino source is the thermal energy, of $\sim 10^{11}$ K, of the newly formed neutron core, which is dissipated via the formation of neutrino-antineutrino pairs of all flavors, [50], the theory being confirmed in 1987, when were detected the flux of neutrino emitted by SN 1987A.

We observe that the Colgate-White theory is in concordance also with the relation (43) of CGT by which we may suppose that- at the considered very high temperatures of the neutronic matter of the collapsed star: $\sim 10^{11}$ K, (which is close to those of particles deconfining experimentally determined:

$T_d \approx 3 \times 10^{12}$ K, [30]), a quantity of neutrons is transformed into γ -rays which- by eq. (43), are partially transformed into neutrins, these radiations being characteristic especially to collapsars, to Ic-hypernovae and to GRB -sources,

(particularly, to pulsars and to magnetars).

- Is is known also- in accordance with the explicative model of CGT, that exists also GRB sources emitting short gamma-rays bursts, of about 2 sec. period, with un-clear origin and cause, because that the gamma bursts are succeeded of X-rays flashes after few minutes or few hours, suggesting the particles emission from a primary object like a neutron star swallowed by a „black hole”.

Another hypothesis considers as cause of the phenomenon a neutron star with „black hole”.

This hypothesis is closest to the CGT’s conclusion looking the generation of a pulsating antigravitic (pseudo) charge of a strong black hole according to eq. (43), by nuclear reactions at $T > 10^{11}K$.

The long and ultra-long gamma-ray bursts are considered also as effect of nuclear matter collapse, particularly-generated by a „black hole” over a „white dwarf”.

-There are also models which considers a newly formed magnetar instead of a BH, as cause of the long GRBs, [51].

This hypothesis may be, partially, in concordance with the CGT’s model because that –according to CGT, it may exist also magnetars formed by a “black hole”, particularly- of Kerr type B-H.

Also, from eq. (47) of CGT looking the particles lifetime, it results that a „black hole” must be enough „cold” for attain a density:

$\rho_{BH} > \rho_n^0 = 4.6 \times 10^{17} \text{kg/m}^3$, (of the impenetrable nucleonic volume), because that a double density: $2\rho_n^0$, for example, may be assimilated with the density of a hexa-quark particle: $3 \times \{q_a^{(2/3)} + q_b^{(-1/3)}\}$, which is un-stable even at $T \rightarrow 10^{10}K$, in accordance with eq. (47).

So, a „black hole” with $\rho_{BH} > \rho_n^0$ must be of „magnetar” type, according to CGT.

5. Conclusions

The possibility to explain the fundamental fields: gravitic, electric, magnetic and nuclear, by equations of the ideal fluids, as being generated by the impulse density of field quanta, may be considered– in our oppinion, a strong argument for the CF-prequantum model of particles of the theory, describing the fermion as ideal un-disturbed chiral soliton cluster in the ground state: $T \rightarrow 0K$, with determined parameters, in a Galileian relativity -like in the scale relativity theory of Nottale [52], which predicts- like in our theory, the natural apparition of some structures by spontaneous self-organising and the existence of a relation between the electron’s mass and charge.

At $T > 0K$, in perturbative conditions, the prequantum particle becomes quantum, as in the case of chiral soliton electron which at $T > 0K$ becomes pseudosperical by spin precession, without the change of its spin value .

The conclusion of CGT regarding the electron’s cold genesis as “freeze” photons confining in a super-strong magnetic field, is indirectly sustained also by the recent experimentally creation of a “super-photon”, i.e. a Bose-Einstein condensate of photons, which emitted yellow

light in the moment of its creation, [53].

This experimental result sustains indirectly also the revised model of photon of CGT, with inertial mass equal with its spinorial mass, in our oppinion.

Also, the possibility to “freeze” light inside a crystal, experimentally proved [54], may be explained by the revised vortexial model of photon, by the hypothesis of ring multiphotons forming in the magnetic field of the electrons or/and of the atoms, (i.e. a $n \cdot h\nu$ ring photon around an e-charge).

The theory shows also that the classic model of nucleon with incorporate electron(s) may explain also the spin and magnetic moment values by the conclusion of a density-dependent degeneration of the electron’s magnetic moment and a null spin of the electronic neutrino, the total spin conservation law being respected only in the moment of the degenerate electron releasing.

This phenomenon invalidates the Yukawa’s hypothesis of the nuclear field mesonic theory, which results as being a formal theory, without prequantum correspondent at $T \rightarrow 0K$.

The conclusion is not contradictory because that a soliton-like particle is an open system in the quantum and subquantum vacuum, according also to the particle’s thermodynamics of de Broglie and explains the fact that- at the proton transformation by K-electron capture, the electron’s spin and its magnetic moment are not transmitted with the normal S_e , μ_B , values to the formed neutron.

So, according to the theory, all elementary particles may be formed „at cold”, by a ‘gravistar’ or by a magnetar type star, in a „cascade vortex” process, as clusters of quasiaelectrons, from the „primordial dark energy” composed of etherons and quantons.

The use of a galieian-like relativity in CGT is justified also by the very low photon speeds obtained in the “stopped light” experiment, [55].

Also, the phenomenological model of expanding Universe, results in CGT with semi-sinusoidal variation of the expansion speed, with the “dark energy” which generates the expansion given by an antigravitic (pseudo)charge of stellary energetic sources with central “black hole”, given by matter \rightarrow energy conversion, with the releasing also of the sinergonic energy.

The considered structure of the particles magnetic moment, with a quantonic vortex inducing the B-induction vector by a sinergonic vortex which is the energetical cause of the magnetic potential A, may explain also microphysically the main magneto-electric and magneto-mechanic phenomena and may be verified also by new possible experiments.

A general field equation compatible with the theory is of Maxwell-Proca type:

$$(\nabla^2 - k_\lambda^2)\Phi - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \Phi = g \cdot \delta[\vec{r} - \vec{r}'(t)]$$

with $k_\lambda \approx m_q \cdot c/\hbar$; $g = -Q/\epsilon$, (m_q - the quantum mass; Q -the electric or electro-gravitic charge) –for the electro-magnetic and the gravitic field and: $k_\lambda^2 \approx (2m_{ni}/\hbar^2) \cdot V_p^0$; $g = 0$, $V_p \sim \Phi$,

with: $V_p = k_n \cdot |\Phi|^2 = V_p^0(o) \cdot e^{-r/\eta}$, for the nuclear field.

The exposed phenomenological approach of CGT does not propose an enough unitary equation of these three basic fields, but explains naturally the particles genesis and their fundamental interactions, by the same basic concept: the “proto- dark energy”, as a vortex cascade mechanism.

A more complete and accuracy approach and the set of possible validation experiments, may be the subject of a further work.

References

- [1] M.Arghirescu - ‘The Cold Genesis of Matter and Fields’, Science Publishing Group, (2015).
- [2] T.H.R. Skyrme, Proc. Roy. Soc. Series A260 (1961) 127 G.S. Adkins, ‘Chiral Soliton’, ed. by G.S. Adkins, Add. - Wesley, (1988); O.L. Battistel, Doctoral thesis, Instituto de Física da Universidad de Sao Paulo, Brazil, (1994).
- [3] E. Mottola, P.O. Mazur, ‘Gravitational Condensate stars: An Alternative to Black Holes’ arXiv: gr-qc/ 0109035 (2001).
- [4] I. Șomărescu - ‘L’électromagnetisme et la gravitoinertie en une theorie unitaire’ International Conf. of Gravitation, G.R.11, Abstract Book, Stockholm, 27 feb. (1986); E.M. Kelly - ‘Vacuum electromagnetic derived from the properties of an ideal fluid’, Nuovo Cimento, 11, 32B, March (1976).
- [5] Georges L. Lesage-‘Lucrece newtonien’, Memoires de l’Academie royale des sciences et belles lettres de Berlin, Berlin (1784) 1-28.
- [6] R. P. Lano- ‘Gravitational Meissner Effect’, arXiv: hep-th/9603077 (1996); N. I. Pallas, L. Sofonea- Book of International Congres of Mathematicians, Helsinki (1978).
- [7] M. Agop, C. Gh. Buzea, B. Ciobanu - ‘On Gravitational Shielding in Electromagnetic Fields’, arXiv: physics/9911011 (1999);
- [8] Hector A. Munera - ‘A semiclassical Model of the Photon based on Objective Reality and Containing Longitudinal Field Components’, Myron W. Evans- ‘Modern Nonlinear Optics’, Ed. John Wiley&Sons Inc., (2002) 335-385.
- [9] G. Hunter, R. L. P. Wadlinger - ‘Photons and Neutrinos as Electromagnetic Solitons’, Physics Essays, Vol. 2 (1989) 158-172.
- [10] R. M. Kiehn - ‘The Falaco Effect, A Topological Soliton’, Talk at 1987 Dynamics Days, Austin, Texas, jan. (1987); R.M. Kiehn-‘Coherent Structures in Fluids are Deformable Topological Torsion Defects’, IUTAM-SIMFLO Conference at DTU, Denmark, May 25-29, (1997)
- [11] G. Hunter, L.P. Wadlinger et. al, ‘Gravitation and Cosmology: From the Hubble Radius to the Planck Scale’, ‘Fundamental Theories of Physics’, Vol. 126, (2002), pp 157-166.
- [12] Weinberger, P., Philosophical Magazine Letters 88 (12), (2008) 897-907.
- [13] A. Cotton, H. Mouton, Ct. r. hebd. Seanc. Acad. Sci. Paris 141, (1905) 317- 349.
- [14] R. D. Chipman, L. D. Jennings, Phys. Rev. 132 (1995) 728; Greenberg, O. W., Yodh, G. B., Phys. Rev. Lett., 32 (1974) 1473.
- [15] K. N. Muhin – ‘Experimental Nuclear Physics’, Vol. 1-2, Ed. ‘Atomizdat’, Moscow (1974).
- [16] I. Sick - ‘On the rms-radius of Proton’, Phys. Lett., B 576 (2003) 62-63.
- [17] G. Adkins, C. Nappi, E. Witten - ‘Static properties of nucleons in the Skyrme model’, Nucl. Phys., B228 (1983) 552-566.
- [18] P. Constantinescu, ‘Hierarchized Systems’, Ed. Acad. R. S. R., Bucharest (1986) 106.
- [19] V. P. Oleinik - ‘Nonlinear QDE for Self-acting Electron’, Nonlinear Math. - Phys. Rev., v.4, No1-2 (1997) 180.
- [20] W. Peschka- ‘Kinetobaric Effect as possible Basis for a New Propulsion Principle’, Raumfahrtforschung, no 18 (1974).
- [21] D. L. Burke et al., ‘Positron production in multiphoton light - by light scattering’, Phys. Rev. Letters 79 (1997) 1626-1629; Physics Today 51, Feb. (1998) 17-18.
- [22] M. Brunetti et al., ‘Gamma-ray bursts of atmospheric origin in the MeV energy range’ Geophys. Research Letters 27, 1 (2000) 1595.
- [23] R. Jastrow-Phys. Review, 81 (1951) 165.
- [24] Y. Yan, R. Tegen- ‘N-N Scattering and Nucleon Quark core’, Science Asia, 27 (2001) 251.
- [25] Patent: US2482773.
- [26] Dan Rădulescu - Bulletin de la Société des Sciences, Cluj (1922).
- [27] The OPERA Collaboration (2011), ‘Measurement of the neutrino velocity with the OPERA detector in the CNGS beam’, ar Xiv: 1109.4897v1.
- [28] L. Wolfenstein- Phys. Lett., B 107 (1981) 77.
- [29] Shull Michael, Stern Alan, Astronomical Journ., Vol. 109, No2 (1995) 690-697.
- [30] Science et Avenir, No. 555, may (1993).
- [31] B. Rossi, D. B. Hall - Phys. Rev., 59 (1941) 223.
- [32] T. Schmidt, Z. Phys., 106 (1937) 358.
- [33] T. Lonnroth, Il Nuovo Cimento 110A (1997) 961.
- [34] W. Bauer, Proc. of 1st Catania Relativistic Ion Studies, Asicastello, Italy (1996) 23.
- [35] A. Săndulescu et. al - Nucl. Physics., 48 (1963) 345.
- [36] Shaw P. E., Davy N., Phys. Rev., Vol. 21, Ser. 2, Iss. 6 (1923) 680-681.
- [37] J. Kristian, A. Sandage, J. Westphal - Astrophys. Journal, 221 (1978) 383.
- [38] W. L. Freedman et al., Astrophys. J. 704 (2009) 1036; arXiv: 0907.4524.
- [39] C. F. C. Brandt, R. Chan, M. F. A. da Silva, P. Rocha, ‘Gravastar with an Interior Dark Energy Fluid and an Exterior de Sitter-Schwarzschild Spacetime’, arXiv: gr-qc/1012.1233v1 (2010).
- [40] Pietronero, L. ‘The Fractal Structure of the Universe: Correlations of Galaxies and Clusters’, Physica A (144): 257, (1987).

- [41] C. Kouveliotou, R. C. Duncan, C. Thompson, 'Magnetars', *Scientific American Rev.*, Febr. (2003) 237.
- [42] I. F. Mirabel- 'Microquasars as sources of high energy phenomena' arXiv: astro-ph/0211085 v1, 5 Nov. (2002).
- [43] 'Scientific American', January, (1999)
- [44] Polnarev, A. G., Khlopov, M. Y. – 'Primordial Black Holes and the Era of Superheavy Particle Dominance in the Early Universe', *Soviet Astronomy Journ.*, V.25 (1981) 406.
- [45] O.B. Zaslavskii, 'Acceleration of particles as universal property of rotating black holes', arXiv: 1007.3678v2 [gr-qc] 15 oct. 2010; O.B. Zaslavskii, 'Acceleration of particles as universal property of rotating black holes', arXiv: 1007.3678v2 [gr-qc] 15 oct. 2010
- [46] Thomsen, D. E.- 'End of the World: You Won't Feel a Thing', *Science News (Society for Science & the Public)*, Jun 20, (1987), 131 (25):391,
- [47] Tingting Liu et al., 'A Periodically varying Luminous Quasar at $z = 2$ from the Pan-Starrs1, Medium Deep Survey: A Candidate Supermassive Black Hole Binary in the Gravitational wave-Driven regime', arXiv:1503.02083v1 [astro-ph. HE]
- [48] Irene, K.-, 'Surprise Quasar Quartet Defies Explanation', *news.discovery.com.*, 14 may, (2015)
- [49] S. A. Colgate & R. H. White- 'The Hydrodynamic Behavior of Supernova Explosions', *The Astrophysical Journal*, 143 (1966) 626 ; doi:10.1086/148549
- [50] A.K. Mann (1997) *Shadow of a star: The neutrino story of Supernova 1987A.* W. H. Freeman. p. 122.ISBN 0-7167-3097-9
- [51] Metzger, B. et al.- 'Proto-Neutron Star Winds, Magnetar Birth, and Gamma-Ray Bursts'. *AIP Conference Proceedings* „SUPERNOVA 1987A: 20 YEARS AFTER”, 937, (2007), pp. 521–525.
- [52] L. Nottale, *Scale Relativity and Fractal Space-Time: Applications to Quantum Physics, Cosmology and Chaotic Systems*, *Chaos, Solitons& Fractals*, Vol.7, No.6 (1996) 887-938.
- [53] Klaers, J., Schmitt, J., Vewinger, F., Weitz, M., *Nature*, 468, 546-548 (2010); Zeeya Merali, 'Chilled light enters a new phase', *Nature*, 24 Nov. (2010).
- [54] Heinze, G, Hubrich, C., Halfmann T., 'Stopped Light and Image Storage by Electromagnetically Induced Transparency up to the Regime of One Minute', *Phys. Rev. Lett.* 111, 038103 (2013);
- [55] L. V. Hau, S. E. Harris, Z. Dutton, C. H. Behroozi, "Ultra-slow, stopped, and compressed light in Bose-Einstein condensates gas", *Nature* 397, 594, (1999); S. E. Harris and L. V. Hau, *Phys. Rev. Lett.* 82, 4611 (1999).