

# Sino-Orbital Cellulitis Due to HSV Type I

Selin Yetkinel<sup>1</sup>, Semai Bek<sup>2</sup>

<sup>1</sup>Agri State Hospital, Department of Neurology, Agri, Turkey

<sup>2</sup>Baskent University Faculty of Medicine, Department of Neurology, Adana Teaching and Medical Research Center, Adana, Turkey

## Email address:

selinyetkinel@gmail.com (S. Yetkinel), semaibek@yahoo.com (S. Bek)

## To cite this article:

Selin Yetkinel, Semai Bek. Sino-Orbital Cellulitis Due to HSV Type I. *International Journal of Ophthalmology & Visual Science*.

Vol. 2, No. 2, 2017, pp. 55-58. doi: 10.11648/j.ijovs.20170202.13

Received: January 24, 2017; Accepted: April 12, 2017; Published: May 17, 2017

**Abstract:** Orbital pseudotumor, is a heterogeneous group of disorders characterised by orbital inflammation and is a rare clinical entity and a diagnosis of exclusion. We report two cases with typical clinical manifestation and magnetic resonance images, but unresponsive to corticosteroid treatment which turned out to be caused by herpes simplex virus type I. Viral etiology must be carefully evaluated, especially in steroid unresponsive recurrent cases presumed to be pseudotumor orbita.

**Keywords:** Orbital Cellulitis, Pseudotumor Orbita, Herpes Simplex Virus, Painful Ophthalmoplegia

## 1. Introduction

Painful ophthalmoplegia is a common clinical presentation for all neurologists and ophthalmologists. There is a long list of differential diagnosis, including vascular, infective, rheumatologic causes and ophthalmologic migraine [1]. Despite improvements in laboratory and scanning techniques, most cases interpreted idiopathic. Extraocular orbital and adnexal inflammation with no identifiable local or systemic cause is often referred to as orbital pseudotumor. It remains a diagnosis of exclusion and is characterized by its chronicity, anatomic location, or histologic subtype.

Corticosteroids are the mainstay of treatment inducing a rapid and dramatic reversal of inflammatory findings. Their mechanism of action is attributable to both anti-inflammatory and immunosuppressive properties. Despite the excellent and rapid response to corticosteroids, failures reported and unusual etiologic causes should be interpreted recurrent steroid unresponsive cases [2]. In this paper we present two cases of sino-orbital cellulites presumed to be orbital pseudotumor, unresponsive to corticosteroid treatment, due to herpes simplex virus type I.

## 2. Case Report

Case 1: A 67-year-old male admitted to our department with major complaints of orbito-frontal pain, ptosis and vision loss on the left side. His complaints began 25 days before admission with diplopia and blurred vision and got

worse gradually. As left sided ptosis and vision loss occurred two days ago, he decided to seek medical assistance. Type II diabetes mellitus was under control with oral antidiabetics for 8 years. His neurological examination revealed complete loss of function in the left II-III-IV and VI cranial nerves. Initial Snellen visual acuity was light perception and relative afferent papillary defect was detected in the affected eye. Retinal examination did not reveal diabetic retinopathy. The examination of the fellow eye was fully normal.




Figure 1. Total ophthalmoplegia on the left side (Case 2).


Case 2: A 51-year-old male admitted to our department with the complaints of orbito-frontal pain, diplopia and ptosis on the left side lasting for three days. His medical history

revealed that he had same symptoms on the left side in 1985 and 1995, and on the right side in 2003. He claimed that his previous complaints had resolved in 2-3 months without any treatment. Neurological examination revealed complete loss of function in left III-IV-VI cranial nerves "Figure 1". Biomicroscopic and fundoscopic examination were fully normal in both eyes. Relative afferent papillary defect was not detected.

Both patients treated with methylprednisolone 1000 mg/day IV for 5 days but no clinical improvement was observed. Biochemical (blood urea nitrogen, creatinine, transaminases, sodium, potassium, erythrocyte sedimentation rate, thyroid stimulating hormone, free T4, angiotensin-converting enzyme level, serum protein electrophoresis), rheumatologic (anti-nuclear antibodies, anti-cardiolipin antibodies, anti-dsDNA, antineutrophil cytoplasmic antibodies), cerebrospinal fluid (CSF) analyses (protein, glucose, microscopy), cerebral magnetic resonance imaging (MRI) and angiography performed for possible etiological causes. Mild elevated CSF protein and leucocytosis was evident. Left sided proptosis was the only pathological finding in both patients. Fatsat turbo spin echo orbital MRI performed to demonstrate possible inflammatory or mass lesion. Orbital MRI revealed contrast enhancement expanding from posteromedial sight of periorbital adipose tissue to optic nerve and cavernous sinus in the first case "Figure 2". Contrast enhancement expanded from retroorbital adipose tissue to maxillary sinus anterior wall in the second case "Figure 3". Preliminary diagnoses were pseudotumor orbita. Biopsy was planned to rule out possible malignancies.



**Figure 2.** Fatsat turbo spin echo MR showed contrast enhancement, spreading from posteromedial sight of periorbital adipose tissue to optic nerve and cavernous sinus (Case 1).



**Figure 3.** Fatsat turbo spin echo MR showed contrast enhancements, spreading from retroorbital adipose tissue to maxillary sinus anterior wall (Case 2).

Neurosurgery and otorhinolaryngology departments together, performed intranasal endoscopic surgery and cytological specimen were obtained from inflammatory tissue passing through lamina papricea. Pathology department reported focal adipose necrosis, chronic inflammatory adipose tissue and findings of mild myositis for both samples. This material was subjected to microscopy, cultures for viruses, microorganisms, and for polymerase chain reaction (PCR) techniques for cytomegalovirus, Epstein-Barr virus, herpes simplex virus (HSV), varicella zoster virus (VZV), and toxoplasmosis. The only test that was positive was PCR for HSV. Sequencing revealed HSV type 1 (all analyses were performed twice).

Patients' symptoms slightly improved after surgery probably due to decompression effect. Both patients were treated with acyclovir 800mg/qid PO for 5 days. The follow up examination revealed that first patient's symptoms regressed partially in three months. Relative afferent papillary defect persisted however visual acuity improved to counting fingers from one meter. Complete recovery was seen in the second case in a month.

### 3. Conclusion

Although the diagnosis of orbital pseudotumor is usually a clinical one, the evaluation should begin with an imaging study. High-resolution orbital computed tomography (CT) may demonstrate soft tissue swelling but is not as sensitive as MRI. Contrast-enhanced MRI with multiple coronal views and fat saturation should be the initial diagnostic study performed, when available [3]. Orbital pseudotumor MRI studies revealed mass lesions that were hypointense to orbital fat on T1-weighted images and isointense or minimally hyperintense to fat on T2-weighted images [4]. Even rarely, orbital pseudotumor may spread to extraorbital areas as seen in our cases with cavernous sinus involvement in the first case and with maxillary sinus involvement in the second case [5, 6, 7].

The etiology of orbital pseudotumor is not clear, but infections (fungal, bacterial, viral), inflammatory syndromes (idiopathic orbital inflammation, idiopathic sclerosing inflammation, Sarcoidosis), autoimmune disorder (Graves' ophthalmopathy, vasculitis), neoplastic processes (primary orbital tumors, metastatic tumors), vascular lesions (carotid cavernous fistulas, cavernous sinus thrombosis) and aberrant wound healings have been put forward as possibilities. In addition orbital pseudotumor has been associated with systemic disorders, including rheumatoid arthritis, Crohn's disease and systemic lupus erythematosus [1, 8-12].

The diagnostic evaluation of a patient with suspected orbital pseudotumor should include a full hematologic work-up with complete blood cell count, electrolytes, sedimentation rate, antinuclear antibodies, anti-dsDNA, antineutrophil cytoplasmic antibodies, angiotensin-converting enzyme level, thyroid stimulating hormone, free T4 and serum protein electrophoresis [2, 13]. Cerebrospinal fluid examination is rarely helpful but should be performed

in questionable cases or when cytology is deemed useful to exclude lymphoma [3]. It is crucial to differentiate between orbital pseudotumor and the disorders that may mimic it. We also performed this extended laboratory work up for our cases but only mild elevated CSF protein and leucocytosis was evident. These laboratory findings did not help in differential diagnosis.

Response to therapy often confirms the diagnosis but in our cases corticosteroid treatment was unresponsive. Biopsy is only indicated if there are rapidly progressive neurologic deficits, a lack of steroid responsiveness, or persistent abnormalities on neuroimaging studies [2]. Chronic inflammatory cells infiltration in various orbital tissues are the main histopathologic features in patients with orbital inflammatory pseudotumor [9]. Chronic inflammation observed in our specimens did not yield any etiological idea. Although the clinical examination and MRI findings were confirmative for orbital pseudo tumor, classical laboratory work up was insufficient to put forward the etiology. Virology study of biopsy specimen was surprising which revealed HSV-DNA was present in both cases, establishing the diagnosis of sino-orbital cellulitis.

Systemic corticosteroids - in the form of oral or intravenous administration- are generally considered mainstay therapy for orbital pseudotumors. Recurrences even after initial corticosteroid treatment have been reported in orbital pseudotumor [15, 16, 17]. Spontaneous remission may also occur [18]. There is a similarity with peripheral facial paralyses because they share the same clinical features; cranial nerve palsy, often dramatic response to corticosteroid therapy, spontaneous remission, usually complete improvement in 2-3 months. MRI findings resemble with different localization [19]. Recent studies support viral etiology and antiviral therapy for Bell's paralyses, sustained as idiopathic perennial [20]. Probable viral etiology seems reasonable with this available data. Ophthalmoplegia associated with herpes virus family, mainly herpes zoster, is not uncommon. As the majority were due to involvement of the ocular motor cranial nerves, they are traditionally interpreted as diseases of III, IV or VI cranial nerves [21, 22, 23, 24]. Ophthalmoplegia due to viral myositis have also been reported [25, 26, 27]. To our knowledge, no prior cases of HSV-induced sino-orbital cellulitis with multiple cranial nerve palsies have been described in the literature. After antiviral treatment, complete resolution was observed in the second case. Insufficient response observed in the first case was probably due to late timing of the proper treatment. There is no particular well-established protocol to guide adjuvant therapy for refractory orbital pseudotumor. Among the available options for the management of refractory orbital pseudotumor is: radiation therapy, cytotoxic agents (Cyclophosphamide and Chlorambucil), immunosuppressants (Methotrexate, Cyclosporine, Azathioprine), IV immunoglobulins, TNF-alpha inhibitor, monoclonal antibody (Infliximab and Adalimumab) and Mycophenolate Mofetil which inhibit denovo purine synthesis and prevent B & T lymphocyte replication [1, 2, 9, 28-32].

Orbital pseudotumor is an unusual inflammatory condition established by typical clinical and imaging findings. Orbital pseudotumor is usually quite responsive to corticosteroids. Corticosteroid unresponsive cases, especially with recurrences, should be interpreted carefully as this type of course is unusual. Biopsy is not always necessary, but it might be helpful to put forward the specific cause. Before initiating any treatment, atypical clinical or imaging findings should prompt biopsy. Although evidence is limited, antiviral therapy should be kept in mind for these atypical cases. These cases showed that we should hesitate in pronouncing the diagnosis of "pseudotumor" before viral evaluation of biopsy specimen if possible.

---

## References

- [1] Mombaerts I, Goldschmeding R, Schlingemann RO, Koornneef AL. What is orbital pseudotumor? *Surv Ophthalmol* 1996; 41: 66-78.
- [2] Rubin PA, Foster CS. Etiology and management of idiopathic orbital inflammation. *Am J Ophthalmol* 2004; 138: 1041-3.
- [3] Kline LB, Hoyt WF. The Tolosa-Hunt syndrome. *J Neurol Neurosurg Psychiatry* 2001; 71: 577-82.
- [4] Atlas SW, Grossman RI, Savino PJ, Sergott RC, Schatz NJ, Bosley TM, Hackney DB, Goldberg HI, Bilaniuk LT, Zimmerman RA. Surface-coil MR of orbital pseudotumor. *AJR Am J Roentgenol* 1987; 148: 803-8.
- [5] Eshaghian J, Anderson RL. Sinus involvement in inflammatory orbital pseudotumor. *Arch Ophthalmol* 1981; 99: 627-30.
- [6] Ishikawa S, Yahikozawa H, Yamazaki M, Kikkawa M, Ikeda S, Hanyu N. Recurrent steroid-responsive trismus and painful ophthalmoplegia. *J Neurol Sci* 1999; 170: 69-71.
- [7] Leibovitch I, Goldberg RA, Selva D. Paranasal sinus inflammation and non-specific orbital inflammatory syndrome: an uncommon association. *Graefes Arch Clin Exp Ophthalmol* 2006; 244: 1391-7.
- [8] Yuen SJ, Rubin PA. Idiopathic orbital inflammation: distribution, clinical features, and treatment outcome. *Arch Ophthalmol*. 2003; 121: 491-499.
- [9] Harris GJ. Idiopathic orbital inflammation: a pathogenetic construct and treatment strategy: The 2005 ASOPRS Foundation Lecture. *Ophthal Plast Reconstr Surg*. 2006; 22: 79-86.
- [10] Brunelle C, Bally S, Provencal J, Lecomte C. Orbital inflammatory pseudotumor secondary to sarcoidosis. *J Fr Ophthalmol*. 2016 Sep; 39 (7): e179-81.
- [11] Leo M, Maggi F, Dottore GR, Casini G, Mazzetti P, Pistello M, Sellari-Franceschini S, Nardi M, Vitti P, Marcocci C, Marinò M. Graves' orbitopathy, idiopathic orbital inflammatory pseudotumor and Epstein-Barr virus infection: a serological and molecular study. *J Endocrinol Invest*. 2016 Dec 16.
- [12] Nakamura A, Miyamura T, Suematsu E. Orbital Inflammatory Pseudotumor Associated with Rheumatoid Arthritis. *Intern Med*. 2016; 55 (11): 1503-4.

- [13] Jacobs D, Galetta S. Diagnosis and management of orbital pseudotumor. *Curr Opin Ophthalmol* 2002; 13: 347-51.
- [14] Yan J, Li Y, Qiu H, Lu G, Wu Z, Lin J, Zhang W. Immunohistochemical study of the presence of mast cells in idiopathic orbital inflammatory pseudotumor: possible role of mast cells in the course of its pathogenesis. *Int Ophthalmol* 2007; 27: 235-9.
- [15] Mombaerts I, Schlingemann RO, Goldschmeding R, Koornneef L. Are systemic corticosteroids useful in the management of orbital pseudotumors? *Ophthalmology* 1996; 103: 521-8.
- [16] Swamy BN, McCluskey P, Nemet A, Crouch R, Martin P, Benger R, Ghabriel R, Wakefield D. Idiopathic orbital inflammatory syndrome: clinical features and treatment outcomes. *Br J Ophthalmol* 2007; 91: 1667-70.
- [17] Wasmeier C, Pfadenhauer K, Rösler A. Idiopathic inflammatory pseudotumor of the orbit and Tolosa-Hunt syndrome--are they the same disease? *J Neurol* 2002; 249: 1237-41.
- [18] Chavis RM, Garner A, Wright JE. Inflammatory orbital pseudotumor. A clinicopathologic study. *Arch Ophthalmol* 1978; 96: 1817-22.
- [19] Kim IS, Shin SH, Kim J, Lee WS, Lee HK. Correlation between MRI and operative findings in Bell's palsy and Ramsay Hunt syndrome. *Yonsei Med J* 2007; 48: 963-8.
- [20] Stjernquist-Desatnik A, Skoog E, Aurelius E. Detection of herpes simplex and varicella-zoster viruses in patients with Bell's palsy by the polymerase chain reaction technique. *Ann Otol Rhinol Laryngol* 2006; 115: 306-11.
- [21] Carroll WM, Mastaglia FL. Optic neuropathy and ophthalmoplegia in herpes zoster oticus. *Neurology* 1979; 29: 726-9.
- [22] Marsh RJ, Cooper M. Ophthalmic herpes zoster. *Eye* 1993; 7: 350-70.
- [23] Pavan-Langston D. Herpes zoster ophthalmicus. *Neurology* 1995; 45: 50-1.
- [24] Womack LW, Liesegang TJ. Complications of herpes zoster ophthalmicus. *Arch Ophthalmol* 1983; 101: 42-5.
- [25] Kawasaki A, Borruat F-X. An unusual presentation of herpes zoster ophthalmicus: orbital myositis preceding vesicular eruption. *Am J Ophthalmol* 2003; 136: 574-5.
- [26] Krasnianski M, Sievert M, Bau V, Zierz S. External ophthalmoplegia due to ocular myositis in a patient with ophthalmic herpes zoster. *Neuromuscular Disorders* 2004; 14: 438 - 441.
- [27] Volpe NJ, Shore JW. Orbital myositis associated with herpes zoster. *Arch Ophthalmol* 1991; 109: 471-2.
- [28] Faktorich EG, Tucker N, O'Brien JM. Thyroid ophthalmopathy. *Curr Opin Ophthalmol*. 1994; 5: 45-52.
- [29] On AV, Hirschbein MJ, Williams HJ, Karesh JW. CyberKnife radiosurgery and rituximab in the successful management of sclerosing idiopathic orbital inflammatory disease. *Ophthalm Plast Reconstr Surg*. 2006; 22: 395-397. [PubMed].
- [30] Garrity JA, Coleman AW, Matteson EL, Eggenberger ER, Waitzman DM. Treatment of recalcitrant idiopathic orbital inflammation (chronic orbital myositis) with infliximab. *Am J Ophthalmol*. 2004; 138: 925-930. [PubMed].
- [31] Hatton MP, Rubin PA, Foster CS. Successful treatment of idiopathic orbital inflammation with mycophenolate mofetil. *Am J Ophthalmol*. 2005; 140: 916-918. [PubMed].
- [32] Wilson MW, Shergy WJ, Haik BG. Infliximab in the treatment of recalcitrant idiopathic orbital inflammation. *Ophthalm Plast Reconstr Surg*. 2004; 20: 381-383.