

Review Article

***Opuntia ficus-indica* as a Source of Bioactive and Nutritional Phytochemicals**

Imen Belhadj Slimen^{1,2,*}, Taha Najjar^{1,2}, Manef Abderrabba²

¹Department of Animal, Food and Halieutic Resources, National Agronomic Institute of Tunisia, Mahragen City, Tunisia

²Laboratory of Materials, Molecules and Applications, Preparatory Institute for Scientific and Technical Studies, La Marsa, Tunisia

Email address:

belhadj_slimen_imen@yahoo.fr (I. B. Slimen)

*Corresponding author

To cite this article:

Imen Belhadj Slimen, Taha Najjar, Manef Abderrabba. *Opuntia ficus-indica* as a Source of Bioactive and Nutritional Phytochemicals. *Journal of Food and Nutrition Sciences*. Vol. 4, No. 6, 2016, pp. 162-169. doi: 10.11648/j.jfns.20160406.14

Received: November 19, 2016; **Accepted:** November 30, 2016; **Published:** December 23, 2016

Abstract: Cacti are known for their minimum water requirement. They grow extensively in arid lands, where they were traditionally used for both human and animal food. *Opuntia ficus-indica*, commonly referred to as prickly pear or nopal cactus, was known as a medicinal plant, owing to its rich composition in polyphenols, vitamins, polyunsaturated fatty acids and amino acids. This makes cactus pear a promising crop for commercial food applications. Recent scientific investigations showed that cactus products may be efficiently used as a source of food additives, mainly fibre, colorants and antioxidants. Tablets, cookies and other forms of fibre derived from cactus cladodes are currently marketed in several American countries. This review details the main functional phytochemicals characterizing different tissues of *Opuntia ficus-indica*.

Keywords: *Opuntia ficus-indica*, Cladodes, Antioxidants, Polyphenols, Betalains

1. Introduction

Cacti are the most conspicuous and characteristic plants of arid and semi-arid regions. *Opuntia ficus-indica* (L.) Mill., commonly called prickly pear or nopal cactus, belongs to the dicotyledonous angiosperm *Cactaceae* family which includes about 1500 species of cactus. Cacti are known for their ability to thrive under environments recognized as stressful for most plant species, and are widely used to prevent soil erosion and to combat desertification [1, 2].

Opuntias have been exploited as a cheap and alternate source of food suitable not only for humans but also for animals. In addition, they have been cultivated as ornamental crops [3]. Two parts of the plant have been used for food: the “nopal” or cladodes and the fruits or the prickly pears. Cladodes are consumed in Mexico as salads [4] whereas fruits are widely eaten fresh, dried or preserved in jams, syrups or processed into candy-like products [4, 5]. *Opuntia* fruits are fleshy and elongated berries, varying in shape, size and color (orange, yellow, red, purple, green, white) and have a consistent number of hard seeds [6].

Opuntia ficus-indica was known to contain several pigments and bioactive molecules having nutritional and medicinal desirable properties [7-17]. Based on the chemical structure of their chromophore, pigments can be classified into (a) Chromophores with conjugated systems, such as carotenoids, anthocyanins, betalains, caramel, synthetic pigments, lakes; and (b) Metal-coordinated porphyrins including myoglobin, chlorophyll, and their derivatives.

Based on this backdrop, the main objective of the present review is to focus on the bioactive molecules from *Opuntia ficus-indica*, their structure-activity relationship, as well as the nutritional value of this plant.

2. Bioactive Phytochemicals and Their Antioxidant Activity

Opuntia ficus-indica was known to be a valuable source of vitamin E, fibers, amino acids, minerals, and antioxidant

molecules (ascorbic acid, flavonoids, carotenoids, betacyanins and betaxanthins) [18-21]. Cactus peel and seeds can be used to prepare cactus oil, peel lipids being enriched in essential fatty acids and liposoluble antioxidants [22]. Cladodes contain vitamins, antioxidants and various flavonoids [23, 24]. Fruits and skin are enriched in betacyanins and betaxanthins [25, 26].

2.1. Polyphenols

“Polyphenols” (or phenolic compounds) is a generic term that refers to more than 8000 compounds widely dispersed throughout the plant kingdom [27]. Polyphenols can be divided into four main classes: flavonoids, phenolic acids, stilbenes, and finally lignan and suberin. The basic structure of some *Opuntia* polyphenols is presented in Figure 1.




Figure 1. Basic structures of some polyphenols from *Opuntia* (a) Basic structure of flavonoids, (b) Basic structure of flavonols, (c) Basic structure of flavones, (d) Basic structure of hydroxycinnamic acids.

Polyphenols are present in different *Opuntia* tissues at various concentrations, as detailed in Table 1.

Table 1. Polyphenol contents in various parts of *Opuntia ficus-indica*.

| Plant part | Molecule | Content (mg/100g) | References |
|------------|---|-------------------|-----------------|
| Flowers | Gallic acid | 1630–4900 | [28-30] |
| | Quercetin 3- <i>O</i> -Rutinoside | 709 | |
| | Kaempferol 3- <i>O</i> -Rutinoside | 400 | |
| | Quercetin 3- <i>O</i> -Glucoside | 447 | |
| | Isorhamnetin 3- <i>O</i> -Robinobioside | 4269 | |
| | Isorhamnetin 3- <i>O</i> -Galactoside | 979 | |
| | Isorhamnetin 3- <i>O</i> -Glucoside | 724 | |
| | Kaempferol 3- <i>O</i> -Arabinoside | 324 | |
| | Total phenolic acid | 48–89 | |
| | Feruloyl-sucrose isomer 1 | 7.36–17.62 | |
| Seeds | Feruloyl-sucrose isomer 2 | 2.9–17.1 | [31] |
| | Sinapoyl-diglucoside | 12.6–23.4 | |
| | Total Flavonoids | 1.5–2.6 | |
| | Total Tannins | 4.1–6.6 | |
| | Total phenolic acid | 45,700 | |
| Peel | Total Flavonoid | 6.95 | [32-34] |
| | Kaempferol | 0.22 | |
| | Quercetin | 4.32 | |
| | Isorhamnetin | 2.41–91 | |
| | Gallic acid | 0.64–2.37 | |
| | Coumaric | 14.08–16.18 | |
| | 3,4-dihydroxybenzoic | 0.06–5.02 | |
| | 4-hydroxybenzoic | 0.5–4.72 | |
| | Ferulic acid | 0.56–34.77 | |
| | Salicylic acid | 0.58–3.54 | |
| Cladodes | Isoquercetin | 2.29–39.67 | [35-38] |
| | Isorhamnetin-3- <i>O</i> -glucoside | 4.59–32.21 | |
| | Nicotiflorin | 2.89–146.5 | |
| | Rutin | 2.36–26.17 | |
| | Narcissin | 14.69–137.1 | |
| | Total phenolic acid | 218.8 | |
| | Quercetin | 9 | |
| | Isorhamnetin | 4.94 | |
| | Kaempferol | 0.78 | |
| | Luteolin | 0.84 | |
| Fruits | isorhamnetin glycosides | 50.6 | [5, 36, 39, 40] |
| | Kaempferol | 2.7 | |

Flavonoids and phenolic acids are the main polyphenols of *Opuntia ficus-indica*. Flavonoids are known for their antioxidant activity. They are able to interact with lipids, proteins and carbohydrates to inhibit their oxidation [41]. Flavonoids can protect from injury caused by free radicals in various ways. One way is the direct scavenging of free radicals according to the following equation (1):



Quercetin and silibin inhibit xanthine oxidase and cytochrome activity, thereby resulting in decreased oxidative injury [42, 43]. Another possible mechanism by which flavonoids act is through interaction with various enzyme systems. When reactive oxygen species are in the presence of iron, lipid peroxidation results. Specific flavonoids, such as quercetin, are known for their iron-chelating and iron-stabilizing properties [44]. Direct inhibition of lipid peroxidation is another protective measure [45]. Another interesting effect of flavonoids on enzyme systems is the inhibition of the metabolism of arachidonic acid [46]. This feature gives flavonoids anti-inflammatory and anti-thrombogenic properties.

2.2. Carotenoids

Carotenoids are lipid-soluble C40 tetraterpenoids synthesized by plants, algae, fungi, yeasts and bacteria. The majority carotenoids are derived from a 40- carbon polyene chain, which could be considered the backbone of the molecule. *Opuntia* carotenoids are generally formed from eight C5 isoprenoid units joined head to tail, except at the center, where a tail-to-tail linkage reverses the order and results in a symmetrical molecule [47]. According to their structure, carotenoids can be classified into (a) carotenes or hydrocarbon carotenoids (such as β -carotene) which contain only carbon and hydrogen atoms (Figure 2), and (b) oxygenated carotenoids which are derivatives of these hydrocarbons or xanthophylls which carry at least one oxygen atom (such as zeaxanthin, lutein, spirilloxanthin, echinenone, and antheraxanthin) [48].

In cladodes, three carotenoids were identified: lutein, β -carotene and α -cryptoxanthin [49]. In young cladodes, concentrations vary between 0.047 and 0.077 mg/100 g [50]. In fruits, carotenoid content ranges from 1.77 to 2.65 mg eq. β -carotene/100 g. The highest values were recorded in the cultivars having orange color [51, 52]. In the peel, total carotenoids content reaches 2.97 mg/100 g [53].




Figure 2. Chemical structure of β -carotene.


Xanthophylls and carotenes were proved to be efficient quenchers of singlet oxygen ($^1\text{O}_2$) and peroxy radicals [54, 55].

This physical quenching is related to the number of conjugated double bonds present in the molecule which determines their lowest triplet energy level. β -Carotene, zeaxanthin, cryptoxanthin, as well as α -carotene were reported to belong to the group of highly active quenchers of $^1\text{O}_2$ since they have triplet energy levels close to that of this radical, thus enabling energy transfer. The most efficient carotenoid is the open ring carotenoid lycopene [56, 57]. Due to their lipophilicity and specific property to scavenge peroxy radicals, carotenoids are thought to play an important role in the protection of cellular membranes and lipoproteins against oxidative damage [58]. They are able to deactivate peroxy radicals and to form a resonance stabilized carbon-centered radical.

2.3. Betalains

Betalains are vacuolar pigments composed of a nitrogenous core structure, betalamic acid. Betalamic acid condenses with imino compounds (*cyclo*-DOPA/ its glucosyl derivatives) or amino acids/derivatives to form violet betacyanins and yellow betaxanthins. Betalains are present in the pulp and the peel of *Opuntia ficus-indica*. Betacyanins and betaxanthins concentrations vary according to the color of the fruit. In addition to neobetanin, betanin, isobetanin, betanidin and indicaxanthin are present in the pulp of *Opuntia* fruits [59-61]. Betanin and indicaxanthin were detected in the peel [61]. Gomphrenin I, portulacaxanthin I, portulacaxanthin III, muscaaurin, (*S*)-serine-betaxanthin, (*S*)-valine-betaxanthin, (*S*)-isoleucine-betaxanthin, (*S*)-Phenylalanine-betaxanthin, Vulgaxanthin I, Vulgaxanthin II, Vulgaxanthin IV and Miraxanthin II were also reported [62-65].

Betalains due their antioxidant activity to their phenolic hydroxy groups (Equation 2) in addition to their imino and tetrahydropyridine groups. The common electronic resonance system supported between the two nitrogen atoms allows creating a stable carbocation upon an electron abstraction.



The redox potential of betanin and indicaxanthin were evaluated respectively at 0.4 V and 0.6 V, indicating that these two pigments are able to donate easily their electrons. At very low concentrations, betanin was demonstrated to inhibit lipid peroxidation and heme decomposition [66]. Betanidin was the most potent antioxidant against peroxy radical and nitric oxide [67]. Moreover, indicaxanthin was shown to be less effective than betanin in radical scavenging reactions [68]. Betanin was also shown to act as an oxidation retarder, and additive effects with α -tocopherol were reported for both betanin and indicaxanthin [69, 70].

3. Vitamins

Opuntia ficus-indica contains considerable amounts of vitamins, mainly ascorbic acid, vitamin B, and α -tocopherol (Table 2). Their concentrations vary among the different plant tissues.

Table 2. Vitamins content in different parts of *O. ficus-indica* (mg/100 g).

| | Pulp | Peels | Cladodes | References |
|----------------------|-------|-------|----------|-------------|
| Ascorbic acid | 1-48 | 59.82 | 7-22 | [6, 71, 53] |
| α -Tocopherol | 84.9 | 1760 | 1.76 | [71,72,73] |
| β -Tocopherol | 12.6 | 222 | NA | [71] |
| γ -Tocopherol | 7.9 | 174 | NA | [71] |
| σ -Tocopherol | 422 | 26 | NA | [71] |
| Total tocopherols | 527.4 | 2182 | 2.18 | [72, 73] |
| Thiamine | NA | NA | 0.14 | [71] |
| Riboflavine | NA | NA | 0.60 | [71] |
| Niacine | NA | NA | 0.46 | [71] |
| Vitamin K1 | 53.2 | 109 | NA | [71] |

NA: Not available

4. Sterols and Fatty Acids

β -sitosterol is the major sterol extracted from *Opuntia* fruits, peel and seeds. Its concentration vary between 6.75 and 21.1 g/kg [22, 73]. Other sterols such as stigmasterol and lanosterol are present in small quantities (Table 3). Chromatographic analyses of total lipids extracted from cactus peels show that palmitic acid (C16:0) and linoleic acid (C18:2) are the major fatty acids (Table 2). In cladodes, palmitic acid, oleic acid (C18:1), linoleic acid and linolenic acid (C18:3) contribute 13.87, 11.16, 34.87 and 32.83% of the total fatty acid content, respectively (Table 3).

Table 3. Sterols and fatty acids from *O. ficus-indica* (g/Kg).

| | Pulp | Peels | Cladodes | References |
|-------------------------|------|-------|----------|------------|
| Sterols | | | | |
| Campesterol | 8.74 | 8.76 | NA | |
| Stigmasterol | 0.73 | 2.12 | NA | |
| Lanosterol | 0.76 | 1.66 | NA | [22, 73] |
| β -Sitosterol | 11.2 | 21.1 | NA | |
| Δ^5 -Avenasterol | 1.43 | 2.71 | NA | |
| Ergosterol | -- | 0.68 | NA | |
| Fatty acids | | | | |
| C12:0 | NA | 7.1 | 13.3 | |
| C14:0 | NA | 19.5 | 19.6 | |
| C16:0 | NA | 231 | 138.7 | |
| C16:1 | NA | 24.8 | 2.4 | [22, 74] |
| C18:0 | NA | 26.7 | 33.3 | |
| C18:1 | NA | 241 | 111.6 | |
| C18:2 | NA | 323 | 348.7 | |
| C18:3 | NA | 92.7 | 332.3 | |
| C20:0 | NA | nd | -- | |
| C22:0 | NA | 5 | -- | |
| C22:1 | NA | -- | -- | |
| C24:0 | NA | 4.1 | -- | |

NA: Not available

5. Sugars

The main carbohydrates reported in *Opuntia* fruits are fructose and glucose in a ratio at about 1:1 [75]. The polysaccharides from cactus pear peel are characterized by sugar constituents typical of pectin with high and medium degrees of esterification of galacturonic acid residues. Cactus pear peels contain a slightly higher content of sucrose (2.85%) and galacturonic acid (2.23%), followed by stachyose (1.81%), mannitol (1.48%), sorbitol (0.71%) and arabinose (0.05%)

[53]. Glucose and galacturonic acid were the main sugars of *Opuntia* cladodes. HPLC analysis revealed the presence of rhamnose, fucose, arabinose, xylose, mannose and galactose at different concentrations [76].

6. Amino Acids

Total proteins in cladodes vary from 4 to 10%. They are represented mainly by glutamine, leucine and lysine. Phytochemical investigation of *Opuntia* fruits revealed a great number of amino acids. The two predominant amino acids are proline and taurine, which represent 46% and 15.78% of the total amino acid content, respectively. Interestingly, the presence of eight essential amino acids was reported (Table 4). Proteins and amino acid contents of cactus pear peels need to be elucidated.

Table 4. Amino acids contents in both cladodes and fruit pulps from *O. ficus-indica* (g/ 100 g).

| | Cladodes | Fruits | References |
|-----------------------------|----------|--------|------------|
| Alanine | 0.6 | 3.17 | |
| Arginine | 2.4 | 1.11 | |
| Asparagine | 1.5 | 1.51 | |
| Asparaginic acid | 2.1 | Trace | |
| Glutamic acid | 2.6 | 2.40 | |
| Glutamine | 17.3 | 12.59 | |
| Glycine | 0.5 | Trace | |
| Cystine | 1.04 | 0.41 | |
| Histidine | 2.0 | 1.64 | |
| Isoleucine | 1.9 | 1.13 | |
| Leucine | 1.3 | 0.75 | |
| Lysine | 2.5 | 0.63 | |
| Methionine | 1.4 | 2.01 | [77, 78] |
| Phenylalanine | 1.7 | 0.85 | |
| Serine | 3.2 | 6.34 | |
| Threonine | 2.0 | 0.48 | |
| Tyrosine | 1.46 | 0.45 | |
| Tryptophane | 1.04 | 0.46 | |
| Valine | 7.72 | 1.43 | |
| α -Aminobutyric acid | Trace | 0.04 | |
| Carnosine | Trace | 0.21 | |
| Citrulline | Trace | 0.59 | |
| Ornithine | Trace | Trace | |
| Proline | Trace | 46.0 | |
| Taurine | Trace | 15.79 | |

7. Minerals

Cactus fruits and peels are rich in magnesium (up to 59 and 195.76 mg/100 g respectively) and calcium (up to 98.4 and 188.58mg/100 g), which makes prickly pears useful in the prevention of osteoporosis and cramps, respectively. In addition, low levels of sodium, potassium, iron and phosphorus were reported [79, 80]. Whilst high levels of calcium, magnesium and potassium are used for energy and to uphold the mineral pool during periods of physical exhaustion, low levels of sodium and chloride are preferred for preventing high blood pressure [81]. Consequently, cactus pear may be used to ensure mineral fortification of diets. Similarly, cladodes contain high levels of calcium, calcium oxalate and magnesium, but at lower concentrations than those registered for fruits (Table 5).

Table 5. Minerals content in various parts of *O. ficus-indica*. (mg/100 g).

| | Fruits | Cladodes | Peels | References |
|------------|-----------|------------|--------|-----------------|
| Mg | 16.1-98.4 | 8.8 | 195.76 | |
| Ca | 12.8-59 | 5.64-17.95 | 188.58 | |
| Ca Oxalate | --- | 11.5-40.3 | --- | |
| Na | 0.6-1.1 | 0.3-0.4 | 183.42 | |
| K | 90-220 | 2.35-55.20 | 63.45 | [4, 53, 77, 78, |
| Fe | 0.4-1.5 | 0.09 | 25.58 | 82, 83] |
| Mn | --- | 0.19-0.29 | 18.00 | |
| Zn | --- | 0.08 | 17.84 | |
| Cu | --- | --- | 9.47 | |
| P | 15-32.8 | --- | --- | |

8. Fibers

Saenz [84] reported the potential use of cactus cladodes as a new source of fiber in human diet. Cactus pear cladodes were shown to have high fiber content and potential health benefits. The moisture content of tender young *O. ficus-indica* pads is about 92%, along with 1–2% protein, and 0.8–3.3% pectin, a soluble fiber included in the 4–6% total fiber [85]. Dried cladodes powder contains about 43% fibre, 28.5% of which are insoluble [86]. Interestingly, it is important to report that fibers are classified into two groups: hydrosoluble and insoluble ones. Soluble fibres include mucilage, gums, pectin, in addition to hemi-celluloses. They ensure the reduction of blood glucose and cholesterol, as well as the stabilization of intestinal food transit. Insoluble fibres are composed mainly of cellulose, lignin, and a large portion of hemicelluloses. They are known for their capacity to retain water, favor ionic exchange, absorption of bile acids, minerals, vitamins and other interaction with microbes [87-89].

9. Conclusions

The therapeutic properties of *Opuntia ficus-indica* are due to its bio-functional phytochemicals and cladodes polysaccharides. Research on cactus pear has been focused mainly on cladodes and fruits. Although there is a scarce scientific work dealing with cactus pear peels, this latter contains higher bioactive molecules and minerals contents than fruits and cladodes. Indeed, cactus fruit peels could be very suitable as a natural food additive, natural colorant and natural antioxidant. However, further investigations are required to optimize the extraction of bioactive phytochemicals from cactus peels, and to describe their application in food industries.

References

[1] Scheinvar L. Taxonomy of utilized Opuntias, in: Barbera G, P Inglese, E Pimienta-Barrios, (editors), Agroecology, cultivation and uses of cactus pear. FAO Plant Production and Protection Paper, Rome, 1995, 20–27.

[2] Le Houerou HN. Utilization of fodder trees and shrubs in the arid and semi-arid zones of West Asia and North Africa. Arid Soil Research and Rehabilitation 2000; 14: 101–135.

[3] Estrada-Luna AA, JDJ Martínez-Hernández, ME Torres-Torres, F Chablé-Moreno. In vitro micropropagation of the ornamental prickly pear cactus *Opuntia lanigera* Salm-Dyck and effects of sprayed GA3 after transplantation to ex vitro conditions. Scientia Horticulturae 2008; 117: 378–385.

[4] Medina EMD, Rodríguez EM, Romero CD. Chemical characterization of *Opuntia dillenii* and *Opuntia ficus-indica* fruits. Food Chemistry 2007; 103: 38–45.

[5] Galati EM, MR Mondello, D Giuffrida, G Dugo, N Miceli, S Pergolizzi, MF Taviano. Chemical characterization and biological effects of sicilian *Opuntia ficus-indica* (L.) Mill. Fruit Juice: Antioxidant and Antiulcerogenic Activity. Journal of Agricultural and Food Chemistry 2003; 51: 4903-4908.

[6] Piga A. Cactus Pear: A Fruit of Nutraceutical and Functional Importance. Journal of the Professional Association for Cactus Development 2004; 9-22.

[7] Agurell S. Cactaceae alkaloids. VIII. N-Methyl-4-methoxyphenethylamine from *Lepidocoryphantha runyonii* (Br. & R.). Experientia 1969; 25:1111–1132.

[8] Knishinsky R. Prickly pear cactus medicine. Healing Arts Press, Rochester, 1971.

[9] Rosemberg H, Paul AG. Biosynthetic production of aberrant alkaloids in *Dolichothele sphaerica* (Cactaceae). Journal of Pharmaceutical Sciences 1973; 62:403–407.

[10] Sahelian R. An anti-inflammatory principle from prickly pear cactus. Fitoterapia 2001; 72(3): 288–290.

[11] Fernández-López JA, Castellar R, Obón JM, Almela L. Screening and mass-spectral confirmation of betalains in cactus pears. Chromatographia 2002; 56:591–595.

[12] Alarcon-Aguilar FJ, Valdes-Arzate A, Xolalpa-Molina S, Banderas-Dorantes T, Jimenez-Estrada M, Hernandez-Galicia E, Roman-Ramos R. Hypoglycemic activity of two polysaccharides isolated from *Opuntia ficus-indica* (prickly pear cactus) and *O. streptacantha*. Proceedings of the Western Pharmacology Society 2003; 46:139–142.

[13] Galati EM, Mondello MR, Monforte MT, Galluzzo M, Miceli N, Tripodo MM. Effect of *Opuntia ficus-indica* (L.) Mill. cladodes in the wound-healing process. Journal of the Professional Association for Cactus Development 2003; 5:1–16.

[14] Oliveira AJB, Da Machado MFPS. Alkaloid production by callous tissue cultures of *Cereus peruvianus* (Cactaceae). Applied Biochemistry and Biotechnology 2003; 104:149–155.

[15] Tesoriere L, Butera D, Am P, Allerga M, Liverea MA. Supplementation with cactus pear (*Opuntia ficus-indica*) fruit decreases oxidative stress in healthy humans: a comparative study with vitamin C. American Journal of Clinical Nutrition 2004; 80(2):391–395.

[16] Zou D, Brewer M, García F, Feugang JM, Wang J, Zang R, Liu H, Zou C. Cactus pear: a natural product in cancer chemoprevention. Nutrition Journal 2005; 4:25–36.

[17] Saleem M, Kim HJ, Han CK, Jin C, Lee YS. Secondary metabolites from *Opuntia ficus-indica* var. Saboten. Phytochemistry 2006; 67(13):1390–1394.

[18] Osorio-Esquivel O, Alicia-Ortiz M, Álvarez VB, Dorantes-Álvarez L, Giusti MM. Phenolics, betacyanins and antioxidant activity in *Opuntia joconostle* fruits. Food Research International 2011, 44: 2160–2168.

- [19] Paiz RC, Juárez-Flores BI, Aguirre RJR, Cárdenas OC, Reyes AJA, García CE, Álvarez FG. Glucose-lowering effect of xoconostle (*Opuntia joconostle* A. Web. Cactaceae) in diabetic rats. *Journal of Medicinal Plants Research* 2010, 4: 2326–2333.
- [20] Schaffer S, Schmitt-Schillig S, Müller WE, Eckert GP. Antioxidant properties of Mediterranean food plant extracts: Geographical differences. *Journal of physiology and pharmacology* 2005, 56 (Suppl. S1): 115–124.
- [21] Stintzing FC, Schieber A, Carle R. Evaluation of colour properties and chemical quality parameters of cactus juices. *European Food Research and Technology* 2003, 216: 303–311.
- [22] Ramadan MF, Mörsel J-T. Oil cactus pear (*Opuntia ficus-indica* L.). *Food Chemistry* 2003, 82: 339–345.
- [23] Lee JC, Kim HR, Kim J, Jang Y-S. Antioxidant property of an ethanol extract of the stem of *Opuntia ficus-indicavar. saboten*. *Journal of Agricultural and Food Chemistry* 2002, 50: 6490–6496.
- [24] Stintzing FC, Carle R. Cactus stems (*Opuntia* spp.): A review on their chemistry, technology, and uses. *Molecular Nutrition & Food Research* 2005, 49: 175–194.
- [25] Stintzing FC, Schieber A, Carle R. Identification of betalains from yellow beet (*Beta vulgaris* L.) and cactus pear [*Opuntia ficus-indica* (L.) Mill.] by high performance liquid chromatography-electrospray ionization mass spectroscopy. *Journal of Agricultural and Food Chemistry* 2002, 50: 2302–2307.
- [26] Impillizzeri G, Piattelli M. Biosynthesis of indicaxanthin in *Opuntia ficus-indica* fruits. *Phytochemistry* 1972, 11: 2499–2502.
- [27] Cartea ME, Francisco M, Soengas P, Velasco P. Phenolic Compounds in Brassica Vegetables. *Molecules* 2011, 16(1): 251–80.
- [28] De Leo M, Abreu MBD, Pawlowska AM, Cioni PL, Braca A. Profiling the chemical content of *Opuntia ficus-indica* flowers by HPLC–PDA–ESI–MS and GC/EIMS analyses. *Phytochemistry Letters* 2010, 3: 48–52.
- [29] Ahmed MS, Tanbouly NDE, Islam WT, Sleem AA, Senousy ASE. Antiinflammatory flavonoids from *Opuntia dillenii* (Ker-Gawl) Haw. flowers growing in Egypt. *Phytotherapy Research* 2005, 19: 807–809.
- [30] Clark WD, Brown GK, Mays RL. Flower flavonoids of *Opuntia* subgenus *Cylindropuntia*. *Phytochemistry* 1980, 19: 2042–2043.
- [31] Chougui N, Tamendjari A, Hamidj W, Hallal S, Barras A, Richard T, Lariat R. Oil composition and characterisation of phenolic compounds of *Opuntia ficus-indica* seeds. *Food Chemistry* 2013, 139: 796–803.
- [32] Kuti JO. Antioxidant compounds from four *Opuntia* cactus pear fruit varieties. *Food Chemistry* 2004, 85: 527–533.
- [33] Moussa-Ayoub TE, El-Samahy SK, Kroh LW, Rohn S. Identification and quantification of flavonol aglycons in cactus pear (*Opuntia ficus indica*) fruit using a commercial pectinase and cellulase preparation. *Food Chemistry* 2011, 124: 1177–1184.
- [34] Jorge AJ, de La Garza TH, Alejandro ZC, Ruth BC, Noé AC. The optimization of phenolic compounds extraction from cactus pear (*Opuntia ficus-indica*) skin in a reflux system using response surface methodology. *Asian Pacific Journal of Tropical Biomedicine* 2013, 3: 436–442.
- [35] Valente LMM, da Paixão D, do Nascimento AC, dos Santos PFP, Scheinvar LA, Moura MRL, Tinoco LW, Gomes LNF, da Silva JFM. Antiradical activity, nutritional potential and flavonoids of the cladodes of *Opuntia monacantha* (Cactaceae). *Food Chemistry* 2010, 123: 1127–1131.
- [36] Bensaadón S, Hervert-Hernández D, Sáyago-Ayerdi SG, Goñi I. By-Products of *Opuntia ficus-indica* as a Source of Antioxidant Dietary Fiber. *Plant Foods for Human Nutrition* 2010, 65: 210–216.
- [37] Gallegos-Infante J-A, Rocha-Guzman N-E, González-Laredo R-F, Reynoso-Camacho R, Medina-Torres L, Cervantes-Cardozo V. Effect of air flow rate on the polyphenols content and antioxidant capacity of convective dried cactus pear cladodes (*Opuntia ficus indica*). *International Journal of Food Sciences and Nutrition* 2009, 60: 80–87.
- [38] Ginestra G, Parker ML, Bennett RN, Robertson J, Mandalari G, Narbad A, Lo Curto RB, Bisignano G, Faulds CB, Waldron KW. Anatomical, Chemical, and Biochemical Characterization of Cladodes from Prickly Pear [*Opuntia ficus-indica* (L.) Mill.]. *Journal of Agricultural and Food Chemistry* 2009, 57: 10323–10330.
- [39] Fernández-López JA, Almela L, Obón JM, Castellar R. Determination of Antioxidant Constituents in Cactus Pear Fruits. *Plant Foods for Human Nutrition* 2010, 65: 253–259.
- [40] Khatabi O, Hanine H, Elothmani D, Hasib A. Extraction and determination of polyphenols and betalain pigments in the Moroccan Prickly pear fruits (*Opuntia ficus indica*). *Arabian Journal of Chemistry* 2013, doi:10.1016/j.arabjc.2011.04.001.
- [41] Jakobek L. Interactions of polyphenols with carbohydrates, lipids and proteins. *Food Chemistry* 2015, 175: 556–567.
- [42] Chang WS, Lee YJ, Lu FJ, Chiang HC. Inhibitory effects of flavonoids on xanthine oxidase. *Anticancer Research* 1993, 13: 2165–70.
- [43] Iio M, Ono Y, Kai S, Fukumoto M. Effects of flavonoids on xanthine oxidation as well as on cytochrome c reduction by milk xanthine oxidase. *Journal of Nutritional Science and Vitaminology (Tokyo)* 1986, 32: 635–42.
- [44] Ferrali M, Signorini C, Caciotti B, Sugerini L, Ciccoli L, Giachetti D, Comporti M. Protection against oxidative damage of erythrocyte membrane by the flavonoid quercetin and its relation to iron chelating activity. *FEBS Letters* 1997, 416: 123–129.
- [45] Sorata Y, Takahama U, Kimura M. Protective effect of quercetin and rutin on photosensitized lysis of human erythrocytes in the presence of hematoporphyrin. *Biochimica et Biophysica Acta* 1984, 799: 313–7.
- [46] Ferrandiz ML, Alcaraz MJ. Anti-inflammatory activity and inhibition of arachidonic acid metabolism by flavonoids. *Agents Actions* 1991, 32: 283–8.
- [47] Eldahshan AO, Singab ANB. Carotenoids. *Journal of Pharmacognosy and Phytochemistry* 2013, 2: 225–234.
- [48] Goodwin TW. *The Biochemistry of the Carotenoids*. Vol. 1: “Plants.” New York: Chapman and Hall, p 203, 1980.

- [49] Jaramillo-Flores ME, Gonzalza-Cruz L, Dorantes-Álvarez L., Gutierrez-Lopez G.F., Hernandez-Sanchez H. Effect to the thermal treatment on the antioxidant activity and content of carotenoids and phenolic compounds of cactus pear cladodes (*Opuntia ficus-indica*). *Food Science and Technology International* 2003, 9: 271–278.
- [50] Hadj Sadok T, Aid F, Bellal M, Abdul Hussain MS. Composition chimique des jeunes cladodes d'*Opuntia ficus-indica* et possibilités de valorisation alimentaire. *Agricultura-Stiinta si practica* 2008, 1: 65-66.
- [51] Chougui N, Louaileche H, Mohedeb S, Mouloudj Y, Hammoui Y, Tamendjari A. Physico-chemical characterisation and antioxidant activity of some *Opuntia ficus-indica* varieties grown in North Algeria. *African Journal of Biotechnology* 2013, 12: 299-307.
- [52] Kuti JO. Antioxidant compounds from four *Opuntia* cactus pear fruit varieties. *Food Chemistry* 2004, 85:527-533.
- [53] El-Said NM, Nagib AI, Rahman ZA, Deraz SF. Prickly pear [*Opuntia ficus-indica* (L.) Mill] peels: Chemical composition, nutritional value, and protective effects on liver and kidney functions and cholesterol in rats. *Functional Plant Science and Biotechnology* 2010, 5: 30-35.
- [54] Young AJ, Lowe GM. Antioxidant and prooxidant properties of carotenoids. *Archives of Biochemistry and Biophysics* 2001, 385: 20–27.
- [55] Baltschun D, Beutner S, Briviba K, Martin HD, Paust J, Peters M, Röver S, Sies H, Stahl W, Steigel A, Stenhorst F. Singlet oxygen quenching abilities of carotenoids. *Liebigs Annals* 1997, 1887–1893.
- [56] Di Mascio P, Kaiser S, Sies H. Lycopene as the most efficient biological carotenoid singlet oxygen quencher. *Archives of Biochemistry and Biophysics* 1989, 274: 532–538.
- [57] Stahl W, Sies H. Antioxidant activity of carotenoids. *Molecular Aspects of Medicine* 2003, 24: 345–351.
- [58] Sies H, Stahl W. Vitamins E and C, beta-carotene, and other carotenoids as antioxidants. *American Journal of Clinical Nutrition* 1995, 62: 1315S–1321S.
- [59] Minale L, Piattelli M, Nicolaus RA. Pigments of centrospermae – IV. On the biogenesis of indicaxanthin and betanin in *Opuntia ficus-indica*. *Phytochemistry* 1965, 4:593–597.
- [60] Piattelli M, Minale L. Pigments of centrospermae – I. Betacyanins from *Phyllocactus hybridus* Hort. and *Opuntia ficus-indica* Mill. *Phytochemistry* 1964, 3:307–311.
- [61] Yeddes N, Chérif JK, Guyot S, Sotin H, Ayadi MT. comparative study of antioxidant power, polyphenols, flavonoids and betacyanins of the peel and pulp of three Tunisian *Opuntia* forms. *Antioxidants* 2013, 2: 37-51.
- [62] Stintzing FC, Schieber A, Carle R. Phytochemical and nutritional significance of cactus pear. *European Food Research and Technology* 2001, 212: 396–407.
- [63] Castellanos-Santiago E, Yahia EM. Identification and Quantification of Betalains from the Fruits of 10 Mexican Prickly Pear Cultivars by High-Performance Liquid Chromatography and Electrospray Ionization Mass Spectrometry. *Journal Agricultural Food Chemistry* 2008, 56: 5758–5764.
- [64] Strack D, Vogt T, Schliemann W. Recent advances in betalain research. *Phytochemistry* 2003, 62: 247–269.
- [65] Stintzing, F. C.; Herbach, K. M.; Mosshammer, M. R.; Carle, R.; Yi, W.; Sellappan, S.; Akoh, C. C.; Bunch, R.; Felker, P. Color, betalain pattern, and antioxidant properties of cactus pear (*Opuntiaspp.*) clones. *Journal Agricultural Food Chemistry* 2005, 53: 442–451.
- [66] Buettner, G. The pecking order of free and antioxidants: lipid peroxidation, α -tocopherol, and ascorbate. *Archives of Biochemistry and Biophysics* 1993, 300: 535-543.
- [67] Taira J, Tsuchida E, Katoh MC, Uehara M, Ogi T. Antioxidant capacity of betacyanins as radical scavengers for peroxy radical and nitric oxide. *Food Chemistry* 2015, 166: 531–536.
- [68] Gandía-Herrero F, Escribano J, García-Carmona F. Purification and antiradical properties of the structural unit of betalains. *Journal of Natural Products* 2012, 75: 1030–1036.
- [69] Tesoriere L, Allegra M, Gentile C, Livrea MA. Betacyanins as phenol antioxidants. Chemistry and mechanistic aspects of the lipoperoxyl radical-scavenging activity in solution and liposomes. *Free Radical Research* 2009, 43(8): 706-717.
- [70] Tesoriere L, Allegra M, Butera D, Gentile C, Livrea, MA. Kinetics of the lipoperoxyl radical scavenging activity of indicaxanthin in solution and in unilamellar liposomes. *Free Radical Research* 2007, 41, 226-233.
- [71] El-Mostafa K, El Kharrassi Y, Badreddine A, Andreoletti P, Vamecq J, Saïd El Kebbjaj M, Latruffe N, Lizard G, Nasser B, Cherkaoui-Malki M. Nopal Cactus (*Opuntia ficus-indica*) as a Source of Bioactive Compounds for Nutrition, Health and Disease. *Molecules* 2014, 19:14879-14901.
- [72] Hassanien MFR, Mörsel JT. Agro-waste products from prickly pear fruit processing as a source of oil. *Fruit Processing* 2003, 13: 242-248.
- [73] Ramadan MF, J-T Mörsel. Recovered lipids from prickly pear [*Opuntia ficus-indica* (L.) Mill] peel: a good source of polyunsaturated fatty acids, natural antioxidant vitamins and sterols. *Food Chemistry* 2003, 83: 447-456.
- [74] Abidi S, Ben Salem H, Vasta V, Priolo A. Supplementation with barley or spineless cactus(*Opuntia ficus indica* f. inermis) cladodes on digestion, growth and intramuscular fatty acid composition in sheep and goats receiving oaten hay. *Small Ruminant Research* 2009, 87: 9–16.
- [75] Kuti JO, Galloway C-M. Sugar composition and invertase activity in prickly pear fruit. *Journal of Food Science* 1994, 59: 387-393.
- [76] Ginestra G, Parker ML, Bennett RN, Robertson J, Mandalari G, Narbad A, Lo Curto RB, Bisignano G, Faulds CB, Wwaldron KW. Anatomical, Chemical, and Biochemical Characterization of Cladodes from Prickly Pear [*Opuntia ficus-indica* (L.) Mill.]. *Journal of Agricultural Food Chemistry* 2009, 57: 10323–10330.
- [77] Feugang JM, Konarski P, Zou D, Stintzing FC, Zou C. Nutritional and medicinal use of Cactus pear (*Opuntia spp.*) cladodes and fruits. *Frontiers in Bioscience* 2006, 11: 2574-2589.
- [78] Sawaya WN, Khalil JK, Al-Mohammad MM. Nutritive value of prickly pear seeds, *Opuntia ficus-indica*. *Plant Food For Human Nutrition* 1983, 33: 91–97.

- [79] Wills RBH, Lim JSK, Greenfield H. Composition of Australian Foods. Tropical and sub-tropical fruit. Food Technology in Australia 1986, 38:118–123.
- [80] Askar A, El-Samahy SK. Chemical composition of prickly pear fruits. Dtsch Lebensm Rundsch 1981, 77:279–281.
- [81] Pszczola DE. Natural colors: Pigments of imagination. Food Technology 1998, 52:70–82.
- [82] Contreras-Padilla M, Pérez-Torrero E, Hernández-Urbiola MI, Hernández-Quevedo G, del Real A, Rivera-Muñoz EM, Rodríguez-García ME. Evaluation of oxalates and calcium in nopal pads (*Opuntia ficus-indica* var. *redonda*) at different maturity stages. Journal of Food Composition Analysis 2011, 24: 38–43.
- [83] Ayadi MA, Abdelmaksoud W, Ennouri M, Attia H. Cladodes from *Opuntia ficus indica* as a source of dietary fiber: Effect on dough characteristics and cake making. Industrial Crops and Products 2009, 30: 40–47.
- [84] Saenz CH. Cladodes: a source of dietary fiber. Journal of the Professional Association for Cactus Development 1997, 2:117–123.
- [85] Brinker FND. Prickly Pear as Food and Medicine. Journal of Dietary Supplements 2009, 6: 362-376.
- [86] Nefzaoui A, Nazareno M, El Mourid M. Review Of Medicinal Uses Of Cactus. CACTUSNET 2007, 11: 3-17.
- [87] Hollingsworth P. Food trends: diversity and choice dominate. Food Technology 1996, 5:40.
- [88] Grijspaardt-Vink C. Ingredients for healthy foods featured at European Expo. Food Technology 1996, 2:30.
- [89] Sáenz C. Utilización agroindustrial del nopal. Boletín de servicios agrícolas de la FAO 2006, 162: 164.