Molecular Cloning, Characterization and Expression Analysis of *MhRAR1* Gene from *Malus Hupehensis*

Zhang Ji-Yu, Guo Zhong-Ren

Institute of Botany, Jiangsu Province and the Chinese Academy of Sciences, Nanjing, China

Email address:

maxzhangjy@163.com (Zhang Ji-Yu), zhongrenguo@cnbg.net (Guo Zhong-Ren)

To cite this article:

Zhang Ji-Yu, Guo Zhong-Ren. Molecular Cloning, Characterization and Expression Analysis of *MhRAR1* Gene from *Malus Hupehensis*. *Journal of Plant Sciences*. Vol. 3, No. 2, 2015, pp. 85-91. doi: 10.11648/j.jps.20150302.17

Abstract: A novel *RAR1* gene, designated *MhRAR1*, was cloned by the methods of RT-PCR and RACE from *Malus hupehensis*. The full length sequence of *MhRAR1* is 1065 bp with an open reading frame of 678 bp, encoding a protein of 225 amino acids. As found in other plant *RAR1* proteins, sequence alignment showed that *MhRAR1* protein contains two CHORD domains and one plant-specific CCCH domain. In addition, the *MhRAR1* contains conserved strings of invariant cysteine and histidine residues within the CHORD domains and CCCH domain. These results suggested that *MhRAR1* protein from *M. hupehensis* might share the similar function with the *Arabidopsis thaliana* *RAR1* and *Hordeum vulgare* *RAR1*, and is an important component of R gene-mediated disease resistance. Phylogenetic analysis revealed that *MhRAR1* was closely related to *Ricinus communis* *RAR1*. The analysis by qRT-PCR revealed that the expression of *MhRAR1* gene was higher in leaves than that in stems and roots. SA, MeJA and ACC treatment induced *MhRAR1* expression in stems and roots, but not in leaves. Expression of *MhRAR1* was weakly induced in *M. hupehensis* after infection with *Botryosphaeria berengeriana*. The cloning and characterization of the *MhRAR1* gene will be useful for further studies of biological roles of *MhRAR1* in plants.

Keywords: *Malus hupehensis*, *MhRAR1*, cDNA Cloning, Expression Pattern

1. Introduction

Plants are frequently attacked by various disease pathogenesis with complicated strategies colonizing their hosts, and have evolved defense mechanisms to protect themselves effectively from microbial pathogens. Plants various resistance proteins (R proteins) activate resistance responses including rapid ion fluxes, generation of reactive oxygen species (ROS), production of antimicrobial compounds, and the accumulation of salicylic acid (SA) which was crucial for the induction of systemic acquired resistance (SAR). These responses are accompanied by the hypersensitive response (HR), localized programmed cell death at the site of pathogen invasion [1].

RAR1 (Required for *Mla12* resistance), which was a crucial component of resistance conferred by many R proteins [2,3], play a key role in disease resistance of plants. *Arabidopsis rar1* mutants are defective in R-protein mediated resistance against several bacterial (*Pseudomonas*) and oomycete (*Peronospora*) pathogens [4,5]. *Rar1* mutants fail to accumulate ROS or mount the HR in barley [6]. In the past several decades, many studies have been done to discover the molecular function of

RAR1 protein in plants. Numbers of researches have been carried out in the model plants such as *Arabidopsis* [4,7,8], barley [3, 9-11], tobacco [12] and so on. However, none of work was done in woody plants especially in fruit trees such as apple.

The plant hormones SA, MeJA and ET are three types of signalling mediators that activate the plant defence response against pathogen attack [13,14]. In general, SA and JA/ET are thought to regulate two different signal transduction pathways in plant disease resistance. SA and JA/ET signalling pathways and the cross-talk between them have been studied in various plants [15-17]. To determine whether exogenous application of these stimuli could induce the expression of *MhRAR1* gene, *Malus hupehensis* were treated with SA, MeJA and ACC.

Malus hupehensis, common names *Hupeh crab* or *tea crabapple*, is a species of flowering plant in the apple genus *Malus* of the family *Rosaceae*, native to Japan and Taiwan and province of China. It is a vigorous deciduous tree growing to 12 m (39 ft) tall and broad, with pink buds opening to fragrant white blossom in Spring, and bright red, cherry-sized

crab-apples in the Autumn. *M. hupehensis* was an apple rootstock that is tolerant to various apple pathogens [18]. Here, we isolated and functionally characterised of a *M. hupehensis* *RAR1* gene. To better understand *MhRAR1* gene functions, we also analysed the expression of *MhRAR1* in response to plant hormones, such as SA, MeJA, and ACC in different tissues of *M. hupehensis*, and in response to biotic stress apple ring spot pathogenesis *Botryosphaeria berengeriana*.

2. Materials and Methods

2.1. Plant Materials and Bacterial Strains

M. hupehensis plants were subcultured in vitro in Murashige and Skoog (MS) medium supplemented with 6-benzylaminopurine (6-BA, 0.5mg/L) and naphthalene acetic acid (NAA, 0.1mg/L) and cultured under a 16-hour light (25°C)/8-hour dark (25°C) cycle. *Escherichia coli* strain DH5 α cells were used for the cloning of *MhRAR1*, and propagation of all the recombinant plasmid vectors. The Pyk2478 vectors were used for the construction of plant expression vectors of the *RAR1* gene.

2.2. Treatment of *M. hupehensis* with SA, MeJA and ACC and Apple Ring Spot Pathogen

Three weeks old *M. hupehensis* in vitro plant leaves were sprayed with 0.1mM SA (Sigma), 0.02mM MeJA (Sigma) and 0.01mM ACC (Sigma) for 4, 12 and 48h, respectively, with water treatment as a control. Each treatment group consisted of three plants. Then, the leaves, stems and roots were harvested and frozen in liquid nitrogen. Tissues were stored at -70°C until use. For apple ring spot pathogen inoculation (*B. berengeriana*), *M. hupehensis* tissue culture plants were rooted after four weeks. Abaxial leaf surfaces were sprayed with freshly collected sporangia propagated in potato dextrose agar (PDA) media and resuspended in water at approximately 1.0×10^6 spores mL^{-1} . Leaves were collected at different time points and immediately frozen in liquid nitrogen.

2.3. Extraction of Total RNA and Clone of *MhRAR1*

PCR was performed with the primers RF1 and RR1 (Table 1) derived from the conserved sequence region of *RAR1* using a complete cDNA library [19] in order to isolate a partial cDNA fragment (554bp) of the *MhRAR1* gene. Then, using a sequence obtained from the initial cDNA fragment, new gene-specific sense and antisense primers were designed for the 5'- and 3'- rapid amplification of cDNA ends (RACE) reactions (Table 1). The missing 3' and 5' sequences were obtained by RACE with SMARTTM RACE cDNA Amplification Kit (Clontech Laboratories, Inc. No. 634902). The conserved, 3' and 5' sequences were assembled using the DNAMAN software. To verify the assembled sequence, PCR was performed using the primers RF3 and RR3 (Table 1) from the complete cDNA library to amplify the open reading frame (ORF) of *MhRAR1*. Genomic DNA was extracted from leaves and treated with RNase I as described by Tong et

al.[20]. *MhRAR1* gene cloning from genomic DNA was performed as above using the primers RF3 and RR3 (Table 1). The PCR products were cloned into a pMD19-T vector (TaKaRa) and sequenced by the Shanghai Invitrogen Biotechnology Company Ltd.

The amino acids sequence of *MhRAR1* was deduced and analyzed using ProtParam tool (<http://cn.expasy.org/tools/protparam.html>). Nucleotide and protein sequence similarity analyses were carried out at the NCBI server using BLAST programs (<http://www.ncbi.nlm.nih.gov/BLAST/>). Amino acid alignment sequences of *RAR1* homologues were downloaded from GenBank and the alignment of the amino acid sequences with that of *MhRAR1* was performed using BioEdit software. A comparative phylogenetic tree was then constructed using MEGA software version 4.0 with the Neighbour-Joining method [21] and 1000 replicates were used for bootstrap test. The tree was then visualised using the tree view package[22].

2.4. Gene Expression Analysis of *MhRAR1* in *M. hupehensis* Using Real-Time Quantitative RT-PCR (qRT-PCR)

Total RNA isolation were performed as described by Cai et al. [23] and DNase I treatment was performed according to the manufacturer's protocol. First strand cDNA was synthesised with the ReverTraAce qPCR RT Kit (TOYOBO, Code No.: FSQ-101) according the manufacturer's instructions. QRT-PCR analyses with a pair of primers (Table 1) were performed to determine the tissue-specific expression (leaf, stem and root) and expression profiles of *MhRAR1* after various treatments. Normal PCR reactions were performed to ensure gene-specific amplification using the primers BRF1 and BRR1 to amplify the *MhRAR1* gene. A single PCR fragment with the expected size was amplified, suggesting that the primers were suitable for qRT-PCR analyses. To confirm the expected fragment of the *MhRAR1* gene, the resulting PCR product was cloned and sequenced. All samples were harvested and three biological replicates were run independently. *M. hupehensis* tubulin was amplified with a pair of primers and used as the housekeeping gene for the qRT-PCR analyses. Templates were the $10 \times$ diluted cDNAs from each sample. The reaction protocol was as follows: 1 μ L of 10-fold diluted cDNA, 0.3 μ L 10pM of each primer (Table 1), 10 μ L SYBR® Premix Ex Taq™ (Perfect Real Time) (TaKaRa Code: DRR041A) and 8.4 μ L of sterile double distilled water. The thermal conditions were: 95°C for 4 min, 95°C for 20 s, 58°C for 20 s and 72°C for 43 s for 40 cycles. Dissociation curves from 55°C to 95°C were generated for each reaction to ensure specific amplification, along with verification by gel electrophoresis. The qRT-PCRs were performed using a 7300 Real-Time PCR System (Applied Biosystems, Foster City, CA, USA). The relative levels of genes respective to control Mhtubulin mRNAs were analysed using the 7300 system software and the $2^{-\Delta\Delta C_t}$ method [24]. Data analyses were conducted using SPSS version 17.0 statistical software. For all analyses, the level of significance between different time points was set at $P < 0.05$.

2.5. Subcellular Localisation of the MhRAR1-EGFP Fusion Protein

Subcellular localisation of MhRAR1 was assayed according to the method of Wang et al.[25]. For generating the MhRAR1-GFP fusion construct, the *MhRAR1* ORF was amplified with a forward primer YXRAF1 containing a

BamH I site and a reverse primer YXRAR1 containing an *EcoR I* site (Table 1). The resulting PCR products were cloned into the pMD19-T simple vector (TaKaRa) and sequenced. The entire ORF of the *MhRAR1* gene was inserted into the plasmid Pyk2784-EGFP which included enhanced green fluorescent protein.

Table 1. Primers used for this paper.

Primers name	Sequence [5'-3']	Usage
RF1	ACTTGCGACGCCATGTTC	Cloning of MhRAR1 conserved sequence
RR1	TCGCAACATTTCCACCCTC	
RF2	AGAGAAAGATAACCACGAGAATGC	3'RACE
RR2	GGAGCAGAAGTGCCTTGATCGTCAGTAAT	5'RACE
RF3	ACACTAGTGGATGGAGGGTC	PCR amplification of MhRAR1 ORF
RR3	GGTACCTTAAGATACCGGGT	
BRF1	TGATATAAACCCAGCCCCAGACG	Expression analysis for MhRAR1
BRR1	CACAGCACTTCCACCCTCTC	
BAF1	AGGTCCATCCATTGTCCACAG	Housekeeping gene
BAR1	TGCCAACCAAACTGACTTCAC	
YXRAF1	ACGGATCCGGATGGAGGGTC	Vector construction of Subcellular localization
YXRAR1	AGAATTACAGATACCGGGT	

3. Results

3.1. Analysis of the MhRAR1 Gene Sequence and the Predicted Protein

Fig. 1. The structure of MhRAR1 gene and protein. (a) The nucleotide and deduced amino acid sequences of cloned MhRAR1. The protein sequence is indicated by one letter code below the nucleotide sequence. * indicate the stop codon. (b) Schematic diagram of MhRAR1 gene. Reseda indicated the exons and White indicated the introns.

A full-length 1,065 bp of *M. hupehensis RAR1* was cloned using the methods of RT-PCR and RACE. It contained an

open reading frame (ORF) of 678 bp encoded a protein of 225 amino acids. The cDNA had a 5'-untranslated region (UTR) of 147 nucleotides, a 3'-UTR of 240 nucleotides including a stop codon (TAA) (Fig. 1a). The calculated molecular mass was 24.90 kDa, and the estimated isoelectric point (pI) of this protein was 7.51. The *MhRAR1* cDNA sequence has been submitted to the NCBI GenBank as accession number FJ593502. The *MhRAR1* genomic DNA sequence from start to stop codon was 2278 bp, and there were five exons and four introns in the complete coding region of *MhRAR1* when the cDNA sequence and genomic DNA sequence were compared (Fig. 1b).

3.2. Multiple Sequences Alignment and Phylogenetic Analysis

A search in the GenBank database, using the BLASTP program, revealed that the predicted MhRAR1 deduced amino acid of *M. hupehensis* exhibited high similarities with those of the *M. domestica* (identities 97 %), *Fragaria vesca subsp. vesca* (identities 84 %), *Prunus mume* (identities 84 %) and *Ricinus communis* (identities 74 %). Similar to *A. thaliana* [4], barley and tobacco [12], there were two cysteine and histidine-rich Zn²⁺-binding domains (designated as CHORD) and one plant-specific conserved Cys residues and a His residue (CCCH) domain in MhRAR1 protein (Fig.2). In addition, the MhRAR1 contains conserved strings of invariant Cys and His residues within the CHORD domains (Fig.2). The amino acid sequences outside the CHORD and CCCH domains are significantly different among plant PAR1 proteins (Fig.2). To determine the phylogenetic relationship between MhRAR1 and other plant RAR proteins, a phylogenetic tree was constructed, revealing that MhRAR1 was closely related to MdRAR1, FvRAR1 and PmRAR1 (Fig. 3).

Fig. 2. Amino acid sequence comparison of the predicted *Malus hupehensis* RARI protein with the other plant species RARI. CLUSTALW produced alignment was formatted using BioEdit software. Conservative substitutions are shaded in dot mark. Two conserved CHORD domains (CHORD-I and II) and one conserved CCCH domain are shown. Triangle (▲) indicate invariant conserved cysteine and histidine residues within the CHORD and CCCH domains. The accession numbers of plant RAR sequences are as follows: *MhRARI*: *Malus hupehensis* RARI (gbACT98264); *RcRARI*: *Ricinus communis* RARI (ref XP002510896); *AtRARI*: *Arabidopsis thaliana* RARI (dbjBAB11239); *NtRARI*: *Nicotiana tabacum* RARI (gbAAM62409); *HvRARI*: *Hordeum vulgare* RARI (gbAAF18432); *TaRARI*: *Triticum aestivum* RARI (gbABN13684); *SaRARI*: *Saccharum hybrid cultivar* RARI (gbACX68653); *GmRARI*: *Glycine max* RARI (gbACI31548).

Fig. 3. Phylogenetic relationships of *MhRARI* with other plant RARI. Phylogenetic tree was constructed using MEGA software version 4.0 with the Neighbour-Joining method (Tamura et al. 2007) and 1000 replicates were used for bootstrap test. The tree was then visualised using the tree view package (Page 1996). The accession numbers of plant RAR sequences are as follows: *MhRARI*: *Malus hupehensis* RARI (gbACT98264); *RcRARI*: *Ricinus communis* RARI (ref XP002510896); *AtRARI*: *Arabidopsis thaliana* RARI (dbjBAB11239); *NtRARI*: *Nicotiana tabacum* RARI (gbAAM62409); *HvRARI*: *Hordeum vulgare* RARI (gbAAF18432); *TaRARI*: *Triticum aestivum* RARI (gbABN13684); *GmRARI*: *Glycine max* RARI (gbACI31548); *MdRARI*: *Malus domestica* RARI (ref XP008376437); *FvRARI*: *Fragaria vesca subsp. vesca* RARI (ref XP004308155); *PmRARI*: *Prunus mume* RARI (ref XP008234345); *VvRARI*: *Vitis vinifera* RARI (ref XP002269178); *SiRARI*: *Setaria italic* (ref XP001267761); *OsRARI*: *Oryza sativa* RARI (gb BAD27802); *CsRARI*: *Cucumis sativus* (ref XP004159991).

3.3. Tissues Expression of *MhRARI* in *M. hupehensis*

QRT-PCR was performed to measure the *MhRARI* gene expression in different tissues of *M. hupehensis*. The dissociation curves were analyzed to determine the amplification specificity for *MhRARI* and *Mhtubulin*. Only one peak existing in the dissociation curves for both *MhRARI* and *Mhtubulin* gene, indicating that the amplifications were specific. The results showed that the mRNA transcripts of *MhRARI* could be detected in all tested tissues and that expression was higher in leaves than that in stems and roots (Fig.4)

Fig. 4. Expression patterns of *MhRARI* in different tissues using qRT-PCR in *M. hupehensis*. *MhTubulin* transcript levels were used to normalise the samples. The mean value and standard deviation were obtained from three independent experiments.

3.4. Expression of the *MhRARI* Gene Induced by SA, MeJA and ACC in Leaves, Stems and Roots of *M. hupehensis*

In order to find out whether exogenous application of these stimuli could induce the expression of *MhRARI* gene, *M. hupehensis* leaves were treated by plant hormones like as SA, MeJA and ACC. The expressions of *MhRARI* gene in different organs of *M. hupehensis* were performed using qRT-PCR (Fig 5). SA, MeJA and ACC could induce the expression of *MhRARI* gene in stems (Fig 5D, E and F) and roots (Fig 5G, H and I), but not in leaves (Fig 5A, B and C).

The amount of height of the expression level varied substantially with both time and tissue. In leaves, the transcript levels had not obviously changed upon treatment with SA (Fig 5A), MeJA (Fig 5B) and ACC (Fig 5C) during the first 48h. In stems after 4, 12 and 48 h, the transcript levels rose 2.03x, 2.64x and 2.75x, respectively, upon treatment with SA, relative to the control treatment (Fig 5D). When stems were treated with MeJA, transcript levels of *MhRARI* rose 0.86x, 2.54x and 1.87x after the same time periods, respectively (Fig 5E). Increases of 0.88x, 1.50x and 1.65x were observed in stems following ACC treatment after 4, 12 and 48 h (Fig 5F). In roots, when compared to measurements at time 0 of the experiment, the accumulation of *MhRARI* gene transcripts was 5.43x, 2.29x and 2.08x with SA (Fig 5G); 2.51x, 2.93x and 2.53x with MeJA (Fig 5H); 4.23x, 8.42x and 2.02x with ACC (Fig 5I), 4, 12 and 48 h post-treatment, respectively.

Fig. 5. Expression patterns of the *MhRAR1* using qRT-PCR in *M. hupehensis* upon treatment with SA (A, D and G), MeJA (B, E and H) and ACC (C, F and I) in leaves (A, B and C), stems (D, E and F) and roots (G, H and I), respectively. *MhTubulin* transcript levels were used to normalise the samples. Mean values and standard deviations were obtained from three independent experiments.

3.5. Induction of the *MhRAR1* Gene with the Fungus *B. Berengeriana*

In this study, we analysed the change in the expression levels of the *MhRAR1* gene in *M. hupehensis* post-infection with apple ring spot pathogen and showed that this pathogen could weakly induce the expression of *MhRAR1* (Fig 6). Levels rose at 6 hours post-inoculation (hpi), remained at the highest level at 12 hpi, and then decreased (Fig 6).

Fig. 6. Expression patterns of *MhRAR1* using qRT-PCR in *M. hupehensis* after inoculation with apple ring spot pathogen (*Botryosphaeria berengeriana* de Not. f. sp. *Parabola Nise Koganezawa et Sakuma*). Mean values and standard deviations were obtained from three independent experiments.

3.6. Subcellular Localisation of *MhRAR1* Protein

To study the subcellular distribution of the *MhRAR1* protein, the coding region of *MhRAR1* without the stop codon was fused to EGFP under the control of the CaMV35S promoter (D35S::MhRAR1-GFP) in the experiment (Fig. 7a). This plasmid, or a control (D35S::GFP), was transformed into onion epidermal cells by the particle bombardment method. It could be observed that the control proteins were uniformly distributed in cells (Fig. 7b, B1 to B3), while the *MhRAR1*-EGFP fusion protein was located in the cell wall

and cell membrane (Fig. 7b, A1 to A3).

Fig. 7. The (a) *MhRAR1* fusion GFP expression vector and (b) subcellular localisation of *MhRAR1* in onion epidermal cells. Onion epidermal cells were transformed with D35S::MhRAR1-EGFP (A1 to A3) or the control plasmid D35S::EGFP (B1 to B3) using the particle bombardment method. The expression and subcellular distribution of proteins were examined under a light microscope (A1, B1) or fluorescent microscope (A2, B2) and then merged (A3, B3). Data shown are representative of three independent experiments. EGFP: enhanced green fluorescent protein.

4. Discussion

RAR1 protein contains two highly similar zinc-binding domains named CHORD-I and CHORD-II that are conserved in eukaryotic phyla except yeast [26]. There are six conserved Cys and three conserved His in CHORD-containing proteins [6]. The tandem organization of CHORD domains and conserved primary structure suggested that these proteins supply important cellular functions. In addition, plant RAR1 proteins have a 20-amino acid motif with three conserved Cys residues and a His residue in CCCH domain which was between CHORD domains I and II [4]. A cDNA sequence named *MhRAR1* gene was isolated from the woody plant *M. hupehensis* in this paper. Analysis of RAR1 protein sequence showed that *MhRAR1* protein has high sequence similarity with other herbaceous plants, such as *A. thaliana*, *N. tabacum*, *T. aestivum* and so on. The same as these plants, *M. hupehensis* *MhRAR1* protein also has two 60 aa long repeat motifs named CHORD domains at the N and C terminus, comprising about 50% of the predicted full length protein. Comparison of deduced protein sequence of *M. hupehensis* RAR1 gene with *R. communis*, *A. thaliana*, *G. max*, *N. tabacum*, *H. vulgare*, *T. aestivum* and *Saccharum* hybrid cultivar showed the same tandem organization of the 60 aa motif as well as similar spacing in each species. And *MhRAR1* contains conserved strings of invariant Cys and His residues within the CHORD domains. In addition, *MhRAR1* protein possesses a CCCH domain which also has three conserved Cys and one His between the CHORD domains. As a result, we suggested that *MhRAR1* protein maybe shared the similar function with the *A. thaliana* AtRAR1 and *H. vulgare*

HvRAR1, and is an important component of R gene-mediated disease resistance.

In this report, *M. hupehensis* leaves were treated by plant hormones like as SA, MeJA and ACC to understand whether exogenous application of these stimuli could induce the expression of *MhRAR1* gene. The results showed that SA, MeJA and ACC could induce the expression of MhRAR1 gene in stem (Fig 5D, E and F) and root (Fig 5G, H and I), but not in leaf (Fig 5A, B and C), suggesting that the expression pattern of *MhRAR1* gene in various tissues under the treatment with SA, MeJA or ACC were different.

Plant innate immune responses were triggered by R genes upon pathogen attack. RAR1, which was engaged by various R genes, is an early union point in a signaling pathway [11]. In plants, RAR1 interacts directly with SGT1 (for suppressor of the G2 allele of *skp1*) and HSP90 [2,8,12,27,28]. Cytosolic HSP90 contains three distinct domains: (1) an N-terminal ATPase domain (N); (2) a substrate binding domain in the middle; (3) the C-terminal end (C) containing a dimerization domain and a MEEVD motif that binds tetratricopeptide repeat (TPR) domains of many cochaperones [8]. Takahashi et al [8] indicated that The CHORD-I domain of RAR1 interacts directly with the N terminus of HSP90. RAR1 physically interacts with SGT1 (suppressor of the G2 allele of *SKP1*) through the C-terminal CHORD-II of RAR1 and the central CS domain of SGT1 [2,29,30]. Azevedo et al [2] state clearly that the interactions among RAR1, SGT1 (suppressor of the G2 allele of *SKP1*), SCF (*Skp1*-Cullin^{DF}-box), and signalosome subunits indicate a link between disease resistance and ubiquitination. In this study, the result of subcellular localisation showed that MhRAR1-EGFP fusion protein was located in the cell wall, cell membrane and cytoplasm, suggesting that MhRAR1 protein could interact with SGT1 and HSP90 in cytoplasm. In future, we will isolate the sequences of SGT1 and HSP90 from *M. hupehensis* to study whether SGT1 and HSP90 act with RAR1.

In plants, a major form of resistance to disease caused by microbial pathogens is by expression of complementary gene pairs in the plant and pathogen, known respectively as resistance (*R*) and avirulence (*avr*) genes [31]. Direct or indirect interaction of their products activates cellular defenses that prevent pathogen colonization of the plant [32]. The most plant *R* gene class encodes predicted cytosolic proteins with a central nucleotide binding (NB) domain and C-terminal Leu-rich repeats (LRRs) [31]. NB-LRR proteins fall into two subclasses based on their different N-terminal motifs. One group possesses an N-terminal coiled-coil (CC) domain. The second group has N-terminal similarity to the cytoplasmic Toll Interleukin-1 Receptor (TIR) domains of human and *Drosophila* Toll-like receptors [31]. Muskett et al [4] reported that RAR is required by TIR- and CC-NB-LRR *R* genes. Barley *RAR1* is a necessary component of *R* gene-mediated resistance to the powdery mildew fungus [32]. In this study, the expression of *MhRAR1* gene was induced by the apple ring spot pathogenesis. We conjecture that MhRAR1 maybe a necessary component of R gene-mediated resistance to the apple ring spot pathogenesis.

5. Conclusion

In this study, a novel RAR1 gene *MhRAR1* was cloned from *M. hupehensis*. The results showed that MhRAR1 protein contains two CHORD domains and one plant-specific CCCH domain. In addition, the MhRAR1 contains conserved strings of invariant cysteine and histidine residues within the CHORD domains and CCCH domain. These results suggested that MhRAR1 protein from *M. hupehensis* might share the similar function with the *Arabidopsis thaliana* RAR1 and *H. vulgare* RAR1, and is an important component of R gene-mediated disease resistance. Phylogenetic analysis revealed that MhRAR1 was closely related to *R. communis* RAR1. The expression of *MhRAR1* gene was higher in leaves than that in stems and roots. SA, MeJA and ACC treatment induced *MhRAR1* expression in stems and roots, but not in leaves. Expression of *MhRAR1* was weakly induced in *M. hupehensis* after infection with *Botryosphaeria berengeriana*. In future studies we will concentrate on introducing the *MhRAR1* gene into the 'Fuji' apple variety to determine if this can enhance resistance to apple plant pathogens, such as apple ring spot, apple powdery mildew and apple canker pathogens.

Acknowledgements

This work was supported by grants from the China Ministry of Science and Technology "863" program # 2011AA100204 and National Natural Science Foundation of China #31401854

References

- [1] Shirasu K., Schulze-Lefert P. 2000. Regulators of cell death in disease resistance. *Plant Molecular Biology* 44: 371–385
- [2] Azevedo C., Sadanandom A., Kitagawa K., Freialdenhoven K., Shirasu A., Schulze-Lefert P. 2002. The RAR1 Interactor SGT1, an Essential Component of R Gene-Triggered Disease Resistance. *Science* 295: 2073-2076
- [3] Bieri S., Mauch S., Shen Q.H., Peart J., Devoto A., Casais C., Francesca C., Schulze S., Steinbiß H.-H., Shirasu K., Schulze-Lefert P. 2004. RAR1 Positively Controls Steady State Levels of Barley MLA Resistance Proteins and Enables Sufficient MLA6 Accumulation for Effective Resistance. *The Plant Cell Online* 16: 3480-3495
- [4] Muskett P.R., Kahn K., Austin M.J., Moisan L.J., Sadanandom A., Shirasu K., Jones J.D.G., Parker J.E. 2002. Arabidopsis RAR1 Exerts Rate-Limiting Control of R Gene-Mediated Defenses against Multiple Pathogens. *The Plant Cell Online* 14: 979-992
- [5] Tornero P., Merritt P., Sadanandom A., Shirasu K., nnes R.W.I., Dangl J.L. 2002. RAR1 and NDR1 Contribute Quantitatively to Disease Resistance in Arabidopsis, and Their Relative Contributions Are Dependent on the R Gene Assayed. *The Plant Cell Online* 14: 1005-1015
- [6] Shirasu K., Lahaye T., Tan M.-W., Zhou F., Azevedo C., Schulze-Lefert P. 1999. A novel class of eukaryotic zinc-binding proteins is required for disease resistance signaling in barley and development in *C. elegans*. *Cell* 99: 355-366

- [7] Shang Y., Li X., Cui H., He P., Thilmony R., Chintamanani S., Zwiesler-Vollick J., Gopalan S., Tang X., Zhou J.M. 2006. RAR1, a central player in plant immunity, is targeted by *Pseudomonas syringae* effector AvrB. *Proceedings of the National Academy of Sciences* 103: 19200-19205
- [8] Takahashi A., Casais C., Ichimura K., Shirasu K. 2003. HSP90 interacts with RAR1 and SGT1 and is essential for RPS2-mediated disease resistance in *Arabidopsis*. *Proceedings of the National Academy of Sciences* 100: 11777-11782
- [9] Tai Y.S. 2007. Interactome of signaling networks in wheat: the protein-protein interaction between TaRAR1 and TaSGT1. *Molecular Biology Reports* 35: 337-343
- [10] Zellerhoff N., Jansen M., Schaffrath U. 2008. Barley Rom1 antagonizes Rar1 function in Magnaporthe oryzae-infected leaves by enhancing epidermal and diminishing mesophyll defence. *New Phytologist* 180: 702-710
- [11] Zhang L., Yang W., Liu D. 2011. TaRAR1 is Required for Lr24-Mediated Wheat Leaf Rust Resistance. *Agricultural Sciences in China* 10: 1732-1738
- [12] Liu Y., Schiff M., Marathe R., Dinesh-Kumar S.P. 2002. Tobacco Rar1, EDS1 and NPR1 NIM1 like genes are required for N mediated resistance to tobacco mosaic virus. *The Plant Journal* 30: 415-429
- [13] Glazebrook J. 2001. Genes controlling expression of defence responses in *Arabidopsis*- 2001 status. *Curr Opin Plant Biol* 4:301-308
- [14] Kim J.C., Lee S.H., Cheong Y.H., Yoo C.M., Lee S.I., Chun H.J., Yun D.J., Hong J.C., Lee S.Y., Lim C.O., Cho M.J. 2001. A novel cold-inducible zinc finger protein from soybean, SCOF-1, enhances cold tolerance in transgenic plants. *Plant J* 25:247-251
- [15] Dong X. 1998. SA, JA, ethylene, and disease resistance in plants. *Curr. Opin. Plant Biol* 1:316-323
- [16] Spoel S.H., Koornneef A., Claessens S.M., Korzelius J.P., Pelt J.A., Mueller M.J., Buchala A.J., Metraux J.P., Brown R., Kazan K., Loon L.C., Dong X.N., Pieterse C.M. 2003. NPR1 modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in the cytosol. *The Plant Cell* 15:760-770
- [17] Leon-Reyes A., Spoel S.H., Lange E.S., Abe H., Kobayashi M., Tsuda S., Millenaar F.F., Welschen R.A., Ritsema T., Pieterse C.M. 2009. Ethylene modulates the role of nonexpressor of pathogenesis-related genes1 in cross talk between salicylate and jasmonate signaling. *Plant Physiology* 149:1797-1809
- [18] Lu Q.N., Jia D.X. 1999. China Fruit Records: Apple Volume, First Ed., China's Agricultural Science & Technology Press, Beijing
- [19] Zhang J.Y., Qu S.C., Dong C., Gao Z.H., Qiao Y.S., Zhang Z. 2010. Utility and construction of full-length cDNA library of *Malus hupehensis* post-introduced with salicylic acid. *Acta Bot Boreal -Occident. Sin* 30 (8):1527-1533
- [20] Tong Z.G., Wang F.R., Zhang Z., Zhao J.B., Zhang K.C., Yan G.H., Zhou Y., Jiang L.J. 2008. A method for DNA extraction from mature leaves of fruit trees. *J. Fruit Sci.* 25, 122-125
- [21] Tamura K., Dudley J., Nei M., Kumar S. 2007. MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596-1599
- [22] Page R.D. 1996. Tree view: an application to display phylogenetic trees on personal computers. *Comput Appl Biosci* 12:357-358
- [23] Cai B.H., Zhang J.Y., Gao Z.H., Qu S.C., Tong Z.G., Mi L., Qiao Y.S., Zhang Z. 2008. An improved method for isolation of total RNA from the leaves of *Fragaria* spp. *Jiangsu Journal of Agricultural Sciences*. 24, 875-877
- [24] Livak K.J., Schmittgen T.D. 2001. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2- $\Delta\Delta$ CT Method. *Methods* 25: 402-408
- [25] Wang Q.J., Xu K.Y., Tong Z.G., Wang S.H., Gao Z.H., Zhang J.Y., Zong C.W., Qiao Y.S., Zhang Z. 2010. Characterization of a new dehydration responsive element binding factor in central arctic cowberry. *Plant Cell, Tissue and Organ Culture (PCTOC)* 101: 211-219
- [26] Wu J., Luo S., Jiang H., Li H. 2005. Mammalian CHORD-containing protein 1 is a novel heat shock protein 90-interacting protein. *FEBS Letters* 579: 421-426
- [27] Thao N.P., Chen L., Nakashima A., Hara S., Umemura K., Takahashi A., Shirasu K., Kawasaki T., Shimamoto K. 2007. RAR1 and HSP90 Form a Complex with Rac/Rop GTPase and Function in Innate-Immune Responses in Rice. *The Plant Cell Online* 19: 4035-4045
- [28] Zhang M., Kadota Y., Prodromou C., Shirasu K., Pearl L.H. 2010. Structural Basis for Assembly of Hsp90-Sgt1-CHORD Protein Complexes: Implications for Chaperoning of NLR Innate Immunity Receptors. *Molecular Cell* 39: 269-281
- [29] Eitas T.K., Nimchuk Z.L., Dangl J.L. 2008. Arabidopsis TAO1 is a TIR-NB-LRR protein that contributes to disease resistance induced by the *Pseudomonas syringae* effector AvrB. *Proceedings of the National Academy of Sciences of the USA*, 105: 6475-6480
- [30] Hubert D.A., He Y., McNulty B.C., Tornero P., Dangl J.L. 2009. Specific Arabidopsis HSP90.2 alleles recapitulate RAR1 cochaperone function in plant NB-LRR disease resistance protein regulation. *Proceedings of the National Academy of Sciences* 106: 9556-9563
- [31] Dangl J.L., Jones J.D. 2001. Plant pathogens and integrated defence responses to infection. *Nature* 411: 826-833
- [32] Austin M.J., Muskett P., Kahn K., Fey B.J., Jones J.D.G., Parker J.E. 2002. Regulatory Role of SGT1 in Early R Gene-Mediated Plant Defenses. *Science* 295: 2077-2080