
Hydrogen Sulphide Improves Iron Homeostasis in Wheat Under Iron-Deficiency

Xia Zhang, Yang Zhang, Lu Zhang, Huijie Zhao, Hua Li*

College of Life Science, Henan Agricultural University, Zhengzhou, China

Email address:

lihuahnd@163.com (Hua Li)

*Corresponding author

To cite this article:

Xia Zhang, Yang Zhang, Lu Zhang, Huijie Zhao, Hua Li. Hydrogen Sulphide Improves Iron Homeostasis in Wheat Under Iron-Deficiency. *Journal of Plant Sciences*. Vol. 5, No. 6, 2017, pp. 170-176. doi: 10.11648/j.jps.20170506.11

Received: October 9, 2017; **Accepted:** October 20, 2017; **Published:** November 24, 2017

Abstract: Hydrogen sulphide (H₂S) is emerging as an important gaseous molecule involved in various plant developmental processes and plant stress responses. In this study, exogenous H₂S donor (sodium hydrosulfide, NaHS) treated wheat plants were used to investigate the role of H₂S in response to iron-deficiency. The results showed that H₂S significantly alleviated leaf chlorosis under iron-deficient conditions, and thus improved photosynthesis. Moreover, H₂S increased the lateral root (LR) number, density and length of wheat seedlings grown in iron-sufficient and deficient culture solution, and promoted phytosiderophores (PSs) secretion from roots simultaneously, which eventually led to an increase in iron uptake. Taken together, these results indicate that H₂S improved iron uptake by regulating root development and PSs secretion, and consequently increased chlorophyll biosynthesis and photosynthesis in plants under iron-deficiency.

Keywords: Hydrogen Sulfide, Iron-Deficiency, Phytosiderophores, Lateral Root, Wheat

1. Introduction

Iron (Fe) is an essential nutrient for all organisms, due to its important role in various cellular metabolic processes. Although the abundance of iron in soils is high, iron availability in a form that can be absorbed and used by plants is limited. It has been estimated that up to 40% of the world's arable land may be iron-deficient [1], and the problem is particularly serious in alkaline soils, as iron availability decreases with increasing soil pH [2]. Iron-deficiency in plants has become a worldwide concern. It usually causes a decrease in chloroplast particles and thylakoid lamellae, disorders in the basal thymoid and matrix thylakoid arrangement, and even leads to chloroplast dissociation. Iron is an important participant in the redox reaction and electron transport in plants, therefore, iron-deficiency usually weakens photosynthesis and respiration, thus affecting plant yield and quality. Iron has been shown to be a limiting factor for biomass in important crop plants, such as tomato [3], spinach [4], and rice [5].

In order to adapt to iron-deficiency conditions, plants usually regulate the processes of iron absorption and transport.

Specifically, under iron-deficiency, *Strategy I* plants secrete protons to the rhizosphere by a plasma membrane H⁺-ATPase, thereby reducing the soil pH and improving iron availability. Subsequently, the ferric-chelate reductase (FCR) on the plasma membrane of root epidermal cells reduces Fe (III) to Fe (II), and then Fe (II) is transported to the cells through a specific Fe transporter. *Strategy II* plants (graminaceous species) increase iron absorption by inducing the release of phytosiderophores (PSs) in roots, which can effectively chelate Fe (III) around the rhizosphere to form Fe³⁺-PS complex. This Fe³⁺-PS complex is then transported into the cell through special membrane transporters (like *YS* and *YSL*) on the root surface [6]. Additionally, the processes of iron absorption and transport are regulated by plant hormones and signal molecules. Exogenous nitric oxide (NO) enhances iron availability and alleviates iron-deficiency symptoms in maize under low iron conditions [7]; Carbon monoxide (CO) facilitates the absorption of iron in *Arabidopsis thaliana* [8]; Auxin promotes the activity of ferric-chelate reductase in soybean [9], cucumber [10] and clover [11]; Ethylene

regulates the expression of iron transporter (IRT1) and ferric-chelate reductase (FRO2) by reducing FIT1 protein degradation [12].

Apart from NO and CO, H₂S is also an important gas signaling molecule [13-15], and its physiological function in plants has been extensively studied in recent years. H₂S regulates the physiological processes of plant stomatal closure, seed germination, root development and senescence [16-19], and it also induces responses to various abiotic stresses (e.g. drought, heavy metals, high temperature, etc.) [20-24]. Moreover, H₂S is also involved in NO, auxin and ethylene signal pathways. For instance, NO enhances the resistance of *Cynodon dactylon* to cadmium (Cd) stress by inducing H₂S production [25]; Auxin regulates lateral root development in tomato and *Arabidopsis thaliana* by inducing H₂S [26-27]. Finally, H₂S also mediates the ethylene-regulated stomatal closure in *Arabidopsis thaliana* [28].

Iron-deficiency leads to plant chlorosis, chlorophyll reduction, and photosynthesis inhibition. Studies have shown that H₂S can enhance rice and spinach photosynthesis under normal conditions [29-30], as well as improve chlorophyll fluorescence of strawberry and wheat when exposed to drought stress [24, 31]. Despite the documented involvement of H₂S in NO, auxin and ethylene signaling pathways, all of which induce responses to iron-deficiency, it is unclear whether H₂S participates in the regulation of plant iron nutrition. However, H₂S has been showed to promote iron uptake in rice roots and leaves under both normal conditions and Cd stress [32]. On the basis of these results, the effects of H₂S on wheat growth, chlorophyll content, photosynthesis, PSs release, and Fe content in wheat under Fe-deficient and sufficient conditions were investigated. Wheat was considered to be research object because it is an important crop and a *Strategy II* plant, which releases PSs under Fe shortage.

2. Methods and Materials

2.1. Plant Growth and Treatments

The seeds of common cultivated wheat varieties Aikang 58 were surface-sterilized and germinated on water soaked filter papers for 48 h, and then grown in 1/2 strength Hoagland nutrient solution. Three days later, the seedlings were transferred to Fe-sufficient (50 mM) and deficient (0 or 1 mM) nutrient solution with or without 0.4 mM NaHS, and the nutrient solution was changed every two days. Therefore, depending on treatment, four groups of wheat seedlings were included in this study: Fe-sufficiency and no NaHS treatment as the control group (+Fe), Fe-sufficiency and NaHS treatment (+Fe+NaHS), Fe-deficiency and no NaHS treatment (-Fe), Fe-deficiency and NaHS treatment (-Fe+NaHS). After two weeks, the third leaves were used for detecting chlorophyll content and photosynthesis (Pn), and all the leaves were used for H₂S and Fe content assays. The

seedlings grew under the conditions of 25/18°C (light/dark), 300 μmol m⁻² s⁻¹ light intensity, 70% humidity and a 14 h photoperiod.

2.2. Determination of Chlorophyll Content and Photosynthesis (Pn)

Total chlorophyll was extracted from leaves with 80% acetone until complete bleaching, and the content of chlorophyll was then calculated from the absorbance at 470, 646, and 663 nm, following Lichtenthaler [33]. Pn was measured using a portable photosynthesis system (Li-6400, NE, USA) on the third fully developed leaf of the seedlings.

2.3. Collection of Root Exudates and Determination of PSs Release

PSs release from seedling roots was analysed by determining PSs content in root washing. After treatment, the roots of wheat seedlings were washed twice with deionized water, and then root systems were transferred to 200 ml deionized water for 6 h. This solution of 200 ml was used as a sample to determine PSs content released from roots according to the Fe-binding assay as described by Reichman and Parker [34].

2.4. Detection of H₂S Content

H₂S quantification was performed as described by Christou [31]. Wheat leaves were powdered with liquid nitrogen, and H₂S was extracted using 100 mM potassium phosphate buffer (pH 7) containing 10 mM ethylene diamine tetraacetic acid (EDTA). After centrifugation of 12,000 g at 4°C for 15min, 100 μl of the supernatant was added to an assay mixture containing 1.88 ml of extraction buffer and 20 μl of 20 mM 5, 5'-dithiobis (2-nitrobenzoic acid). The absorbance was measured at 412 nm after a 2 min incubation time at room temperature. NaHS was used for standard curve.

2.5. Iron Quantification

Harvested seedlings were washed twice in 5 mM CaSO₄ and 10 mM EDTA solution and then dried at 70°C. The samples with a weight of at least 0.2 g were digested completely in 70% HNO₃ at 120°C. Finally, the solution was diluted to a certain volume with distilled deionized water. Iron was quantified with inductively coupled plasma spectrometry (Perkin Elmer Optimal 2100DV).

2.6. Statistical Analysis

All data were collected from at least three independent experiments. For photosynthesis measurement, at least six leaves were used. For other analyses, the results were the mean of three replicated treatments. Differences of the means among treatments were compared using Duncan's multiple range tests at 0.05 probability level.

3. Results

3.1. Phenotype, Chlorophyll Content and Photosynthesis in Wheat Leaves

Iron-deficiency affected chlorophyll synthesis, and resulted in leaf chlorosis. However, exogenous sodium hydrosulfide (NaHS, an H₂S donor) treatment significantly alleviated plant

chlorosis and increased chlorophyll content under iron-deficiency (Figure 1). Naturally, the reduction of chlorophyll caused a decline in photosynthesis under Fe-deficiency. Wheat seedlings under Fe-deficiency showed a 67.2% reduction in photosynthesis compared to the control group. However, that reduction was limited under treatment with H₂S, dropping to 51.4% of that in the control.




Figure 1. Effect of NaHS on phenotype (a), chlorophyll content (b) and photosynthesis (c) of iron-deficient wheat leaves. Wheat seedlings were grown in Fe-sufficient (50 mM) and deficient (0 mM) nutrient solution with or without 0.4 mM NaHS treatment for two weeks. After that, the third leaves were used to assay. Values are the means \pm standard deviation (SD). Different letters denote significant differences between treatments ($P < 0.05$).

3.2. Lateral Root (LR) Formation

Fe-deficiency usually causes changes in plant lateral roots. Therefore, several parameters of wheat lateral root (LR) formation, namely lateral root number, density and length were investigated. As shown in Figure 2, under iron-deficiency for two weeks, the number, density and length of LR increased significantly. Exogenously applied NaHS

further exacerbated this increase under both Fe-deficient and sufficient conditions. In comparison with the no-NaHS treatment group, treatment with NaHS increased the number, density and length of lateral roots. The increase was of 40.4%, 61.4%, 57.2%, respectively, under Fe-sufficiency, and of 32.7%, 263.0%, 60.3%, respectively, in Fe-deficient conditions.




Figure 2. Effect of NaHS on root phenotype (a), lateral root number (b), density (c) and length (d) of iron-deficient wheat plants. Wheat seedlings were grown in Fe-sufficient (50 mM) and deficient (0 mM) nutrient solution with or without 0.4 mM NaHS treatment for two weeks. After that, the third leaves were used to assay. Values are the means \pm standard deviation (SD). Different letters denote significant differences between treatments ($P < 0.05$).

3.3. Fe Content and Phytosiderophores (PSs) Release

Fe content significantly decreased in leaves and roots under Fe-limited conditions, while NaHS treatment improved iron accumulation under both Fe-sufficient and limited conditions (Table 1). In comparison with the no-NaHS treatment group, NaHS application promoted iron accumulation in leaves (1.73 times) and roots (1.43 times) under Fe-sufficiency. Iron accumulation increased also under Fe-limited conditions both in leaves (1.51 times) and roots (1.90 times).

The secretion of PSs from roots promotes iron absorption and, therefore, the content of PSs released by roots increased significantly under conditions of iron-deficiency, as expected.

The increase was up to 1.75 times compared to the control group, and up to 2.28 times under exposure to NaHS. Even under Fe-sufficient conditions, NaHS also increased PSs secretion by 1.42 times compared to the control group (Table 1).

Table 1. Effect of NaHS on Fe content and PSs release in iron-deficient wheat plants. Values are the means \pm standard deviation (SD). Different letters denote significant differences between treatments ($P < 0.05$).

Treatment	Fe content (mg g ⁻¹ DW)		PSs release (nmol g ⁻¹ RFW)
	Leaf	Root	
+Fe	0.081 \pm 0.006b	3.131 \pm 0.394b	12.350 \pm 3.445c
+Fe+NaHS	0.140 \pm 0.029a	4.486 \pm 0.412a	20.371 \pm 2.372b
-Fe	0.069 \pm 0.007c	0.216 \pm 0.038d	19.039 \pm 3.072b
-Fe+NaHS	0.104 \pm 0.019ab	0.411 \pm 0.079c	31.353 \pm 5.868a

3.4. H₂S Content in Shoots and Roots

H₂S concentration was determined in both shoots and roots of wheat plants grown in Fe-sufficient and limited culture solutions (Figure 3). The H₂S content increased in leaves (145%) and decreased in roots (50%) under Fe-limited conditions compared to the Fe-sufficient group. In comparison with the no-NaHS treatment group, exogenously applied NaHS increased H₂S content in leaves and roots under both Fe-limited and sufficient conditions. The increase was of 21.2% in leaves and 253.6% in roots under Fe-sufficiency, and of 41.9% and 307.1% respectively, in Fe-limited conditions.




Figure 3. H₂S content in iron-deficient wheat leaf (a) and root (b). Wheat seedlings were grown in Fe-sufficient (50 mM) and deficient (1 mM) nutrient solution with or without 0.4 mM NaHS treatment for two weeks. After that, all the leaves and roots were used to assay respectively. Values are the means \pm standard deviation (SD). Different letters denote significant differences between treatments ($P < 0.05$).

4. Discussion

Iron is one of the most important elements for plant growth, especially for chloroplast development, photosynthesis, respiration and DNA synthesis. Leaf yellowing is a typical phenotype caused by Fe-deficiency and first exhibits itself in young leaves. This was also shown in this study, where the content of chlorophyll in the youngest leaf (the third leaf) of wheat seedlings significantly decreased under Fe-deficiency conditions (Figure 1a, b). As chlorophyll is the basis of photosynthesis in plants, it is not surprising that, with

chlorophyll content decreased, Fe-deficiency naturally caused a significant decline in photosynthesis (Figure 1c). However, exogenous NaHS treatment significantly increased chlorophyll synthesis and photosynthetic rate in wheat leaves under Fe-deficiency, thus alleviating the effects of Fe-deficiency (Figure 1). These findings that H₂S can regulate chlorophyll synthesis and photosynthetic rate are supported by previous studies. For example, H₂S enhances photosynthesis through promoting chloroplast biogenesis, photosynthetic enzyme expression, and thiol redox modification in *Spinacia oleracea* seedlings [29]; H₂S can regulate plant growth and photosynthesis under Cr and Pd stress in barley and cotton, respectively [35-36]; A treatment with low concentration of NaHS can improve photosynthesis in rice by increasing its stomatal aperture and density [30].

Plants induce a variety of morphological and physiological changes to adapt to Fe-limited conditions [37]. One of those changes is the increase of lateral root (LR) development, which has been demonstrated in *Arabidopsis* and tomato [38-39]. Jin et al. [40] showed that the increase of root branching contributes to enhanced activity of the root ferric chelate reductase under iron-deficiency. Another study showed that NO acts downstream of auxin in regulating Fe-deficiency-induced root branching that enhances Fe-deficiency tolerance in tomato [41]. Recently, the role of H₂S in lateral root development has also been investigated. NaHS treatment induced pepper lateral root formation, while hypotaurine (H₂S scavenger) showed an adverse effect [42]. H₂S inhibits primary root, LR and root hair elongation but promotes LR initiation in *Arabidopsis thaliana* seedlings [27]. Further, auxin depletion-induced inhibition of lateral root formation is rescued by NaHS [26]. In this study, NaHS treatment significantly increased lateral root number, density and length under both Fe-deficiency and sufficiency conditions. These results, combined with previous findings, lead to the conclusion that the function of H₂S on improving wheat Fe-deficiency was related to the regulation of lateral root development.

Phytosiderophores (PSs) secretion from roots is known to occur under Fe-deficiency for *Strategy II* plants. Studies on grain crops and other grasses have shown that Fe-deficiency tolerance was strongly correlated to PSs content, which may facilitate iron absorption by forming ferric-phytosiderophore [43-46]. The results in this study lend support to this idea, as the increased PSs secretion induced by H₂S corresponded to higher iron accumulation in wheat leaves and roots under both Fe-deficient and sufficient conditions (Table 1).

H₂S, as a signal molecule, is known to regulate plant growth and development, and also induce responses to various stresses. In this study, results showed that H₂S content differed under Fe-limited and sufficient conditions (Figure 3). Low iron induced H₂S production in wheat leaves but decreased it in roots, while PSs secretion from roots increased under Fe-limited conditions. Moreover, the increase of H₂S content caused by exogenous NaHS in leaves was also accompanied by the increase of PSs secretion in wheat roots. Previous studies have proposed that plants' Fe-deficiency response was

related to sensors which are localized in leaves and roots [2, 47]. The results of this study, combined, suggest that H₂S concentration in shoots might play a major role in response to Fe-deficiency, potentially by being a sensor in the leaves that responds to Fe-deficiency.

5. Conclusion

The findings of this study showed that H₂S plays an important role in the response of wheat plants to Fe-deficiency. In particular, H₂S can improve PSs secretion and iron uptake, consequently increasing chlorophyll biosynthesis and photosynthesis. Although this study provides important insights into the role of H₂S in plants' response to Fe-deficiency, further studies are needed to understand the regulatory mechanism of H₂S on PSs secretion and iron uptake.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (31501238), The National Key Research and Development Program of China (2017YFD0300400), Scientific Research Foundation of the Higher Education Institutions of He'nan Province, China (15A180040) and Science and Technology Innovation Fund of Henan Agricultural University (KJCX2015A13).

References

- [1] Vose PB (1982) Iron nutrition in plants: a world overview. *J Plant Nutr* 5:233-249.
- [2] Schmidt (2003) Iron solutions: acquisition strategies and signaling pathways in plants. *Trends Plant Sci* 8:188-193.
- [3] Jin CW, Du ST, Chen WW, Li GX, Zhang YS, Zheng S J (2009) Elevated carbon dioxide improves plant iron nutrition through enhancing the iron-deficiency-induced responses under iron-limited conditions in tomato. *Plant Physiol* 150:272-280.
- [4] Jin CW, Liu Y, Mao QQ, Wang Q, Du S T (2013a). Mild Fe-deficiency improves biomass production and quality of hydroponic-cultivated spinach plants (*Spinacia oleracea* L.). *Food chem* 138:2188-2194.
- [5] Takahashi M, Nakanishi H, Kawasaki S, Nishizawa NK, Mori S (2001) Enhanced tolerance of rice to low iron availability in alkaline soils using barley nicotianamine aminotransferase genes. *Nat. Biotechnol* 19:466-469.
- [6] Walker EL and Connolly EL (2008) Time to pump iron: iron-deficiency-signaling mechanisms of higher plants. *Curr Opin Plant Biol* 11:530-535.
- [7] Graziano M, Beligni M V, Lamattina L. Nitric oxide improves internal iron availability in plants. *Plant Physiol*. 2002, 130: 1852-1859.
- [8] Kong WW, Zhang LP. Guo K, Liu ZP, Yang ZM (2010) Carbon monoxide improves adaptation of *Arabidopsis* to iron deficiency. *Plant Biotechnol J* 8:88-99.
- [9] Li C, Zhu X, Zhang F (2000) Role of shoot in regulation of iron deficiency responses in cucumber and bean plants. *J Plant Nutr* 23:1809-1818.
- [10] Li X, Li C (2004) Is ethylene involved in regulation of root ferric reductase activity of dicotyledonous species under iron deficiency. *Plant Soil* 261:147-153.
- [11] Zheng SJ, Tang CX, Arakama Y, Masaoka Y (2003) The responses of red clover (*Trifolium pratense* L.) to iron deficiency: a root Fe (III) chelate reductase. *Plant Sci* 164:679-687.
- [12] Lingam S, Mohrbacher J, Brumbarova T, Potuschak T, Fink-Straube C, Blondet E, Genschik P, Bauer P (2011) Interaction between the bHLH transcription factor FIT and ETHYLENE INSENSITIVE3/ETHYLENE INSENSITIVE3-LIKE1 reveals molecular linkage between the regulation of iron acquisition and ethylene signaling in *Arabidopsis*. *Plant Cell* 23: 1815-1829.
- [13] Wang R (2002) Two's company, three's a crowd: can H₂S be the third endogenous gaseous transmitter? *FASEB J* 16:1792-1798.
- [14] Olson KR (2009) Is hydrogen sulfide a circulating "gasotransmitter" in vertebrate blood? *BBA-Bioenergetics* 1787:856-863.
- [15] Tan BH, Wong PTH, Bian JS (2010) Hydrogen sulfide: a novel signaling molecule in the central nervous system. *Neurochem Int* 56:3-10.
- [16] Lisjak M, Teklić T, Wilson ID, Wood M, Whiteman M, Hancock JT (2011) Hydrogen sulfide effects on stomatal apertures. *Plant Signal Behav* 6:1444-1446.
- [17] Jin Z, Xue S, Luo Y, Tian B, Fang H, Li H, Pei Y (2013b) Hydrogen sulfide interacting with abscisic acid in stomatal regulation responses to drought stress in *Arabidopsis*. *Plant Physiol Bioch* 62:41-46
- [18] Dooley FD, Nair SP, Ward PD (2013) Increased growth and germination success in plants following hydrogen sulfide administration. *Plos one* 8:e62048
- [19] Xie Y, Zhang C, Lai D, Sun Y, Samma MK, Zhang J, Shen W (2014) Hydrogen sulfide delays GA-triggered programmed cell death in wheat aleurone layers by the modulation of glutathione homeostasis and heme oxygenase-1 expression. *J Plant Physiol* 171:53-62.
- [20] Zhang H, Hu LY, Hu KD, He YD, Wang SH, Luo JP (2008) Hydrogen sulfide promotes wheat seed germination and alleviates oxidative damage against copper stress. *J Integr Plant Biol* 50:1518-1529.
- [21] Zhang H, Hu LY, Li P, Hu KD, Jiang CX, Luo JP (2010a) Hydrogen sulfide alleviated chromium toxicity in wheat. *Biol Plantarum* 54:743-747.
- [22] Zhang H, Jiao H, Jiang CX, Wang SH, Wei ZJ, Luo JP, Jones RL (2010b) Hydrogen sulfide protects soybean seedlings against drought-induced oxidative stress. *Acta Physiol Plant* 32:849-857.
- [23] Li ZG, Yang SZ, Long WB, Yang GX, Shen ZZ (2013) Hydrogen sulphide may be a novel downstream signal molecule in nitric oxide-induced heat tolerance of maize (*Zea mays* L.) seedlings. *Plant Cell Environ* 36:1564-1572.

- [24] Li H, Gao MQ, Xue RL, Wang D, Zhao HJ (2015) Effect of hydrogen sulfide on D1 protein in wheat under drought stress. *Acta Physiol Plant* 37:225.
- [25] Shi H, Ye T, Chan Z (2014) Nitric oxide-activated hydrogen sulfide is essential for cadmium stress response in bermudagrass (*Cynodon dactylon* (L). Pers.). *Plant Physiol Biochem* 74: 99-107.
- [26] Fang T, Cao Z, Li J, Shen W, Huang L (2014) Auxin-induced hydrogen sulfide generation is involved in lateral root formation in tomato. *Plant Physiol Bioch* 76:44-51.
- [27] Jia H, Hu Y, Fan T, Li J (2015) Hydrogen sulfide modulates actin-dependent auxin transport via regulating ABPs results in changing of root development in *Arabidopsis*. *Sci Rep-UK* 5: srep08251.
- [28] Hou ZH, Wang LX, Liu J, Hou LX, Liu X (2013) Hydrogen Sulfide Regulates Ethylene-induced Stomatal Closure in *Arabidopsis thaliana*. *J Integr Plant Biol* 55:277-289.
- [29] Chen J, Wu FH, Wang WH, Zheng CJ, Lin GH, Dong XJ, He JX, Pei ZM, Zheng HL (2011) Hydrogen sulphide enhances photosynthesis through promoting chloroplast biogenesis, photosynthetic enzyme expression, and thiol redox modification in *Spinacia oleracea* seedlings. *J Exp Bot* 62: 4481-4493.
- [30] Duan B, Ma Y, Jiang M, Yang F, Ni L, Lu W (2015) Improvement of photosynthesis in rice (*Oryza sativa* L.) as a result of an increase in stomatal aperture and density by exogenous hydrogen sulfide treatment. *Plant Growth Regul* 75:33-44.
- [31] Christou A, Manganaris GA, Papadopoulos I, Fotopoulos V (2013) Hydrogen sulfide induces systemic tolerance to salinity and non-ionic osmotic stress in strawberry plants through modification of reactive species biosynthesis and transcriptional regulation of multiple defence pathways. *J Exp Bot* 64:1953-1966.
- [32] Mostofa MG, Rahman A, Ansary MMU, Watanabe A, Fujita M, Tran LSP (2015) Hydrogen sulfide modulates cadmium-induced physiological and biochemical responses to alleviate cadmium toxicity in rice. *Sci Rep* 5:14078.
- [33] Lichtenthaler HK (1987) Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Method Enzymol* 148:350-382.
- [34] Reichman SM, Parker DR (2007) Critical evaluation of three indirect assays for quantifying phytosiderophores released by the roots of Poaceae. *Eur J Soil Sci* 58:844-853.
- [35] Ali S, Farooq MA, Hussain S, Yasmeen T, Abbasi GH, Zhang G (2013) Alleviation of chromium toxicity by hydrogen sulfide in barley. *Environ Toxicol Chem* 32:2234-2239.
- [36] Bharwana SA, Ali S, Farooq MA, Ali B, Iqbal N, Abbas F, Ahmad MSA (2014) Hydrogen sulfide ameliorates lead-induced morphological, photosynthetic, oxidative damages and biochemical changes in cotton. *Environ Sci Pollut R* 21:717-731.
- [37] Marschner H, Römheld V, Kissel M (1986) Different strategies in higher plants in mobilization and uptake of iron. *J Plant Nutr* 9:695-713.
- [38] Moog PR, Kooij TA, Brüggemann W, Schiefelbein JW, Kuiper PJ (1995) Responses to iron deficiency in *Arabidopsis thaliana*: the turbo iron reductase does not depend on the formation of root hairs and transfer cells. *Planta* 195:505-513.
- [39] Dasgan HY, Römheld V, Cakmak I, Abak K (2002) Physiological root responses of iron deficiency susceptible and tolerant tomato genotypes and their reciprocal F₁ hybrids. *Plant Soil* 241:97-104.
- [40] Jin CW, Chen WW, Meng ZB, Zheng SJ (2008) Iron Deficiency-induced increase of root branching contributes to the enhanced root ferric chelate reductase activity. *J Integr Plant Biol* 50:1557-1562.
- [41] Jin CW, Du ST, Shamsi IH, Luo BF, Lin X Y (2011) NO synthase-generated NO acts downstream of auxin in regulating Fe-deficiency-induced root branching that enhances Fe-deficiency tolerance in tomato plants. *J Exp Bot* 62:3875-3884.
- [42] Xue YF, Zhang M, Qi ZQ, Li YQ, Shi, Chen J (2016) Cinnamaldehyde promotes root branching by regulating endogenous hydrogen sulfide. *J Sci Food Agr* 96:909-914.
- [43] Gries D, Brunn S, Crowley DE, Parker DR (1995) Phytosiderophore release in relation to micronutrient metal deficiencies in barley. *Plant Soil* 172:299-308.
- [44] Chaignon V, Di Malta D, Hinsinger P (2002) Fe-deficiency increases Cu acquisition by wheat cropped in a Cu-contaminated vineyard soil. *New Phytol* 154:121-130.
- [45] Meda AR, Scheuermann EB, Prechsl UE, Erenoglu B, Schaaf G, Hayen H, Weber G, von Wirén N (2007) Iron acquisition by phytosiderophores contributes to cadmium tolerance. *Plant Physiol* 143:1761-1773.
- [46] Neelam K, Tiwari VK, Rawat N, Tripathi SK, Randhawa GS, Dhaliwal HS (2010) Identification of *Aegilops* species with higher production of phytosiderophore and iron and zinc uptake under micronutrient-sufficient and-deficient conditions. *Plant Genet Resour-C* 8: 132-141.
- [47] Bocchini M, Bartucca ML, Ciancaleoni S, Mimmo T, Cesco S, Pii Y, Albertini E, Del Buono D (2015) Iron deficiency in barley plants: phytosiderophore release, iron translocation, and DNA methylation. *Front Plant Sci* 6.