
Metal complexes of Schiff bases derived from 2-thiophenecarboxaldehyde and mono/diamine as the antibacterial agents

Mohammad Nasir Uddin^{1,*}, Didarul Alam Chowdhury¹, Md. Moniruzzman Rony¹,
Md. Ershad Halim²

¹Department of Chemistry, University of Chittagong, Chittagong-4331, Bangladesh

²Department of Chemistry, University of Dhaka, Dhaka, Bangladesh

Email address:

nasircu72@gmail.com (M. N. Uddin)

To cite this article:

Mohammad Nasir Uddin, Didarul Alam Chowdhury, Md. Moniruzzman Rony, Md. Ershad Halim. Metal Complexes of Schiff Bases Derived from 2-Thiophenecarboxaldehyde and Mono/Diamine as the Antibacterial Agents. *Modern Chemistry*. Vol. 2, No. 2, 2014, pp. 6-13. doi: 10.11648/j.mc.20140202.11

Abstract: The Schiff bases, formed by condensation of 2-thiophenecarboxaldehyde with 2-aminothiophenol (LH) and propane-1,2-diamine (L₁), and their complexes of Ni(II), Cu(II), Zn(II) and Cd(II) have been prepared and characterized by elemental analysis, and magnetic and spectroscopic measurements. Infrared and NMR spectra of the complexes agree with the coordination to the central metal atom through the nitrogen of the azomethine (-HC=N-) group and the sulfur atom of the thiophene ring. Magnetic susceptibility data coupled with electronic suggest octahedral structure for the complexes. Conductance measurements suggest the non-electrolytic nature of the complexes. Stoichiometry of complexes has been suggested as [M(L)₂] and [ML₁Cl₂]. The Schiff base and its metal chelates have been screened for their in vitro antibacterial activity against four human pathogenic bacteria. Ligands show moderately whereas some of metal chelates show highly antibacterial activity against them.

Keywords: Metal Complexes, Schiff Bases, 2-Thiophenecarboxaldehyde, 2-Aminothiophenol, Propane-1, 2-Diamine, Antibacterial Activity

1. Introduction

Schiff base ligands can be synthesized from an amine and a carbonyl compound by nucleophilic addition forming a hemi-aminal group followed by dehydration to generate an imine compound [1]. Schiff bases, an important class of ligands plays an important role in the development of coordination chemistry as they can easily form stable complexes with most of the transition metals [2]. Metal chelates of the Schiff bases derived from salicylaldehyde and amino acids have been shown to be important class of compounds in elucidating the mechanism of transamination reaction in biological systems [3]. Metal complexes of the Schiff bases possess numerous applications including antibacterial, antifungal and other biological applications, as well as clinical, analytical and industrial in addition to their important roles in catalysis [4].

A large number of Schiff base complexes of various transition metal ions with a variety of donor atoms have

been reported. Synthesis, spectroscopic and electrochemical properties of some complexes of a new symmetric bidentate Schiff base ligand of bis ((3-(2-nitrophenyl) allylidene) propane-1,2-diamine (L) with a general formula of MLX₂ (M= Cd(II) and Hg(II); X= Cl-, Br-, I-, SCN- and NO₃-) are reported [5]. Several new dioxouranium(VI) complexes of Schiff bases, derived from salicylaldehyde, substituted salicylaldehydes, 2-hydroxy-1-naphthaldehyde and 2-aminothiophenol, have been prepared and characterized on the basis of their elemental analyses, IR and electronic absorption spectra and magnetic susceptibility measurements. The results suggest that each Schiff base is a bivalent anion with tridentate ONS donors from the phenolic oxygen, azomethine nitrogen and thiophenolic sulfur [6]. Some dioxouranium (VI) bis-complexes of ligands (LH); benzoylhydrazine (BH), salicyloylhydrazine (SaH), nicotinoylhydrazine (NH) and

their Schiff bases with acetone; isopropylbenzoylhydrazine (IPBH), isopropylsalicyloylhydrazine (IPSalH), isopropyl-nicotinoylhydrazine (IPNH) in their both keto and enol forms have been prepared [7].

Most of the tetradentate Schiff base ligands are formed by the condensation of β -diketones or *o*-hydroxyaldehydes or ketones with diamines. These ligands with ONNO donor atom set are well known to coordinate with various metal ions and this has attracted the interest of many authors [8]. The titanium(IV) complexes of the unsymmetrical Schiff base ligands of ethylenediamine and salicylaldehyde, *o*-hydroxyacetophenone, *o*-hydroxynaphthaldehyde have been prepared and characterized when unsymmetrical ligands are synthesized through *in situ* partial displacement of the symmetrical bis-Schiff bases [9].

Metal complexes [Fe(II), Co(II), Ni(II), Cu(II), Zn(II), or Cd(II)] of the Schiff bases formed by condensation of 2-thiophenecarboxaldehyde with 2-aminopyridine, *N*-(2-thienylmethylidene)-2-aminopyridine have been prepared and characterized [10]. Co(II), Ni(II) and Cu(II) complexes of three Schiff bases, derived from the condensation reaction of thiophene-2-carbaldehyde and DL-alanine, DL-valine and DL-phenylalanine, have been synthesized and characterized by their analytical and spectral data [3]. (TCPA 4) Copper (II) and Iron (III) complexes of Schiff base ligand namely (Z)-3-hydroxy-4-((5-methylthiophen-2-yl) methyleneamino) benzoic acid derived from the condensation of 5-methyl-2-thiophene carboxaldehyde with 4-amino-3-hydroxy benzoic acid were synthesized [4]. Mg(II), Ca(II) and Ba(II) metal chelates of the Schiff base moiety, formed from condensation of 2-thiophenecarboxaldehyde with 2-chlorophenyl hydrazine, have been prepared and characterized by elemental analysis and infrared and ¹H-NMR spectroscopic measurements [11]. CoII, NiII and CuII complexes of Schiff bases formed by condensation of 2-thiophenecarboxaldehyde and propylamine, *N*-[2-thienylmethylidene]-1-propanamine, or ethylamine, *N*-[2-thienylmethylidene]ethanamine, have been prepared and characterized [12]. The present work has been an attempt to study preparation, characterization and antibacterial activities some metal complexes using new tridentate and tetradentate Schiff base ligand having thiophenecarboxaldehyde moiety.

2. Experimental

2.1. Chemicals

2-aminothiophenol, 1,2-propane-di-amine, 2-thiophenecarboxaldehyde were obtained from M/S. E. Merk (Germany). Methanol, chloroform, N, N-dimethyl formamide (DMF), dimethylsulphoxide (DMSO) and carbon-tetrachloride were obtained from Aldrich Chemical Company Ltd. Ethyl alcohol was obtained from Carew and

company Ltd. Perchloric acid, nitric acid, sulphuric acid and disodium salt of ethylenediaminetetraacetic acid were obtained from BDH Chemicals Ltd. All chemicals except solvent were used as received. Solvents were dried by standard methods and distilled under an inert atmosphere.

2.2. Preparation of Schiff Base Ligands

2.2.1. 2-Thiophenecarboxaldehyde -2-Aminothiophenol, TPCA-ATPH (LH)

2-aminothiophenol contains one $-NH_2$ group attached to the aromatic ring with a sulfhydryl group (-SH) in 2 position. An active carbonyl group reacts with $-NH_2$ group of 2-Aminothiophenol yielding a Schiff base product containing azomethine ($>C=N-$) function. A hot solution (60 °C) of 2-aminothiophenol (10 mmol) in 25 mL of ethanol was mixed with a hot solution (60°C) of 2-thiophenecarboxaldehyde (10 mmol) in the same solvent and the reaction mixture was left under reflux for one hr. The solid product formed was separated by filtration, purified by crystallization from ethanol and washed with diethyl ether and then dried in a vacuum desiccator over anhydrous calcium chloride.

2.2.2. 2-Thiophenecarboxaldehyde-1, 2-Propanediamine, TPCA-Pn-TPCA (L₁)

1,2-propanediamine contains two amino ($-NH_2$) groups attached with propane group at 1,2- positions. An active carbonyl group reacts with $-NH_2$ group of 1, 2-propanediamine yielding a Schiff base product containing azomethine ($>C=N-$) function. 2-thiophenecarboxaldehyde (30 mmol) was dissolved in 60 mL of ethanol and 1,2-propanediamine (15 mmol) was added drop wise to this solution with continuous stirring. The solution was refluxed for about one hour. Then it was cooled, allowed to stand in a deep freeze for cooling over some days (3-4). Brown product was formed after cooling. The product was then filtered off, washed with ethanol and dried in a desiccator over anhydrous calcium chloride.

The color, yield and melting point of the prepared ligands are given in Table 1.

2.3. Synthesis of Metal Complexes

The metal complexes of the Schiff bases, LH and L₁, were prepared by the addition of a hot solution (60°C) of the appropriate metal chloride, nitrate, or acetate (1 mmol) in an ethanol-water mixture (1:1, 25 mL) to the hot solution (60°C) of the Schiff bases (2 mmol LH and 1 mmol L₁) in the same solvent (25 mL). The resulting mixture was stirred under reflux for 1 h whereupon the complexes precipitated. They were collected by filtration, washed with a 1:1 ethanol-water mixture and diethyl ether. The color, Yield, melting point and % of element content and elemental analysis of the prepared complexes are given in Table 1.

Table 1. Color, % of Yield, M. P. and % of Metal contents of the prepared Schiff bases and their metal complexes

Complexes	Color	Yield (%)	M.P. (°C)	% of Metal (Calc.)	Elemental analysis		
					%C	%H	%N
TPCA-ATP	Yellow	80	180	-	59.97 (60.27)	4.32 (4.11)	6.19 (6.39)
[Ni(TPCA-ATP) ₂]	Red	88	>250	14.79 (15.5)	50.07 (49.72)	3.55 (3.77)	5.55 (5.27)
[Cu(TPCA-ATP) ₂]	Black	86	>250	12.63 (11.5)	52.09 (52.85)	3.01 (3.2)	5.35 (5.61)
[Zn(TPCA-ATP) ₂]	Brown	80	>300	25.7 (25.55)	52.21 (52.69)	3.58 (3.19)	5.53 (5.59)
[Cd(TPCA-ATP) ₂]	Yellow Brown	87	>250	20.37 (20.92)	-	-	-
TPCA-pn-TPCA	brown	85	160	-	-	-	-
[Ni(TPCA-pn-TPCA)Cl ₂]	Brown	85	>300	14.91 (13.75)	39.63 (38.25)	4.06 (4.28)	7.11 (6.58)
[Cu(TPCA-pn-TPCA)Cl ₂]	Brown	85	>300	15.93 (14.85)	39.14 (40.81)	4.02 (4.25)	7.03 (5.95)
[Zn(TPCA-pn-TPCA)Cl ₂]	Light Brown	85	>250	16.31 (17.01)	-	-	-
[Cd(TPCA-pn-TPCA)Cl ₂]	Brown	88	>250	25.13 (25.51)	34.87 (33.95)	3.58 (3.18)	6.26 (5.99)

2.4. Test for Antibacterial Evaluation

Synthesized complexes were screened by the agar well diffusion method for their antibacterial activity against different types of bacteria, gram positive- *Bacillus cereus* and gram negative- *Salmonella typhi*, *pseudomonas aeruginosa*, *Escherichia coli* regarded as pathogen to man. All media and bacteria suspension were prepared using a suitable method. The in-vitro evaluation of antimicrobial activities was performed according to the diffusion technique [13]. The bacteria were grown in nutrient broth at 37°C for 24 hours. The complexes were tested using diffusion on solid media. Sterile (5 mm) diameter sensitivity paper disc were impregnated with concentration of dimethylformamide (DMF) and their bimetallic complexes at concentration of 50 µg cm⁻³ and placed in the nutrient agar. The plates were then incubated for 24 hours. The results were recorded by measuring the growth inhibition (% zones of inhibition) surrounding the disc.

2.5. Analytical Methods


The analyses of the metal contents of the prepared complexes were obtained by Atomic Absorption Spectrophotometer (model Thermo Scientific ICE-3000) from B.C.S.I.R. Laboratory, Chittagong, Bangladesh. The Infrared spectra of the prepared ligands and complexes were obtained by FTIR spectrophotometer (Model-883, Perkin-Elmer) using KBr as the matrix in the range 400-4000 cm⁻¹ from research laboratory of Department of Chemistry, University of Chittagong. Electronic absorption spectra were run on Shimadzu UV-Visible Recording Spectrophotometer (Model-1800) using 1 cm cell in the research laboratory of the Department of Chemistry, University of Chittagong. An electrothermal melting point apparatus was used for the determination of the melting or decomposition points of the complexes and the ligands. Using N, N-dimethylformamide (DMF) as the solvent the solutions of the complexes (of the order of 10⁻³ M) were

used for conductivity measurements. Conductivity measurement was performed on a Philips Conductivity Meter (Model-HI 9255). Determination of conductivity of an electrolytic solution involves measuring of the electrical resistance of that solution at a particular temperature, usually 25°C. Magnetic susceptibility values of some of the prepared complexes were determined using the Magnetic Susceptibility Balance in the research laboratory of the Department of Chemistry, University of Chittagong. Elemental analysis data and mass spectra were obtained from Shiumyozu's Laboratory, Department of Applied Chemistry at Kyushu University, Japan, by Ultra light Performance Mass Spectrometer (Model- JMS-HX 11 OA).


3. Results and discussion

3.1. Schiff Bases Characterization

The tridentate Schiff base TPCA-ATPH (L_H) was prepared by the condensation of 2-thiophenecarboxaldehyde (TPCA) and 2-aminothiophenol (ATP) in their 1:1 molar ratio. Another tetradentate Schiff base TPCA-pn-TPCA (L₁) was prepared by the condensation of 1,2-propanediamine (pn) with 2-thiophenecarboxaldehyde (TPCA) in their 1:2 molar ratio. Both ligands were obtained after reflux and then on cooling as brownish precipitates. The Schiff bases were subjected to elemental analyses. The results of elemental analyses (C, H, N) with molecular formulae and the melting points are presented in Table 1. The results obtained are in good agreement with those calculated for the suggested formulae and the melting points are sharp, indicating the purity of the prepared Schiff bases. The structure of these Schiff bases is also confirmed by mass, IR and ¹H NMR spectra. The mass spectrum of L_H shows a well-defined parent peak at m/z = 218 (M⁺), with a relative intensity = 15%. And that of L₁ shows a well-defined parent peak at m/z = 264, with a relative intensity = 5%. The parent ion and the fragments obtained



Scheme 2. The representative diagram for the preparation of some Schiff base ligand, (TCPA-ATPH) and its metal (M) complexes



Scheme 3. The representative diagram for the preparation of some Schiff base ligand, (TCPA-pn-TCPA) and its metal (M) complexes

3.2.2. Elemental Analyses of the Complexes

The results of elemental analyses, as shown in Table 1, are in good agreement with those required by the proposed formulae. The formation of these complexes may proceed according to the following equations given in scheme 2, 3.

3.2.3. Infrared Spectra

The infrared spectra for the present compounds taken in the range 400-4000 cm^{-1} help to indicate regions of absorption vibrations. The main stretching modes are for $\nu_{\text{S-H}}$, $\nu_{\text{C=N}}$, $\nu_{\text{C=C}}$, $\nu_{\text{C-N}}$, $\nu_{\text{C-S}}$, $\nu_{\text{M-N}}$ and $\nu_{\text{M-S}}$. The IR data of the spectra of Schiff base ligands (LH and L_1) and their complexes are presented in Table 2. The IR spectra of the complexes were compared with those of the free ligands in order to determine the coordination sites that may be involved in chelation. The position and/or the intensities of these peaks are expected to change upon chelation. The peak between 1620 and 1580 cm^{-1} have been assigned to $\nu_{\text{C=N}}$ band [8]. This band shows a downward shift by about 25-35 cm^{-1} in the spectra of all the metal complexes, indicating the participation of the azomethine nitrogen in coordination with metals [5, 8,9]. Upon comparison, it was determined that the $\nu(\text{C=N})$ stretching vibration is found in the free ligands at 1604 and 1633 cm^{-1} for the LH and L_1 ligands, respectively. This band was shifted to higher or lower wave numbers in the complexes, indicating the participation of the azomethine nitrogen in coordination (M-N) [10]. The bands appearing between 1520 and 1600 cm^{-1} have been assigned to aromatic or thiophene $\nu_{\text{C=C}}$. The shifted band in many cases is coincident with the C=C band, which then shows greater intensity or broadening. The $\nu_{\text{C=N}}$ modes are often mixed with higher frequency $\nu_{\text{C=C}}$ (near 1600 cm^{-1}) and are seen as a strong band [5,9]. The bands appearing at 1315-1400 cm^{-1} (specially the highest frequency ones near 1400 cm^{-1}) have been assigned to $\nu_{\text{C-N}}$ mode [8,9]. It was observed that the symmetric vibrations of C-S, which appeared as a band near 701 cm^{-1} in the ligand spectrum, has been shifted to lower frequencies after complexation. Similarly a weak band of S-H, which appeared at 2650 cm^{-1} in the case of ligand, has been disappeared in the spectrum of all complexes [6]. However, the participation of the SH group in chelation is ascertained from the shift of the $\nu_{\text{asym(C-S)}}$ and $\nu_{\text{sym(C-S)}}$ from 706 cm^{-1} and 758 cm^{-1} to lower or higher wave numbers in the spectra of the complexes [14]. The sharp IR ligand bands at 824 cm^{-1} assigned as $\nu_{\text{(C-S-C)}}$ of thiophene moiety, for the ligand, shifted to 827-828 cm^{-1} for metal complexes [3]. The $\nu_{\text{(M-S)}}$ bands appeared at 467-468 (thiophene) and 428-431 cm^{-1} (thiophenol) for metal complexes [6]. Spinu and Kriza [12] have prepared Schiff base complexes who observed the bands at 563-567 cm^{-1} in have been assigned to $\nu_{\text{(M-N)}}$ mode and the $\nu_{\text{(M-S)}}$ bands appeared at 428-431 cm^{-1} for the complexes. New bands at 507-619 cm^{-1} in LH and L_1 metal complexes, have been assigned to $\nu(\text{M-N})$ mode. The $\nu(\text{M-S})$ bands appeared at 420-430 cm^{-1} (thiophene) and 440-469 cm^{-1} (thiophenol) for LH metal complexes [6,12]. Therefore, from the IR spectra, it is

concluded that the LH ligand behaves as a uninegatively tridentate ligand coordinated to the metal ions via deprotonated thiophenol S, azomethine N, and thiophene S;

whereas L₁ behaves as a neutral tetradentate ligand coordinated to the metal ions via azomethine N and thiophene S.

Table 2. Infrared spectral bands of Schiff bases and their metal complexes

Complexes	$\nu_{C=N}$	$\nu_{C=C}$	ν_{C-N}	ν_{C-S-C}	ν_{C-S} (sym)	ν_{C-S} (asym)	ν_{M-N}	ν_{M-S} (-ol)	ν (M-S) (-phene)
TPCA-ATPH	1604(s)	1550 m	1425(sh)	893(vs)					
Ni(TPCA-ATP) ₂	1635(m) 1645(m)	1581m	1386(w) 1338(m)	802(s)	755s	710w	617(m) 507(w)	469m	430m
Cu(TPCA-ATP) ₂	1680(w) 1640(ms)	1562m	1435(ms) 1420ms)	859(vs)	750s	705s	579(w)	440(w)	422(w)
Zn(TPCA-ATP) ₂	1625(m) 1616(ms)	1575s	1458(ms) 1388(ms)	813(s)	760m	715s	615(s)	468m	430m
Cd(TPCA-ATP) ₂	1635(ms)	1562m	1436(vs)	859(s)	758m	709m	472(w)	456m	420(w)
TPCA-pn-TPCAH ₂	1633(vs)	1550ms	1425(sh)	898(vs)					
[Ni(TPCA-pn-TPCA)Cl ₂]	1650(m) 1640(m)	1577s	1388(s) 1338(m)	827sh	763m	725s	509(m)		428m
[Cu(TPCA-pn-TPCA)Cl ₂]	1635(vs) 1613(s)	1568m	1388(ms)	828sh	753s	705s	615(s)		429m
[Zn(TPCA-pn-TPCA)Cl ₂]	1635(vs)	1581s	1338(ms)	831(s)	745s	712w	649(s)		431m
[Cd(TPCA-pn-TPCA)Cl ₂]	1635(w)	1581s	1388(m)	827(m)	758m	709m	617(ms)		430m

3.2.4. ¹H NMR Spectra

The ¹H NMR spectra of the free ligands and their diamagnetic Zn(II) complex shows that the SH signal appeared in the spectrum of the LH ligand at 3.36 ppm and completely disappeared in the spectrum of its Zn(II) complex, indicating that the SH proton is removed by chelation with the Zn(II) ion. The peaks at 8.71 and 7.09 ppm characteristic to the azomethine and thiophene protons are down fielded to 8.30 and 6.87 ppm respectively in the complexes supporting well binding of the azomethine groups of ligand to metal ions. The signals for aromatic hydrogens at 7.91-7.79 ppm are down fielded to 7.52-7.39 ppm.

3.2.5. Electronic Spectra

The absorption spectra for 2.5×10^{-3} M solution of LH and L₁ ligands in absolute ethanol ranging from 200 to 700 nm against the same solvent as a blank. LH gives 3 sharp bands at 216, 278 and 337 nm and L₁ shows 3 bands at 253, 316 and 402 nm. These bands can be attributed to $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ transitions within the Schiff bases. Being a d¹⁰ system, the present Cd(II) and Zn(II) complexes show no d-d transition in the visible region, and are colored only through their intense charge transfer absorptions tailing in from the ultraviolet. Besides, the other bands are due to intra ligand transitions. The present Ni(II) complexes show spectral bands around 400-550 nm which may be attributed to the d-d transition. The high energy peaks are tentatively assumed as due to $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$ and $n \rightarrow \sigma^*$ transitions. Electronic spectra of the Cu(II) complex displayed bands at 286-563 417nm which may be assigned to $\pi \rightarrow \pi^*$ transition of thiophene ring, L → Cu charge transfer, $n \rightarrow \pi^*$ transition of azomethine group and d-d transitions.

3.2.6. Magnetic Measurement

From the experimental values of magnetic measurements of the prepared complexes, it is revealed that Cd(II) and Zn(II) complexes of the Schiff bases are diamagnetic, thus indicating the d¹⁰ electronic configuration of Cd(II) and Zn(II). The Ni(II) complexes reported herein are high-spin with room temperature magnetic moment values of 3.69-4.12 B.M., which are in the normal range observed for octahedral Ni(II) complexes ($\mu_{eff} = 2.9-3.3$ B.M.). The Cu(II) complex was found to be paramagnetic as expected for d⁹ configuration. The magnetic moment value is 1.71-1.72 BM is very close to the spin only value 1.73, which support that a octahedral geometry of Cu(II) complex is most likely.

3.2.7. Conductance Measurement

The chelates were dissolved in DMF and the molar conductivities of 10^{-3} M of their solutions at 25 ± 2 °C were measured. The molar conductivity values of the metal chelates of LH and L₁ ligands ($\Lambda_m = 10-29 \Omega^{-1} \text{ mol}^{-1} \text{ cm}^2$) indicate that these complexes are non-electrolytes. Based on these data, it is clear that chelates are considered as 1: 2 [M(L)₂], indicating the nonionic nature of the bonding of the complex nucleus. The +2 oxidation states of M(II) is satisfied by two uni-negatively charged tridentate (TPCA-ATPH) ligands as expected. Since the Schiff base ligand (TPCA-pn-TPCA) contains no charge, so +2 oxidation states of M(II) are satisfied by two negatively charged monodentate Cl⁻ ligands in the complexes of [M(L)₁Cl₂] residing in coordination sphere.

Table 3 shows the electronic spectral data, magnetic moments and molar conductivity for the prepared complexes.

Table 3. The electronic spectra, magnetic moments and conductivity data for the prepared complexes

No.	Complexes	Medium	Spectral bands (nm)	μ_{eff} (B.M)	$\lambda_{\text{m}}^{\circ}$ $\text{ohm}^{-1} \text{cm}^2 \text{mol}^{-1}$
	TPCA-ATPH		337, 278, 216		
1	[Ni(TPCA-ATP)]		550, 453, 426	4.12(para)	23
2	[Cu(TPCA-ATP) ₂]		563, 433, 388,	1.71 (para)	21
3	[Zn(TPCA-ATP)]		353, 318, 286,	0.47 (Dia.)	29
4	[Cd(TPCA-ATP) ₂]	DMF	333, 281, 273,	0.68 (Dia.)	15
	TPCA-pn-TPCA		402, 316, 263		
5	[Ni(TPCA-pn-TPCA)Cl ₂]		540, 438, 352	3.69 (para)	15
6	[Cu(TPCA-pn-TPCA)Cl ₂]		560, 322, 286	1.72 (para)	14
7	[Zn(TPCA-pn-TPCA)Cl ₂]		281, 273	0.51 (Dia.)	10
8	[Cd(TPCA-pn-TPCA)Cl ₂]		317, 271	0.55 (Dia.)	10

3.3. Structural Interpretation

The structures of the complexes of Schiff bases LH and L₁, with Ni(II), Cu(II), Zn(II), and Cd(II) ions were confirmed by elemental analyses, UV-Vis, mass, IR, ¹H NMR, molar conductance and magnetic analysis data. Therefore, from the IR spectra, it is concluded that LH behaves as a uni-negatively tridentate ligand, coordinated to the metal ions via azomethine N, thiophene S and deprotonated thiophenol S. L₁ behaves as a neutral terdentate ligand, coordinated to the metal ions via azomethine N, and thiophene S. From the molar conductance data, it was found that the M(II) chelates are considered as 2:1 (L:M) composition with L ligand. While complexes with the L₁ ligand are considered as 1:1 composition. Non-electrolytic nature is satisfied by two negatively charged monodentate Cl⁻ ligands. After coordination, all signals in ¹H NMR spectra are down fielded. On the basis of the above observations and from the magnetic measurements, octahedral geometries are suggested for the investigated complexes with the formulae, [M(L)₂] and [M(L₁)Cl₂].

3.4. Stability Constants

The stability constants of the Ni(II), Cu(II), Zn(II) and Cd(II) complexes with HL are determined potentiometrically using the method described by Bjerrum [15]. The complex-forming abilities of the metal ions are frequently characterized by stability orders. The order of stability constants was found to be Cd²⁺ < Ni²⁺ < Cu²⁺ > Zn²⁺ for divalent metal ions. The stability of Cu(II) complex is considerably more as compared to other metals ions. Under the influence of the ligand field, Cu(II) (3d⁹) will receive some extra stabilization due to tetragonal distortion of the octahedral symmetry in its complexes. The Cu(II) complex will be further stabilized due to the Jahn-Teller effect.

3.5. Effect of Chemicals on the Bacterial Growth

In the present work, Nickel(II), Copper(II), Zinc(II) and

Cadmium(II) complexes were selected for antibacterial activity against four human pathogenic bacteria. Antibacterial activity of complexes was studied on one Gram positive and three Gram-negative bacteria by disc diffusion method. The selected bacteria can cause different fatal diseases; viz. *Salmonella typhi* causes typhoid, each of *Escherichia coli* and *Bacillus cereus* cause gastroenteritis. The inhibition zones of the test organisms for different complexes are represented in Figure 1. The results are graphically shown in Figures 2, 3 and for the complexes of the respective ligand. Both ligands themselves are found to be moderately effective against tested bacteria. Most cases chelation increases their antibacterial activity than free ligands. Nickel(II) complexes of LH are effective against the Gram-positive and Gram-negative bacteria except *Escherichia coli* whereas that of L₁ are effective for both *Escherichia coli* and *pseudomonas aeruginosa*. Copper(II) complexes of ligand LH exhibit effective antibacterial activity against all bacteria except *Bacillus cereus* whereas that of L₁ are quite ineffective to almost all bacteria. Zinc(II) complexes of ligand LH are moderately effective to all bacteria except *Bacillus cereus* whereas that of L₁ are more effective for both *Escherichia coli* and *pseudomonas aeruginosa*. All the cadmium(II) complexes are quite effective against the *Escherichia coli* and *Salmonella typhi* and ineffective against *pseudomonas aeruginosa*. The higher inhibition zone of metal complexes than those of the ligands can be explained on the basis of overtone's concept and Chelation theory. On chelation, the polarity of the metal ion will be reduced to greater extent due to the overlap of the ligand orbital and partial sharing of the positive charge of the metal ions with donor groups. Further, it increases the delocalization of π -electrons over the whole chelation ring and enhances the penetration of the complexes into lipid membranes and blocking of the metal bonding sites in the enzymes of microorganisms. There are other factors which also increase the activity are solubility, conductivity and bond length between the metal and ligand.




Fig. 1. Inhibition zone against *Escherichia coli* by the complexes R_3 - $[Zn(TPCA-ATP)_2]$ and R_8 - $[Cd(TPCA-pn-TPCA)Cl_2]$.




Fig. 2. The graphical presentation of the antibacterial activity against tested bacteria by the complexes of ligand, TPCA-ATP (HL).




Fig. 3. The graphical presentation of the antibacterial activity against tested bacteria by the complexes of ligand, TPCA-pn-TPCA (L1).

4. Conclusion

The new Schiff bases were derived by the condensation of 2-thiophenecarboxaldehyde with 2-aminothiophenol (L_H) and propane-1,2-diamine (L₁). The complexes of the types [M(L)₂] with tridentate Schiff base (L_H) ligand and [M(L₁)Cl₂] with tetradentate Schiff base ligand (L₁) were prepared [M = Ni(II), Cu(II), Zn(II) and Cd(II)]. The analytical data confirmed the metal : ligand stoichiometry of the complexes. Conductivity measurements indicate their non-electrolytic nature. The magnetic measurements along with electronic spectral data of the prepared complexes support the electronic configuration of the central metal ion consistent with the +2 oxidation states of Ni(II), Cu(II), Zn(II) and Cd(II) ions. The synthesized metal complexes, in comparison to the uncomplexed Schiff base ligand, were screened for their antibacterial activity against pathogenic bacteria species. The activity of the some Schiff base complexes became more pronounced when coordinated with the metal ions.

Acknowledgement

Authors are pleased to acknowledge the Department of Microbiology, University of Chittagong, Bangladesh for cordial cooperation in collecting bacterial strains and performing antibacterial tests.

References

- [1] M Vikotora; K Rudolfa; L Molbank. M354 Open Access Publication, 2004.
- [2] A Ahmed; MA El-Sherif; T Eldebss. *Spectrochimica Acta Part A*, 2011, 79, 1803-1814.
- [3] N Sari; P Gürkan. *Z. Naturforsch.* 2004, 59b, 692 – 698.
- [4] K Jamuna; BR Naik; B Sreenu; K Sessaiah. *J. Chem. Pharm. Res.*, 2012, 4(9), 4275-4282.
- [5] M Montazerozohori; M Sedighipoor; S Joohari. *Int. J. Electrochem. Sci.*, 2012, 7, 77– 88.
- [6] DA Chowdhury; MN Uddin; F Hoque. *Chiang Mai J. Sci.* 2010, 37(3), 443-450.
- [7] DA Chowdhury; MN Uddin; MAH Sarker; *Chiang Mai J. Sci.* 2008, 35(3), 483-494.
- [8] DA Chowdhury; MN Uddin; GK Chowdhury. *The Chitt. Uni. J. of Sci.* 2006, 30(1), 85-88.
- [9] MN Uddin; DA Chowdhury; K Hossain, *J. Chinese Chem. Soc.*, 2012, 59(12), 1520-1527.
- [10] C S[□]Inu; M Pleniceanu; C Tigae. *Turk J Chem*, 2008, 32, 487– 493.
- [11] OB Ibrahim; MA Mohamed; MS Refat. *Int. J. of Innovative Res. in Sci., Eng. and Tech.* 2013, 2(11), 6355-6570.
- [12] C Spinu; A Kriza; *Acta Chim. Slov.* 2000, 47, 179-185.
- [13] MN Uddin; DA Chowdhury; MT Islam; F Hoque. *Orbital Elec. J. Chem.* 2012, 4(4), 273-287.
- [14] AP Mishra; A Tiwari; R Jain. *Int. J. Res. Pharm. Sci.* 2011, 3(2), 186-191.
- [15] J Bjerrum. "Metal Amine Formation in Aqueous Solution, Haase, Copenhagen, 1941.