
Modular cone metric spaces

Saeedeh Shamsi Gamchi, Asadollah Niknam

Department of Pure Mathematics, Ferdowsi University of Mashhad, P.O. Box 1159-91775, Mashhad, Iran

Emails address:

saeedeh.shamsi@gmail.com (S. S. Gamchi), dassamankin@yahoo.co.uk (A. Niknam)

To cite this article:

Saeedeh Shamsi Gamchi, Asadollah Niknam. Modular Cone Metric Spaces. *Pure and Applied Mathematics Journal*. Vol. 2, No. 6, 2013, pp. 191-196. doi: 10.11648/j.pamj.20130206.14

Abstract: In this paper a generalization of a modular metric which is also a generalization of cone metric, is introduced and some of its topological properties are studied. Next, a fixed point theorem in this space is proved and finally by an example, it is proved that the fixed point result of the paper "Ch. Mongkolkeha, W. Sintunavarat, P. Kumam, Fixed point theorems for contraction mapping in modular metric spaces, *Fixed Point Theory Appl.* (2011)" is not true.

Keywords: Ordered Spaces, Modular Cone Metric, Fixed Point Theorem

1. Introduction and Preliminaries

The theory of modular spaces was initiated by Nakano [11] in 1950 in connection with the theory of order spaces and reddened and generalized by Musielak and Orlicz [13] in 1959. In 2008, Chistyakov [5] introduced the notion of modular metric spaces generated by F-modular and developed the theory of this spaces, on the same idea he defined the notion of a modular on an arbitrary set and developed the theory of metric spaces generated by modular such that called the modular metric spaces in 2010 [6].

Ordered normed spaces and cones have applications in applied mathematics and optimization theory [7]. Replacing the real numbers, as the codomain of metrics, by ordered Banach spaces one may obtain a generalization of metric spaces. Such generalized spaces called cone metric spaces, were introduced by Rzepecki [12]. Recently, some authors have studied cone metric spaces over solid topological vector spaces and fixed point theorems in such spaces. (see [1], [8]).

In this paper, which is split into two parts, our aim is to develop the theory of cone metric spaces called modular cone metric spaces. In the first part, the notion of modular cone metric space is introduced and some properties of such spaces are investigated. Also we define convex modular cone metric which takes values in $C_{\mathbb{R}}(Y)$ where Y is a compact Hausdorff space. Then a fixed point theorem is proved for contractions in these spaces. In the second part of this paper, we make a remark on paper [10] and it will be proved that their fixed point result in modular metric spaces is not true.

Let E be a topological vector space (TVS, for short) with its zero vector θ . A nonempty subset P of E is called a convex cone if $P + P \subseteq E$ and $\lambda P \subseteq P$ for all $\lambda \geq 0$. A convex cone P is said to be pointed if $P \cap (-P) = \{\theta\}$. For a given convex cone P in E , a partial ordering \preceq on E with respect to P is defined by $x \preceq y$ if and only if $y - x \in P$. We shall write $x < y$ if $x \preceq y$ and $x \neq y$, while $x \ll y$ will stand for $y - x \in \text{int}P$, where $\text{int}P$ denotes the topological interior of P .

In the sequel, we will need the following useful lemmas:

Lemma 1.1.

[9] Let P be a cone in E . Then:

(i) If $\theta \preceq a_n \rightarrow \theta$, then for each $c \in \text{int}P$, there exists $N \in \mathbb{N}$ such that for every $n > N$, $a_n \ll c$.

(ii) For every $c_1, c_2 \in \text{int}P$, there exists $c \in \text{int}P$ such that $c \ll c_1$ and $c \ll c_2$.

(iii) For every $a \in P$ and $c \in \text{int}P$, there exists $n_0 \in \mathbb{N}$ such that $a \ll n_0 c$.

The nonlinear scalarization function $\xi_e: E \rightarrow \mathbb{R}$ is defined as follows:

$$\xi_e(y) = \inf\{r \in \mathbb{R}: y \in re - P\},$$

for all $y \in E$.

Lemma 1.2.

[2] Let $r \in \mathbb{R}$ and $y \in E$, then

(i) $\xi_e(y) \leq r \Leftrightarrow re - y \in P$,

(ii) $\xi_e(y) < r \Leftrightarrow re - y \in \text{int}P$,

(iii) $\xi_e(\cdot)$ is positively homogeneous and continuous on E ,

- (iv) If $y_1 \in y_2 + P$, then $\xi_e(y_2) \leq \xi_e(y_1)$,
- (v) $\xi_e(y_1 + y_2) \leq \xi_e(y_1) + \xi_e(y_2)$.

Definition 1.3.

[8] Let X be a nonempty set and $d: X \times X \rightarrow E$ be a mapping that satisfies:

(CM1) For all $x, y \in X$, $d(x, y) \geq \theta$ and $d(x, y) = \theta$ if and only if $x = y$,

(CM2) $d(x, y) = d(y, x)$ for all $x, y \in X$,

(CM3) $d(x, y) \leq d(x, z) + d(z, y)$ for all $x, y, z \in X$.

Then d is called a topological vector space cone metric (TVS cone metric, for short) on X and (X, d) is said to be a topological vector space cone metric space.

Definition 1.4.

[6] Let X be a nonempty set. A function $w: (0, \infty) \times X \times X \rightarrow [0, \infty]$ is said to be a modular metric on X , if it satisfies the following three axioms:

(i) for given $x, y \in X$, $w_\lambda(x, y) = 0$ for all $\lambda > 0$ if and only if $x = y$,

(ii) $w_\lambda(x, y) = w_\lambda(y, x)$ for all $x, y \in X$ and $\lambda > 0$,

(iii) $w_{\lambda+\mu}(x, y) \leq w_\lambda(x, z) + w_\mu(z, y)$ for all $x, y, z \in X$ and $\lambda, \mu \in (0, \infty)$.

If, instead of (i), the function w satisfies only

(i') $w_\lambda(x, x) = 0$ for all $\lambda > 0$ and $x \in X$,

then w is said to be a pseudo modular on X , and if

$$X_w^* = X_w^*(x_0) = \{x \in X: \exists \lambda = \lambda(x) > 0, \text{ such that } w_\lambda(x, x_0) < \infty\}$$

are said to be modular spaces (around x_0).

If w is a convex modular on X , then according to [6], Section 3.5 and Theorem 3.6, the two modular spaces coincide, $X_w = X_w^*$.

Definition 1.6.

[10] Let w be a modular metric on X , $x \in X_w$ and $\{x_n\}$ be a sequence in X_w . Then

(1) $\{x_n\}$ is said to be modular convergent to x if for every $\lambda > 0$, $w_\lambda(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.

(2) $\{x_n\}$ is said to be a modular Cauchy sequence if $\lambda > 0$, $w_\lambda(x_n, x_m) \rightarrow 0$ as $n, m \rightarrow \infty$

(3) X_w is called a complete modular metric space if every modular Cauchy sequence is modular convergent.

Theorem 1.7.

[3] Let w be a convex modular metric on X and X_w^* be a complete modular metric space. Suppose that $T: X_w^* \rightarrow X_w^*$ is a mapping which satisfies the following condition:

$$w_{k\lambda}(Tx, Ty) \leq w_\lambda(x, y) \text{ for all } 0 < \lambda \leq \lambda_0.$$

where $\lambda_0 > 0$ and $0 < k < 1$ are constant. Then T has a fixed point. If in addition, the modular w assumes only finite values on X_w^* , then the fixed point of T is unique.

w satisfies (i') and

(i'') for given $x, y \in X$, if there exists $\lambda > 0$ such that $w_\lambda(x, y) = 0$, then $x = y$.

The function w is called a strict modular on X .

A modular (pseudomodular, strict modular) w on X is said to be convex if, instead of (iii), for all $\lambda, \mu > 0$ and $x, y, z \in X$ it satisfies the inequality

$$w_{\lambda+\mu}(x, y) \leq \frac{\lambda}{\lambda+\mu} w_\lambda(x, z) + \frac{\mu}{\lambda+\mu} w_\mu(z, y).$$

Note that the conditions (i) to (iii) imply that for all $y, z \in X$ and $\lambda > 0$, $w_\lambda(y, z) \geq 0$. If $w_\lambda(x, y)$ does not depend on $x, y \in X$, then by (i) $w \equiv 0$. Now if $w_\lambda(x, y) = w(x, y)$ is independent of $\lambda > 0$, then axioms (i)-(iii) mean that w is a metric on X .

The essential property of a pseudo modular w on X (cf. [6], Section 2.3) is that, for any given $x, y \in X$, the function $0 < \lambda \mapsto w_\lambda(x, y)$ is decreasing on $(0, \infty)$.

Definition 1.5.

[6] Let w be pseudo modular on X , the sets

$$X_w = X_w(x_0) = \{x \in X: w_\lambda(x, x_0) \rightarrow 0 \text{ as } \lambda \rightarrow \infty\}$$

and

2. Convergence in Modular Cone Metric Spaces

In the following, unless otherwise specified, we always suppose that E is a locally convex Hausdorff TVS with its zero vector θ , P a proper closed and convex pointed cone in E with $\text{int}P \neq \emptyset$, $e \in \text{int}P$ and \leq the partial ordering with respect to P .

Throughout this paper functions $w: \text{int}P \times X \times X \rightarrow E$ will be written as $w(c, x, y) = w_c(x, y)$ for all $c \in \text{int}P$ and $x, y \in X$.

Definition 2.1.

Let X be a nonempty set. A function $w: \text{int}P \times X \times X \rightarrow E$ is said to be a modular cone metric on X , if it satisfies the following three axioms:

(i) For given $x, y \in X$, $w_c(x, y) = \theta$ for all $c \in \text{int}P$ if and only if $x = y$,

(ii) $w_c(x, y) = w_c(y, x)$, for all $x, y \in X$ and $c \in \text{int}P$,

(iii) $w_{c_1+c_2}(x, y) \leq w_{c_1}(x, z) + w_{c_2}(z, y)$ for all $x, y, z \in X$ and $c_1, c_2 \in \text{int}P$.

(X, E, P, w) is called a modular cone metric space.

Note that the conditions (i) to (iii) imply that for all $y, z \in X$ and $c \in \text{int}P$, $\theta \leq w_c(y, z)$. Indeed, by setting $x = y$ and $c_1 = c_2 = c$ in (iii), for all $y, z \in X$ one gets

$$\theta = w_{2c}(y, y) \leq 2w_c(y, z).$$

If $w_c(x, y)$ does not depend on $x, y \in X$, then by (i) $w \equiv 0$. Now if $w_c(x, y) = w(x, y)$ is independent of $c \in \text{int}P$, then axioms (i)-(iii) mean that w is a cone metric on X .

For given $x, y \in X$, the function $\theta \ll c \mapsto w_c(x, y)$ is decreasing on $\text{int}P$. In fact, if $\theta \ll c_1 \ll c_2$, then

$$w_{c_2}(x, y) \leq w_{c_2 - c_1}(x, x) + w_{c_1}(x, y) = w_{c_1}(x, y).$$

By the following example, we may construct a class of examples of modular cone metric spaces using a cone

$$w_{c_1 + c_2}(x, y) = \varphi(c_1 + c_2)d(x, y) \leq \varphi(c_1 + c_2)d(x, z) + \varphi(c_1 + c_2)d(z, y) \leq \varphi(c_1)d(x, z) + \varphi(c_2)d(z, y) = w_{c_1}(x, z) + w_{c_2}(z, y)$$

Now we define some convergence concepts in this space. It will be proved that these definitions are compatible with some topology on X constructed by a modular cone metric.

Definition 2.3.

Let (X, E, P, w) be a modular cone metric space, $x \in X$ and $\{x_n\}$ be a sequence in X . Then

(1) $\{x_n\}$ is said to be modular cone convergent to x and we denote it by $x_n \xrightarrow{w} x$ if for every $c \gg \theta$ there exists a positive integer N such that for all $n > N$, $w_c(x_n, x) \ll c$.

(2) $\{x_n\}$ is said to be a modular cone Cauchy sequence if for every $c \gg \theta$ there exists a positive integer N such that for all $m, n > N$, $w_c(x_n, x_m) \ll c$.

(3) (X, w) is called a complete modular cone metric space if every modular cone Cauchy sequence is convergent.

Let E be a Banach space and P be a cone in E . P is called normal if there exists a constant $K > 0$ such that for all $a, b \in P$, $a \leq b$ implies that $\|a\| \leq K \|b\|$. The least positive number satisfying the above inequality is called the normal constant of P .

In the next theorem some equivalent condition for convergence is proved.

Theorem 2.4.

Let E be a Banach space and P be a normal cone with normal constant K . Suppose that (X, E, P, w) is a modular cone metric space and $\{x_n\}$ is a sequence in X . Then,

(i) $x_n \xrightarrow{w} x$ if and only if for each $c \in \text{int}P$, $\|w_c(x_n, x)\| \rightarrow 0$ as $n \rightarrow \infty$.

(ii) $\{x_n\}$ is modular cone Cauchy if and only if for each $c \in \text{int}P$, $\|w_c(x_n, x_m)\| \rightarrow 0$ as $n, m \rightarrow \infty$.

Proof. Let $x_n \xrightarrow{w} x$. Fix $c \in \text{int}P$. Then for every $0 < \varepsilon < 1$, there exists a positive integer N such that for all $n > N$, $w_{\varepsilon c}(x_n, x) \ll \varepsilon c$. On the other hand, so for each n we have $w_c(x_n, x) \leq w_{\varepsilon c}(x, y)$, since $\varepsilon c \ll c$. Hence, $w_c(x_n, x) \ll \varepsilon c$, for each $n > N$. Therefore, normality of P implies that $\|w_c(x_n, x)\| \leq K \varepsilon \|c\|$, for each $n > N$. For proving its converse, let for each $c \in \text{int}P$, $\|w_c(x_n, x)\| \rightarrow 0$ as $n \rightarrow \infty$. For a given $c \in \text{int}P$, by assumption we have $w_c(x_n, x) \rightarrow \theta$ as $n \rightarrow \infty$. Applying Lemma 1.1 there is a positive integer N such that for all $n > N$, $w_c(x_n, x) \ll c$. Thus the proof of (i) is complete.

For proving (ii) let $\{x_n\}$ be a modular cone Cauchy

metric.

Example 2.2.

Let (X, d) be a cone metric space and $\varphi: \text{int}P \rightarrow (0, \infty)$ be a decreasing function. Define $w_c(x, y) = \varphi(c)d(x, y)$. It is easy to see that w is a modular cone metric on X . Indeed, by the properties of a cone metric d axioms (i)-(ii) of Definition 2.1 are satisfied. On the other hand, for all $x, y, z \in X$ and $c_1, c_2 \in \text{int}P$, we have:

sequence and fix $c \in \text{int}P$. Similar to (i) one can show that $\|w_c(x_n, x_m)\| \rightarrow 0$ as $n, m \rightarrow \infty$, for each $c \in \text{int}P$. Conversely, let $w_c(x_n, x_m) \rightarrow \theta$ as $n, m \rightarrow \infty$, for each $c \in \text{int}P$. For given $c \in \text{int}P$, there exists $\varepsilon > 0$ such that $c - y \in \text{int}P$ for each $y \in B_\varepsilon(\theta)$. For this $\varepsilon > 0$, there is a positive integer N such that for all $n, m > N$, $\|w_c(x_n, x_m)\| < \varepsilon$. Therefore, for all $n, m > N$,

$$c - w_c(x_n, x_m) \in \text{int}P,$$

that means $w_c(x_n, x_m) \ll c$ for all $n, m > N$. Thus the proofs complete. \square

Now we are going to construct a topology on X , where (X, E, P, w) is a modular cone metric space. For any $x \in X$, $c \in \text{int}P$, set $B_w(x, c) = \{y \in X: W_c(x, y) \ll c\}$.

Theorem 2.5.

Let (X, E, P, w) be a modular cone metric space. Then

$$\tau_w = \{U \subset X: \forall x \in U \exists c \in \text{int}P \text{ s.t. } B_w(x, c) \subset U\}$$

forms a Hausdorff topology on X .

Proof. Trivially $\emptyset, X \in \tau_w$.

Also for any $U, V \in \tau_w$ and $x \in U \cap V$, then $x \in U$ and $x \in V$, one may find $c_1, c_2 \in \text{int}P$ such that $x \in B_w(x, c_1) \subset U$ and $x \in B_w(x, c_2) \subset V$. By Lemma 1.1, there exists $c \in \text{int}P$ such that $c \ll c_1$ and $c \ll c_2$. Now suppose that $y \in B_w(x, c)$, so we have $w_{c_1}(x, y) \leq w_c(x, y) \ll c \ll c_1$, hence, $y \in B_w(x, c_1)$. Similarly we have $y \in B_w(x, c_2)$. Thus

$$B_w(x, c) \subset B_w(x, c_1) \cap B_w(x, c_2) \subset U \cap V.$$

Therefore $U \cap V \in \tau_w$.

Let $U_\alpha \in \tau_w$ for each $\alpha \in \Delta$, and let $x \in \bigcup_{\alpha \in \Delta} U_\alpha$, then $\exists \alpha_0 \in \Delta$ such that $x \in U_{\alpha_0}$. Hence, $x \in B(x, c) \subset U_{\alpha_0}$, for some $c \in \text{int}P$. That is $\bigcup_{\alpha \in \Delta} U_\alpha \in \tau_w$. Thus τ_w is a topology on X .

Now we prove that this topology is a Hausdorff topology. Suppose that $x, y \in X$ and $x \neq y$. So by the property (i) of Definition 2.1 there exists $c \in \text{int}P$ such that $w_{c_0}(x, y) \neq \theta$. In contrary, we assume that for each $c \in \text{int}P$,

$$B_w(x, c) \cap B_w(y, c) \neq \emptyset.$$

Hence, for each $n \in \mathbb{N}$, there is a $z_n \in X$ such that

$$z_n \in B_w(x, \frac{c_0}{2n}) \cap B_w(y, \frac{c_0}{2n}).$$

$$w_{c_0}(x, y) \leq w_{\frac{c_0}{n}}(x, y) \leq w_{\frac{c_0}{2n}}(x, z_n) + w_{\frac{c_0}{2n}}(x, z_n) \ll \frac{c_0}{2n} + \frac{c_0}{2n} = \frac{c_0}{n}$$

Therefore for each $n > 1$, $\frac{c_0}{n} - w_{c_0}(x, y) \in P$. But P is closed, so

$$\lim_{n \rightarrow \infty} (\frac{c_0}{n} - w_{c_0}(x, y)) = -w_{c_0}(x, y) \in P.$$

This is a contradiction since $P \cap (-P) = \{\theta\}$. So the proof is complete. \square

One can easily see that the collection $\{B_w(x, c) : x \in X, c \in \text{int}P\}$ forms a basis for τ_w under which the above definitions of convergent and Cauchy sequences are fully justified.

Theorem 2.6.

Let (X, E, P, w) be a modular cone metric space. Then

- (i) (X, E, P, w) is first countable.
- (ii) A self-map $f : X \rightarrow X$ is continuous at $x \in X$ if and only if whenever $x_n \rightarrow x$, we have $f(x_n) \rightarrow f(x)$, as $n \rightarrow \infty$.

Proof. Let $x \in X$. Fix $c \in \text{int}P$. We show that $\beta_x = \{B_w(x, \frac{c}{n}) : n \in \mathbb{N}\}$ is a local base at x . Let U be an open set such that $x \in U$. Therefore, there is a $c_1 \in \text{int}P$ such that $B_w(x, c_1) \subset U$. On the other hand there exists a positive integer n , such that $\frac{c}{n} \ll c_1$. Now suppose that $y \in B_w(x, \frac{c}{n})$,

$$W_{\lambda_1 + \lambda_2}^e(x, y) = \xi_e(w_{\lambda_1 e + \lambda_2 e}(x, y)) \leq \xi_e(w_{\lambda_1 e}(x, z)) + \xi_e(w_{\lambda_2 e}(z, y)) = W_{\lambda_1}^e(x, z) + W_{\lambda_2}^e(z, y).$$

Theorem 2.8.

Let (X, E, P, w) be a modular cone metric space, $e \in \text{int}P$, $x \in X$ and $\{x_n\}$ be a sequence in X . Then

- (i) $x_n \rightarrow x$ if and only if for every $\lambda > 0$, $W_{\lambda}^e(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- (ii) $\{x_n\}$ is a w -Cauchy sequence if and only if for every $\lambda > 0$, $W_{\lambda}^e(x_n, x_m) \rightarrow 0$ as $n, m \rightarrow \infty$.

Proof. (i) Let $x_n \rightarrow x$. Fix $\lambda > 0$. For a given $\epsilon > 0$ if $\lambda > \epsilon$, then for $c = \epsilon e$, there is a positive integer N such that for each $n > N$, $w_{\epsilon e}(x_n, x) \ll \epsilon$. On the other hand, so $w_{\lambda e}(x_n, x) \leq w_{\epsilon e}(x_n, x) \ll \epsilon e$, since $\epsilon e \ll \lambda e$. Hence by the property (iv) of Lemma 1.2, for each $n > N$ we have

$$W_{\lambda}(x_n, x) \leq W_{\epsilon}(x_n, x) = \xi_e(w_{\epsilon e}(x_n, x)) < \xi_e(\epsilon) = \epsilon.$$

Now suppose that $\lambda \leq \epsilon$. For $c = \lambda e$, there is a positive integer N such that for each $n > N$, $w_{\lambda e}(x_n, x) \ll \lambda e$. Hence, for each $n > N$ we have

$$W_{\lambda}(x_n, x) = \xi_e(w_{\lambda e}(x_n, x)) < \xi_e(\lambda e) = \lambda < \epsilon.$$

For its converse, let for each $\lambda > 0$, $W_{\lambda}^e(x_n, x) \rightarrow 0$. Fix $c \in \text{int}P$. By Lemma 1.1, there exists a positive real number λ_0 such that $\lambda_0 e \ll c$. So, there is a positive integer N

Thus for each $n > 1$, we have

so we have $w_{c_1}(x, y) \leq w_{\frac{c}{n}}(x, y) \ll \frac{c}{n} \ll c_1$, hence, $y \in B_w(x, c_1)$. Thus $B_w(x, \frac{c}{n}) \subset B_w(x, c_1) \subset U$.

The second part is trivial using (i). \square

The following useful theorem shows that we may construct a family of modular metrics using a modular cone metric and the mapping ξ_e .

Theorem 2.7.

Let (X, E, P, w) be a modular cone metric space and $e \in \text{int}P$. Then $W^e : (0, \infty) \times X \times X \rightarrow [0, \infty)$ which is defined by $W_{\lambda}(x, y) = \xi_e(w_{\lambda e}(x, y))$ is a modular metric on X .

Proof. Let $x, y \in X$. If $x = y$, then $w_{\lambda e}(x, y) = \theta$ for each $\lambda > 0$, so $W_{\lambda}^e(x, y) = \xi_e(w_{\lambda e}(x, y)) = 0$ for each $\lambda > 0$. Now, let for each $\lambda > 0$, $W_{\lambda}^e(x, y) = 0$. So $w_{\lambda e}(x, y) = \theta$ for each $\lambda > 0$. On the other hand for each $c \in \text{int}P$ there exists a $\lambda > 0$ such that $\lambda e \ll c$, hence $w_c(x, y) \leq w_{\lambda e}(x, y) = \theta$. Therefore, $w_c(x, y) = \theta$, for each $c \in \text{int}P$. That means $x = y$. So we prove that $x = y$ if and only if $W_{\lambda}^e(x, y) = 0$, for each $\lambda > 0$.

It is easy to see that $W_{\lambda}^e(x, y) = W_{\lambda}^e(y, x)$, for all $\lambda > 0$ and $x, y \in X$.

Also if $x, y, z \in X$ and $\lambda_1, \lambda_2 > 0$, then

such that for each $n > N$,

$$\xi_e(w_{\lambda_0 e}(x_n, x)) = W_{\lambda_0}(x_n, x) < \lambda_0.$$

Thus by property (ii) of Lemma 1.2 and the fact that the function $\theta \ll c \mapsto w_c(x, y)$ is decreasing on $\text{int}P$, we have

$$w_c(x_n, x) \leq w_{\lambda_0 e}(x_n, x) \ll \lambda_0 e \ll c,$$

for each $n > N$. The proof of (ii) is similar to (i). \square

3. A Fixed Point Theorem for Contractions in Convexmodular Cone Metric Spaces

In this section we suppose that $E = C_{\mathbb{R}}(Y)$ where Y is a compact Hausdorff space. E with the sup-norm is a real Banach space. Put $P = \{f \in C_{\mathbb{R}}(Y) : f(x) \geq 0 \text{ for any } x \in Y\}$. It is clear that P is a cone in E . Thus the partial order on E with respect to P is of the form

$$f \leq g \Leftrightarrow f(x) \leq g(x) \forall x \in Y.$$

Note that $\text{int}P \neq \emptyset$. Indeed, $1 \in \text{int}P$.

Lemma 3.1.

If $f \in \text{int}P$, then f is invertible.

Proof. Let $f \in \text{int}P$. Then there exists $\epsilon > 0$ such that

$$B_\epsilon(f) = \{g \in C_{\mathbb{R}}(Y) : \|g - f\|_\infty < \epsilon\} \subset P.$$

If, in contrary, there is a $t_0 \in Y$ such that $f(t_0) = 0$, then by defining $g(t) = f(t) - \frac{\epsilon}{2}$, it is clear that $g \in B_\epsilon(f)$ and hence $g \in P$. But $g(t_0) < 0$ and this is a contradiction. Thus $0 \notin \text{rang}(f)$ and so f is invertible. \square

Now we introduce the concept of convex modular cone metric spaces.

Definition 3.2.

Let X be a nonempty set. A function $w: \text{int}P \times X \times X \rightarrow E$ is said to be a convex modular cone metric on X , if it satisfies the following three axioms:

- (i) For given $x, y \in X$, $w_c(x, y) = \theta$, for all $c \in \text{int}P$, if and only if $x = y$,
- (ii) $w_c(x, y) = w_c(y, x)$, for all $x, y \in X$ and $c \in \text{int}P$,
- (iii) $w_{c_1+c_2}(x, y) \leq \frac{c_1}{c_1+c_2} w_{c_1}(x, z) + \frac{c_2}{c_1+c_2} w_{c_2}(z, y)$, for all $x, y, z \in X$ and $c_1, c_2 \in \text{int}P$.

(X, w) is called a convex modular cone metric space.

Example 3.3.

Let (X, d) be a cone metric space. Define $w_c(x, y) = \frac{d(x, y)}{c}$. It is easy to see that w is a convex modular cone metric on X .

Note that every convex modular cone metric space is a modular cone metric space.

If instead of (i), the function w satisfies in (i') and (i''), then it is called strict convex modular cone metric.

(i') $w_c(x, x) = \theta$ for all $c \in \text{int}P$ and $x \in X$,

(i'') Given $x, y \in X$, if there exists $c \in \text{int}P$ such that $w_c(x, y) = \theta$, then $x = y$.

Remark 3.4.

Let (X, w) be a convex modular cone metric space and $1 \in E = C_{\mathbb{R}}(Y)$ be the unit constant function. It is easy to see that $W_\lambda^1(x, y) = \xi_1(w_{\lambda 1}(x, y))$ is a convex modular metric on X and $W_\lambda^1(x, y) < \infty$ so $X_w^* = X$.

Theorem 3.5.

Let (X, w) be a complete convex modular cone metric space. Suppose that there exists a constant number $k \in (0, 1)$ and $c_0 \in \text{int}P$ such that a mapping $T: X \rightarrow X$ satisfies the following condition:

$$w_{kc}(Tx, Ty) \leq w_c(x, y), \text{ for all } \theta \ll c \ll c_0.$$

Then T has a unique fixed point.

Proof. Take $\lambda_0 = \|c_0\|_\infty$. For all $0 < \lambda < \lambda_0$ by the above inequality for convex modular metric $W_\lambda^1(x, y) = \xi_1(w_{\lambda 1}(x, y))$ we have

$$W_{k\lambda}^1(x, y) = \xi_1(w_{k\lambda 1}(x, y)) \leq \xi_1(w_{\lambda 1}(x, y)) = W_\lambda^1(x, y).$$

(X_w^*, W) is complete since $X_w^* = X$ and (X, w) is complete. Hence, the proof is complete by Theorem 1.7. \square

4. A Note on the Paper [10]

Recently, in the paper "Ch. Mongkolkeha, W. Sintunavarat, P. Kumam, Fixed point theorems for contraction mapping in modular metric spaces, Fixed Point Theory and Applications, (2011)", the authors have studied and introduced some fixed-point theorems in the framework of a modular metric space. We will first state the main result that the authors have proved in that paper and then by constructing an example we will show that the result is not true.

Definition 4.1.

(Definition 3.1.[10]) Let w be a modular metric on X and X_w be a modular metric space induced by w and $T: X_w \rightarrow X_w$ be an arbitrary mapping. A mapping T is called a contraction if for each $x, y \in X_w$ and for all $\lambda > 0$ there exists $0 < k < 1$ such that

$$w_\lambda(Tx, Ty) \leq kw_\lambda(x, y).$$

Theorem 4.2.

(Theorem 3.2.[10]) Let w be a modular metric on X and X_w be a modular metric space induced by w . If X_w is a complete modular metric space and $T: X_w \rightarrow X_w$ is a contraction mapping, then T has a unique fixed point in X_w . Moreover, for any $x \in X_w$, iterative sequence $\{T_n x\}$ converges to the fixed point.

Now by the following example we show that the above theorem is not valid.

Example 4.3.

Let $X = \{(a, 0) \in \mathbb{R}^2 : \frac{1}{2} \leq a \leq 1\} \cup \{(0, b) \in \mathbb{R}^2 : \frac{1}{2} \leq b \leq 1\}$. Defined the mapping $w: (0, \infty) \times X \times X \rightarrow [0, \infty)$ by

$$w_\lambda((a_1, 0), (a_2, 0)) = \frac{4|a_1 - a_2|}{3\lambda}$$

$$w_\lambda((0, b_1), (0, b_2)) = \frac{|b_1 - b_2|}{\lambda},$$

and

$$w_\lambda((a, 0), (0, b)) = \frac{4a}{3\lambda} + \frac{b}{\lambda} = w_\lambda((0, b), (a, 0)).$$

We note that if we take $\lambda \rightarrow \infty$, then we see that $X = X_w$ and also it is easy to see that if $\{x_n\}$ is a Cauchy sequence in X_w then we just have one of the following assertions:

(1) There exists a positive integer N such that for each $n > N$, $x_n \in \{(a, 0) \in \mathbb{R}^2 : \frac{1}{2} \leq a \leq 1\}$.

(2) There exists a positive integer N such that for each $n > N$, $x_n \in \{(0, b) \in \mathbb{R}^2 : \frac{1}{2} \leq b \leq 1\}$.

Indeed, it follows from the fact that for $(a, 0), (0, b) \in X$,

$$w_\lambda((a, 0), (0, b)) = \frac{4a}{3\lambda} + \frac{b}{\lambda} \geq \frac{7}{3\lambda} = w_\lambda\left(\left(\frac{1}{2}, 0\right), \left(0, \frac{1}{2}\right)\right).$$

Thus X_w is a complete modular metric space. Now we define a mapping $T: X_w \rightarrow X_w$ by

$$T((a, 0)) = (0, a)$$

and

$$T((0, b)) = \left(\frac{b}{2}, 0\right)$$

Simple computations show that

$$w_\lambda(T((a_1, b_1)), T((a_2, b_2))) \leq \frac{3}{4} w_\lambda((a_1, b_1), (a_2, b_2)),$$

for all $(a_1, b_1), (a_2, b_2) \in X_w$. But T does not have any fixed point in X_w .

Acknowledgements

The first author would like to thank her advisor, Dr. Mohammad Janfada, for his valuable comments on this paper.

References

- [1] H. Cakalli, A. Sonmez, C. Genc, On an equivalence of topological vector space valued cone metric spaces and metric spaces. *Appl. Math. Lett.* 25 (2012) 429-433.
- [2] G.Y. Chen, X.X. Huang, X.Q. Yang, *Vector Optimization*. Springer-Verlag, Berlin, Heidelberg. (2005).
- [3] V. Chistyakov, A fixed point theorem for contractions in modular metric spaces. *arXiv: 1112.5561v1*. (2011)
- [4] V. Chistyakov, Metric modulars and their application. *Dokl. Math.* 73 (1)(2006) 32-35.
- [5] V. Chistyakov, Modular metric spaces generated by F-modulars, *Folia Math.* 14 (2008) 3-25.
- [6] V. Chistyakov, Modular metric spaces, I: Basic concepts. *Nonlinear Anal.* 72 (2010) 1-14.
- [7] K. Deimling, *Nonlinear Functional Analysis*. Springer-Verlag, Berlin. (1985).
- [8] W.S. Du, A note on cone metric fixed point theory and its equivalence. *Nonlinear Anal.* 72. (2010) 2259-2261.
- [9] S. Jankovic, Z. Golubovic, S. Radenovic, Compatible and weakly compatible mappings in cone metric spaces. *Math. Comp. Modelling.* 52(2010) 1728-1738.
- [10] Ch. Mongkolkeha, W. Sintunavarat, P. Kumam, Fixed point theorems for contraction mapping in modular metric spaces. *Fixed Point Theory Appl.* (2011)
- [11] H. Nakano, *Modular Semi-Ordered Linear Spaces*. In *Tokyo Math Book Ser, vol. 1. Maruzen Co., Tokyo.* (1950).
- [12] B. Rzepecki, On fixed point theorems of Maia type. *Publications de l'Institut Mathématique*, 28 (42)(1980) 179-186.
- [13] J. Musielak, W. Orlicz, On modular spaces. *Studia Math.* 18 (1959) 591-597.