



# Measurement of Different Heavy Metals Concentration in Roadside Dust in the Vicinity of Gujrat, Pakistan

Nusrat Bashir<sup>1</sup>, Mujahid Farid<sup>1</sup>, Rashid Saeed<sup>1</sup>, Hafiz Muhammad Tauqeer<sup>1</sup>, Shafaqat Ali<sup>2</sup>, Muhammad Rizwan<sup>2</sup>, Ayesha Siddiqua<sup>1</sup>, Rasham Sallah-Ud-Din<sup>1</sup>

<sup>1</sup>Department of Environmental Sciences, University of Gujrat, Hafiz Hayat Campus, Gujrat, Pakistan

<sup>2</sup>Department of Environmental Sciences and Engineering, Government College University, Faisalabad, Pakistan

## Email address:

mujahid726@yahoo.com (M. Farid)

## To cite this article:

Nusrat Bashir, Mujahid Farid, Rashid Saeed, Hafiz Muhammad Tauqeer, Shafaqat Ali, Muhammad Rizwan, Ayesha Siddiqua, Rasham Sallah-Ud-Din. Measurement of Different Heavy Metals Concentration in Roadside Dust in the Vicinity of Gujrat, Pakistan. *Science Journal of Chemistry*. Vol. 5, No. 4, 2017, pp. 51-57. doi: 10.11648/j.sjc.20170504.12

Received: December 18, 2016; Accepted: March 7, 2017; Published: August 29, 2017

**Abstract:** Due to rapid urbanization and industrialization in many countries of the world, heavy metals are continuously emitted into the terrestrial environment and posing great threat to human health. The present study was conducted to measure the concentration of six heavy metals, chromium (Cr), nickel (Ni), cobalt (Co), zinc (Zn), cadmium (Cd) and lead (Pb) in roadside dust of five major roads in Gujrat city. The dust samples were collected from five different locations in the vicinity of Gujrat. The results showed that the average concentrations of Cr, Ni, Co, Zn, Cd, and Pb in the dust samples were higher than their permissible limits. The concentrations of Cu, Zn, Cd and Pb were much higher than those in the background value of Gujrat soil. Results clearly concluded that the heavy traffic and transportation is the major responsible source of these heavy metals in the road dust of Gujrat city along with the transportation of sewage sludge, surface runoff from agricultural land and discharge of sewage water from drains.

**Keywords:** Industrialization, Heavy Metal, Traffic, Chromium, Lead, Sludge

## 1. Introduction

Pollution is the most severe problem in developing countries now a day [36, 16]. Continuous increasing population leads the pollution by using transport and industrialization. In urban areas, traffic is more congested as compared to rural areas [30]. Roads are considered as the second largest non-point source of pollution in all media of environment [44, 15]. Developed countries could combat or minimize traffic pollution impacts by using different advanced technologies and recourses because they have substantial economic parameters as compared to developing countries [18]. Anthropogenic activities, paints, petrol, aerosols are the major sources of traffic pollution [3].

In Pakistan about 38.6% is urban population [42]. Due to rapid urbanization the roads are loaded with high burden of traffic which cause more pollution in urban areas. Due to the deficiency of resources, poverty, lack of awareness and government's perfunctory; Pakistan is struggling to control

traffic pollution.

The major concern of time is the emission of heavy metals from transportation and other means. These heavy metals has ability to circulate in all media of environment by bio-magnifications and bioaccumulation. These not just pollute the soil but also accumulate in food plants [32, 19, 24] and intake by human and animals during food consumption and cause health related problems [35, 1]. Traffic pollution has emerged as a serious issue due to potential hazard on human health and nearby environment. Road dust could easily be inhaled, ingested or absorbed by human beings due to its small particle size [8, 9]. These persistent heavy metals cause potential threat to human health and safety along with environment. In small concentrations, some heavy metals are necessary for plants and human health but in excessive amount these are toxic [15, 5, 2]. People living or working near roadways, are more vulnerable to health related problems with respect to the people living in remote areas. The severe health problems which they face are asthma, cardiac diseases, lungs disorder (i.e. asthma, bronchitis,

Chronic obstructive pulmonary disease (COPD), lung cancer, (Pneumonia) premature deaths and brain disorder. Heavy metals incorporate in bio-geochemical cycle and disturbed the natural environment [40].

Recent studies reported that natural weathering of building material, industrial emissions, tire abrasion, mining processes, fuel burning at homes, metallurgical processes, brick kilns and heating systems inside the vehicles are also responsible for producing different heavy metals as a byproduct [43] but burning of gasoline produce more percentage of heavy metals.

Climatic conditions, traffic frequency and density, urbanization, temperature profile and topography of that area are the effective factors for the distribution and

contamination of road dust [22].

Keeping In view the above described scenario the present study was planned to estimate the concentration of six selected heavy metals at five different locations of Gujrat City, Pakistan.

## 2. Material and Methods

### 2.1. Study Area

The geographical location of Gujrat city is “32°40’N latitude and “74°4’E” longitude. The location of study area is shown in figure 1.




Figure 1. Map of study area.

Soil samples were collected from five selected sites of Gujrat city. Samples were collected in dry season because rainfall may keep wet the soil present on roads and green belts. The samples were collected in three replicates from each location. Soil samples at 0-10cm depth of surface soil were collected in polythene bags with the approximately weight of 500 grams. Samples were collected during peak hours of traffic i.e. in between 7:30am-9:30am in the morning and 2:00Pm – 5:30pm in the evening. During sample collection, efforts were made to avoid other contamination, organic waste, debris material, effluents, and industrial wastes and manure objects that might disguise the outcome of motor vehicle discharge. All cores taken for a given sample were collected in a clean bucket and mixed thoroughly. As crushing is easier at right moisture level, the soil was passed through 2- 3 mm sieve and air dried. Samples were preserved at room temperature (25°C) for further analysis of heavy metal concentration.

### 2.2. DTPA Preparation

1.97g of DTPA (Diethylenetriamine Penta acetic Acid), 1.1g of anhydrous CaCl<sub>2</sub> and 14.92 g of TEA (tri-ethanol amine) were dissolved in approximately 800 ml of distilled

water. DTPA requires a sufficient time on hate plate, along with magnetic stirrer to dissolve, and then prepare volume. The pH was adjusted at 7.3 with 1:1 HCl or 1:1 NH<sub>4</sub>OH while stirring [29].

### 2.3. Method for Soil Analysis

25 g soil sample was taken and 50 ml of DTPA solution added to it. Then it was shaken continuously for 2 hours on horizontal shaker and then filtered to a flask. A blank solution (containing all reagents except soil) was run with samples as blank. Read each element concentration on concentration mode by Atomic Absorption Spectrophotometer (novAA 350, based in Jena, Germany) (At least 5 standards were prepared for each element with a range as following in the table 1. [15].

Table 1. Standards for heavy metals.

Serial No.	Heavy Metals	Parameters/ Standards
1	Lead (Pb)	0.5, 1.0, 1.5, 2.0, 2.5 ppm
2	Cobalt(Co)	0.5, 1.0, 1.5, 2.0, 2.5 ppm
3	Chromium (Cr)	0.5, 1.0, 1.5, 2.0, 2.5 ppm
4	Cadmium (Cd)	0.5, 1.0, 1.5, 2.0, 2.5 ppm
5	Nickel (Ni)	0.5, 1.0, 1.5, 2.0, 2.5 ppm
6	Zinc (Zn)	0.5, 1.0, 1.5, 2.0, 2.5 ppm

## 2.4. Calculations Required

Heavy Metal (ppm) = AAS reading x dilution factor

## 2.5. Statistical Analysis


Different procedures of statistical analysis were used to analyze data. All statistical analysis were done using SPSS (Statistical Package for Social Sciences).


## 3. Results

Concentration of different heavy metals was measured at five selected locations in the vicinity of Gujrat city. All samples were in positive value for heavy metal contamination in traffic congested areas of Gujrat. At location (A) the concentration of Pb was higher as compared to other heavy metals such as  $Pb > Zn > Cd > Cr > Co > Ni$  while Similar trend was found at location (B) where Ni concentration was slightly equal to the concentration of Co while Zn concentration was not significantly different from Pb.


The location (C) has maximum value of Pb followed by Zn, Cd, Cr, Co and Ni respectively. Cobalt found in lesser quantity as compared to other heavy metals at location (D). The sequence can be written as  $Co < Ni < Cd < Zn < Cr < Pb$ . Quite similar trend found at location (E) but the concentration of Cd was lesser while Zn concentration was higher as compared to location (D) such as  $Co < Cd < Ni < Cr < Zn < Pb$ .

The comparison of single heavy metal concentration at all locations is given in Figure 3. Almost all samples were in positive value of selected heavy metals while Pb concentration was found higher at all locations and Kachari Chowk holds the maximum concentration of Pb while other locations were not significantly different from each other. Cobalt concentrations were found lesser at all locations and were not significantly different to each other. Zinc holds second highest concentration at all locations but not significantly different from each other except Kachari and GTS chowk. Cr and Cd were found almost in same range of concentrations at all locations. Ni concentration was lesser at Fawara chowk while found highest at Kachari chowk.





**Figure 2.** The concentration of lead (Pb), cobalt (Co), chromium (Cr), cadmium (Cd), nickel (Ni) and zinc (Zn) at Fawara Chowk (A), Jail Chowk (B), Katchari Chowk (C), GTS Chowk (D) and Service Morr Chowk (E). Values are means of three replicates  $\pm$  S. D. Different small letters on bars depicts that values are significantly different from each other at  $p \leq 0.005$ .



**Figure 3.** The concentration of lead (Pb), cobalt (Co), chromium (Cr), cadmium (Cd), nickel (Ni) and zinc (Zn) at Fawara Chowk, Jail Chowk, Kachari Chowk, GTS Chowk and Service Morr Chowk. Values are means of three replicates  $\pm$  S. D. Different small letters on bars depicts that values are significantly different from each other at  $p \leq 0.005$ .

### 4. Discussion

Assessment of heavy metals is becoming an important tool

to determine the quality of traffic pollution at a specific point, both in rural and urban environments. The concentration of heavy metal is more in traffic congested areas. Traffic volume is increasing by public dependence more on their

private transport. This dependence or leasing of private transport cause more transport, more fuel consumption and eventually more traffic pollution [38]. In developed countries there is proper legislation for monitoring and tuning of vehicles on regular basis. This constant increase in number of automobiles lead to more traffic load on roads and cause traffic jams due to which more pollutants released into the atmosphere. The seasonal changes affect the contamination of heavy metals in road dust such as, heavy metal concentration is more in normal to dry season, while in rainy season these spread/thresh away along with water and diluted [33, 25].

The current study involved the monitoring of different heavy metals, from road dust during the peak hours of traffic. The traffic volume changes along with school and office timings. i.e. from 7:30am to 9:30am in the morning and 2:30pm -5:30pm in the evening. Traffic stampede was found to be increasing gradually and a bit higher concentration of heavy metals was recorded. In current study the most elevated level of lead was at location L3 Kachari Chowk (Fig 2 & 3). Contamination level of Pb is at elevated levels in metropolitan areas [23] when large number of vehicles passed at the timing of rush hours [12]. The heavy metals emitted from automobiles exhaust dispersed in the atmosphere and then accumulated in the surrounding environment, plant, fruits (which kept by vendors in open air) [21, 37]. Further these heavy metals taken up by human beings and biomagnified in food chain and food web [39, 31]. Approximately 3600 automobiles passing through this point which includes, rickshaws, motor bikes, cars, buses/mini buses, hiace/bolan etc. Lead also entered into the atmosphere by wind erosion, during sludge transportation and sludge extraction. A study was conducted by yang. (2011) that Pb contamination was due to over urbanization and industrialization but in current study this might be due to congested traffic pollution [45].

In current study the level of Co contamination was also least as compared to other selected heavy metals. General source are mining and industrial wastes, vehicle emissions, lead-acid batteries, fertilizers, paints, treated woods, plastics floating in rivers and drains [20, 34]. Cobalt can accumulate to toxic levels in the liver, kidney, pancreas, and heart, as well as the skeleton and skeletal muscles [6]. Cobalt has been found to produce tumors in animals and is likely a human carcinogen as well. Occupational exposure to cobalt powder has been linked to vision, asthma and hearing problems [41].

The adjacent areas of Gujrat city are more congested in tannery industry such as Sialkot and Gujranwala. The waste water coming from these cities through drains and also transportation of the tannery through the Gujrat city is the major source of Cr in roadside dust. Aydin et al. (2006) reported that elevated level of Cr was due to industrialization but in current study it might be due to tannery industry and its transportation by roads [14]. These are carcinogenic in nature and affects public health [27, 11].

Automobile emissions are the major source of Cd contamination [26]. Inhalation exposure of Cd can occur

from road dust. Prolonged exposure to Cd can cause severe toxicity in humans. Mostly Cd affects the kidney and lungs of human beings [4].

According to Eisler (2007) nickel contamination mainly originates from industrial and traffic pollution [17]. According to Doung and Lee. (2011) atmospheric dispersion and frequently brake use were additional factors in nickel contamination [13]. Nickel moves along with food web by bio-accumulation and bio-magnification that cause several diseases such as skin allergy, lung and cardiovascular diseases [10, 25].

At L2 (jail chowk) number of hospitals and pharmacies are more as compared to other locations, so that's why zinc is released from hospital waste and pharmaceutical waste (Figure 3). Approximately 372 automobiles passing through this point per hour which includes cars, motor bikes, rickshaws, hiace and buses etc. Zn released in atmosphere and affects human health and also causes damages to environment [28].

## 5. Conclusion

The results of present study concluded that the concentration of selected heavy metals, Pb, Co, Cr, Cd, Ni and Zn were found in roadside dust of Gujrat City which clearly indicated that the heavy traffic and transportation is the major responsible source of these heavy metals along with other sources. There is need to develop and implement the proper legislation for the monitoring and maintaining of automobile vehicles and transportation of waste material.

## References

- [1] Adrees M, Ali S, Rizwan M, Ibrahim M, Abbas F, Farid M, Zia-ur-Rehman M, Irshad MK, Bharwana SA (2015a) The effect of excess copper on growth and physiology of important food crops: A review. *Environmental Science and Pollution Research* 22(11): 8148-8162. doi: 10.1007/s11356-015-4496-5.
- [2] Adrees M, Ali S, Rizwan M, Zia-ur-Rehman M, Ibrahim M, Abbas F, Farid M, Qayyum MF, Irshad MK (2015b) Mechanisms of silicon-mediated alleviation of heavy metal toxicity in plants: A review. *Ecotoxicology and Environmental Safety* 119: 186-197. doi: 10.1016/j.ecoenv.2015.05.011.
- [3] Akhter MS, Madany IM (1993) Heavy metals in street and house dust in Bahrain. *Water, Air, and Soil Pollution* 66(1), 111-119. doi: 10.1007/BF00477063.
- [4] Al-Fatlawi SM, Al-Alwani M, (2012) Heavy metals pollution of roadside dust sample with different traffic volumes at Hilla city. *The Iraqi Journal for Mechanical And Material Engineering* 12(4): 660-672.
- [5] Ali S, Bharwana SA, Rizwan M, Farid M, Kanwal S, Ali Q, Ibrahim M, Gill RA, Khan MD (2015) Fulvic acid mediates chromium (Cr) tolerance in wheat (*Triticum aestivum* L.) through lowering of Cr uptake and improved antioxidant defense system. *Environmental Science and Pollution Research* 22(14): 10601-10609. doi: 10.1007/s11356-015-4271-7.

- [6] Apostoli P, Catalani S, Zaghini A, Mariotti A, Poliani PL, Vielmi V, Semeraro F, Duse S, Porzionato A, Macchi V, Padovani A, Rizzetti MC, de Caro R. (2013) High doses of cobalt induce optic and auditory neuropathy. *Experimental and Toxicologic Pathology* 65(6): 719-727. doi: 10.1016/j.etp.2012.09.006.
- [7] Benin AL, Sargent JD, Dalton M, Roda S. (1999) High concentrations of heavy metals in neighborhoods near ore smelters in northern Mexico. *Environmental Health Perspectives* 107(4): 279-284.
- [8] Bharwana SA, Ali S, Farid M, Jamshaid M, Azeem M, Shakoor MB, Ahmad R, Hannan F. (2014) Noise and ambient air pollution monitoring and its effects on local community in Faisalabad, Pakistan. *Academia Journal of Environmental Sciences* 2(2): 15-28.
- [9] Bharwana SA, Ali S, Farid M, Hannan F, Ahmad R, Tauqeer HM, Baloch IH, Iqbal M. (2015) Work related injuries in small scale metal press industries of Shahdra Town, Lahore, Pakistan. *The Holistic Approach to Environment* 5(1): 3-11.
- [10] Chashschin VP, Artunina GP, Norseth T. (1994) Congenital defects, abortion and other health effects in nickel refinery workers. *Science of The Total Environment* 148(2): 287-291. doi: 10.1016/0048-9697(94)90405-7.
- [11] Chervona Y, Arita A, Costa M. (2012) Carcinogenic metals and the epigenome: understanding the effect of nickel, arsenic, and chromium. *Metallomics* 4(7): 619-627. doi: 10.1039/c2mt20033c.
- [12] Cholak J, Schafer LJ, Yeager D. (1968) The air transport of lead compounds present in automobile exhaust gases. *American Industrial Hygiene Association Journal* 29(6): 562-568. doi: 10.1080/00028896809343056.
- [13] Duong TTT, Lee BK. (2011) Determining contamination level of heavy metals in road dust from busy traffic areas with different characteristics. *Journal of Environmental Management* 92(3): 554-562. doi: 10.1016/j.jenvman.2010.09.010.
- [14] Duzgoren-Aydin NS, Aydin A. (2006) Chemical and mineralogical heterogeneities of weathered igneous profiles: implications for landslide investigations. *Natural Hazards and Earth System Science*, 6(2): 315-322.
- [15] Ehsan S, Ali S, Noureen S, Farid M, Shakoor MB, Aslam A, Bharwana SA, Tauqeer HM (2013) Comparative assessment of different heavy metals in urban soil and vegetables irrigated with sewage/industrial waste water. *Ecoterra Journal of Environmental Research and Protection* 35: 37-53.
- [16] Ehsan S, Ali S, Noureen S, Mehmood K, Farid M, Ishaque W, Shakoor MB, Rizwan M. (2014) Citric acid assisted phytoremediation of Cd by *Brassica napus L.* *Ecotoxicology and Environmental Safety*. 106: 164-172.
- [17] Eisler R. (2007) *Eisler's encyclopedia of environmentally hazardous priority chemicals*. Elsevier, Amsterdam, The Netherlands. 986p.
- [18] Ezeala-Harrison F. (2015) Environmental Quality in economic development in low-income countries: Application of an EQT model using cross-country sample data. *Journal of Sustainable Development* 8(2): 122-133 doi: 10.5539/jsd.v8n2p122.
- [19] Farid M, Ali S, Ishaque W, Shakoor MB, Niazi NK, Bibi I, Dawood M, Gill RA, Abbas F (2015) Exogenous application of thylenediamminetetraacetic acid enhanced phytoremediation of cadmium by *Brassica napus L.* *International Journal of Environmental Science and Technology* 12(12): 3981-3992. doi: 10.1007/s13762-015-0831-0.
- [20] Gill HS, Grammatopoulos G, Adshead S, Tsialogiannis E, Tsiroidis E. (2012) Molecular and immune toxicity of CoCr nanoparticles in MoM hip arthroplasty. *Trends in Molecular Medicine* 18(3): 145-155. doi: 10.1016/j.molmed.2011.12.002.
- [21] Habiba U, Ali S, Farid M, Shakoor MB, Rizwan M, Ibrahim M, Abbasi GH, Hayat T, Ali B. (2015) EDTA enhanced plant growth, antioxidant defense system, and phytoextraction of copper by *Brassica napus L.* *Environmental Science and Pollution Research* 22(2): 1534-1544. doi: 10.1007/s11356-014-3431-5.
- [22] Hidalgo J, Masson V, Pigeon G. (2008) Urban-breeze circulation during the CAPITOU experiment: numerical simulations. *Meteorology and Atmospheric Physics* 102(3-4): 243-262. doi: 10.1007/s00703-008-0345-0.
- [23] Kabadayi F, Cesur H. (2010) Determination of Cu, Pb, Zn, Ni, Co, Cd, and Mn in road dusts of Samsun City. *Environmental Monitoring and Assessment* 168(1-4): 241-253. doi: 10.1007/s10661-009-1108-1.
- [24] Kanwal U, Ali S, Shakoor MB, Farid M, Hussain S, Yasmeen T, Adrees M, Bharwana SA, Abbas F. (2014) EDTA ameliorates phytoextraction of lead and plant growth by reducing morphological and biochemical injuries in *Brassica napus L.* under lead stress. *Environmental Science and Pollution Research*, 21(16), 9899-9910.
- [25] Khaliq A, Ali S, Hameed A, Farooq MA, Farid M, Shakoor MB, Mahmood K, Ishaque W, Rizwan M (2015) Silicon alleviates nickel toxicity in cotton seedlings through enhancing growth, photosynthesis and suppressing Ni uptake and oxidative stress. *Archives of Agronomy and Soil Science* 62(5): 633-647. doi: 10.1080/03650340.2015.1073263.
- [26] Kim ND, Fergusson JE (1994) The concentrations, distribution and sources of cadmium, copper, lead and zinc in the atmosphere of an urban environment. *Science of The Total Environment* 144(1-3): 179-189. doi: 10.1016/0048-9697(94)90437-5
- [27] Léonard A, Lauwerys RR (1980) Carcinogenicity and mutagenicity of chromium. *Mutation Research/Reviews in Genetic Toxicology* 76(3): 227-239. doi: 10.1016/0165-1110(80)90018-4.
- [28] Li X, Poon CS, Liu PS (2001) Heavy metal contamination of urban soils and street dusts in Hong Kong. *Applied Geochemistry* 16(11-12): 1361-1368. doi: 10.1016/S0883-2927(01)00045-2.
- [29] Lindsay WL, Norvell WA (1978) Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil science society of America Journal*, 42(3), 421-428. doi: 10.2136/sssaj1978.03615995004200030009x.
- [30] Liu R, Guan CQ (2005) Mode biases of urban transportation policies in China and their implications. *Journal of Urban Planning and Development*. 131(2): 58-70. doi: 10.1061/(ASCE)0733-9488(2005)131: 2(58).
- [31] Lokeshwari H, Chandrappa GT (2006) Impact of heavy metal contamination of Bellandur Lake on soil and cultivated vegetation. *Current Science* 91(5): 622-627.

- [32] Muchuweti M, Birkett JW, Chinyanga E, Zvauya R, Scrimshaw MD, Lester JN (2006) Heavy metal content of vegetables irrigated with mixtures of wastewater and sewage sludge in Zimbabwe: Implications for human health. *Agriculture, Ecosystems & Environment* 112(1): 41-48. doi: 10.1016/j.agee.2005.04.028.
- [33] Rahman SH, Khanam D, Adyel TM, Islam MS, Ahsan MA, Akbor MA (2012) Assessment of heavy metal contamination of agricultural soil around Dhaka Export Processing Zone (DEPZ), Bangladesh: Implication of seasonal variation and indices. *Applied Sciences* 2(3): 584-601. doi: 10.3390/app2030584.
- [34] Rizo OD, Castillo FE, López JA, Merlo MH (2011) Assessment of heavy metal pollution in urban soils of Havana city, Cuba. *Bulletin of Environmental Contamination and Toxicology* 87(4): 414-419. doi: 10.1007/s00128-011-0378-9.
- [35] Rizwan M, Ali S, Ibrahim M, Farid M, Adrees M, Bharwana SA, Zia-ur-Rehman M, Qayyum MF, Abbas F (2015) Mechanisms of silicon-mediated alleviation of drought and salt stress in plants: a review. *Environmental Science and Pollution Research* 22(20): 15416-15431. doi: 10.1007/s11356-015-5305-x.
- [36] Shakoor MB, Ali S, Farid M, Farooq MA, Tauqeer HM, Iftikhar U, Hannan F, Bharwana SA (2013) Heavy metal pollution, a global problem and its remediation by chemically enhanced phytoremediation: A review. *Journal of Biodiversity and Environmental Sciences* 3(3): 12-20.
- [37] Shakoor MB, Ali S, Hameed A, Farid M, Hussain S, Yasmeen T, Najeeb U, Bharwana SA, Abbasi GH (2014) Citric acid improves lead (Pb) phytoextraction in *Brassica napus* L. by mitigating Pb-induced morphological and biochemical damages. *Ecotoxicology and Environmental Safety*. 109: 38-47.
- [38] Shapiro RJ, Hassett KA, Arnold FS (2002) Conserving energy and preserving the environment: The role of public transportation. Report by American Public Transportation Association. 39p. Available at: <https://www.apta.com/resources/reportsandpublications/Documents/shapiro.pdf> [access date: 09.08.2016].
- [39] Sharma RK, Agrawal M, Marshall FM (2008) Heavy metal (Cu, Zn, Cd and Pb) contamination of vegetables in urban India: A case study in Varanasi. *Environmental Pollution* 154(2): 254-263. doi: 10.1016/j.envpol.2007.10.010.
- [40] Victoria A, Cobbina SJ, Dampare SB, Duwiejuah AB (2014) Heavy Metals Concentration in Road Dust in the Bolgatanga Municipality, Ghana. *Journal of Environment Pollution and Human Health* 2(4): 74-80. doi: 10.12691/jephh-2-4-1.
- [41] Walters GI, Moore VC, Robertson AS, Burge CBSG, Vellore AD, Burge PS (2012) An outbreak of occupational asthma due to chromium and cobalt. *Occupational Medicine* 62(7): 533-540. doi: 10.1093/occmed/kqs111.
- [42] Worldmeters (2016) Population of Pakistan, Retrieved 27 August 2016, from <http://www.worldometers.info/world-population/pakistan-population/>.
- [43] Wei B, Jiang F, Li X, Mu S (2010) Contamination levels assessment of potential toxic metals in road dust deposited in different types of urban environment. *Environmental Earth Sciences* 61(6): 1187-1196. doi: 10.1007/s12665-009-0441-6.
- [44] Yan X, Gao D, Zhang F, Zeng C, Xiang W, Zhang M (2013) Relationships between heavy metal concentrations in roadside topsoil and distance to road edge based on field sources observations in the Qinghai-Tibet Plateau, China. *International Journal of Environmental Research and Public Health* 10(3): 762-775. doi: 10.3390/ijerph10030762.
- [45] Yang Z, Lu W, Long Y, Bao X, Yang Q (2011) Assessment of heavy metals contamination in urban topsoil from Changchun City, China. *Journal of Geochemical Exploration* 108(1): 27-38. doi: 10.1016/j.gexplo.2010.09.006.