Synthesis of Schiff Bases Compounds from Oxamic Hydrazide: Spectroscopic Characterization, X-ray Diffraction Structure and Antioxidant Activity Study

Fatou Faye¹, Amadou Guèye¹, Papa Samba Camara², Aïssatou Alioune Gaye¹, Farba Bouyagui Tamboura², Nathalie Gruber³, Mohamed Gaye^{1,*}

¹Department of Chemistry, University Cheikh Anta Diop, Dakar, Senegal

²Department of Chemistry, University Alioune Diop, Bambey, Senegal

³Institut Le Bel, University of Strasbourg, Strasbourg, France

Email address:

mohamedl.gaye@ucad.edu.sn (M. Gaye)

*Corresponding author

To cite this article:

Fatou Faye, Amadou Guèye, Papa Samba Camara, Aïssatou Alioune Gaye, Farba Bouyagui Tamboura, Nathalie Gruber, Mohamed Gaye.

Synthesis of Schiff Bases Compounds from Oxamic Hydrazide: Spectroscopic Characterization, X-ray Diffraction Structure and Antioxidant Activity Study. *American Journal of Applied Chemistry*. Vol. 9, No. 1, 2021, pp. 6-12. doi: 10.11648/j.ajac.20210901.12

Received: January 2, 2021; Accepted: January 11, 2021; Published: January 22, 2021

Abstract: The compounds (*E*)-2-amino-*N'*-(1-(2-hydroxyphenyl)ethylidene)-2-oxoacetohydrazide (I) and (*E*)-*N'*-(2-hydroxy-3-methoxybenzylidene)-2-amino-2-oxoacetohydrazide (II) were synthesized by the 1:1 ratio condensation reaction of oxamic hydrazide and 2-hydroxyacetophenone or o-vanillin respectively. The two compounds were characterized by physico-chemical analyses, elemental analysis, FTIR, ¹H and ¹³C NMR spectroscopies techniques. The structure of the compound (I) was determined by single-crystal X-ray diffraction study. The compound (I) (C₁₀H₁₁N₃O₃) crystallises in the triclinic space group *P*-1 with the following unit cell parameters: *a* = 7.0399 (5) Å, *b* = 8.6252 (8) Å, *c* = 9.5474 (9) Å, *a* = 81.730 (3)°, *b* = 72.738 (3)°, *g* = 67.450 (3)°, *V* = 510.99 (8) Å³, *Z* = 2, *T* = 173 (2) K, *m* = 0.11 mm⁻¹, *D*_{calc} = 1.438 g/cm³, *R*_{int} = 0.028, *R*_{sigma} = 0.073. The oxamic hydrazide moiety of the molecule is slightly twisted as reflected by the torsion angles values of 177.2 (2)° [N1-N2-C9-C10], -171.3 (3)° [N2-C9-C10-N3], -4.6 (4)° [O2-C9-N2-N1] and 8.4 (4)° [O3-C10-C9-N2]. The intramolecular hydrogen bond O1(phenol)-H1...N1(hydrazide) which close in *S* (6) ring stabilized the conformation. The intermolecular hydrogen bonds, C3-H3...O1ⁱ(phenol) (i: -x+1, -y, -z+1), N3(amide)-H3A...O3ⁱⁱ(amide) (ii: -x+1, -y+2, -z) and N3(amide)-H3B...O2ⁱⁱⁱ(hydrazide) (iii: -x+1, -y+1, -z) lead to the formation of sheets parallel to *ac* plane. Compounds (I) and (II) showed antioxidant activities less than 10% inhibition of DPPH.

Keywords: Oxamic Hydrazide, 2-hydroxyacetophenone, O-vanillin, Antioxidant, X-ray

1. Introduction

Potentially ditopic Schiff bases prepared from oxamic or thioxamic hydrazide have been widely reported in the literature by chemists and in particular those interested in coordination chemistry [1-6]. Oxamic hydrazide has two different arms. The hydrazide arm condenses more easily than the amide arm with carbonyl [7-10]. The use of these ligands has made it possible to prepare coordination compounds with atypical structures [11-13]. The derivatives of oxamic hydrazide have been used for the synthesis of

nucleoside [14, 15]. Oxamic hydrazide have been used as co-reactant in electroluminescence [16, 17]. A wide variety of heterocyclic molecules with good medicinal properties are obtained starting from these oxamic precursors [18]. They are used as antibacterial agent, [19] anticonvulsant, [8] anti-inflammatory [20], antituberculosis [18] and anticancer agents [4, 21]. Schiff's bases are synthesized from oxamic hydrazide and used for the preparation of complexes with transition metal and lanthanide ions [9, 22-25]. The properties of these complexes are evaluated in superoxide catalysis [2, 3] and magnetism [26]. It is in this perspective that we have studied these types of ligands and reported their

transition metal complexes [7, 27]. Continuing our work in this field, we obtained ligands (I) and (II). In the present study, we report the spectroscopic study of the two compounds and the structure of (I) obtained by X-ray diffraction.

2. Material and Methods

2.1. Materials and Physical Methods

Oxamic hydrazide, 2'-hydroxyacetophenone, o-vanillin, cyclohexanol and 1,1-diphenyl-2-picrylhydrazyl (DPPH.) were of analytical reagent grade and were obtained from Sigma-Aldrich Company. All used solvents were of UV spectroscopic quality. The elemental analyses of C, H and N were recorded on a VxRio EL Instrument. FT-IR spectra were recorded in the region of 4000–400 cm⁻¹ using a Perkin Elmer Spectrum Two FT-IR spectrometer. The UV-Visible spectra were recorded on a Perkin Elmer Lambda UV-Vis spectrophotometer. The ¹H and ¹³C NMR spectra were recorded in DMSO-d₆ on a Bruker 500 MHz spectrometer at room temperature using TMS as an internal reference.

2.2. Free Radical Scavenging Antioxidant Assay

Antioxidant capacities of compound (I) were measured according to Akhtar et al. [28] method with modifications. The methanol solution of 3.8 mL DPPH• was added to test compounds (200 µL) at different concentrations. The mixture was shaken vigorously and incubated in dark for 30 min at room temperature. After the incubation time, the absorbance of the solution was measured at 517 nm by using UV-vis spectrophotometer Perkin two. The DPPH• radical scavenger effect was calculated using the following equation:

$$\text{Scavenging activity (\% control)} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100$$

where A_{control} is the absorbance of the control reaction and A_{sample} is the absorbance of the test compound. The tests were carried out in triplicate. Trolox was used as positive control.

2.3. Synthesis of (E)-2-amino-N'-(1-(2-hydroxyphenyl)ethylidene)-2-oxoacetohydrazide (I)

Oxamic hydrazide (2.0307 g, 19.7 mmol) was suspended in 20 mL of cyclohexanol. 2'-hydroxyacetophenone (2.6821 g, 19.7 mmol) previously dissolved in 20 mL of cyclohexanol was added. The mixture was heated at reflux for 24 hours. On cooling, a white solid was collected by filtration, washed with 2 x 10 mL of ethanol, and dried in the open air. The precipitate was recrystallized in DMF and the solution was left at room temperature. Slow evaporation of the solvent gave colorless crystals after one week. Yield: 95%. Anal. Calc. for [C₁₀H₁₁N₃O₃] (%): C, 54.29; H, 5.01; N, 19.00. Found: C, 54.27; H, 5.03; N, 19.03. FT-IR (n, cm⁻¹): 3383(OH), 3293 (NH), 3222 (NH₂), 1702 (CONH), 1652 (CONH₂), 1606 (C=N), 1562 (C_{Ar}=C_{Ar}), 1466 (C_{Ar}=C_{Ar}), 1411 (C_{Ar}=C_{Ar}), 1250 (C-O_{phenol}). ¹H RMN δ: 2.424 (S, 3H, CH₃), 6.921–8.056 (m, 4H, H_{Ar}), 8.396 (Ar-OH), 9.940 (S,

1H, CONH), 11.428 (S, 1H, HNCOH), 12.925 (S, 1H, HNCOH). RMN ¹³C (dmso-d₆, d (ppm)): 162.232 (C=O), 159.198 (HN-COH), 158.840 (C=N), 157.840 (C_{Ar}-OH), [132.344, 129.416, 119.607, 119.231, 117.812] (C_{Ar}), 14.677 (-CH₃),

2.4. Synthesis of (E)-N'-(2-hydroxy-3-methoxybenzylidene)-2-amino-2-oxoacetohydrazide (II)

Oxamic hydrazide (2.037 g, 19.7 mmol) was suspended in 20 mL of cyclohexanol. o-vanillin (2.9973 g, 19.7 mmol) previously dissolved in 20 mL of cyclohexanol was added. The mixture was heated at reflux for six hours. On cooling, a solid was collected by filtration, washed with 2 x 10 mL of ethanol and dried in the open air. The precipitate was recrystallized in DMF and the solution was left at room temperature. Slow evaporation of the solvent gave colorless precipitate after one week. Yield: 97%. Anal. Calc. for [C₁₀H₁₁N₃O₄] (%): C, 50.63; H, 4.67; N, 17.71. Found: C, 50.60; H, 4.69; N, 17.75. FT-IR (n, cm⁻¹): 3375 (N-H), 3221 (O-H), 1704 (CONH), 1656 (C=O), 1603 (C=N), 1560 (C_{Ar}=C_{Ar}), 1464 (C_{Ar}=C_{Ar}), 1409 (C_{Ar}=C_{Ar}), 1252 (C-O_{phenol}), 1230 (C-O_{ether}), 1080 (N-N), 837 (C-H_{Ar}), 669 (C-H_{Ar}). RMN ¹H (dmso-d₆, d (ppm)): 3.80 (S, 3H, -OCH₃), [6.82–7.10] (m, 3H, H_{Ar}), 7.97 (S, 1H, CONH), 8.30 (S, 1H, Ar-OH), 8.75 (S, 1H, HC=N), 10.75 (S, 1H, HNCOH), 12.36 (S, 1H, HNCOH). RMN ¹³C (dmso-d₆, d (ppm)): 161.51 (C=O), 156.54 (HN-COH), 150.46 (C=N), 147.87 (C_{Ar}-OH), 147.20 (C_{Ar}-OCH₃), [120.65, 119.03, 118.72, 113.96] (C_{Ar}), 55.76 (-OCH₃).

I: R₁ = -CH₃, R₂ = H and t = 24 hours; II: R₁ = H, R₂ = -OCH₃ and t = 6 hours

Figure 1. Synthetic scheme of (I) and (II).

2.5. Crystal Structure Determination

Crystals suitable for single-crystal X-ray diffraction, of the reported compound, was grown by slow evaporation of DMF solution of the compound. Details of the crystal structure solution and refinement are given in Table 1. Diffraction data were collected using a Bruker APEX-II CCD diffractometer with graphite monochromatized MoK α radiation ($\lambda = 0.71073$ Å). All data were corrected for Lorentz and polarization effects. The structure was solved and refined using the Bruker SHELXTL Software Package [29]. All the structures were refined on F^2 by a full-matrix least-squares procedure using anisotropic displacement parameters for all non-hydrogen atoms [30]. H atoms of the NH groups was located in the Fourier difference maps and refined without restraints. Other H atoms were geometrically optimized and refined as riding on their carriers with $U_{iso}(H) = 1.2U_{eq}(C)(1.5 \text{ for } CH_3 \text{ group})$. Molecular graphics were generated using ORTEP-3 [31].

3. Result and Discussion

3.1. General Study

The synthesis of Schiff bases usually takes place in simple alcohols such as methanol, ethanol or propanol. In the synthesis of Schiff bases from oxamic hydrazide, the subject of our study, the use of these solvents leads to excessively long reaction times. In fact, for the condensation of oxamic hydrazide with carbonyl compounds, it is necessary to heat to temperature high enough to shorten the reaction time. Cyclohexanol which has a high boiling point (161.8° C) is suitable to prepare compounds (I) and (II) with short time reaction (Figure 1).

The results of elemental analysis agree with the expected formulas for the two compounds. The solid-state infrared spectrum of (I) reveals a broad band around 3383 cm⁻¹ attributed to the OH stretching vibration and another band around 3293 cm⁻¹ indicating the presence of NH. These two bands are present in the spectrum of compound (II) at 3375 cm⁻¹ and 3221 cm⁻¹ respectively [32, 33]. The band due to the C=N group formed after the condensation reaction between the oxamic hydrazide and the appropriate carbonyl is pointed at 1606 cm⁻¹ for (I) and at 1603 cm⁻¹ for (II). The stretching vibrations due to C=O of the oxamic unit were noted at 1702 cm⁻¹ and 1652 cm⁻¹ for (I) and at 1704 cm⁻¹ and 1656 cm⁻¹ for (II) [34, 35]. The shift of the second band towards the low frequencies is justified by the strong resonance of the oxalate group. The additional bands in the range [1570–1405 cm⁻¹] are due to the aromatic groups.

The ¹H NMR spectra of the compounds, in DMSO-d₆ solution, are recorded. Compound (I) gives two signals characteristic of iminolisation. Indeed, the single signal designating the moiety [–C(=O)–NH₂] does not appear on the spectrum. The two signals at 11.428 ppm and 12.925 ppm assigned respectively to HN=C–OH and HN=C–OH are indicative of the iminolisation of the amide function of the Schiff base. The same phenomenon is observed for

compound (II). The corresponding signals are pointed at 10.750 ppm and 12.360 ppm, respectively. These observations are confirmed by ¹³C NMR spectra in DMSO-d₆. Compound (I) gives a signal at 159.198 ppm corresponding to the iminol carbon atom HN=C–OH. This signal is identified at 156.540 ppm in the spectrum of (II). The signals due to the hydrazide carbon atoms are at 162.232 ppm and 161.510 ppm for (I) and (II) respectively. This behavior is observed in amide–iminol tautomerism [36].

Table 1. Crystal data and structure refinement for compound (I).

Chemical formula	C ₁₀ H ₁₁ N ₃ O ₃
<i>Mr</i>	221.22
Crystal shape/Color	Prismatic/colorless
Crystal system, space group	Triclinic, <i>P</i> –1
<i>T</i> (K)	173 (2)
<i>a</i> (Å)	7.0399 (5)
<i>b</i> (Å)	8.6252 (8)
<i>c</i> (Å)	9.5474 (9)
<i>a</i> (°)	81.730 (3)
<i>b</i> (°)	72.738 (3)
<i>g</i> (°)	67.450 (3)
<i>V</i> (Å ³)	510.99 (8)
<i>Z</i>	2
<i>D</i> _{calc} (g cm ⁻³)	1.438
<i>F</i> (000)	232
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	0.11
Crystal size (mm)	0.12 x 0.10 x 0.10
<i>T</i> _{min} , <i>T</i> _{max}	0.982, 0.991
<i>h</i>	–7 → 8
<i>k</i>	–11 → 11
<i>l</i>	–12 → 12
Diffractometer	Bruker APEX-II CCD
Absorption correction	Multi-scan <i>SADABS</i>
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	9791, 2063, 1634
<i>R</i> _{int}	0.028
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)]	0.073
<i>wR</i> (<i>F</i> ²)	0.227
<i>S</i>	1.09
No. of parameters/restraints	147/0
$\Delta\rho_{max}$, $\Delta\rho_{min}$ (e Å ⁻³)	0.44, –0.29

3.2. Crystal Structure

The DMF solution of compound C₁₀H₁₁N₃O₃ which was left for slow evaporation for two weeks gave colorless crystals suitable for X-ray analysis. The compound crystallizes in the triclinic group *P*–1. The molecular structure with the atomic–labelling scheme is shown in figure 2. The crystal structure solution and refinement are given in Table 1. Selected bond distances and angles are listed in Table 2. The asymmetric unit contains one organic molecule. The –NH₂ of the amid group is in *anti*–position with respect to the hydrazide group across C9–C10, while the carbonyl groups are in *trans*–position with respect to each other, across C9–C10. The compound assumes an *E* configuration across the C7=N1 bond. The ((2–hydroxyphenyl)–2–ethylidene) imino moiety adopts a *cis* conformation with the oxygen atom O2 of the carbonyl through the C9–N2 bond. The bond lengths values of 1.221 (3) Å [C9–O2] and 1.234 (3) Å [C1–O3] are indicative of a double character as

observed for similar compounds which crystallize in their keto forms [37, 7]. Those distances are comparable to the values found for a derivative which has an oxalate group [38]. The bond lengths values of 1.321 (4) Å [C10–N3] and 1.340 (4) Å [C9–N2] bonds are in the normal range observed for single C–N bonds [39]. The oxamic hydrazide fragment N1/N2/C9/O2/C10/O3/N3 is planar with a maximum deviation from the least-squares plane of –0.132 (2) Å for the N3 atom. The oxamic hydrazide moiety of the molecule is slightly twisted as reflected by the torsion angles values of 177.2 (2)° [N1–N2–C9–C10], –171.3 (3)° [N2–C9–C10–N3], –4.6 (4)° [O2–C9–N2–N1] and 8.4 (4)° [O3–C10–C9–N2].

The crystal packing of compound (I) is stabilized by intramolecular O(phenol)–H···N(hydrazide) and intermolecular N(amide)–H···O(amide), N(amide)–H···O(hydrazide) and C–H···O(phenol) hydrogen bonds. The intramolecular hydrogen bond O1(phenol)–H1···N1(hydrazide) which close in *S* (6) ring stabilized the conformation. Intermolecular hydrogen bonds, C3–H3A···O1ⁱ(phenol) (i: –x+1, –y, –z+1) N3(amide)–

H3A···O3ⁱⁱ(amide) (ii: –x+1, –y+2, –z) and N3(amide)–H3B···O2ⁱⁱⁱ(hydrazide) (iii: –x+1, –y+1, –z) lead to the formation of sheets parallel to *ac* plane (Figure 3, Table 3).

Table 2. Selected geometric parameters (Å, °).

C2–O1	1.362 (4)	C9–O2	1.221 (3)
C7–N1	1.286 (4)	C10–O3	1.234 (3)
N1–N2	1.371 (3)	C10–N3	1.321 (4)
C9–N2	1.340 (4)		
N1–C7–C1	114.7 (2)	O2–C9–C10	122.9 (2)
N1–C7–C8	123.5 (3)	N2–C9–C10	111.5 (2)
C1–C7–C8	121.8 (3)	O3–C10–N3	126.3 (3)
O2–C9–N2	125.6 (2)	O3–C10–C9	119.6 (2)

Table 3. Hydrogen-bond geometry (Å, °).

D–H···A	D–H	H···A	D···A	D–H···A
C3–H3···O1 ⁱ	0.95	2.51	3.395 (4)	154.9
O1–H1···N1	0.84	1.79	2.521 (3)	144.3
N3–H3A···O3 ⁱⁱ	0.88	2.11	2.946 (3)	158.6
N3–H3B···O2 ⁱⁱⁱ	0.88	2.23	3.022 (3)	150.1

Symmetry codes: (i) –x+1, –y, –z+1; (ii) –x+1, –y+2, –z; (iii) –x+1, –y+1, –z.

Table 4. Antioxidant activity of (I) and (II) at different concentration of DPPH.

Concentration (μM)	[DPPH] = 0.1014 mM			[DPPH] = 0.0507 mM		
	Trolox	(I)	(II)	Trolox	(I)	(II)
50	3.92±0.04	1.84±0.08	1.58±0.4	3.35±0.05	4.84±0.11	4.58±0.07
100	8.42±0.06	1.32±0.11	2.63±0.02	11.60±0.012	4.32±0.23	5.63±0.09
200	18.34±0.13	3.42±0.13	3.95±0.05	23.45±0.11	6.42±0.15	6.95±0.06
300	27.22±0.22	5.53±0.08	6.05±0.05	51.03±0.13	8.53±0.18	9.05±0.11
400	42.10±0.18	6.32±0.09	5.26±0.09	81.70±0.09	9.32±0.12	8.26±0.15
500	51.67±0.13	6.32±0.05	4.74±0.06	93.30±0.08	9.32±0.09	7.74±0.07

Figure 2. The crystal structure of the compound (I). Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small sphere.

Figure 3. Chain formed by hydrogen bonds in the title compound (I) along *ac* plane.

3.3. Antioxidant Activity

DPPH[•] is a stable free radical which becomes a stable molecule when it accepts an electron or hydrogen radical. DPPH[•] radical scavenging is a method widely used to evaluate the antioxidant activity of compounds [40, 41]. The capacity of scavenging DPPH[•] radical of the two compounds (I) and (II) have been screened (Table 4). The Figure 4 shows the plots of DPPH[•] free radical scavenging activity (%) for the compounds (I) and (II). For compound (I), the scavenging activity increases with increasing the concentration in the range tested (50–500 mmol/L) for the two DDPH[•] initial concentrations. The scavenging activity of (I) varies, for the highest DPPH[•] (0.1014 mM) concentration, in the range 1.84±0.08 – 6.32±0.05% and between 4.84±0.11 and 9.32±0.09% for the lowest DPPH[•] concentration (0.0507 mM). This activity is due to the NH or OH groups which can react with DPPH[•] radical by the typical H–abstraction reaction to form a stable radical. Radical scavenging activity of compound (II) [(1.58±0.15 – 4.74±0.11% for the highest DPPH[•] concentration), (1.58±0.15 – 4.74±0.11% for the lowest DPPH[•] concentration)] is slightly lower than that observed for compound (I) in the concentration range screened (Figure 4). Comparatively to the scavenging activity of Trolox (3.92±0.04 – 51.67±0.13% for the highest DPPH[•] concentration), (3.35±0.05 – 93.30±0.08% for the lowest

DPPH concentration)), the values observed for compounds (I) and (II) are lower than those of Trolox in concentration

range investigated (50–500 mM). The scavenging activity of compounds (I) and (II) are low and do not exceed 10%.

Figure 4. Antioxidant activity of (I) and (II): (a) [DPPH] = 0.1014 mM and (b) [DPPH] = 0.0507 mM.

4. Conclusion

The compounds (*E*)-2-amino-*N'*-(1-(2-hydroxyphenyl)ethylidene)-2-oxoacetohydrazide (I) and (*E*)-*N'*-(2-hydroxy-3-methoxybenzylidene)-2-amino-2-oxoacetohydrazide (II) were successfully synthesized condensation reaction of oxamic hydrazide and 2-hydroxyacetophenone or o-vanillin respectively. The structures of the compounds were confirmed by elemental analysis and spectroscopic techniques (FT-IR, ^1H and ^{13}C NMR). The molecular structure of the (I) was also determined using X-ray crystallography technique. Compounds (I) and (II) showed low antioxidant activity of about 10% in the screened concentration range [50–500 ppm].

Supplementary Materials

CCDC-2049222 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via <https://www.ccdc.cam.ac.uk/structures/>, or by e-mailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 (0) 1223-336033.

References

- [1] Čobeljić, B., Pevec, A., Turel, I., Spasojević, V., Milčić, M., Mitić, D., Sladić, D. and Anđelković, K. (2014). Analysis of the structures of the Cu (I) and Cu (II) complexes with 3-acetylpyridine and thiocyanate. *Polyhedron*, 69: 77–83.
- [2] Gutman, C. T. and Brunold, T. C. (2012). Spectroscopic and computational studies of a small-molecule functional mimic of iron superoxide dismutase, iron 2,6-diacetylpyridinebis (semioxamazine). *Inorganic Chemistry*, 51 (23): 12729–12737.
- [3] Gutman, C. T., Guzei, I. A. and Brunold, T. C. (2013). Structural, spectroscopic, and computational characterization of the azide adduct of Fe^{III} (2,6-diacetylpyridinebis (semioxamazine)), a functional analogue of iron superoxide dismutase. *Inorganic Chemistry*, 52 (15): 8909–8918.
- [4] Jarestan, M., Khalatbari, K., pouraei, A., Sadat Shandiz, S. A., Beigi, S., Hedayati, M., Majlesi, A., Akbari, F. and Salehzadeh, A. (2020). Preparation, characterization, and anticancer efficacy of novel cobalt oxide nanoparticles conjugated with thiosemicarbazide. *3 Biotech*, 10 (5): 230.
- [5] Haba, P., Sow, M. M., Sarr, M., Thiam, I. E., Diaw, M. and Gaye, M. L. (2020). Syntheses, characterization and X-ray crystal structure of polymeric heteronuclear oxo-bridged Fe/Na assembled with salen-type Schiff base and dicyanamide. *Sciences Journal of Chemistry*, 8 (2): 20–27.
- [6] Li, S., Khan, M. H., Wang, X., Cai, M., Zhang, J., Jiang, M., Zhang, Z., Wen, X., Liang, H. and Yang, F. (2020). Synthesis of a series of novel In (III) 2,6-diacetylpyridine bis (thiosemicarbazide) complexes: structure, anticancer function and mechanism. *Dalton Trans*, 49 (47): 17207–17220.
- [7] Kane, C. H., Thiam, E. I., Tamboura, F. B., Gaye, M. and Retailleau, P. (2012). Bis{2-Amino-2-oxo-*N'*-(1*E*)-1-(pyridin-2-yl-*kN*)ethylidene]acetohydrazido- k^2N' , O' }nickel (II). *Acta Crystallographica E*, 68 (5), m553.
- [8] Dimmock, J. R., Vashishtha, S. C. and Stables, J. P. (2000). Anticonvulsant properties of various acetylhydrazones, oxamoylhydrazones and semicarbazones derived from aromatic and unsaturated carbonyl compounds. *European Journal of Medicinal Chemistry*, 35 (2): 241–248.
- [9] Sumar, M., Ivanovic-Burmazovic, I., Hodzic, I. and Andjelkovic, K. (2002). Pentagonal-bipyramidal Mn (II) and Zn (II) complexes with 2',2'''-(2,6-pyridindiyldiethylidene)dioxamohydrazide. *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry*, 32 (4): 721–737.
- [10] Quacqyaegens, F., Desseyn, H. O., Bracke, B. and Lenstra, A. T. H. (1990). Vibrational analysis of different isomers of oxalyldihydrazide and the crystal structure of the “trans-trans” form. *Journal of Molecular Structure*, 238: 139–157.
- [11] Quacqyaegens, F., Hofmans, H. and Desseyn, H. O. (1987). The vibrational spectra of the Ni (II) and Cu (II) complexes with oxamic hydrazide. *Spectrochimica Acta Part A*, 43 (4): 531–537.
- [12] Quacqyaegens, F., Perlepes, S. P. and Desseyn, H. O. (1987). The vibrational spectra of Pd (II) complexes with planar monosubstituted oxamides. *Spectrochimica Acta Part A*, 43 (7): 917–923.

- [13] Perlepes, S. P., Quaeys, F. J. and Desseyn, H. O. (1990). Synthesis, spectroscopy and thermal properties of the nickel (II), palladium (II) and copper (II) complexes of oxalylhydrazide. *Transition Metal Chemistry*, 15 (2): 132–140.
- [14] Huynh-Dinh, T., Igolen, J., Bisagni, E., Marquet, J. P. and Cuvier, A. (1977). Synthesis of C-nucleosides. Part 14. 5 (3)-glycosyl-1,2,4-triazole-3 (5) carboxamides as analogues of ribavirin. *Journal of Chemical Society Perkin Trans 1*, 7, 761–764.
- [15] Chudinov, M. V., Konstantinova, I. D., Ryzhova, O. I., Esipov, R. S., Yurkevich, A. M., Shvets, V. I. and Miroshnikov, A. I. (2005). A new effective method for the synthesis of 1,2,4-triazole-3-carboxamide and ribavirin derivatives. *Pharmaceutical Chemistry Journal*, 39 (4): 212–215.
- [16] Lan, Y., Yuan, F., Tadesse Haile Fereja, T. H., Lou, B., Han, S., Li, J. and Xu, G. (2019). Electrochemiluminescence of 3,4,9,10-perylene-tetracarboxylic acid/oxamic hydrazide and its application in the detection of tannic acid. *Analyst*, 144 (15): 4449–4720.
- [17] Chen, M., Ning, Z., Chen, K., Zhang, Y. and Shen, Y. (2020). Recent advances of electrochemiluminescent system in bioassay. *Journal of Analysis and Testing*, 4 (2): 57–75.
- [18] Narang, R., Narasimhan, B. and Sharma, S. (2012). A review on biological activities and chemical synthesis of hydrazide derivatives. *Current Medicinal Chemistry*, 19 (4): 569–612.
- [19] Hakobyan, S., Boily, J.-F. and Ramstedt, M. (2014). Proton and gallium (III) binding properties of a biologically active salicylidene acylhydrazide. *Journal of Inorganic Biochemistry*, 138: 9–15.
- [20] Altowyan, M. S., Ali, M., Soliman, S. M., Al-Majid, A. M., Islam, M. S., Yousuf, S., Choudhary, M. I., Ghabbour, H. A. and Barakat, A. (2020). Synthesis, computational studies and biological activity of oxamohydrazide derivatives bearing isatin and ferrocene scaffolds. *Journal of Molecular Structure*, 1202: 127372.
- [21] Mohan, M., Agarwal, A. and Jha, N. K. (1988). Synthesis, characterization, and antitumor properties of some metal complexes of 2,6-diacetylpyridine bis (N-4-azacyclic thiosemicarbazones). *Journal of Inorganic Biochemistry*, 34 (1): 41–54.
- [22] Ivanovic-Burmazovic, I., Hamza, M. S. A. and van Eldik, R. (2002). A detailed mechanistic study of the substitution behavior of an unusual seven-coordinate iron (III) complex in aqueous solution. *Inorganic Chemistry*, 41 (20): 5150–5161.
- [23] El-Toukhy, A. (1991). Stoichiometry, Products and kinetics of oxidation of neutral 2,6-diacetylpyridine-bis (N-4-azacyclic thiosemicarbazone) Cobalt (II) complexes by dioxygen in aprotic solvents. *Inorganica Chimica Acta*, 180 (1): 85–91.
- [24] de Sousa, G. F., Deflon, V. M., Gambardella, M. T. do P., Francisco, R. H. P., Ardisson, J. D. and Niquet, E. (2006). X-Ray crystallographic and Mössbauer spectroscopic applications in dependence of partial quadrupole splitting, [R], on the C-Sn-C angle in seven-coordinated diorganotin (IV) complexes. *Inorganic Chemistry*, 45 (11): 4518–4525.
- [25] Aljahdali, M., El-Sherif, A. A., Shoukry, M. M., Hosny, W. M. and Abd-Elmoghny, M. G. (2013). Complex formation equilibria of unusual seven coordinate Fe (III) complexes with DNA constituents. *Journal of Solution Chemistry*, 42 (8): 1663–1679.
- [26] Liu, G.-F., Filipović, M., Heinemann, F. W. and Ivanović-Burmazović, I. (2007). Seven-coordinate iron and manganese complexes with acyclic and rigid pentadentate chelates and their superoxide dismutase activity. *Inorganic Chemistry*, 46 (21): 8825–8835.
- [27] Gaye, P. A., Sy, A., Sarr, A. D., Gaye M. and Besnard, C. (2011). 1- (Thiophen-2-yl)ethanone thiosemicarbazone. *Acta Crystallographica E*, 67 (5): o1168.
- [28] Akhtar, P., Yaakob, Z., Ahmed, Y., Shahinuzzaman, M. and Hyder, M. K. M. (2018). Total phenolic contents and free radical scavenging activity of different parts of *Jatropha* species. *Asian Journal of Chemistry*, 30 (2): 365–370.
- [29] Sheldrick, G. M. (2015). It *SHELXT*—Integrated space-group and crystal-structure determination. *Acta Crystallographica A*, 71 (1): 3–8.
- [30] Sheldrick, G. M. (2015). Crystal structure refinement with It *SHELXL*. *Acta Crystallographica C*, 71 (1): 3–8.
- [31] Farrugia, L. J. (2012). It WinGX and It *ORTEP* for Windows: An Update. *Journal of Applied Crystallography*, 45 (4): 849–854.
- [32] Singh, A. K., Pandey, O. P. and Sengupta, S. K. (2013). Synthesis, spectral and antimicrobial activity of Zn (II) complexes with Schiff bases derived from 2-hydrazino-5-[substituted phenyl]-1,3,4-thiadiazole and benzaldehyde / 2-hydroxyacetophenone / indoline-2,3-dione. *Spectrochimica Acta Part A*, 113: 393–399.
- [33] Saghatforoush, L. A., Hosseinpour, S., Bezpalko, M. W. and Kassel, W. S. (2019). X-Ray crystal structural and spectral studies of copper (II) and nickel (II) complexes of functionalized bis (thiosemicarbazone) ligands and investigation of their electrochemical behavior. *Inorganica Chimica Acta*, 484: 527–534.
- [34] Tripathi, G. N. R. and Katon, J. E. (1979). Vibrational spectra and structure of crystalline oxalyl hydrazide and semioxamizide. *Journal of Molecular Structure*, 54: 19–29.
- [35] Mashevskaya, M. S., Kolla, V. É., Nazmetdinov, F. Ya., Plaksina, A. N. and Semenova, Z. N. (1991). Synthesis of oxalic acid arylidenedihydrazides and their biological activity. *Pharmaceutical Chemistry Journal*, 25 (2), 83–85.
- [36] Chandra, S. and Sharma, A. K. (2009). Nickel (II) and Copper (II) complexes with Schiff base ligand 2,6-diacetylpyridine bis (carbohydrazone): synthesis and IR, Mass, ¹H NMR, electronic and EPR spectral studies. *Spectrochimica Acta Part A*, 72 (4): 851–857.
- [37] Karimian, R., Hosseini-Monfared, H., Bikas, R., Arslan, N. B., Kazak, C. and Koroglu, A. (2012). (*E,E*)-*N*-4-[(2-benzoylhydrazin-1-ylidene)methyl]benzylidenebenzo hydrazide. *Acta Crystallographica E*, 68 (5): o1433.
- [38] Tai, X.-S., Yin, J. and Kong, F.-Y. (2007). Crystal structure of 2-carboxybenzaldehyde furan-2-carbohydrazide methanol hemisolvate, C₁₃H₁₀N₂O₄·0.5CH₃OH. *Zeitschrift für Kristallographie—New Crystal Structures*, 222 (4): 401–402.
- [39] Monfared, H. H., Bikas, R. and Mayer, P. (2010). (*E*)-3-Hydroxy-*N'*-(2-hydroxybenzylidene)-2-naphthohydrazide. *Acta Crystallographica E*, 66 (1), o236–o237.

- [40] Taha, Z. A., Ajlouni, A. M., Momani, W. A. and Al-Ghzawi, A. A. (2011). Syntheses, characterization, biological activities and photophysical properties of lanthanides complexes with a tetradentate Schiff base ligand. *Spectrochimica Acta Part A*, 81 (1): 570–577.
- [41] Foti, M. C., Daquino, C. and Geraci, C. (2004). Electron–transfer reaction of cinnamic acids and their methyl esters with the DPPH• radical in alcoholic solutions. *Journal of Organic Chemistry*, 69 (7): 2309–2314.