



Aloe vera Leaf Extract as a Potential Growth Enhancer for Populus Trees Grown Under *in vitro* Conditions

Fadia El Sherif^{1,2}

¹Department of Biological Sciences, Faculty of Science, King Faisal University, Al-Ahsaa, the Kingdom of Saudi Arabia

²Department of Horticulture, Faculty of Agriculture, Suez Canal University, Ismailia, Egypt

Email address:

Felsherif@kfu.edu.sa

To cite this article:

Fadia El Sherif. *Aloe vera* Leaf Extract as a Potential Growth Enhancer for *Populus* Trees Grown Under *in vitro* Conditions. *American Journal of Plant Biology*. Vol. 2, No. 3, 2017, pp. 101-105. doi: 10.11648/j.ajpb.20170203.13

Received: May 18, 2017; Accepted: May 31, 2017; Published: July 5, 2017

Abstract: Natural plant extracts are a cost-effective and environmental friendly alternative to synthetic plant growth regulators and phytohormones. The present *in vitro* study investigated the promotory activity of various concentrations (0, 10, 20 and 40 mL/L) of *Aloe vera* leaf extract (ALE) on the growth of hybrid aspen (*Populus tremula* L. × *Populus tremuloides* "Michx") clone T89 and aspen (*Populus tremula* L.) clone W52. The extract isolated from *Aloe vera* leaves increased the plant height and weight, number of shoots, leaves and roots, and the root length, as well as mineral concentrations of both *Populus* clones. The rooted plants were acclimatized in the greenhouse with 90% survival. The results showed that ALE is an efficient alternative source to improve the growth of both *Populus* clones under study.

Keywords: *Aloe vera* Leaf Extract, Male Trees, Aspen, Hybrid Aspen, *in vitro*, Shoot Tip

1. Introduction

Plant hormones and growth regulators have been used in agriculture and tissue culture for many decades. The search for natural compounds to replace the use of hormones and PGRs is becoming popular because of the high cost of hormones and the risk of toxicity in plants, animals, and humans due to overdose [1]. Natural plant extracts, which are rich in plant hormones, can be used to improve and stimulate growth of other plant species. Among these is the extract from the leaves of *Aloe vera* plants. *Aloe vera* is an important medicinal plant from the Liliaceae family with African origin. It is a succulent herb which grows in many countries [2]. Its large leaf parenchymatic cells contain liquid of yellow latex and clear gel [3, 4], which is rich in essential amino acids, mono- and polysaccharides, lignin, macronutrients, micronutrients, vitamins, gibberellins and salicylic acid [5, 6, 7, 8, 9, 10]. *Aloe* leaf extract (ALE) has been used to improve the vegetative growth of *Abelmoschus esculentus*, *Oenothera biennis* and *Majorana hortensis* [8, 11, 12, 13] and also as a natural plant growth regulator in *Majorana hortensis* and *Salvia officinalis* [8, 14, 15, 16]. Since the *Aloe vera* extract contains plant hormones such as gibberellin, therefore, we may be able to use it as a source of hormone instead of

synthetic growth regulators and purified natural hormones. *In vitro* culture provides controlled environmental conditions for plant culture and shortens experimental time frames.

Populus trees is a dioecious plant, native to the Northern hemisphere. It is a rapid-growing tree commonly used for wood and fuel, as well as a model organism for the study of biological functions of trees [17, 18]. Male *Populus* trees tend to have longer internodes, higher dry weight and heavier wood compared with female clones [19, 20].

The present work aimed to study the effect of various concentration of *Aloe* leaf extract (ALE) on vegetative growth of *Populus* clones under *in vitro* conditions.

2. Materials and Methods

2.1. Establishment of *in-vitro* *Populus* Plantlets

Plant material was obtained from the Experimental Farm of Ismailia Agriculture Research Station, Ismailia Governorate, Egypt. The experiments were conducted from March 1st, 2016 until October 1st, 2016 at the Tissue Culture Lab, Department of Biological Science, Faculty of Science, King Faisal University, Saudi Arabia. The male aspen (*Populus tremula* L.) clone W52 and the hybrid aspen (*Populus tremula* L. × *Populus tremuloides* "Michx") clone

T89 were used in this experiment. Shoot tips of 0.5-1.0 cm length, excised from the respective poplar trees (four years old), were sterilized to establish the *in vitro* plantlets [21].

2.2. Preparation of Aloe Leaf Extracts (ALE)

Aloe vera leaves cut from the plant (5 years old) which grown in The Agricultural-Veterinarian Training and Research Station, King Faisal University, were cold pressed using a stainless steel drum to obtain the gel solution of *Aloe* leaves. The extract was then filter-sterilized. Determination of mineral and phytohormone contents in the *Aloe vera* leaves extract were performed according to [22] and [10], respectively (Table 1 and 2).

Table 1. Determination of minerals in *Aloe vera* extract according to Rawe (1966).

Mineral	Result mg/100ml f. w
Nitrogen	81
Phosphorus	7
Potassium	61
Iron	0.3
Zinc	0.02
Manganese	0.03
Calcium	40
Copper	0.004
Magnesium	14
Sodium	51

Table 2. Determination of phytohormones (GA3, IAA and ABA) in *Aloe vera* extract according to Shyamal et al. (1990).

Phytohormone/ Nutrient	Result	Unit
GA3	16	mg/100gm f. w
IAA	0.6	mg/100gm f. w
ABA	3.1	mg/100gm f. w
Total carbohydrate	10	(%)
Glucose	3	g/100g
Protien	1.0	mg/g
Cholesterol	19	mg/g

2.3. Effect of Different Aloe Leaf Extract (ALE)

Concentrations on Growth of Two *Populus* Clones

Eight week old *in vitro* shoot tips (0.5-1.0 cm in length) were obtained from established *in vitro* cultures of male aspen *P. tremula* clone W52 and hybrid aspen *P. tremula* × *P. tremuloides* clone T89 from *in-vitro* grown plantlets. A single shoot tip explant was cultured into a 40 mL capacity jar containing 10 mL of half strength MS basal salts and vitamins medium [23] supplemented with 2.0% (w/v) sucrose and 6.0 gL⁻¹ agar. NaOH (0.1 N) or HCl (0.1 N) solution was used to adjust the medium pH to 5.7. The medium was autoclaved for 20 min at 121°C. Various concentrations of filter-sterilized *Aloe vera* extract (0, 10, 20 and 40 mL/L) were added to the medium after the autoclaved medium reached 47°C. Each treatment contained 20 replicate jars. After 8 weeks of culture, growth parameters such as plant height and weight, number of shoots, leaves and roots, as well as the longest root, were recorded. Some of the plantlets were transferred to pots with a moist mixture of sand and perlite (1:1) for acclimatization. Potted plants were grown in

a growth chamber and irrigated with a fine mist of water for 4 weeks. The percentage of plants surviving was determined after 4 weeks.

2.4. Mineral Composition

Plantlet samples from both *Populus spp.* after 8 weeks of culturing on MS medium supplemented with *Aloe vera* leaf extracts were dried at 70 °C for 24 h. The samples were ground and digested with sulphuric acid [24]. Nitrogen determination was carried out using the modified micro-Kjedahl method, as described by [25] The percentage of phosphorus was estimated calorimetrically according to the method of [26]. The Potassium (K) content was determined by using Atomic Absorption Flame Photometry according to [27]

2.5. Statistical Design and Analysis

The experiment was set up in a complete randomized design with 20 replicates per treatment. Data were statistically analyzed using ANOVA/MANOVA of Statistic 6 software [28]. The significance of differences among means was analyses using the Least Significant Test (L. S. D) at p = 0.05.

3. Result and Discussion

3.1. Effect of (ALE) on Vegetative Growth

3.1.1. Effect on Plant Height (cm)

ALE at the concentration of 10 and 20 mL/L increased the plant height of both *Populus* clones as compared with control treatment (Table 3). The ALE concentrations of 10 mL/L produced the highest plant height as (10.0 and 11.86 cm) for hybrid *P. tremula* L. x *P. tremuloides* "Michx" and *P. tremula* L. respectively (Figure 1). In contrast, the ALE at 40 mL/L significantly reduced the plant height of both clones as compared with the control treatment. The used of ALE extract in the *in-vitro* culture of *Populus* clones increase the plant height, Data in Table (2) showed that ALE extract contained the GA3 (16 mg/100gm f. w), higher growth observed could be due to GA3 found in ALE. Many reports have described that ALE increases cell membrane permeability, oxygen uptake, respiration and photosynthesis, root and cell elongation and ion transport and effects on different factor of plant growth [12, 13, 16].

3.1.2. Effect on Fresh Weight of Plant (gm)

ALE at 10 and 20 mL/L increased the fresh weight of both *Populus* clones as compared with control treatment (Table 3). However, the concentration of ALE at 40 mL/L decreased the plant fresh weight. The highest plant fresh weight of 1.32 and 1.02 g were observed in 10 mL/L ALE treatment for hybrid *P. tremula* L. x *P. tremuloides* "Michx" and *P. tremula* L. respectively. The enhancing effect of ALE on plantlet fresh weight is another indication of the plant growth enhancement effect of ALE on *in vitro* cultured *Populus* plantlets. ALE has been reported to stimulate assimilation of major and minor elements, activates enzyme, and enhances membrane

permeability, protein synthesis and biomass production [29].

3.1.3. Shoot Number Per Explant

ALE concentration at 10 and 20 ml/L produced higher shoot number per explant as compared to control treatment for both *Populus* clones (Table 3). The concentration of 10 ml/L gave the highest shoot/explant as 4.5 and 4.0

shoot/explant for the hybrid *P. tremula* L. x *P. tremuloides* “Michx” and *P. tremula* L. respectively (Figure 1). High concentration of ALE (40 ml/L) significantly decreased the shoot number per explants in the case of *P. tremula* as compared to control treatment. The use of ALE as a natural plant growth regulator has been reported by [13, 16].

Table 3. Effect of (ALE) on plant height (cm), Plant Fresh weight (g) and number of shoot /explant of two *Populus* clones, the hybrid aspen clone T89 (*Populus tremula* L. x *Populus tremuloides* Michx.) and clone W52 (*Populus tremula* L.) after eight weeks from *in vitro* culture.

Aloe leaf extract treatments (ml/L)	Plant height (cm)		Plant Fresh weight (g)		No. of shoots (n)	
	Clone T89	Clone W52	Clone T89	Clone W52	Clone T89	Clone W52
Control	7.67 b*	10.00 ab	0.51bc	0.56 bc	2.56 bed	1.86 cd
10	10.00 ab	11.86 a	1.32 a	1.02 ab	4.50 a	4.00 ab
20	9.00 ab	10.19 ab	0.73 bc	0.85 abc	4.57 a	3.63 abc
40	4.00 c	4.25 c	0.44 bc	0.28 c	2.75 abcd	1.50 d

*Means followed by the same letter within a column are not significantly different at 0.05 level of probability according to L. S. D. test




Figure 1. Effects of ALE extract on growth of *Populus* hybrid aspen (*Populus tremula* L. x *Populus tremuloides* Michx) clones T89 (A) and aspen (*Populus tremula* L.) clone W52 (B) after eight weeks growth on half MS medium supplemented with 10 ml/L ALE.

Table 4. Effect of (ALE) on number of leaves (n), root length (cm), number of roots and plant height (cm) of two *Populus* clones, the hybrid aspen clone T89 (*Populus tremula* L. x *Populus tremuloides* Michx.) and clone W52 (*Populus tremula* L.) after 8 weeks from *in vitro* culture.

Aloe leaf extract treatments (ml/L)	No. of leaves (n)		No. of roots (n)		Root length (cm)	
	Clone T89	Clone W52	Clone T89	Clone W52	Clone T89	Clone W52
Control	23.89 bcd*	14.86 d	3.22 bc	4.71 abc	7.00 ab	5.29 bc
10	34.50 a	25.14 abc	6.33 a	6.14 a	9.33 a	5.07 bc
20	30.57ab	21.13 ab	6.14 a	5.63 ab	6.00 bc	4.56 bc
40	27.25 abc	16.75 cd	3.00 bc	2.50 c	2.75 c	4.13 bc

*Means followed by the same letter within a column are not significantly different at 0.05 level of probability according to L. S. D. test

3.1.6. Ex-vitro Acclimatization

In vitro regenerated plantlets of both *Populus* clones showed 90 % survival when transferred to soil and did not display any phenotypic variation during subsequent vegetative development.

3.1.4. Number of Leaves / Explant

All ALE concentration increased the number of leaves per explant than that of control in both populus clones (Table 4). The highest numbers of leaves/ explant were 34.5 and 25.14 for hybrid *P. tremula* L. x *P. tremuloides* “Michx” and *P. tremula* L. respectively. These were observed on plantlets grown on media with 10 ml/L of ALE. The obtained results are in line with the findings of [8, 14].

3.1.5. Root Number Per Explants and Root Length Per Explant

ALE at 10 and 20 ml/L increase the number of root per explant for both *Populus* clones (Table 4, Figure 1 a and b). At 40 ml/L ALE concentration, the root number of the two *Populus* clones decreased compared to control treatment. Interestingly, ALE at 10 ml/L only increased the root length of the hybrid *P. tremula* L. x *P. tremuloides* “Michx”. As for *P. tremula* L., decrease root length was observed with the increase of ALE concentration. The results showed that ALE at (10 and 20 ml/l) could be considered as an alternative auxin-enriched *in vitro* rooting medium at rooting stage for both clones as ALE containing IAA (0.6 mg/100gm f. w) Table (2).

3.2. Effect of ALE leaf extract Treatment on Chemical Contents of Leaves

3.2.1. Total Nitrogen (N) Percentage in the Leaves

The nitrogen percentage increased in parallel with ALE concentration in both *Populus* clones (Figure 2a). At ALE

concentrations of 40ml/L, highest nitrogen percentages were observed, with (3.09 and 3.20 %) for hybrid *P. tremula* L. x *P. tremuloides* “Michx” and *P. tremula* L. respectively. Lowest nitrogen percentage was observed in control treatment, with 0.96 and 1.19 % for hybrid *P. tremula* L. x *P. tremuloides* “Michx” and *P. tremula* L. respectively.

3.2.2. Total Phosphorus (P) Percentage in the Leaves

All ALE concentrations increased the phosphorus concentration in the leaves of two *Populus* clones as compared to control (Figure 2b). The higher percentages of phosphorus were (0.37 and 0.43%) for the plants treated with 10 or 40ml/L in the case of *P. tremula* L. x *P. tremuloides* “Michx” and *P. tremula* L. respectively.

3.2.3. Total Potassium (K) Percentage in the Leaves




Figure 2. Effect of (ALE) on the percentage of Nitrogen (N) (a), Phosphorus (P) (b) and Potassium (K) (c) in the leaves of *Populus* hybrid aspen (*Populus tremula* L. x *Populus tremuloides* Michx) clone T89 and aspen (*Populus tremula* L.) clone W52 after 8 weeks from *in vitro* culture.

All concentrations of ALE tended to increase the potassium percentage in relative to the control treatments as for both *Populus* clones (Figure 2c). The potassium percentage were (1.90 and 1.95%) for hybrid *P. tremula* L. x *P. tremuloides* “Michx” and *P. tremula* L. in the control treatment. The highest potassium percentage as (4.05 and 3.55%) were recorded by using ALE at concentrations of 10 or 40ml/L in the case of *P. tremula* L. x *P. tremuloides* “Michx” and *P. tremula* L. respectively. Increasing N, P and K in leaves of both *Populus* clones treated with ALE supported the hypothesis that ALE stimulates plant growth by

assimilation of major and minor elements. Similar increase in major elements in plant leaves as result of ALE treatments was also reported by [7, 9, 30].

4. Conclusion

Results showed that *Aloe vera* leaf extract (ALE) increased the growth (plant height, fresh weight, number of leaves and roots, root length), multiplication (number of shoot per explants), and major nutrients (nitrogen, phosphorus and potassium) contents of leaves in both *Populus* clones. The best ALE concentration was 10 ml/L, which gave the highest recorded value for all the above mentioned parameters. In summary, the results have suggested that ALE is an efficient alternative source to improve the growth of both *Populus* clones under study.

Acknowledgements

The author would like to thank Dr. Salah Khattab and Dr. Yun-Kiam Yap, Department of Biological Sciences, Faculty of Science, King Faisal University, Al-Hassa, Kingdom of Saudi Arabia, for their kind help and assistance.

References

- [1] Cutler HG, Schneider BA. Plant Growth Regulator Handbook. LaGrange, GA: Plant Growth Regulator Society of America. 1990.
- [2] Pandey A, Singh S. *Aloe Vera*: A Systematic Review of its Industrial and Ethno-Medicinal Efficacy. International Journal of Pharmaceutical Research and Allied Sciences. 2016; 5: 21-33.
- [3] Surjusha A, Vasan i R, Saple D. *Aloe vera*: a short review,”Indian Journal of Dermatology. 2008; 53: 163–166.
- [4] Ni Y, Turner D, Yates KM, Tizard I. Isolation and characterization of structural components of *Aloe vera* L. leaf pulp. Int Immunopharmacol 2004; 4: 1745–55. doi: 10.1016/j.intimp.2004.07.006 PMID: 15531291.
- [5] Chatterjee p, Bodhisattwa C, Subhangkar N. Significant Pharmacological Activities of *Aloe vera* plant. Min. J. Phar. Med. Sci. 2013; 3: 21-24.
- [6] Raman RPV, Rita R S, Mondal B C, Singh SK. Effect of *Aloe vera* and clove powder supplementation on carcass characteristics, composition and serum enzymes of Japanese quails. Vet. World. 2013; 8: 664-668.
- [7] Sahu K P, Giri D D, Singh R, Pandey P, Gupta S, Shrivastava A K, Kumar A, Pandey D K. Effect of *Aloe vera* on some annual plants Sci. Res. Pharmacology and Pharmacy. 2013; 4: 599-610.
- [8] Hamouda A M A, Hendi D M G, Abu-El-Leel O F A. Improving basil growth, yield and oil production by *Aloe vera* extract and active dry yeast. Egypt Journal Horticulture. 2012; 39: 45-71.
- [9] Rajeswari R, Umadevi M, Rahale S C, Pushpa R, Selvavenkadesh K P, Kumar S, Bhowmik D. *Aloe vera*: The Miracle Plant Its Medicinal and Traditional Uses in India. J. of Pharmacognosy and Phytochemistry. 2012; 4: 118-124.

- [10] Shyamal KN, Lok MP, Charles WP. Dynamics of endogenous cytokinins during the growth cycle of a hormone- autotrophic genetic tumor line of Tobacco. *Plant Physiology*. 1990; 94: 1084-1089.
- [11] Ahmed S K, Hammam KHA, Amer A A. Effect of bio-fertilization and some plant extracts on the growth, yield and chemical constituents of basil plants. *Journal Plant Production Mansoura Univ*. 2014; 5: 193-210.
- [12] El-Shayeb N S A. Physiological studies on *Oenothera biennis* bio- fertilizer and plant extracts. PhD. Thesis, Hort. Dept. Fac. Agric. Benha Univ. 2009.
- [13] Padmaja C K, Kowsalya B, Seethalakshmi C. Efficacy of *Aloe vera* leaf powder as biostimulant in enhancing the growth and yield of Lady's Finger (*Abelmoschus esculentus* L.). *Research on Crops*. 2007; 8: 395-397.
- [14] Mady A. Effect of certain medicinal plant extracts on growth, yield and metabolism of some medicinal and aromatic and plants. (M. Sc. Thesis) Fac. Science Al-Azher Univ. 2008.
- [15] Nour El-Din T. Physiological studies on marjoram plants (*Majorana hortensis*). M. Sc. Thesis of Ornamental Horticulture Department Faculty of Agriculture at Moshtohor Zagazig University (Banha Branch). 2005.
- [16] Dongzhi L, Tsuzuki E, Sugimoto Y, Yanjun D, Matsuo M, Terao H Allelopathic effects of aqueous *Aloe vera* leaf extract on selected crops. *Statistical Allelopathy Journal*. 2004; 13: 67-74.
- [17] Xiao X, Yang F, Zhang S, Korpelainen H, Li C. Physiological and proteomic responses of two contrasting *Populus cathayana* populations to drought stress. *Physiologia Plantarum*. 2009; 136: 150-168.
- [18] Taylor G A I L. *Populus*: Arabidopsis for forestry. Do we need a model tree?" *Annals of Botany*. 2002; 90: 681-689.
- [19] Khosla P K, Deol G S. Growth response of Male and Female clones of *Populus Ciliata* "Wall." Ex royle to nitrogenfertilizer. *Proc. Indian main. Sci. Acad*. 1984; 50: 603-606.
- [20] Khattab S. Effect of Different Media and Growth Regulators on the *in vitro* Shoot Proliferation of Aspen, Hybrid Aspen and White Poplar Male Tree and Molecular Analysis of Variants in Micropropagated Plants. *Life Science Journal*. 2011; 8: 177-184.
- [21] Jiang C 1, Zhaopu L, Qingsong Z. Direct Regeneration of Plants Derived from *in vitro* Cultured Shoot Tips and Leaves of Poplar (*Populus ×euramericana* 'Neva'). *Journal of Life Sciences*. 2015; 9: 366-372.
- [22] Rawe GJ. Food analysis by atomic absorption spectroscopy varian. Tech. Tron. Australia, USA Switzerland. 1966.
- [23] Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant physiology*. 1962; 13: 473-497.
- [24] Piper OS. Soil and plant analysis. In. University of Adelaide, Adelaide, Australia. 1947.
- [25] Jackson ML. Soil Chemical Analysis. Prentice Hall of India, Private Limited; New Delhi. 1967.
- [26] Murphy J, Riley JP. A modified single-solution method for the determination of phosphorus in natural water: *Analytica Chimica Acta*. 1962; 27: 31-36.
- [27] Wilde SA, Corey RB, Lyer JG, Voight GK. Soil and plant analysis for Tree Culture, 3 edn. Oxford and IBM Puplicing Co, New Delhi. 1985.
- [28] Statsoft Inc. STATISTICA for Windows (software-system fur Datenanalyse) Version 6. <http://www.statsoft.com>. 2001.
- [29] Sheikh F A, Singh R P P, Lehana P. Effect of high Voltage on the resistance of *Aloe vera* leaves. (*IJRTE*). 2013; 2: 42-45.
- [30] Hamman H J. Composition and applications of *Aloe vera* leaf Gel. *Molecules*. 2008; 13: 1599-1616.