

Study on Present Status of Fish Biodiversity in Wetlands of Sylhet District, Bangladesh

Mohammed Ariful Islam¹, Md. Jahidul Islam¹, Sanzib Kumar Barman¹, Farjana Morshed¹,
Sabiha Sultana Marine²

¹Department of Aquatic Resource Management, Faculty of Fisheries, Sylhet Agricultural University, Sylhet, Bangladesh

²Department of Fisheries Technology and Quality Control, Faculty of Fisheries, Sylhet Agricultural University, Sylhet, Bangladesh

Email address:

arif.fisheries@gmail.com (M. A. Islam)

To cite this article:

Mohammed Ariful Islam, Md. Jahidul Islam, Sanzib Kumar Barman, Farjana Morshed, Sabiha Sultana Marine. Study on Present Status of Fish Biodiversity in Wetlands of Sylhet District, Bangladesh. *Agriculture, Forestry and Fisheries*. Vol. 4, No. 6, 2015, pp. 296-299.
doi: 10.11648/j.aff.20150406.20

Abstract: The study was conducted to identify the present status of fish biodiversity in the wetlands of Sylhet district for a period of 12 months from January 2014 to December 2014. It was done by questionnaire interviews (QI) of fishers, focus group discussions (FGD), and key informant interviews (KII) and secondary data collection. During the study period, a total of 58 fish species under 21 families were recorded. The species availability status was remarked in four categories and obtained as 24 commonly available, 16 moderately available, 18 rarely available species. Highest number of commonly available species was found in October to December and lowest number of commonly available species was observed in March to April. Among 54 threatened fish species listed by IUCN Bangladesh, about 30 species were found 10-15 years ago in those wetlands but only 23 were found during the study period. It is revealed that there has been gradual reduction of fish diversity in the wetlands of the area of Sylhet district and average fish catch per fisherman per day was also reduced. Community based fisheries management, fishing gears maintenance, sanctuary establishment and management, implementation of fish acts and regulations, stocking of fish fingerling in the open waters, dredging of beels and raising public awareness can play a great role in conserving fish biodiversity.

Keywords: Biodiversity Reduction, Species Availability, Threatened Species, Conservation, Questionnaire Interview

1. Introduction

Aquatic biodiversity has enormous economic and aesthetic value and is largely responsible for maintaining and supporting overall environmental health (Hossain, 2012). Wetlands are one of the world's key natural resources (Khan et al., 1994). It is the transition between land and water and is the most productive ecosystems in the world. The biodiversity of the wetland ecosystem is variable in the world; it encompasses the range of living things, the degree of genetic variation, and the wealth of different habitats within a particular ecosystem.

The Ramsar Convention (1971) has defined wetlands as – “areas of marsh, fen, peat land, or water, whether natural or artificial, permanent or temporary, with water that is static or flowing, fresh, brackish or salt, including areas of marine water the depth of which at low tide does not exceed six meters.” Bangladesh is a home to at least 265 freshwater fish

species (Rahman, 2005). Huge number of wetlands in various forms viz. rivers, haors, boars, beels, pond, ditch etc. support these large number of fish species. Among them, wetlands of Sylhet district are so important that contributes a huge amount of fish to the people of the country. The major wetlands in Sylhet district are Hailkar, Jilkar, Patharchauli, Jainkar, Chauldhani, Balai, Muria, Erali and Damrir haors.

At present time, reduction in the abundance and kinds of fish species from the inland waters or wetlands of Bangladesh is a burning issue in the country (Galib et al. 2009, Imteazzaman and Galib 2013). However, a total of 54 fish species of Bangladesh have been declared threatened by IUCN (IUCN Bangladesh 2000). All these findings clearly indicate the need for water body specific detailed biodiversity studies which is essential to assess the present status of fish biodiversity and sustainable management of a body of water (Galib et al. 2013a; Imteazzaman and Galib 2013). Though such type of research efforts are much common in neighbor

countries like India (Dahanukar *et al.* 2012, Kharat *et al.* 2012, Baby *et al.* 2010, Jadhav *et al.* 2011, Patra 2011, Johnson and Arunachalam 2009, Heda 2009, Saha and Bordoloi 2009) but very few in Bangladesh.

To the best knowledge of the authors no previous research work has been conducted on fish biodiversity of Wetlands of Sylhet Districts, Bangladesh. So, this study will be very significant for the assessment of present status of fish biodiversity of Wetlands of Sylhet district, Bangladesh.

2. Methodology

2.1. Description of Study Area

Wetland fish species diversity was recorded in Sylhet district in Bangladesh from January to December in 2014. The selected wetlands were Hailkar, Jilkar, Patharchauli, Jainkar, Chauldhani, Balai, Muria, Erali and Damrir haors. The geographical location of the Sylhet district is an area of 3452.07 km² (1332.85 sq. miters); it is bounded by Sunamgonj district on the west, Moulvibazar district on the south and Habogonj district on the south-west. The sites of the study are illustrated in Figure 1.




Figure 1. Location of wetlands at different upazilas of Sylhet district.

2.2. Data Collection

The study was based on field survey method where an appropriate questionnaire was prepared and used for collecting data from villagers of the surrounding study area under Sylhet district. During collection of data, both primary and secondary sources were considered to interpret the results.

Primary data were collected from 40 randomly selected fishermen through questionnaire interviews (QI) and focus

group discussions (FGD) where Upazila Fisheries Officer (UFO), union parishad chairman & members, leaders of the fisher community, fish market leaders, fish traders, fry traders and community people of the selected wetlands area were also present. The secondary information was collected from upazila fisheries office under Sylhet district, district fisheries office of Sylhet, books, journals and others. After collecting data, it was cross-checked through key informant interviews (KII) with Upazila Fisheries Officer (UFO), District Fisheries Officer (DFO), school teachers, local leaders and NGO workers in the study area. Finally data were analyzed by using Microsoft office excels 2010.




Figure 2. The design of the present study involved the steps.

3. Results and Discussion

According to the statement of local fishermen, a total of 58 fish species under 21 families were recorded. The recorded fish species with accurate taxonomy were identification by the cross-matching of definite fish characters from the text book, named “Freshwater Fishes of Bangladesh” (A. K. Ataur Rahman, Department of Fisheries, Matshya Bhaban, Dhaka). The recorded fish species are represented in the following table with their biodiversity status and IUCN status.

Table 1. List of recorded fish species with their status.

Sl. No.	Family	Local Name	Common Name	Scientific Name	Biodiversity status	IUCN Status
1	Cyprinidae	Rohu	Indian Major Carp	<i>Labeo rohita</i>	CA	NO
2	Cyprinidae	Carpio	Common Carp	<i>Cyprinus carpio</i>	CA	EX
3	Cyprinidae	Kalibaush	Black Rohu	<i>Labeo calbasu</i>	CA	EN
4	Bagridae	Bujuri	Long Bled Catfish	<i>Mystus tengra</i>	CA	NO
5	Bagridae	Tengra	Striped Dwarf Catfish	<i>Mystus vittatus</i>	CA	NO
6	Siluridae	Boal	Freshwater Shark	<i>Wallago attu</i>	CA	NO

Sl. No.	Family	Local Name	Common Name	Scientific Name	Biodiversity status	IUCN Status
7	Siluridae	Modhu Pabda	Butter Catfish	<i>Ompok pabda</i>	CA	EN
8	Siluridae	Pabda	Pabo Catfish	<i>Ompok pabo</i>	CA	EN
9	Clariidae	Magur	Walking Catfish	<i>Clarius batrachus</i>	CA	NO
10	Heteropneustidae	Shing	Stinging Catfish	<i>Heteropneustes fossilis</i>	CA	NO
11	Cyprinidae	Tit Punti	Ticto Barb	<i>Puntius ticto</i>	CA	VU
12	Cyprinidae	Teri Punti	One Spot Barb	<i>Puntius terio</i>	CA	NO
13	Cyprinidae	Jat Punti	Spot Fin Swamp Barb	<i>Puntius sophore</i>	CA	NO
14	Channidae	Taki	Spotted Snakehead	<i>Channa punctatus</i>	CA	NO
15	Channidae	Shol	Snakehead Murrel	<i>Channa striatus</i>	CA	NO
16	Mastacembelidae	Guchi Baim	Striped Spiny Eel	<i>Macrognathus pancalus</i>	CA	NO
17	Mastacembelidae	Tara Baim	One Striped Spiny Eel	<i>Macrognathus aculeatus</i>	CA	VU
18	Anabantidae	Baro Khalisha	Striped Gourami	<i>Colisa fasciatus</i>	CA	NO
19	Anabantidae	Chota Khalisha	Honey Gourami	<i>Colisa chuno</i>	CA	NO
20	Anabantidae	Koi	Climbing Perch	<i>Anabas testudineus</i>	CA	NO
21	Ambassidae	Gol Chanda	Indian Glass Fish	<i>Parambassis ranga</i>	CA	VU
22	Nandidae	Meni	Mud Perch	<i>Nandus nandus</i>	CA	VU
23	Cobitidae	Gutum	Guntea Loach	<i>Lepidocephalichthys guntea</i>	CA	NO
24	Palaemonidae	Sada Icha	Prawn	<i>Macrobrachium</i> sp.	CA	NO
25	Cyprinidae	Catla	Indian Major Carp	<i>Catla catla</i>	MA	NO
26	Cyprinidae	Mrigal	Indian Major Carp	<i>Cirrhinus cirrhosus</i>	MA	NO
27	Cyprinidae	Grass Carp	Grass Carp	<i>Ctenopharyngodon idella</i>	MA	EX
28	Cyprinidae	Silver Carp	Silver Carp	<i>Hypophthalmichthys molitrix</i>	MA	EX
29	Cyprinidae	Mola	Carplet	<i>Amblypharyngodon mola</i>	MA	NO
30	Cyprinidae	Darkina	Flaying Barb	<i>Esomus danricus</i>	MA	DD
31	Channidae	Cheng	Asiatic Snakehead	<i>Channa orientalis</i>	MA	VU
32	Bagridae	Golsha	Long Whiskered Catfish	<i>Mystus cavasius</i>	MA	VU
33	Channidae	Gozar	Giant Snakehead	<i>Channa marulius</i>	MA	EN
34	Mastacembelidae	Baro Baim	Two-track Spiny Eel	<i>Mastacembelus armatus</i>	MA	EN
35	Sybranchidae	Cuchia	Gangetic Mud Eel	<i>Monopterusuchia</i>	MA	VU
36	Ambassidae	Lamba Chanda	Elongated Glass Perchlet	<i>Chanda nama</i>	MA	VU
37	Gobiidae	Bele	Bar Eyed Goby	<i>Glossogobius giuris</i>	MA	NO
38	Notopteridae	Foli	Bronze Featherback	<i>Notopterus notopterus</i>	MA	VU
39	Cobitidae	Bou Rani	Bengal Loach	<i>Botio dario</i>	MA	EN
40	Tetraodontidae	Potka	Ocellated Puffer fish	<i>Tetraodon cutcutia</i>	MA	NO
41	Cyprinidae	Goniya	Kuria Labeo	<i>Labeo gonius</i>	RA	EN
42	Cyprinidae	Bata	Minor Carp	<i>Labeo bata</i>	RA	EN
43	Cyprinidae	Dhela	Cotio	<i>Osteobrama cotio</i>	RA	EN
44	Cyprinidae	Chela	Finescale Razorbelly Minnow	<i>Chela phulo</i>	RA	NO
45	Cyprinidae	Shar Punti	Olive Barb	<i>Puntius sarana</i>	RA	CR
46	Bagridae	Rita	Rita	<i>Rita rita</i>	RA	CR
47	Bagridae	Ayre	Long Whiskered Catfish	<i>Mystus aor</i>	RA	VU
48	Bagridae	Gagla	Menoda Catfish	<i>Hemibagrus menoda</i>	RA	NO
49	Schilbeidae	Gharua	Gharua Bacha	<i>Clupisoma garua</i>	RA	CR
50	Schilbeidae	Batashi	Indian Potasi	<i>Pseudeutropius atherinoides</i>	RA	NO
51	Schilbeidae	Kajoli	Gangetic Ailia	<i>Ailia coilia</i>	RA	NO
52	Pangasidae	Thai Pangus	Sutchi Catfish	<i>Pangasius hypophthalmus</i>	RA	EX
53	Clupeidae	Chapila	Indian River Shad	<i>Gaduasias chapra</i>	RA	NO
54	Anabantidae	Lal Khalisha	Dwarf Gourami	<i>Colisa lalia</i>	RA	NO
55	Ambassidae	Lal Chanda	Indian Glass Perchlet	<i>Parambassis lala</i>	RA	EN
56	Cichlidae	Tilapia	Mozambique Tilapia	<i>Oreochromis mossambicus</i>	RA	EX
57	Hemiramphidae	Ekthute	Congaturi Halhbeak	<i>Hyporhamphus limbatus</i>	RA	NO
58	Palaemonidae	Golda	Prawn	<i>Macrobrachium rosenbergii</i>	RA	NO

CA=Commonly available, MA=Moderately available, RA=Rarely available.

CR=Critically endangered, EN=Endangered, VU=Vulnerable, NO=Not threatened, DD=Data deficient and EX=Exotic species.

4. Recommendation

- Fish sanctuary should be established in the selected wetlands area before breeding season.
- No fishing in the fish sanctuary in a defined time (several months) should be ensured.
- Overfishing should be prohibited in the wetlands area.
- Banded fishing gears (specially banded fishing nets)

should be prohibited in the selected areas.

- Community based fisheries management (CBFM) should be established in the haor areas.
- Alternative earning source of the people of haor areas should be provided during banded season of fishing.
- Increasing awareness among the much people of surrounding wetland areas.

5. Conclusion

Wetlands of Sylhet district are generally considered as a highly diversified zone of Bangladesh mainly for its rich aquatic biodiversity. According to the statement of the respondent fishermen, the total fish biodiversity is reducing drastically in the wetlands of Sylhet district due to lack of proper management. As a consequence, wetland ecosystem protection is important for species conservation and the protection of a sustainable environment.

References

- [1] Baby F, Tharian J, Ali A and Raghavan R (2010). A checklist of freshwater fishes of the New Amarambalam Reserve Forest (NARF), Kerala, India. *Journal of Threatened Taxa* 2(12): 1330-1333.
- [2] Dahanukar N, Paingankar M, Raut RN and Kharat SS (2012). Fish fauna of Indrayani River, northern Western Ghats, India. *Journal of Threatened Taxa* 4(1): 2310-2317.
- [3] Galib SM, Naser SMA, Mohsin ABM, Chaki N and Fahad MFH (2013a). Fish diversity of the River Choto jamuna, Bangladesh Present status and conservation needs. *International Journal Biodiversity and Conservation* 5(6): 389-395. DOI: 10.5897/IJBC2013.0552.
- [4] Hossain, M., 2012. Biodiversity of Threatened Fish Species of Choto Jamuna River in Badalgachhi Area under Naogaon District. MS Thesis, Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh, pp: 30-53.
- [5] IUCN Bangladesh (2000). Red book of threatened fishes of Bangladesh, IUCN- The world conservation union. xii+116 pp.
- [6] Jadhav BV, Kharat SS, Raut RN, Paingankar M and Dahanukar N (2011). Freshwater fish fauna of Koyna River, northern Western Ghats, India. *Journal of Threatened Taxa* 3(1): 1449-1455.
- [7] Johnson JA and Arunachalam M (2009). Diversity, distribution and assemblage structure of fishes in streams of southern Western Ghats, India. *Journal of Threatened taxa* 1(10): 507-513.
- [8] Patra AK (2011). Catfish (Teleostei: Siluriformes) diversity in Karala River of Jalpaiguri District, West Bengal, India. *Journal of Threatened Taxa* 3(3): 1610-1614.
- [9] Rahman, AKA (2005). Freshwater Fishes of Bangladesh, second edition. Zoological Society of Bangladesh, University of Dhaka, Dhaka, Bangladesh, 263 pp.
- [10] Saha S and Bordoloi S (2009). Ichthyofaunal diversity of two beels of Goalpara District, Assam, India. *Journal of Threatened Taxa* 1(4): 240-242.
- [11] Rahman, A. K. A. (1989). Freshwater Fishes of Bangladesh. Zoological Society of Bangladesh. Department of Zoology, University of Dhaka. Dhaka-1000.
- [12] Khan, M. S., Haq, S., Rahman, A. A., Rashid S. M. A. and Ahmed. H., 1994. Wetlands of Bangladesh. Bangladesh Centre for Advanced Studies & Nature Conservation Movement.