

Morphological and genetic variation of *Thrips parvispinus* (Thysanoptera: Thripidae) in chili plantation (*Capsicum annum* L.) in the lowland and highland of Jambi Province, Indonesia

Asni Johari^{1,*}, Siti Herlinda², Yulia Pujiastuti², Chandra Irsan², Dewi Sartiami³

¹Graduate School of Science Education, Jambi University, Jl. Raden Mattaher No. 16 Jambi

²Department of Plant Pests and Diseases, Faculty of Agriculture, Sriwijaya University, Jl. Raya Palembang- Prabumulih Km 32, Ogan Ilir, Indralaya 30662

³Laboratory of Biosystematics, Plant Protection Department, Bogor Agricultural University, Jl. Meranti, Campus IPB, Bogor 16680

Email address:

johari_asni@yahoo.com (A. Johari)

To cite this article:

Asni Johari, Siti Herlinda, Yulia Pujiastuti, Chandra Irsan, Dewi Sartiami. Morphological and Genetic Variation of *Thrips parvispinus* (Thysanoptera: Thripidae) in Chili Plantation (*Capsicum annum* L.) in the Lowland and Highland of Jambi Province, Indonesia.

American Journal of BioScience. Special Issue: Chemical Biology. Vol. 2, No. 6-1, 2014, pp. 17-21.

doi: 10.11648/j.ajbio.s.2014020601.14

Abstract: *Thrips parvispinus* Karny (Thysanoptera: Thripidae) are dangerous pests for chili (*Capsicum annum* L.). Thrips can be found in lowland to highland. The damage caused by thrips can reach 60% and this damage will be greater if the thrips also be vectors of plant virus diseases. The research aims to analyze morphological and genetic variation of *T. parvispinus* located in the lowlands and in the highlands of Jambi province, Indonesia. *T. parvispinus* were collected in 2012 from chili plantation in 19 villages spread over 5 districts in the lowlands and 26 villages spread over 2 districts in the highlands. The results showed that *T. parvispinus* were found at each research site. The size and color of the body *T. parvispinus* have variations with long, medium and short criteria. Variation in body size of *T. parvispinus* in lowland: length of 1.42 ± 0.065 mm, medium 1.32 ± 1.15 mm and short 0.122 ± 0.051 mm. Variation in body size of *T. parvispinus* in the highlands: length of 1.46 ± 0.035 , medium of 1.35 ± 0.051 and short of 1.22 ± 0.089 . *T. parvispinus* with long body size and dark brown color predominantly are found in all survey sites, is 80.53% in the lowlands and 75.91% in the highlands. The results also showed that the insects having different body sizes belong to the same *T. parvispinus* species.

Keywords: *T. parvispinus*, Morphology, Color, DNA

1. Introduction

T. parvispinus is a dangerous insect pests that attack and damage chili plants. These pests damage plants by sucking and whittling (Kalshoven 1981). The damage, which was caused by thrips in Bandung and Bogor Regency of Indonesia ranged 10-46%. Yield losses due to attack *T. parvispinus* on chili plantation could reach 22.8% (Sastrosiswojo 1991). Plant damage, which was caused by thrips will be greater if the thrips was to be vectors of plant virus diseases (Ullman & German, 1995; Marullo and Mound, 2002).

T. parvispinus is originating from Southeast Asia with regional distribution starting from Thailand, Malaysia to

New Guinea and northern Australia. Currently these thrips are also found in Greece. According to Mound (2006) the inclusion of *T. parvispinus* in Greece it's because taken along accidentally by the *Gardenia* sp. flower from Indonesia.

T. parvispinus are thrips that become major pests for vegetable plants in Indonesia (Sastrosiswojo 1991). According to Vos (1994) *T. parvispinus* is a major pest for cultivating chili in Java. Mound (2006) reported that one of the thrips that are so damaging species in Southeast Asia is *T. parvispinus*. Talekar (1991) reported that the species of *T. parvispinus* predominantly are found in Indonesia and most destructive chili plants. Prabaningrum (2007) reported that only *T. parvispinus* are found in paprika plant in Bandung

regency, West Java. The result of preliminary survey found that morphological *T. parvispinus* are varied. However, research of morphology of *T. parvispinus* is still very limited. Therefore, aims of this study were to identify and analyze morphological variations and some other characteristics of *T. parvispinus* that attack chili plants in the lowlands and in the highlands province of Jambi, Indonesia. This research can increase knowledge about of *T. parvispinus*.

2. Methods

The survey was conducted between February and August 2012 at the height of 1-200 meters above sea level in the lowlands and 800-1750 meters above sea level in the highlands. Research sites had temperatures in the range 23-32°C in the lowlands and 17-29°C in the highlands, the air relative humidity in the lowlands ranged between 60-98% and highlands ranged 58% -95%.

Exploration thrips was done by collecting thrips imago of 150 plants at each location by tracing a line transect (Khan 2006). The thrips imago collected were put into a plastic vial which contains 70% alcohol volume of 50 ml to prevent damage before it's made slides for identification. The microscopic slides were made for identification of collected thrips species. Identification was conducted in the department of plant protection Biosystematics Laboratory IPB Bogor, Indonesia.

Identification was done by observing, recording and documenting morphological characteristics such as, antenna, ocelli, tassels and wings color, metanotum,

abdomen and other vital parts binocular compound microscope 40x magnifications. Identification of species using identification keys which made by; Mound and Kibby (1998); Moritz *et al.* (2001); Mound (2006).

DNA extraction from thrips was performed by method of Goodwin *et al.* (1994).

3. Results and Discussion

The results of identification of *T. parvispinus* specimen are showed in Table 1 and Fig. 1. The morphological features of *T. parvispinus* are as follows: (1) body color was brown to dark brown; color of head and thorax was brighter than the abdomen, (2) head had strong reticulation patterns and big eyes that have pigment, (3) compound eyes had not elongated shape, (4) antenna consisted of seven segments, while the second and third segments had a fork-shaped sensory organs, (5) sample does not have ocellar setae 1, ocellar setae 2 is shorter than ocellar setae 3, (6) pronotum had 2 pairs of long setae posteroangular and 3 pairs of setae posteromarginal were shorter, (7) at the metanotum there was no sensilla campaniform, (8) at Tergit VIII there was a comb (comb microtrichia), (9) at the V-VIII there was a ctenidia at part of lateral and at Tergit VIII ctenidia which was located behind the spiracles, (10) length of wing was more than half the length of the abdomen, (11) dark wings or shaded, with a pale base, (12) at the first and second wings vein front there was a complete line of seta (Fig. 1). Sequences analysis of DNA fragments from three specimens (A, B and C) were found approximately 655 nucleotides with 99.38% consider identical with *T. Parvispinus*.

Table 1. Plateau place found *T. parvispinus* in chilli plantation at lowland and highland in Jambi Province

Family	Species	Altitude (mdpl)
Lowland		
Thripidae	<i>Thrips parvispinus</i>	8, 10, 17, 20, 21, 22, 24, 25, 26, 27, 29, 31, 34, 54, 56, 60, 65, 126
Highland		
Thripidae	<i>Thrips parvispinus</i>	837, 847, 984, 1.054, 1.071, 1.079, 1.104, 1.391, 1.416, 1.416, 1.445, 1.471, 1.473, 1.480, 1.498, 1.507, 1.508, 1.510, 1.517, 1.558, 1.560, 1.607, 1.623, 1.639, 1.710, 1.713




Figure 1. Morphology of *T. parvispinus* a. female imago, b. Ocellar, c. metanotum, d. antenna, e. wings, f. tergite VIII

T. parvispinus which were found in its own body size varies. They were long, medium and short (Figure 2). The average body length of *T. parvispinus* found in the lowlands was the size of the body length of 1.42 ± 0.065 mm, medium of 1.32 ± 1.15 mm and short of 1.22 ± 0.051

mm. In the highlands of *T. parvispinus* body size length of 1.46 ± 0.035 mm, medium was 1.35 ± 0.051 mm and short of 1.22 ± 0.089 mm (Table 2). Differences in body length of *T. parvispinus* long category, medium and short were statistically significantly different.




Figure 2. Body size variation of long *T. parvispinus* (1.42 mm) (a), medium (1.32 mm) (b) and short (1.17 mm) (c) at mulberry 4x

Table 2. Body size of imago of *T. parvispinus*

Parameters	Body size, mm		
	Long, dark brown	Medium, dark brown	Short, dark brown
Lowland			
Long of body	1.42 ± 0.065	1.32 ± 0.051	1.15 ± 0.122
Long of abdomen	0.85 ± 0.041	0.79 ± 0.049	0.68 ± 0.088
Long of wings	0.72 ± 0.031	0.69 ± 0.042	0.65 ± 0.046
Long of antenna	0.28 ± 0.010	0.27 ± 0.008	0.26 ± 0.015
Width of head	0.16 ± 0.015	0.16 ± 0.012	0.15 ± 0.015
Width of thorax	0.30 ± 0.024	0.30 ± 0.027	0.29 ± 0.029
Highland			
Long of body	1.46 ± 0.035	1.35 ± 0.051	1.22 ± 0.089
Long of abdomen	0.89 ± 0.043	0.83 ± 0.064	0.73 ± 0.077
Long of wings	0.75 ± 0.014	0.73 ± 0.031	0.70 ± 0.046
Long of antenna	0.29 ± 0.013	0.28 ± 0.010	0.27 ± 0.011
Width of head	0.16 ± 0.014	0.15 ± 0.010	0.15 ± 0.007
Width of thorax	0.30 ± 0.017	0.28 ± 0.021	0.29 ± 0.027

Explanation: Mean of the result measurement from 165 trips imago (adult thrips)

Body color of *T. parvispinus* was found vary (Table 3). Body color begins to brown to dark brown. *T. parvispinus* size and body length dark brown are dominant species found than *T. parvispinus* medium-sized body and a short. The results of Prabaningrum research (2007) found that the color and body length of *T. parvispinus* varied. Mound and Collins (2000) found a morphological variation in *T. parvispinus*. To the variations in body length of *T. parvispinus* were performed electrophoresis testing. The Testing used 1% gel of agarose gel PCR thrips with primer pairs ITS 2 thrips. The test results showed that *T. parvispinus* with varying body length was still classified into the same species.

Table 3. The amount and percentage of *T. parvispinus* at lowland and highland in Jambi Province

<i>T. parvispinus</i>	Amount	Percentage (%)
lowland		
long, dark brown	4,691	80.53
long, brown	325	5.58
Medium, dark brown	336	5.77
Medium, brown	31	0.53
Short, dark brown	426	7.31
Short, brown	16	0.27
Highland		
long, dark brown	9,161	75.91
long, brown	455	3.77
Medium, dark brown	1,013	8.39
Medium, brown	202	1.67
Short, dark brown	994	8.24
Short, brown	244	2.02




Figure 3. The result of electrophoresis with 1% gel agarose gel PCR Thrips in pairs primer ITS 2, row 1 (long *T. parvispinus* , 2 (medium *T. parvispinus* , 3 (short *T. parvispinus*): amplified, and 4 (male individu) : not amplified, row M: marker 100 bp (Sigma).

T. parvispinus which were found with different morphological variations, as identified by the key identification made Mound & Kibby (1998); Moritz *et al.* (2001); Mound (2006) have the same specific characteristics. Specific characteristics that are owned by *T. parvispinus* are long, medium and short, that are, the antenna consists of 7 segments, the second and third segments have fork-shaped sensory organ, there have no ocellar setae 1 and ocellar setae 2 is shorter than ocellar setae 3, at the pronotum there are 2 pairs of long posteroangular setae and 3 pairs of posteromarginal setae, on metanotum there is no sensilla campaniform, on Tergit VIII contained no comb, on Tergit V-VIII are lateral ctenidia and at Tergit VIII ctenidia located behind the spiracles, dark wing color or shaded with pale base and the first vein and second front wing there are complete row of seta.

Electrophoresis used 1% gel agarose gel PCR thrips with primer pairs ITS 2 thrips showed (Fig. 3) the primer pair ITS 2 (Hillis and Dixon 1991) 28 Z 5 'AGACTCCTGGTCCGTGTTTC 3' and P1 5 'ATCACTCGGCTCGTGGATCG 3' was able to identify the species of *T. parvispinus* long good-sized, medium, or short with 150 bp DNA fragment size. This ITS 2 primer determines the site of DNA replication is begun. Each of any genetic material can be replicated in large quantities by determining a pair of primers flanked the sequences DNA desired. The way of PCR working is started from binding primer which has known its sequences. Then the enzyme of polymerase DNA will extend the oligonucleotide (primer). Each of reaction in the PCR is repeated after denaturation step so there was amplification (multiplication) happened exponentially (Stansfield *et al.* 2003).

On these PCR thrips there were three temperature or stages of incubation which repeated 35 times. A replay of these stages is called a cycle. The first stage is called denaturation, where the two strands of the target DNA molecules are separated (denatured) by its heat. DNA at temperature of 95°C can break the hydrogen bonds between the bases and produce two separated DNA strands. The second stage is called annealing (annealing), where the two primers will hybridize to become complementary sequences on single DNA strands. The primers mean here are a single-stranded synthetic of DNA sequences and short (length 20-30 bases). Primers were chosen of that sort in order to one primer is complementary to one end of the desired gene on one strand. Meanwhile, the second primer is complementary to the other end on the other DNA strand. Primer will form a hydrogen bond (stick) with complementary sequences so with the result that form a stable double-stranded molecule. Annealing temperature used in PCR thrips is 60°C. At the third stage, extension or elongation, primer is extended by DNA polymerase at 72°C.

4. Conclusions

T. parvispinus found in highlands and lowlands of Jambi province, Indonesia, were investigated. The same species of

T. parvispinus can have different morphology and other characteristics. The size and color of the body *T. parvispinus* had variations with long, medium and short criteria. The results also showed that the insects having different body sizes belong to the same *T. parvispinus* species.

Acknowledgements

Authors would like to thank Bert Vierbergen in HC Wageningen Laboratory of Entomology department Netherlands who helped in sequencing DNA of *T. parvispinus*.

References

- [1] Berger D, Walters R, Gotthard K. 2008. What limits insect fecundity? Body size-and temperature-dependent egg maturation and oviposition in a butterfly. *Functional Ecology* 22:523-529.
- [2] Hills DM, Dixon T. 1991. Ribosomal DNA: Molecular evolution and phylogenetic interference. *The Quarterly review of biology* 66: 411-453.
- [3] Khan I, Din S, Khalil SK, Rafi MA. 2006. Survey of predatory coccinellids (Coleoptera:Coccinellidae) in the Chitral, District, Pakistan. *Journal of Insect Science* 7(7):13-21
- [4] Lewis T. 1997. *Trips as Crop Pests*. CAB International. International University Press. p. 15-65.
- [5] Marullo R. and L Mound. 2002. Trips and Tospovirus. *Proceedings of the international symposium on Thysanoptera.Australian national insect collection*. Canberra. p.391
- [6] Moritz G, D Morris, and L Mound. 2001. *Trips ID. Pest Trips of the world. An interactive identification and information system*. CSIRO Publishing Canberra.
- [7] Mound L. 2006. *Thysanoptera Biology and Identification*. CSIRO Entomology Canberra Australia
- [8] Mound LA & Kibby G. 1998. *Thysanoptera. An Identification Guide*. CSIRO Entomology, Canberra Australia; Natural History Museum, London, UK. CAB International
- [9] Mound LA, Collins DW. 2000. A Southeast Asian pest species newly recorded from Europe: *Thrips parvispinus* (Thysanoptera:Thripidae), its confused identity and potential quarantine significance. *Eur J Entomol* 97:197-200
- [10] Murai T. and Satoshi Toda. 2002. Variation of Trips tabaci in colour and size. Trips and Tospoviruses. *Proc. Of the 7 th International Symposium on Thysanoptera*.
- [11] Palmer JM, L Mound and GJ du Heaume. 1989. *CIE guides to insect of importance to man 2. Thysanoptera*. C.R. Betts. CAB International Inst. British Museum Natural History p.72
- [12] Prabaningrum L, dan Yayuk R, Suhardjono. 2007. Identifikasi spesies trips (Thysanoptera) pada tanaman paprika (*Capsicum annum* var. Grossum) di Kabupaten Jawa Barat. *Jurnal Hortikultura* 17 (3) p. 270-276

- [13] Sastrosiswojo S. 1991. Trips on vegetables in Indonesia. In: N.S. Talekar. Trips in Southeast Asia. *Proc. Regional Consultation Workshop Bangkok, Thailand*, 13 March 1991. AVRDC, Taiwan, ROC. p 12-17.
- [14] Stansfield W, Colome JS, Cano RJ. 2003. Schaums's Easy Outlines Molecular and Cell Biology. McGraw-Hills.
- [15] Talekar NS. 1991. Trips in Southeast Asia. *Proc. Regional Consultation Workshop Bangkok, Thailand*, 13 March 1991. AVRDC, Taiwan, ROC. p 61-67.
- [16] Ullman DE, German TL, Sherwood JL & Westcot DM. 1995. Trips transmission of tospoviruses: Future possibilities for management. In B.L. Parker, M. Skinner and T. Lewis. *Trips Biology and Management*. p. 3-16. NATO ASI Series. Series A: Life Sciences Vol 276. Plenum Press, New York.
- [17] Vos JGM, S Sastrosiswojo, TS Uhan, dan Setiawati. 1991. Trips on hot pepper in Java, Indonesia. In: N.S. Talekar. *Trips in Southeast Asia*. P.18-28. *Proc. Regional consultation workshop*. Bangkok, Thailand, 13 March 1991. AVRDC, Taiwan, ROC.
- [18] Vos JGM. 1994. *Integrated Crop Management of Hot Pepper (Capsicum spp.) In Tropical Lowlands*. Wageningen Agricultural University, The Netherlands. p 188.