



Short Communication: Medication Adherence: Most Important but Mostly Avoided

Abdul Kader Mohiuddin

Faculty of Pharmacy, Dhaka University, Dhaka, Bangladesh

Email address:

trymohi@yahoo.co.in

To cite this article:

Abdul Kader Mohiuddin. Short Communication: Medication Adherence: Most Important but Mostly Avoided. *American Journal of Nursing and Health Sciences*. Vol. 4, No. 1, 2023, pp. 6-11. doi: 10.11648/j.ajnhs.20230401.12

Received: November 16, 2022; **Accepted:** December 9, 2022; **Published:** March 3, 2023

Abstract: The indices of patients' health outcomes have historically included recurrence of symptoms, number of emergency visits, hospitalization and re-admission rates, morbidity, and mortality. Even patients from developed nations have trouble staying on top of their drug compliance. When it comes to improper medicine use, there is an odd parallel between underdeveloped, emerging nations and the so-called developed world in the West. The key factor influencing whether patients stick to their treatment plan is their understanding and perception of the disease. Medication adherence was already a difficult and complex health behavior prior to the COVID-19 pandemic and the disruptions it brought. The promptness of disease diagnosis and management, the cost of treatment, the availability of health insurance, and medication adherence are all factors that providers, who play a significant role in the healthcare system, can affect. Beyond the availability and accessibility of healthcare, the ability of patients to follow doctors' treatment advice greatly contributes to patient outcomes. Unfortunately, there has been a lot of medication non-adherence, which has led to worsened health conditions, higher healthcare costs, and higher healthcare spending. Care providers in clinic and pharmacy settings can set up routines to support adherence to medication and educate and encourage patients about services like distance-accessible technologies and online ordering of medications.

Keywords: Patient Compliance, Healthcare Denials, Medication Adherence, Elderly Patient Care, Treatment Failure

1. Introduction

According to the WHO, noncompliance with treatment regimens causes major problems in patients, particularly those with chronic illnesses. "Right administration" depends on at least 5 right factors--right patient, right drug, right time, right dose and right route [1]. "Medicines simply will not work if you don't take it right"—This simple fact is not understood by most people around the world, and as a result, more than half of chronic disease patients in the developed world do not take their medicine correctly, according to WHO [2]. Patients suffering from chronic diseases may have a particularly difficult time adhering because their medications must frequently be taken for an extended period of time, sometimes for the rest of their lives. Patients may struggle to stick to treatment regimens for a variety of reasons, and the CDC estimates that medication non-adherence accounts for 30 to 50% of chronic disease treatment failures. Poor adherence can lead to treatment failure, worsening symptoms, and health deterioration [3].

2. Non-Adherence in the So-Called Developed Countries

In the United Kingdom, up to 50% of medicines are not taken as prescribed, and 60% of NHS patients were unable to receive the appropriate treatment within 18 weeks [4-6]. In patients with chronic diseases, noncompliance with medications leads to poorer health outcomes, higher healthcare costs, increased hospitalizations, and even higher mortality rates [7]. Medication non-adherence alone accounts for at least 10% of hospitalizations in the United States, 250,000 hospitalizations in Australia, and 1.1 million hospital days in France [8-10]; causes \$300 billion in annual medical costs in the United States and \$125 billion in the European Union; and causes more than 1,25,000 premature deaths in the United States and 2,00,000 deaths in the European Union [8, 11, 12]. Furthermore, two-thirds of medication-related hospitalizations in Australia are potentially avoidable [9]. According to a recent Canadian study, 30% of patients stop taking their

medicine before it is recommended, and 25% do not fill their prescription or take less than prescribed [13]. Medication non-adherence attributed to \$679-\$898 more preventable spending among patients who had at least one preventable encounter [14]. However, pharmaceutical companies worldwide lost \$637 billion in potential sales due to non-adherence, with \$250 billion lost in the United States alone last year [15].

3. Misuse of Antibiotics

More than half of all antibiotics sold in the world are sold without a prescription, and the CDC reports that 30-50% of antibiotics prescribed in hospitals are inappropriate or unnecessary [16, 17]. According to a recent Lancet study funded by the Bill and Melinda Gates Foundation and the Wellcome Trust, nearly 5 million deaths worldwide in 2019 were caused by bacterial resistance, which is expected to double by 2050 [18]. In South Asia, nearly 70% of hospitalized patients received one or more antibiotics, whereas 100% of ICU patients received antibiotics [19, 20]. However, 70% to 80% of COVID-19 patients received various antibiotics for COVID-19 treatment [21-23]. The antibiotics most commonly prescribed were azithromycin, ceftriaxone, amoxicillin, metronidazole, and amoxicillin-clavulanic acid [24]. In addition, it has been reported that about 90% of patients with COVID-19 are being unnecessarily treated with antibiotics and close to 100% of these prescriptions were empiric [25].

4. Abuse of NSAIDs in Patients with COVID-19, Dengue, and Chikungunya

NSAIDs are responsible for at least 650,000 hospitalizations, 165,000 deaths, and 30% of ADR-related hospital admissions worldwide each year [26, 27]. Overuse of this class of drugs can result in kidney damage, and their side effects can be three to four times more severe in patients with kidney disease [28]. Many studies have found that these drugs are widely abused in Dengue, Chikungunya, and COVID-19 patients. It is even more important to keep the body hydrated than to reduce the fever with pain relievers, especially in Dengue or COVID-19 patients. Excessive use of Paracetamol syrup or suppositories in children can cause gastric irritation, which can lead to vomiting and hospitalization. With a few exceptions, most hospitalizations or ICU admissions among those patients could be avoided simply by halting dehydration at residence with saline and fruit juice or simply by drinking more water [29].

5. A New Era of Uncontrolled Use of Prescription Only and Recreational Drugs

Sleep disturbances are reported by approximately 40% of COVID-19 patients. Benzodiazepines increase the risk of

delirium in COVID-19 patients, depress the system in patients with compromised breathing functions, and are contraindicated with some anti-viral medications [30, 31]. Surprisingly, benzodiazepine dispensing increased dramatically in Canada between 2020 and 2021, while abuse of similar drugs more than doubled in Italy [32]. According to the American Journal of Public Health [33], approximately 300 metric tons of morphine-type analgesics are used worldwide each year, with less than 1% distributed to low- and middle-income countries. As a result, the developed world retains their misuse and associated side effects. Prior to the US midterm elections, an announcement from authorities on "simple possession of cannabis" to thousands of convicted citizens exploded recreational drug abuse in both the US and the EU [34, 35].

6. Negative Attitude Towards COVID-19 Vaccine

A cross-sectional study of 259 school leaders in Hong Kong carried out during the COVID-19 pandemic between April 2021 and February 2022 shows that more than 50% of participants had limited health literacy, which was strongly associated with a negative attitude towards vaccination, confusion about COVID-19-related information and secondary symptoms [36]. Earlier, a US-based study in 2020 concluded that two-thirds of the Americans will not get the COVID-19 vaccine when it is first available, while 25% report that they do not have any intention to get vaccinated at any time [37]. In India, vaccine hesitancy was high in Tamil Nadu, more than 40% and willingness for vaccine uptake was found to be close to 90% in Kerala [38, 39]. Another vaccine hesitancy survey by University College London, UK finds mistrust among 16% respondents, and 23% were confused [40].

7. Medical Cost and Low-Health-Literacy: The Two Major Barriers of Adherence among Diabetes Patients

A strange similarity can be found in under-developed, developing countries and the so-called developed world in the West or the Middle-East when it comes to not taking medicine properly. According to a WHO report, only half of patients in developed countries adhere to treatment guidelines for chronic diseases, which is much less in developing countries [41]. Several studies among diabetic patients in South Asian countries have shown that nearly half of patients do not adhere to their prescribed medication and are at risk of acute and long-term complications, resulting in increased hospitalization rates and medical costs [42, 43]. "Medical costs are barriers to adherence to proper clinical guidelines for chronic diseases in poor countries"-- although discussed in many forums but forgetfulness, confusion about the duration required for medication use and mistrust about

the overall efficacy of medication are among the reasons for non-adherence to diabetes management protocols in Middle Eastern countries [44]. Health literacy and medication adherence are strongly associated. Poor glycemic control due to low-health-literacy among diabetes patients reported to both South-East Asian and Middle Eastern countries [45-51].

8. Humanitarian Crisis: Poor BP Control Among Cardiac Patients

A recent study by the American Heart Association revealed that patients with high blood pressure do not follow treatment guidelines because of--(a) suboptimal dosing or prescribing the wrong medication (b) lack of insurance or lack of health care access and (c) patient failure to comply prescribed medication or other lifestyle guidelines [52]. Among hypertensive patients, less than 50% have persistent control over BP, even though more patients have received treatment over time. Furthermore, inadequate BP control was reported among those with elevated total cholesterol, LDL, and uric acid levels in both high, low and middle income countries [53]. Humanitarian crisis is associated with increased short-term and long-term cardiac morbidity and mortality and increases in BP [54]. For example, hypertensive patients with diabetes mellitus were twice as likely to exhibit poor BP control, found in war-torn Palestine [55]. Also, a US-based survey on re-settled Rohingya refugees from Myanmar shows a higher trend of chronic diseases like diabetes, hypertension and obesity [56].

9. Superstitions: An Elephant in the Room

Epilepsy and schizophrenia still seen in most countries

of the world as an evil spirit --although two-thirds of patients can become seizure-free with adequate treatment, poor adherence to proper guidelines is a major problem for effective recovery [57, 58]. In a study conducted in India, 60% of the patients believed in luck and superstition with regard to illnesses [59]. Superstitions also reported in close to 40% men and 70% women in Northern Germany [60]. In Africa, 70% of people turn to indigenous treatments such as charms and witchery to treat their illness [61]. Surprisingly, more than 40% of Americans believe in spiritual treatments and researchers found that 73% of addiction treatment programs in the USA include a spirituality-based element [62, 63]. Phobia was the cause of insulin refusal among 60% diabetic patients, despite physician recommendations—found in a study conducted in South Iran [64].

10. Pediatric and Geriatric Complications to Non-Adherence

Due to multiple physical complications and additional medication burden, three-quarters of geriatric persons worldwide are unable to adhere to appropriate long-term treatment regimens [65]. Patients over the age of 65 who take at least five medications are at an increased risk of mild cognitive impairment, memory loss, falls, frailty, impairment, and death, while ADRs are estimated to account for 5% to 28% of acute geriatric medical admissions [66, 67]. For children, common non-adherences are related to family routines, child-raising issues, and to social issues such as poverty. Long-term disease conditions like asthma, cystic fibrosis, HIV, diabetes, inflammatory bowel disease and juvenile arthritis—are attributable to around 60% of non-adherence among children [68-70].

Table 1. Several identified reasons for non-adherence to treatment guidelines for chronic diseases [7, 71-73].

No.	Status	Factors
1	Patient's socio-economic status	Low health literacy, lack of family or social support network, unstable living or homelessness, financial insecurity
2	Treatment-related	Complexity and duration of treatment procedures, frequent changes in medication regimen, lack of immediate results, real or perceived unpleasant side effects, interference with lifestyle
3	Health system-related	High treatment costs, limited health system for patient education and follow-up, doctor-patient relationship, patient trust in health care, long waits, lack of patient information materials
4	Patient-related	Visual-hearing and cognitive impairment, mobility and dexterity, psychological and behavioral factors, perceived risk of disease susceptibility, superstitions and stigmatization by disease, etc

11. Conclusion

Finally, it can be stated that patients' knowledge and interpretation of the disease are the primary factors influencing their adherence to the treatment regimen. Health-care providers should explore more effective health-education methods for identifying patients' attitudes toward disease, medicine trust, psychological stressors, and increasing adherence to medication.

Abbreviations

Adverse drug Reactions (ADRs)
Non-steroidal anti-inflammatory drugs (NSAIDs)
The Centers for Disease Control and Prevention (CDC)
World Health Organization (WHO)

Conflict of Interest

The author declares that he has no competing interests.

Acknowledgements

I'm thankful to Khulud Alosaimi, Department of Pharmacology and Toxicology, Faculty of Medicine, Umm Al-Qura University, Makkah for his valuable time to audit my paper and for his thoughtful suggestions. I'm also grateful to seminar library of Faculty of Pharmacy, University of Dhaka and BANSDOC Library, Bangladesh for providing me books, journal and newsletters.

References

- [1] Grissinger M. (2010). The Five Rights: A Destination Without a Map. *Pharmacy and Therapeutics*, 35 (10), 542.
- [2] Brown MT, Bussell JK. Medication adherence: WHO cares? *Mayo Clin Proc.* 2011 Apr; 86 (4): 304-14. doi: 10.4065/mcp.2010.0575.
- [3] Center for Drug Evaluation and Research. (n.d.). Why you need to take your medications as prescribed or instructed. U.S. Food and Drug Administration. Retrieved November 5, 2022, from <https://www.fda.gov/drugs/special-features/why-you-need-to-take-your-medications-prescribed-or-instructed>
- [4] Barnett NL Medication adherence: where are we now? A UK perspective *European Journal of Hospital Pharmacy: Science and Practice* 2014; 21: 181-184. <https://doi.org/10.1136/ejpharm-2013-000373>
- [5] Campbell, Denis, and Pamela Duncan. "Record 6.8m People Waiting for Hospital Treatment in England." *The Guardian*, 8 Sept. 2022, www.theguardian.com/society/2022/sep/08/waiting-lists-for-routine-hospital-treatment-in-england-break-record.
- [6] Andrews, Luke. "60% of Patients Waiting 18 Wks for Treatment at Worst-performing Trust." *Mail Online*, 7 Mar. 2022, www.dailymail.co.uk/news/article-10577387/60-NHS-patients-waited-18-weeks-treatment-worst-performing-hospital-trust.html.
- [7] Mohiuddin, A. K. "Chapter 14. Patient Compliance". *The Role of the Pharmacist in Patient Care: Achieving High Quality, Cost-Effective and Accessible Healthcare Through a Team-Based, Patient-Centered Approach*, Universal-Publishers, 2020, pp. 250-270. <https://www.universal-publishers.com/book.php?method=ISBN&book=1627343083>
- [8] Cutler RL, Torres-Robles A, Wiecek E, Drake B, Van der Linden N, Benrimoj SI, Garcia-Cardenas V. Pharmacist-led medication non-adherence intervention: reducing the economic burden placed on the Australian health care system. *Patient Prefer Adherence.* 2019; 13: 853-862 <https://doi.org/10.2147/PPA.S191482>
- [9] Lim, Renly, et al. "The Extent of Medication-Related Hospital Admissions in Australia: A Review from 1988 to 2021." *Drug Safety*, vol. 45, no. 3, 28 Jan. 2022, pp. 249–257. <https://doi.org/10.1007/s40264-021-01144-1>.
- [10] "Medication Nonadherence: Medicine's Weakest Link." *Wolters Kluwer, Health/Experts Insight*, 13 Feb. 2020, <https://www.wolterskluwer.com/en/expert-insights/medication-nonadherence-medicines-weakest-link>.
- [11] Kardas, Przemysław et al. "Reimbursed medication adherence enhancing interventions in 12 european countries: Current state of the art and future challenges." *Frontiers in pharmacology* vol. 13 944829. 11 Aug. 2022, doi: 10.3389/fphar.2022.944829.
- [12] van Boven JF, Tsiligianni I, Potočnjak I, Mihajlović J, Dima AL, Nabergoj Makovec U, Ágh T, Kardas P, Ghiciuc CM, Petrova G, Bitterman N, Kamberi F, Culig J, Wettermark B. European Network to Advance Best Practices and Technology on Medication Adherence: Mission Statement. *Front Pharmacol.* 2021 Oct 11; 12: 748702. doi: 10.3389/fphar.2021.748702.
- [13] Bonsu KO, Young S, Lee T, Nguyen H, Chitsike RS. Adherence to Antithrombotic Therapy for Patients Attending a Multidisciplinary Thrombosis Service in Canada - A Cross-Sectional Survey. *Patient Prefer Adherence.* 2022 Jul 26; 16: 1771-1780. doi: 10.2147/PPA.S367105.
- [14] Zhang Y, Flory JH, Bao Y. Chronic Medication Nonadherence and Potentially Preventable Healthcare Utilization and Spending Among Medicare Patients. *J Gen Intern Med.* 2022 Nov; 37 (14): 3645-3652. doi: 10.1007/s11606-021-07334-y.
- [15] Bulik, Beth Snyder. "Nonadherence Costs Pharma \$600B-plus in Annual Sales: Study." *Fierce Pharma*, 22 Nov. 2016, <https://www.fiercepharma.com/marketing/non-adherence-costs-healthcare-system-patient-outcomes-and-pharma-bottom-line>.
- [16] Bahta M, Tesfamariam S, Weldemariam DG, Yemane H, Tesfamariam EH, Alem T, Russom M. Dispensing of antibiotics without prescription and associated factors in drug retail outlets of Eritrea: A simulated client method. *PLoS One.* 2020 Jan 24; 15 (1): e0228013. doi: 10.1371/journal.pone.0228013.
- [17] "Improve Antibiotic Use." *Centers for Disease Control and Prevention*, Centers for Disease Control and Prevention, 9 Mar. 2022, <https://www.cdc.gov/sixteenthai/index.htm>.
- [18] Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet.* 2022 Feb 12; 399 (10325): 629-655. doi: 10.1016/S0140-6736(21)02724-0.
- [19] Rawson TM, Moore LSP, Zhu N, et al. Bacterial and fungal coinfection in individuals with coronavirus: a rapid review to support COVID-19 antimicrobial prescribing. *Clin Infect Dis.* 2020; 71: 2459-2468.
- [20] Clancy CJ, Nguyen MH. Coronavirus disease 2019, superinfections, and antimicrobial development: what can we expect? *Clin Infect Dis.* 2020; 71: 2736-2743.
- [21] Daria, Sohel, and Md Rabiul Islam. "Indiscriminate Use of Antibiotics for COVID-19 Treatment in South Asian Countries is a Threat for Future Pandemics Due to Antibiotic Resistance." *Clinical pathology (Thousand Oaks, Ventura County, Calif.)* vol. 15 2632010X221099889. 18 May. 2022, doi: 10.1177/2632010X221099889.
- [22] Langford B. J., So M., Raybardhan S., Leung V., Soucy J. R., Westwood D., Daneman N., MacFadden D. R. Antibiotic prescribing in patients with COVID-19: Rapid review and meta-analysis. *Clin. Microbiol. Infect.* 2021; 27: 520–531. doi: 10.1016/j.cmi.2020.12.018.
- [23] Cong W., Poudel A. N., Alhusein N., Wang H., Yao G., Lambert H. Antimicrobial Use in COVID-19 Patients in the First Phase of the SARS-CoV-2 Pandemic: A Scoping Review. *Antibiotics.* 2021; 10: 745. doi: 10.3390/antibiotics10060745.

- [24] Kamara, Ibrahim Franklyn et al. "Antibiotic Use in Suspected and Confirmed COVID-19 Patients Admitted to Health Facilities in Sierra Leone in 2020-2021: Practice Does Not Follow Policy." *International journal of environmental research and public health* vol. 19, 7 4005. 28 Mar. 2022, doi: 10.3390/ijerph19074005.
- [25] Usman M, Farooq M, Hanna K. Environmental side effects of the injudicious use of antimicrobials in the era of COVID-19. *Sci Total Environ.* 2020; 745: 141053. doi: 10.1016/j.scitotenv.2020.141053.
- [26] Kasciūškevičiūtė S, Gumbrevičius G, Vendzelytė A, Ščiupokas A, Petrikonis K, Kaduševičius E. Impact of the World Health Organization Pain Treatment Guidelines and the European Medicines Agency Safety Recommendations on Nonsteroidal Anti-Inflammatory Drug Use in Lithuania: An Observational Study. *Medicina (Kaunas)*. 2018 May 11; 54 (2): 30. doi: 10.3390/medicina54020030.
- [27] Davis, Abigail, and John Robson. "The Dangers of NSAIDs: Look Both Ways." *British Journal of General Practice*, vol. 66, no. 645, 2016, pp. 172–173. <https://doi.org/10.3399/bjgp16x684433>.
- [28] Lucas GNC, Leitão ACC, Alencar RL, Xavier RMF, Daher EF, Silva Junior GBD. Pathophysiological aspects of nephropathy caused by non-steroidal anti-inflammatory drugs. *J Bras Nefrol.* 2019 Jan-Mar; 41 (1): 124-130. doi: 10.1590/2175-8239-JBN-2018-0107.
- [29] Mohiuddin, Abdul Kader. "Taking Medicine in the Right Way: Most Important but Most Neglected." *Cases*, vol. 1, no. 1, 13 Nov. 2022, pp. 1–3., <https://doi.org/https://doi.org/10.35702/cases.10003>. <https://www.jcases.org/articles/taking-medicine-in-the-right-way-most-important-but-most-neglected.pdf>
- [30] Jahrami H, BaHammam AS, Bragazzi NL, Saif Z, Faris M, Vitiello MV. Sleep problems during the COVID-19 pandemic by population: a systematic review and meta-analysis. *J Clin Sleep Med.* 2021 Feb 1; 17 (2): 299-313. doi: 10.5664/jcsm.8930.
- [31] Ostuzzi G, Papola D, Gastaldon C, Schoretsanitis G, Bertolini F, Amaddeo F, Cuomo A, Emsley R, Fagiolini A, Imperadore G, Kishimoto T, Michencigh G, Nosé M, Purgato M, Dursun S, Stubbs B, Taylor D, Thornicroft G, Ward PB, Hiemke C, Correll CU, Barbui C. Safety of psychotropic medications in people with COVID-19: evidence review and practical recommendations. *BMC Med.* 2020 Jul 15; 18 (1): 215. doi: 10.1186/s12916-020-01685-9.
- [32] Sarangi, Ashish et al. "Benzodiazepine Misuse: An Epidemic Within a Pandemic." *Cureus* vol. 13, 6 e15816. 21 Jun. 2021, doi: 10.7759/cureus.15816.
- [33] Bhadelia A, De Lima L, Arreola-Ornelas H, Kwete XJ, Rodriguez NM, Knaul FM. Solving the Global Crisis in Access to Pain Relief: Lessons From Country Actions. *Am J Public Health.* 2019 Jan; 109 (1): 58-60. doi: 10.2105/AJPH.2018.304769.
- [34] Lopez, German. "Marijuana Majority/ Americans Support Marijuana Legalization, but Many of Their Political Leaders Do Not." *The New York Times*, 23 Nov. 2022, <https://www.nytimes.com/2022/11/23/briefing/legal-weed-marijuana.html>. Accessed 30 Nov. 2022.
- [35] Oltermann, Philip. "Germany Announces Plan to Legalise Cannabis for Recreational Use." *The Guardian*, 26 Oct. 2022, <https://www.theguardian.com/world/2022/oct/26/germany-to-legalise-cannabis-for-recreational-use>. Accessed 30 Nov. 2022.
- [36] Shahid, Rabia et al. "Impact of low health literacy on patients' health outcomes: a multicenter cohort study." *BMC health services research* vol. 22, 1 1148. 12 Sep. 2022, doi: 10.1186/s12913-022-08527-9.
- [37] Alam, Md Moddassir et al. "Public Attitude Towards COVID-19 Vaccination: Validation of COVID-Vaccination Attitude Scale (C-VAS)." *Journal of multidisciplinary healthcare* vol. 15 941-954. 29 Apr. 2022, doi: 10.2147/JMDH.S353594.
- [38] Danabal, K. G. M., Magesh, S. S., Saravanan, S. et al. Attitude towards COVID-19 vaccines and vaccine hesitancy in urban and rural communities in Tamil Nadu, India – a community based survey. *BMC Health Serv Res* 21, 994 (2021). <https://doi.org/10.1186/s12913-021-07037-4>
- [39] Leelavathy, Manju et al. "Attitude towards COVID-19 vaccination among the public in Kerala: A cross sectional study." *Journal of family medicine and primary care* vol. 10, 11 (2021): 4147-4152. doi: 10.4103/jfmpe.jfmpe_583_21.
- [40] Paul, Elise et al. "Attitudes towards vaccines and intention to vaccinate against COVID-19: Implications for public health communications." *The Lancet regional health. Europe* vol. 1 (2021): 100012. doi: 10.1016/j.lanepe.2020.100012.
- [41] Chauke GD, Nakwafila O, Chibi B, Sartorius B, Mashamba-Thompson T. Factors influencing poor medication adherence amongst patients with chronic disease in low-and-middle-income countries: A systematic scoping review. *Heliyon.* 2022 Jun 15; 8 (6): e09716. doi: 10.1016/j.heliyon.2022.e09716.
- [42] Chong E, Wang H, King-Shier KM, Quan H, Rabi DM, Khan NA. Prescribing patterns and adherence to medication among South-Asian, Chinese and white people with Type 2 diabetes mellitus: a population-based cohort study. *Diabet Med.* 2014. August 11 10.1111/dme.12559.
- [43] Sohal T, Sohal P, King-Shier KM, Khan NA. Barriers and Facilitators for Type-2 Diabetes Management in South Asians: A Systematic Review. *PLoS One.* 2015 Sep 18; 10 (9): e0136202. doi: 10.1371/journal.pone.0136202.
- [44] Alsairafi ZK, Taylor KM, Smith FJ, Alattar AT. Patients' management of type 2 diabetes in Middle Eastern countries: review of studies. *Patient Prefer Adherence.* 2016 Jun 10; 10: 1051-62. doi: 10.2147/PPA.S104335.
- [45] Almigbal, Turkey H et al. "Association of health literacy and self-management practices and psychological factor among patients with type 2 diabetes mellitus in Saudi Arabia." *Saudi medical journal* vol. 40, 11 (2019): 1158-1166. doi: 10.15537/smj.2019.11.24585.
- [46] Nair, Satish Chandrasekhar et al. "Health literacy in a high income Arab country: A nation-wide cross-sectional survey study." *PloS one* vol. 17, 10 e0275579. 5 Oct. 2022, doi: 10.1371/journal.pone.0275579.
- [47] Hashim, Saman Agad et al. "Association of Health Literacy and Nutritional Status Assessment with Glycemic Control in Adults with Type 2 Diabetes Mellitus." *Nutrients* vol. 12, 10 3152. 15 Oct. 2020, doi: 10.3390/nu12103152.
- [48] Hussein, Shaimaa H et al. "Association of health literacy and other risk factors with glycemic control among patients with type 2 diabetes in Kuwait: A cross-sectional study." *Primary care diabetes* vol. 15, 3 (2021): 571-577. doi: 10.1016/j.pcd.2021.01.011.

- [49] Khatiwada, Bhushan, et al. "Prevalence of and Factors Associated with Health Literacy among People with Noncommunicable Diseases (Ncds) in South Asian Countries: A Systematic Review." *Clinical Epidemiology and Global Health*, vol. 18, 3 Nov. 2022, p. 101174. <https://doi.org/10.1016/j.cegh.2022.101174>.
- [50] Rajah R., Hassali M. A. A., Murugiah M. K. A systematic review of the prevalence of limited health literacy in Southeast Asian countries. *Public Health*. 2019; 167: 8–15. doi: 10.1016/j.puhe.2018.09.028.
- [51] Saleh, Ariyanti, et al. "The Relationships among Self-Efficacy, Health Literacy, Self-Care and Glycemic Control in Older People with Type 2 Diabetes Mellitus." *Working with Older People*, vol. 25, no. 2, 4 May 2021, pp. 164–169. <https://doi.org/10.1108/wwop-08-2020-0044>.
- [52] Choudhry NK, Kronish IM, Vongpatanasin W, Ferdinand KC, Pavlik VN, Egan BM, Schoenthaler A, Houston Miller N, Hyman DJ; American Heart Association Council on Hypertension; Council on Cardiovascular and Stroke Nursing; and Council on Clinical Cardiology. Medication Adherence and Blood Pressure Control: A Scientific Statement From the American Heart Association. *Hypertension*. 2022 Jan; 79 (1): e1-e14. doi: 10.1161/HYP.0000000000000203.
- [53] Elnaem, Mohamed Hassan et al. "Disparities in Prevalence and Barriers to Hypertension Control: A Systematic Review." *International journal of environmental research and public health* vol. 19, 21 14571. 6 Nov. 2022, doi: 10.3390/ijerph192114571.
- [54] Keasley, James et al. "A systematic review of the burden of hypertension, access to services and patient views of hypertension in humanitarian crisis settings." *BMJ global health* vol. 5, 11 (2020): e002440. doi: 10.1136/bmjgh-2020-002440.
- [55] Alawneh, Issa S et al. "The Prevalence of Uncontrolled Hypertension among Patients Taking Antihypertensive Medications and the Associated Risk Factors in North Palestine: A Cross-Sectional Study." *Advances in medicine* vol. 2022 5319756. 25 Aug. 2022, doi: 10.1155/2022/5319756.
- [56] Rahman, Ayesha, et al. "Non-Communicable Diseases Risk Factors among the Forcefully Displaced Rohingya Population in Bangladesh." *PLOS Global Public Health*, vol. 2, no. 9, 27 Sept. 2022, <https://doi.org/10.1371/journal.pgph.0000930>.
- [57] Lossius, M. I., Alfstad, K. Å., Aaberg, K. M., & Nakken, K. O. (2017). Seponering av Antiepileptika Ved anfallsfrihet – når og hvordan? *Tidsskrift for Den Norske Legeforening*, 137 (6), 451–454. <https://doi.org/10.4045/tidsskr.16.0957>
- [58] Yang, D. *Zhonghua shen jing jing shen ke za zhi = Chinese journal of neurology and psychiatry* vol. 25, 4 (1992): 215-8, 253.
- [59] Banerjee S, Varma RP. Factors affecting non-adherence among patients diagnosed with unipolar depression in a psychiatric department of a tertiary hospital in Kolkata, India. *Depress Res Treat*. 2013; 2013: 809542.
- [60] Graeupner D, Coman A. The dark side of meaning-making: How social exclusion leads to superstitious thinking. *Journal of Experimental Social Psychology*. 2017; 69: 218–22.
- [61] Puckree T, Mkhize M, Mgobhozi Z, Lin J. African traditional healers: What health care professionals need to know. *Int J Rehabil Res*. 2002; 25: 247–51.
- [62] Taher, Mohammad et al. "Superstition in health beliefs: Concept exploration and development." *Journal of family medicine and primary care* vol. 9, 3 1325-1330. 26 Mar. 2020, doi: 10.4103/jfmpc.jfmpc_871_19.
- [63] Grim, Brian J, and Melissa E Grim. "Belief, Behavior, and Belonging: How Faith is Indispensable in Preventing and Recovering from Substance Abuse." *Journal of religion and health* vol. 58, 5 (2019): 1713-1750. doi: 10.1007/s10943-019-00876-w.
- [64] Mirahmadizadeh, Alireza et al. "Factors Affecting Insulin Compliance in Patients with Type 2 Diabetes in South Iran, 2017: We Are Faced with Insulin Phobia." *Iranian journal of medical sciences* vol. 44, 3 (2019): 204-213.
- [65] Félix IB, Henriques A. Medication adherence and related determinants in older people with multimorbidity: a cross sectional study. *Nurs Forum*. 2021; 56: 834–843. doi: 10.1111/nuf.12619.
- [66] Chippa, Venu, and Kamalika Roy. "Geriatric Cognitive Decline and Polypharmacy." *National Library of Medicine, StatPearls Publishing*, Jan. 2022, <https://www.ncbi.nlm.nih.gov/books/NBK574575/>.
- [67] Varghese D, Ishida C, Haseer Koya H. Polypharmacy. *National Library of Medicine, StatPearls Publishing*, Jan. 2022, <https://www.ncbi.nlm.nih.gov/books/NBK532953/>.
- [68] Santer, Miriam et al. "Treatment non-adherence in pediatric long-term medical conditions: systematic review and synthesis of qualitative studies of caregivers' views." *BMC pediatrics* vol. 14 63. 4 Mar. 2014, doi: 10.1186/1471-2431-14-63.
- [69] Al-Hassany, Linda et al. "Assessing methods of measuring medication adherence in chronically ill children-a narrative review." *Patient preference and adherence* vol. 13 1175-1189. 22 Jul. 2019, doi: 10.2147/PPA.S200058.
- [70] Wu, Y Y et al. *Zhonghua er ke za zhi = Chinese journal of pediatrics* vol. 60, 11 (2022): 1191-1195. doi: 10.3760/cma.j.cn112140-20220110-00036.
- [71] Jin J, Sklar GE, Min Sen Oh V, Chuen Li S. Factors affecting therapeutic compliance: A review from the patient's perspective. *Ther Clin Risk Manag*. 2008 Feb; 4 (1): 269-86. doi: 10.2147/tcrm.s1458.
- [72] Wilder, Marcee E et al. "The Impact of Social Determinants of Health on Medication Adherence: a Systematic Review and Meta-analysis." *Journal of general internal medicine* vol. 36, 5 (2021): 1359-1370. doi: 10.1007/s11606-020-06447-0.
- [73] Kardas, Przemyslaw et al. "Determinants of patient adherence: a review of systematic reviews." *Frontiers in pharmacology* vol. 4 91. 25 Jul. 2013, doi: 10.3389/fphar.2013.00091.