

**Research paper**

Phenolic Profile and Biological Properties of *Momordica charantia*

Zermina Khalid¹, Syeda Mona Hassan¹, Shahzad Sharif Mughal^{1, *}, Syed Khurram Hassan², Huma Hassan³

¹Department of Chemistry, Lahore Garrison University, Lahore, Pakistan

²Institute of Quality and Technology Management, University of the Punjab, Lahore, Pakistan

³Department of Chemical Engineering, NFC Institute of Engineering and Fertilizer Research, Faisalabad, Pakistan

Email address:

shezi1130@gmail.com (S. S. Mughal)

*Corresponding author

To cite this article:

Zermina Khalid, Syeda Mona Hassan, Shahzad Sharif Mughal, Syed Khurram Hassan, Huma Hassan. Phenolic Profile and Biological Properties of *Momordica charantia*. *Chemical and Biomolecular Engineering*. Vol. 6, No. 1, 2021, pp. 17-29. doi: 10.11648/j.cbe.20210601.13

Received: June 22, 2020; **Accepted:** November 2, 2020; **Published:** March 26, 2021

Abstract: *M. charantia* is an important medicinal plant belongs to family *cucurbitaceae*. It originates from India, Malasiya and is widely spread all over tropical, subtropical and warm temperate regions of the world. This research work has been designed to evaluate the antioxidant, antimicrobial and toxicological potential of *M. charantia*. The antifungal and antioxidant components of *M. charantia* leaves, seeds and peels were extracted by using four solvent systems (80% methanol, 80% ethanol, 100% methanol and 100% ethanol) and leaves presented maximum extract yield (22.7 g/100g DW) in 80% methanolic solvent system. Phytochemical analysis of *M. Charantia* leaves, seeds and peels extracts performed in terms of total phenolic and total flavonoid contents, showed that 80% methanolic leaves extract offered highest total phenolic contents (47.1 mg GAE/g DW), whereas 80% ethanolic leaves gave maximum total flavonoid contents (67.3 mg CE/g DW). The phenolic contents were also analysed by HPLC. Antioxidant activity was determined by DPPH radical scavenging activity and measure of reducing power. Results revealed that 80% methanolic leaves extract showed highest radical scavenging activity and reducing potential. Antimicrobial activity of *M. charantia* leaves, seeds and peels was investigated by Disc Diffusion Method and Minimum Inhibitory Concentration (MIC). Results showed that 80% methanolic extract of leaves exhibited highest antibacterial and antifungal potential against *P. multocida* (30 mm DIZ) and *A. paraciticus* (28 mm DIZ), respectively. Cytotoxicity analysis was performed on BHK-21 cell by adopting the MTT assay. The cytotoxicity activity of the 80% methanolic extract of leaves was evaluated by noticing the cell survival percentage (53.4%). Overall results of the present study showed that 80% methanolic leaves extracts of *M. charantia* possesses very good antioxidant, antimicrobial and cytotoxic properties.

Keywords: *Momordica charantia*, General Description, Medicinal Properties

1. Introduction

Momordica charantia commonly known as bitter melon or bitter gourd is tropical or subtropical climber of the family of *cucurbitaceae* [1, 5]. It is mostly available in China, Malaysia, India and tropical Africa. All parts of the plant, including the fruit, taste is very bitter, as it contains a bitter compound called *momordicin* that is believed to have a stomachic effect [15, 19]. Bitter gourd has been used as folk medicine to cure toothache, diarrhea, furuncle, diabetes, dysmenorrhea, eczema,

emmenagogue, galactagogue, gout, jaundice, kidney (stone), leprosy, leucorrhea, piles, pneumonia, psoriasis, rheumatism and scabies [3]. The Latin name *Momordica* means "to bite," referring to the jagged edges of the leaves, which appear as if they have been bitten [35].

The last few decades several hundred studies that have been carried with *M. charantia*. *M. charantia* possess antidiabetic, antiviral, antitumor, antileukemic, antibacterial, anthelmintic, antimutagenic, anti-mycobacterial, antioxidant, antiulcer, anti-inflammatory, hypocholesterolemic, hypotriglyceridemic,

hypotensive, immunostimulant and insecticidal properties [18]. Traditionally, it is believed to have hypoglycemic effect. *M. charantia* containing food dishes are not popular in the Western world due to its taste [46, 47]. Although the different parts of plant *M. charantia* have been used as food and drug but the fruit is the most important part. Depending on the maturity stages, *M. charantia* seeds (BGS) have been found as a rich source of oil (18.1-37.6%) and protein (28-30%) [6, 69]. Previous investigations have shown that the fruits and leaves of *M. charantia* had rich phenolics and exhibited a high antioxidant activity. Fresh bitter melon is also used as a nourishing food, as it contains: 93.8% water, 0.9% protein, 0.1% lipid, 3.3% dietary fiber, 20 kJ energy per 100 g, 0.6% ash, and a small quantity, 0.05%, of vitamin C [70]. The nutritional value is less due to low levels of proteins and carbohydrates, as compared to other member of *Cucurbitaceae* with best nutritional value, might be due to the phosphorus, iron, and ascorbic acid content [17]. *M. charantia* plant is used as medicine and is a good nutritious source of phytochemicals [38]. The residue of alcoholic ether extract of *M. charantia* leaves, is reported to possess hypoglycemic activity. P-insulin (the pure protein) of *M. charantia* fruits plant part is also tested as in crystalline form [34].

Medicinal value of bitter melon has been attributed to its high antioxidant properties due in part to phenols, flavonoids, isoflavones, terpenes, anthroquinones and glucosinolates [41]. Alteration in the total antioxidant content and activity due to different processing methods is of scientific importance as it has a direct impact on dietary nutrition. Blanching is a treatment of vegetables to inactivate enzymes such as polyphenol oxidase, catalase, peroxidase, lipogenase and chlorophyllase [7, 8]. The antioxidant enzymes (AOE) include superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx) and indirectly glutathione reductase (GR). Their role as protective enzymes are well known and have been extensively investigated both in vivo and in vitro in model systems [54]. Studies have reported that phenolics have potent antioxidant and free radical-scavenging activities. Whole bitter melon (flesh, aril and seeds) has been shown to be a good source of phenolic compounds and one study demonstrated that the flesh, aril and seeds all had very high antioxidant activity [64].

Many dietary phytochemicals, particularly phenolic compounds, have demonstrated antioxidant characteristics in various disease states such as diabetes, liver disease, cardiovascular disease and cancer. Consequently, the commercial development of plants as sources of antioxidants for health and nutritional purposes is of great interest worldwide [30]. Medicinal value of bitter melon has been attributed to its high antioxidant properties due in part to phenols, flavonoids, isoflavones, terpenes, anthroquinones, and glucosinolates [41].

Colon rectal cancer is the second leading cause of cancer related deaths and the third most commonly occurring noncutaneous carcinoma in the United States of America [36]. *M. charantia* is rich in various biologically active chemicals including triterpenes, proteins, and steroids. Triterpenes of *M.*

charantia has the ability to inhibit the enzyme guanylate cyclase that is thought to be linked to the cause of psoriasis [20]. In addition, guanylate cyclase is one of the important enzymes, necessary for the growth of leukemia and other cancer cells. In addition to these biologically active triterpenes, *M. charantia* proteins like momordin, alpha- and beta momorcharin and cucurbitacin B were also tested for possible anticancerous effects [26].

The anti-cancer properties of *M. charantia* are recently elucidated. Many researchers have found that treatment of *M. charantia* related products in a number of cancer cell lines induces cell cycle arrest and apoptosis without affecting normal cell growth [57]. The role of free radicals and active oxygen in treating chronic diseases including cancer, aging and atherosclerosis has been recognized [37]. Most recently, the same research group extended the antitumor potential of *M. charantia* crude extract to prostate cancer based on results from both in vitro and in vivo studies. It was observed that crude *M. charantia* extracts impaired cell cycle progression and inhibited xenograft proliferation [15]. The medicinal values of *M. charantia* have bioactive phytochemical constituents with non nutritive chemicals that produce definite physiological effects on human body and protect them from various diseases. In *M. charantia* primary metabolites are common sugars, proteins and chlorophyll while secondary metabolites are alkaloids, flavonoids, tannins and so on [10]. The aims of this study are to find the phenolic contents present in *M. charantia* its potential antimicrobial and biological effects by using DPPH and MIC methods [11].

2. Material and Method

2.1. Chemicals and Standard Compounds

Butylatedhydroxytoluene (BHT) (99.0%), DPPH, catechin, gallic acid and Folin-Ciocalteu reagents were brought from Sigma Chemicals Co. (St, Louis, MO, USA). All standard antibiotic and culture media were bought from Oxoid Ltd., Immunoaffinity column (AflaTest® WB VICAM, USA) (Hampshire, UK).

2.1.1. Collection of Plant Materials

M. charantia leaves, seeds and peels were obtained from the vicinity of Lahore Garrison University, Lahore, Pakistan.

2.1.2. Preparation of Extract

For extraction of bioactive compounds, four solvent systems (100% methanol, 80% methanol, 100% ethanol and 80% ethanol) were being used. In this regard powdered seeds and leaves (20g) were extracted with 200mL in an orbital shaker for 8 hours at room temperature (Gallenkamp, UK). In order to separate the extract from residue whatman No. 1 filter paper was used. The resulting residues were extracted two times with the same solvent system. The drying of extracts was done at temperature of 45°C and their yield was calculated by weighing extracts. The extracts were kept in a refrigerator at 4°C for further analysis [20]. The ground samples were kept in polythene bags at 4°C until for further analysis.

2.1.3. Determination of Total Phenolic and Flavonoid Contents

By using Folin-Ciocalteu reagent total phenolic contents were determined [71]. The plant crude extract of 50 mg was mixed with 0.5 mL of Folin-Ciocalteu reagent and then 7.5 mL deionized water was added. The mixture was kept at room temperature for 10 min. Then 1.5 mL of 20% sodium carbonate (w/v) was added. At 40°C the mixture was in a water bath for 20 min and then cooled in an ice bath. Absorbance was measured at 755 nm using a spectrophotometer (U-2001, Hitachi Instruments Inc., Tokyo, Japan). Using gallic acid calibration curve the amounts of TP were calculated. The results were expressed as mg GAE/g DW. In thrice time all samples were analysed. The result was measured in average. All the results are reported on dried weight basis. Total flavonoid contents were determined by following the procedure of Dewanto *et al.* (2002). The aqueous extract of one milliliter of containing 0.01 g/mL of dry matter was placed in a 10 mL volumetric flask. Then 5 mL of distilled water was added followed by 0.3 mL of 5% NaNO₂. Then after 5 min, 0.6 mL of 10% AlCl₃ was added. After 5 min, 2 mL of 1 M NaOH was added. Then volume was made up with distilled water. After this the absorbance was measured at 510 nm. TF amounts were expressed as mg Catechin equivalent (CE)/g DW. All samples were analysed thrice and results were averaged [20, 24].

2.2. Analysis of Phenolics by High Performance Liquid Chromatography (HPLC)

2.2.1. HPLC Sample Extraction

With slight modifications Extraction/hydrolysis of phenolics was carried out following the method of [65]. Permanently acidified methanol (25 mL) containing 1% (v/v) HCl and 0.5 mg mL⁻¹ TBHQ was added to each plant material (5 g). HCl (1.2 M, 5 mL) was added and the mixture was stirred at 90°C under reflux for 2 h. The extract was cooled to room temperature. Then centrifuged at 1500g (5000 rpm) for 10 min. Upper layer was taken and sonicated for 5 min, to remove air. Before injecting into HPLC, the final extract was filtered through a 0.45mL (Millipore) filter.

Using high-performance liquid chromatography the phenolic analysis was performed. An HPLC (model LC-10A, Shimadzu, Kyoto, Japan), CTO-10A column oven, equipped with two LC-10 AS pumps, Rheodyne injector, SCL-10A system control unit, SPD-10A UV-vis detector, and data acquisition class LC-10 software was used. A 20µL volume of the filtered sample was injected into an analytical Supelco (Supelco Inc., Supelco Park, Bellefonte, PA, USA) ODS reverse phase (C18) column (250×4.6 mm; 5µm particle size). Two solvent systems, A: contained water and Acetic acid (94:6 v/v) and B: contained 100% acetonitrile, were used. The chromatographic separation was performed by gradient elution of the mobile phase (0-15min= 15%B, 15-30=45%B, 30-45min=100%B) at a flow rate of 1.0 mL min⁻¹ at room temperature. Detection was performed at a wavelength of 280nm. By comparing their retention times identification of phenolics compound was carried out with authentic standards

(Sigma Chemicals Co., St Louis, MO, USA). Quantitative determination was carried out by using calibration curves of the standards [20].

2.2.2. DPPH Radical Scavenging Assay

By using 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical then this way free radical scavenging activity was determined this described by [60]. This solution of DPPH (33mg/L) was prepared in methanol. This type of solution then this way the absorbance was taken at 0 min. Then the solution of extract (250µg/mL) were prepared. Then 5mL of methanolic solution of DPPH was added in 1mL of extract solution. For 30 minutes the mixture was left in the dark. Then absorbance was measured at 517nm using a spectrophotometer. Then methanol used as blank. Free radical scavenging activity was expressed as percentage inhibition and calculated by using the following formula:

$$\text{Inhibition (\%)} = \frac{\text{Control Absorbance} - \text{Sample Absorbance}}{\text{Control Absorbance}} \times 100$$

Control Absorbance

Where control absorbance is absorbance of methanolic solution of DPPH taken at zero minute [60].

2.2.3. Reducing Power Determination

With slight modification then the reducing power of the plant leave extracts was determined according to the procedure described by [12]. Concentrated extract (2.5-10.0 mg) was mixed with sodium phosphate buffer (5.0 mL, 0.2 M, pH 6.6) and potassium ferricyanide (5.0 mL, 1.0%). Then the mixture was incubated at 50°C for 20 min. Then 5 mL of 10% trichloroacetic acid was added. Then the mixture was centrifuged at 980 g for 10 min at 5°C in a refrigerated centrifuge (CHM-17; Kokusan Denki, Tokyo, Japan). The upper layer of the solution (5.0 mL) was decanted and diluted with 5.0 mL of distilled water and ferric chloride (1.0 mL, 0.1%). Then absorbance was checked at 700 nm using spectrophotometer (U-2001, Hitachi Instruments Inc., Tokyo, Japan). All samples were analysed thrice and results averaged [41].

2.2.4. Evaluation of Antimicrobial Potential

The extracts of leaves and seeds were tested individually against a panel of microorganisms which may included five fungal strains (*Aspergillus parasiticus*, *Aspergillus flavus*, *Fusariumoryzae*, *Fusariumtritichum*, *Aspergillus oryzae*) and three bacterial strains (*Escherichia coli*, *Pasturellamultocida* and *Staphylococcus aureus*) obtained from the Fungal Bank, University of Punjab, Lahore. Fungal strains were cultured overnight at 28°C in Potato Dextrose agar (Oxoid Hampshire, UK), however the bacterial strain were cultured at 37°C in nutrient agar (Oxoid Hampshire, UK). The slants of microbial strains were stored at 4°C. Antimicrobial potential of plants extracts were determined by using the disc diffusion and microdilution broth assays [22, 25].

2.3. Disc Diffusion Assay

Antimicrobial activity of leaves, seeds and peels of *M.*

charantia was tested against fungal strains (*Aspergillus flavus*, *Aspergillus paraciticus*, *Fusariumoryzae*, *Fusariumtritichum*, *Aspergillus oryzae*) and bacterial strains (*Escherichia coli*, *Pasturellamultocida* and *Staphylococcus aureus*) by previously adopted method (NCCLS, 2004) with little modifications. The solution of Potato dextrose agar (PDA) was prepared and then autoclaved. Under laminar air flow, then about 20mL PDA solution was transferred in sterilized petri plate. Sterilized discs (6mm) of wicks sheet impregnated with 50 μ L of particular plant extract were placed on the agar plates. To equate the activity with standard antibiotics, as positive reference Fluconazol (30 μ g/disc) (Oxoid) and Rifampicin (30 μ g/disc) (Oxoid) were used as for fungal and bacterial strains respectively. As a negative control disc without samples were used. Standard disc and test discs and were placed in separate petri dishes. The plates were incubated at 28°C for 48h for fungal growth and 37°C for 24h for bacterial growth. Antimicrobial activity was evaluated by measuring the diameter of inhibition zones (mm) by zone reader [44].

2.4. Microdilution Broth Method

The method reported by [37] then this way MIC of plant extracts was evaluated. Briefly, in the first row 100 μ L of plant extract was transferred into 96 well microliter plates. To entirely other wells, 50 μ L of Sabarouraud dextrose broth and nutrient broth was added for fungal and bacterial strains, respectively [42]. Two-fold serial dilutions were performed using a micropipette such that each well had 50 μ L of the test material in serially descending concentrations. Finally, 10 μ L of microbial suspension was added to each well. Each plate had a row of negative control, a row of positive control of Fluconazole and Rifampicin for antifungal and antibacterial activities, respectively. The plates were prepared in triplicate and incubated at 28°C for 48h for fungi and 37°C for 24h for bacteria. By ELISA reader the absorbance was measured at 620 nm. At the lowest concentration then no growth was taken as the MIC value [51, 52, 53].

2.5. Toxicological Analysis

2.5.1. Cytotoxicity Assay

By adopting the MTT assay Cytotoxicity of plant extracts was evaluated then by using baby hamster kidney cells (BHK-21) as described by Freshney and Frame (1982), though 10% DMSO was used as a positive control. Solutions of the tested materials were evaluated for cytotoxic potential. The BHK-21 cells were revived using DMEM (Sigma-Aldrich, Germany) media as described by Freshney (1998). Then transferred into 40 ml cell culture flasks (Karrel Flasks, Corning, USA), which were then incubated for 72 h to get the confluent monolayer of cells. One hundred microliters of cell suspension (105 cells/ml) was dispensed into each well of 96-well plates (Corning, USA) and incubated at 37°C for 72 h. Media on the confluent monolayer of cells was regularly changed. Then 100 μ L of the respective sample concentrations was added in triplicate, which was then incubated at 37°C for

48 h. Finally, the growth medium was removed. Then wells were washed with PBS and replenished with fresh media. One hundred microliters of 0.5% MTT solution was added to each well. Then each plates were incubated for ~4 h. The MTT solution was then removed, and plates were incubated at 37°C for 2 h. Thenthis after adding 5% DMSO to each well. Optical density was measured at 570 nm by an ELISA reader (Type355, Model 2005–05, Thermo, China) [6].

2.5.2. Statistical Analysis

By performing all experiments in triplicate (n=3), the data was described, mean \pm SD. Data analyzed at 5% significant level by using Minitab 2000 Version 13.2 statistical software (Minitab Inc. Pennsylvania, USA). As mean values at 95% confidence interval data of antifungal activity are presented. Significant differences of mean were calculated by using LSD [54, 55, 56].

3. Result and Discussion

3.1. Extracts Yields

The extraction yields from leaves and seeds and peels of using different solvent systems. Comparatively, 80% methanol exhibited higher extraction yields from leaves (22.7%). The extraction ability of different solvent systems for recovering extractable components from leaves followed the order: 80% methanol > 80% ethanol > absolute methanol > absolute ethanol. Yields of extract from seeds followed the order: absolute methanol > 80% ethanol > 80% methanol > absolute ethanol. Yield of extract from peels followed the order: 80% methanol > 80% ethanol > absolute ethanol > absolute methanol [57].

3.2. Total phenolic and Total Flavonoid Contents

The total phenolic content from leaves, seeds and peels of using different solvent systems are presented in Table 1. Relatively, 80% methanol showed higher extraction yields from leaves (47.1%). The extraction capacity of different solvent systems for recovering extractable components from leaves followed order: 80% methanol > 80% ethanol > absolute methanol > absolute ethanol. Yields of extract from seeds followed the order: 80% methanol > 80% ethanol > absolute methanol > absolute ethanol. Yield of extract from peels followed the order: 80% methanol > 80% ethanol > absolute ethanol > absolute methanol [57, 58]. The total flavonoid content from leaves, seeds and peels of using different solvent systems are presented in Table 2. Moderately, 80% methanol exhibited higher extraction yields from leaves (67.3%). The extraction skill of different solvent systems for recovering extractable components from leaves followed the order: 80% ethanol > 80% methanol > absolute ethanol > absolute methanol. Yields of extract from seeds followed the order: 80% methanol > 80% ethanol > absolute methanol > absolute ethanol. Yields of extract from peels followed the order: 80% ethanol > 80% methanol > absolute ethanol > absolute methanol [59].

Table 1. Total phenolic contents (mg GAE/g DW) of *M. charantia* seeds, peels and leaves extracts.

Sr. no	Solvent System	Total phenolic contents (mg GAE/g DW)		
		Leaves	Seeds	Peels
1	80% Methanol	47.1±0.56 ^a	36.2±0.42 ^{ab}	36.1±0.49 ^a
2	80% Ethanol	42.2±0.55 ^{ab}	35.3±0.46 ^{bc}	34.3±0.44 ^b
3	100% Methanol	40.3±0.50 ^b	34.5±0.59 ^{bc}	30.2±0.32 ^d
4	100% Ethanol	35.5±0.41 ^{bc}	27.1±0.45 ^d	32.1±0.45 ^c

Values are mean ± SD of three samples analyzed individually in triplicate at $p < 0.05$. Superscripts alphabets within the column depicted significant difference among different medicinal plants. Subscripts alphabets within the rows depicted significant difference among different plant parts.

Table 2. Total flavonoid contents (mg CE/g DW) of *M. charantia* seeds peels and leaves extracts.

Sr. no	Solvent System	Total flavonoid contents (mg CE/g DW)		
		Leaves	Seeds	Peels
1	80% Methanol	64.1±0.56 ^{ab}	57.4±0.42 ^a	61.2±0.49 ^{ab}
2	80% Ethanol	67.3±0.55 ^a	54.3±0.46 ^b	63.3±0.44 ^a
3	100% Methanol	57.2±0.50 ^c	49.2±0.59 ^{bc}	52.1±0.32 ^d
4	100% Ethanol	62.1±0.41 ^{bc}	45.1±0.45 ^c	57.4±0.45 ^b

Values are mean ± SD of three solvents analyzed individually in triplicate at $p < 0.05$. Superscripts alphabets within the column depicted significant difference among different solvents system. Subscripts alphabets within the rows depicted significant difference among different plant parts.

3.3. Phenolics Detected by HPLC

The HPLC examination of medicinal plant leaves revealed the presence of polyphenolic compounds like gallic acid, chlorogenic acid, caffeic acid, *p*-coumaric acid, *m*-coumaric acid, 4-hydroxy, 3-methoxy benzoic acid, ferulic acid, syringic acid and vanillic acid [50]. Phenolic compounds of *M. charantia* of different plant parts of leaves, seeds and peels are given in table 3. The total phenolic profile of *M. charantia* leaves revealed the presence of phenolic compounds like gallic acid (23.44ppm), chlorogenic acid (66.73 ppm) and *p*-coumaric acid is (53.21ppm). However *m*-coumaric acid (17.21ppm) and 4-hydroxy, 3-methoxy benzoic acid (403.7ppm) were found to be present in *M. charantia* peels. Moreover, syringic acid (56.34 ppm) and vanillic acid (301.9 ppm) were present in seeds of *M. charantia* [60, 61].

3.4. Antioxidant Activity

A large number of medicinal plants have been investigated for their antioxidant properties. The natural antioxidants, either in the form of extracts or their chemical constituents are very important to prevent the destructive actions caused by oxidative stress [53].

3.5. DPPH Radical Scavenging Activity

Antioxidant property of the natural products is due to radical scavenging ability of phytoconstituents such as flavonoids, polyphenols, tannins, and phenolic terpenes [20]. Antioxidant activity of these products can be evaluated by

using DPPH radical scavenging assay which is widely used to test the ability of compounds that act as free radical scavengers or hydrogen donors [6]. DPPH is a stable nitrogen-centered free radical, its color changes from violet to yellow upon reduction by either the process of hydrogen or electron donation. Substances which are capable to perform this type of reaction can be considered as antioxidants and therefore radical scavengers. It was also found that with increasing the DPPH radical scavenging activity of extract with concentration, antioxidant activity also increases [14]. So, DPPH (1, 1-diphenyl-2-picryl-hydrazyl-hydrate) free radical method is an antioxidant assay which based on phenomenon of electron transfer that produces a violet colour in ethanol solution. This free radical, stable at room temperature, is reduced in the presence of an antioxidant molecule, give rise to colorless solution [15, 16]. Proton radical scavenging action is important mechanism for oxidation. The reduction capability of DPPH radicals was evaluated by the decrease in its absorbance at 517 nm, suggesting that antioxidant activity of plant extract is due to its proton donating ability [9, 10]. The hydrogen donating ability of the antioxidants molecules contributes to its free radical scavenging nature which is an important quality of antioxidants [52]. The DPPH radical assay has been used as a quick, reliable and easy method in order to investigate the general antioxidant activity of pure compounds as well as plant extracts [4]. It is a rapid method for screening of many samples for radical scavenging potential and is independent of sample polarity [13, 14].

Table 3. Total Phenolic profile of *M. charantia* seeds, peels and leaves.

Plant Parts	Total Phenolics (ppm)			
	Gallic acid	Caffeic acid	4-hydroxy, 3-methoxy benzoic acid	<i>p</i> -coumaric acid
Leaves	23.44±0.04	ND	ND	53.21±0.05
Peels	ND	ND	403.7±0.04	ND
Seed	ND	ND	ND	ND

Table 3. Continued.

Plant Parts	Total Phenolics (ppm)					Total Phenolics
	Ferulic acid	m-coumaric acid	Chlorogenic acid	Syringic acid	Vanillic acid	
Leaves	ND	ND	66.73±0.01	ND	ND	166.54±0.04
Peels	ND	17.21±0.07	ND	ND	ND	420.91±0.05
Seed	ND	ND	ND	56.34±0.04	301.9±0.03	358.24±0.03

The DPPH radical scavenging activity from leaves, seeds and peels of using different solvent systems are presented in Table 4. Somewhat, 80% methanol exhibited higher extraction yields from leaves (71.2%). The extraction ability of different solvent systems for recovering extractable components from leaves followed the order: 80% methanol >


absolute ethanol > 80% ethanol > absolute methanol. Yields of extract from seeds followed the order: 80% ethanol > 80% methanol > absolute ethanol > absolute methanol. Yields of extract from peels followed the order: 80% methanol > 80% ethanol > absolute methanol > absolute ethanol [15, 16].

Table 4. DPPH radical scavenging activity of *M. Charantia* seeds, peels and leaves extract.

Sr. no	Solvent System	DPPH (%) radical scavenging activity		
		Leaves	Seeds	Peels
1	80% Methanol	71.2±0.56 ^a	48.2±0.42 ^c	63.2±0.49 ^{a_b}
2	80% Ethanol	65.3±0.55 ^{b_c}	51.3±0.46 ^{b_c}	60.3±0.44 ^{b_b}
3	100% Methanol	61.2±0.50 ^{c_a}	44.2±0.59 ^c	58.1±0.32 ^{b_c}
4	100% Ethanol	67.5±0.41 ^{b_a}	46.1±0.45 ^{d_c}	53.2±0.45 ^{c_b}

Values are mean ± SD of three samples analyzed individually in triplicate at $p < 0.05$. Superscripts alphabets within the column depicted significant difference among different solvents. Subscripts alphabets within the rows depicted significant difference among different plant parts.

Reducing power

Figure 1. Reducing power of *M. charantia* seeds, peels and leaves extracts.

Reductive abilities of the plant extracts can work as an indicator of their potential antioxidant activities [66, 67]. The antioxidant ability of phenolic compounds is generally due to their redox properties. Then this allows them to react as a reducing agent, an electron donor and an oxygen quencher. Study on medicinal plants and vegetables revealed that plant possesses the antioxidant activity and are capable of applying the protective effects against oxidative stress in biological systems [63, 64]. Antioxidants have the ability to donate electron to reactive radicals. Then they neutralize them into stable and nonreactive species [44, 45]. In this assay, the presence of reducers (i.e., antioxidants) causes the reduction

of the Fe^{3+} to the ferrous form. So reducing power can be measured by donating electron and reducing $\text{Fe}^{3+}(\text{CN})_6$ into $\text{Fe}^{2+}(\text{CN})_6$. The product of prussian blue color was formed that can monitor the Fe^{2+} concentration at the wavelength of 700 nm [2]. Then this was observed that higher absorbance value showed more reducing potential of medicinal plants [43, 44]. Hence, activity of reducing power increases with increasing the concentration of extracts [54, 55]. The result of the present investigation presented in Table 4 showed that leaves of *M. charantia* showed maximum reducing power [38].

3.6. Antimicrobial Activity

The antimicrobial activity of leaves and seeds extract of *M. charantia* was determined by disc diffusion and microdilution broth methods against a panel of micro-organisms [39].

3.7. Antibacterial Activity by Disc Diffusion Method

The antibacterial activity of leaves seeds and peels extracts of *M. Charantia* was determined by disc diffusion method against a panel of bacterial strains (*P. multocida*, *E. coli* and *S. aureus*). The antibacterial potential of leaves, seeds and peels

extracts of *M. Charantia* varied significantly ($P < 0.05$) against bacterial strains as presented in table 5. The aqueous methanolic leaves extract was found to be the most effective against *P. Multocida* with DIZ of 30mm. However, absolute alcoholic leaves extracts exhibited no activity against *S. aureus*. The aqueous methanolic seeds extract showed highest DIZ (22mm) against *E. coli*. Whereas, all seed extracts showed no appreciable activity against *S. aureus*. Rifampicin was used as standard antibiotic. The Rifampicin showed DIZ (32mm, 29 mm and 20mm) against *P. multocida*, *E. coli* and *S. aureus* respectively [50, 51, 52].

Table 5. Antibacterial activity of *M. charantia* seeds, peels and leaves extracts

Plant parts	Microorganism	DIZ (mm)			
		80% Ethanol	Absolute ethanol	80% methanol	Absolute methanol
Leaves	<i>E. coli</i>	25±1.30 ^b	20±1.66 ^{bc}	27±2.45 ^{bc}	23±2.0 ^b
	<i>P. multocida</i>	29±1.23 ^a	23±1.35 ^b	30±0.45 ^a	25±1.2 ^a
	<i>S. aureus</i>	13±1.78 ^c	Nil	14±1.51 ^c	Nil
Seeds	<i>E. coli</i>	22±0.67 ^b	17±0.98 ^a	22±1.63 ^c	21±1.73 ^a
	<i>P. multocida</i>	21±0.55 ^c	16±1.25 ^b	17±1.95 ^d	15±1.22 ^b
	<i>S. aureus</i>	Nil	Nil	Nil	Nil
Peels	<i>E. coli</i>	21±0.67 ^d	15±0.98 ^b	20±1.63 ^c	19±1.73 ^a
	<i>P. multocida</i>	22±0.55 ^c	18±1.25 ^a	15±1.95 ^d	18±1.22 ^b
	<i>S. aureus</i>	Nil	Nil	Nil	Nil

Values are mean ± SD of three samples analyzed individually in triplicate at $p < 0.05$. Superscripts within the same column depicted significant difference among different fungal strains, while Subscripts within the same row indicated significant difference ($p < 0.05$) between solvent systems.




Figure 2. Antibacterial activity of *M. charantia* seeds, peels and leaves extracts.

3.8. Antifungal Activity Disc Diffusion Method


The antifungal activity of leaves, seeds and peels extracts of different parts of *M. Charantia* was determined by disc diffusion method against *A. parasiticus*, *A. flavus*, *F. oryzae*, *F. tritichum*, *A. Oryzae* [23-25]. The zones of inhibition are presented in table 6. It was concluded from the data that 80% ethanolic extract of leaves part showed strong activity against *A. Parasiticus* (28 mm

zone of inhibition) nearly equal to control (27 mm) while 80% ethanolic leaves extract of peels part showed lowest activity against *A. oryzae* (9 mm zone of inhibition) [68]. It was observed that 80% ethanolic extracts of leaves are best antifungal source. Fluconazol was used as standard antibiotic [40]. The Fluconazol showed DIZ (31mm, 30 mm, 29mm, 28mm and 18mm) against *F. tritichum*, *A. parasiticus*, *A. flavus*, *A. oryzae* and *F. oryzae* respectively [21, 26, 27].

Table 6. Antifungal activity of *M. charantia* leaves, seeds and peels extract.

Plant parts	Microorganism	DIZ (mm)			
		80% Ethanol	Absolute ethanol	80% methanol	Absolute methanol
Leaves	<i>parasiticus</i>	28±1.13 ^{ab} _a	20±1.73 ^c _c	22±1.40 ^{bc} _b	16±2.07 ^c _d
	<i>flavus</i>	24±1.27 ^b _a	21±1.64 ^c _c	17±2.02 ^{cd} _d	22±1.85 ^{ab} _b
	<i>F. oryzae</i>	Nil	Nil	Nil	Nil
	<i>F. tritichum</i>	12±2.04 ^{cd} _d	13±1.81 ^{ef} _c	24±2.08 ^{ab} _a	18±2.09 ^{bc} _b
	<i>oryzae</i>	19±1.06 ^{cd} _{ab}	13±1.64 ^{ef} _{bc}	12±1.50 ^f _{cd}	12±2.05 ^d _{cd}
Seeds	<i>A. parasiticus</i>	20±1.31 ^{bc} _b	17±0.98 ^{ab} _c	22±1.63 ^{de} _a	15±2.02 ^d _d
	<i>flavus</i>	18±1.02 ^{ce} _{bc}	16±1.25 ^{cd} _{de}	12±1.95 ^h _f	17±2.08 ^c _{cd}
	<i>F. oryzae</i>	Nil	Nil	Nil	Nil
	<i>F. tritichum</i>	15±1.39 ^{ce} _c	15±2.07 ^f _c	17±1.55 ^g _b	24±1.69 ^a _a
	<i>oryzae</i>	10±1.95 ^f _d	16±1.54 ^{cd} _b	18±2.03 ^e _a	12±1.89 ^{de} _c
Peels	<i>parasiticus</i>	21±1.33 ^{ab} _{bc}	16±0.95 ^b _f	20±1.66 ^{bc} _{cd}	19 ^{cd} ±2.07 ^d _{de}
	<i>flavus</i>	20±1.01 ^{bc} _a	14±1.20 ^{bc} _c	13±1.94 ^f _d	18±2.09 ^{de} _b
	<i>F. oryzae</i>	Nil	Nil	Nil	Nil
	<i>F. tritichum</i>	18±1.36 ^{cd} _{bb}	13±2.04 ^e _d	16±1.54 ^e _{cd}	29±1.67 ^{bc} _{ab}
	<i>oryzae</i>	9±1.93 ^e _d	17±1.59 ^a _b	19±2.04 ^{ce} _a	16±1.87 ^{ef} _c

Values are mean ± SD of three samples analyzed individually in triplicate at $p < 0.05$. Superscripts indicated significant difference ($p < 0.05$) between fungal strains while subscripts alphabets within the row depicted significant difference among different solvent.

**Figure 3.** Antifungal activity of *M. Charantia* leaves, seeds and peels extract.

3.9. Minimum inhibitory Concentration (MIC) of *M. Charantia* Seeds, Peels and Leaves Extracts Against Bacterial Strains

MIC values of leaves, seeds and peels of *M. charantia* against bacterial strains are given in table 7. Data showed that the least MIC value was obtained from the absolute methanolic extract of seeds against *P. multocida*, showing its

greater potential against *P. Multocida* [62]. While data showed that the least MIC value was obtained from the 80% ethanolic extract of leaves against *E. coli*, showing its lowest potential against *E. coli*. Seeds extracts were found to be ineffective against *S. aureus*. Rifampicin was used as standard antibiotic. The Rifampicin showed MIC (38µg/ml, 30µg/ml and 40µg/ml) against *P. Multocida*, *E. coli* and *S. aureus* respectively [28, 29, 30].

Table 7. Minimum inhibitory concentration (MIC) of *M. charantiaseeds*, peels and leaves extracts ageist bacterial strains.

Plant parts	Microorganism	MIC (µg/ml)			
		80% Ethanol	Absolute Ethanol	80% Methanol	Absolute Methanol
Leaves	<i>E. coli</i>	34±1.04 ^{de} _d	45±1.08 ^{ef} _c	55±2.08 ^{cc} _b	67±2.09 ^{bc} _a
	<i>P. multocida</i>	40±0.87 ^{cd} _{cd}	53±1.26 ^{de} _{bc}	69±1.86 ^{ab} _{ab}	75±1.97 ^a _a
	<i>S. aureus</i>	45±1.32 ^{bc} _b	Nil	73±1.55 ^{ab} _a	Nil
Seeds	<i>E. coli</i>	49±0.67 ^b _d	65±0.98 ^{bc} _b	62±1.63 ^b _c	71±1.77 ^b _a
	<i>P. multocida</i>	58±0.55 ^a _d	69±1.25 ^a _{bc}	73±1.95 ^a _b	85±2.05 ^{ab} _{ab}
	<i>S. aureus</i>	Nil	Nil	Nil	Nil
Peels	<i>E. coli</i>	42±0.67 ^{bc} _d	61±0.98 ^{ef} _c	66±1.63 ^{bc} _b	70±1.77 ^{cd} _a
	<i>P. multocida</i>	54±0.55 ^{ab} _d	63±1.25 ^{cd} _c	76±1.95 ^{ab} _b	80±2.05 ^{bc} _a
	<i>S. aureus</i>	Nil	Nil	Nil	Nil

Values are mean ± SD of three samples analyzed individually in triplicate at $p < 0.05$. Superscripts indicated significant difference ($p < 0.05$) between fungal strains while subscripts alphabets within the row depicted significant difference among different solvent.




Figure 4. Minimum inhibitory concentration (MIC) of *M. charanti* seeds, peels and leaves extracts against bacterial strains.

3.9.1. Minimum Inhibitory Concentration (MIC) of *M. Charantia* Seeds, Peels and Leaves Extracts against Fungal Strain

The minimum inhibitory concentration of the *M. charantia* extracts was determined against the test organisms. The minimum inhibitory concentrations are presented in the Table 8. From the data, it was concluded that 80% ethanolic extract of leaves part showed lowest minimum inhibitory

concentration (34 µg/ml) against *A. Paraciticus* while the absolute methanolic extract of peels part showed highest concentration (106 µg/ml) against *A. oryzae*. Fluconazol was used as standard antibiotic [48, 49]. The Fluconazol showed MIC (39 µg/ml, 30 µg/ml, 32 µg/ml, 46 µg/ml and 47 µg/ml) against *F. tritichum*, *A. parasiticus*, *A. flavus*, *A. oryzae* and *F. oryzae* respectively [31, 32, 33].

Table 8. Minimum inhibitory concentration (MIC) of *M. charanti* seeds, peels and leaves extracts against fungal strains.

Plant parts	Microorganism	MIC (mm)			
		80% Ethanol	Absolute ethanol	80% methanol	Absolute methanol
Leaves	<i>parasiticus</i>	34±1.13 ^{de} _d	36±2.73 ^{fg} _c	45±1.44 ^{ef} _b	47±2.08 ^{ef} _a
	<i>flavus</i>	38±1.27 ^{de} _f	42±2.67 ^{fg} _{cd}	48±2.05 ^{ef} _{bc}	52±1.85 ^{ce} _{ab}
	<i>F. oryzae</i>	Nil	Nil	Nil	Nil
	<i>F. tritichum</i>	46±2.07 ^{cd} _f	53±2.89 ^{ef} _e	67±2.45 ^{de} _d	78±2.09 ^{bc} _c
	<i>oryzae</i>	53±1.67 ^{cd} _d	57±2.24 ^{cd} _e	72±2.47 ^{cd} _b	82±1.93 ^{bc} _a
Seeds	<i>parasiticus</i>	55±1.33 ^c _d	64±2.07 ^{cd} _c	77±2.41 ^{cd} _{bc}	88±2.00 ^{bc} _{ab}
	<i>flavus</i>	62±1.09 ^{bc} _d	67±2.54 ^c _{cd}	79±2.59 ^{cd} _b	91±2.09 ^{bc} _a
	<i>F. oryzae</i>	Nil	Nil	Nil	Nil
	<i>F. tritichum</i>	74±1.41 ^a _e	75±2.09 ^{bc} _d	89±1.53 ^{bc} _{cd}	99±1.67 ^{ab} _{bc}
	<i>oryzae</i>	68±1.94 ^b _d	79±1.77 ^{ab} _c	92±2.04 ^b _{bb}	105±1.87 ^a _{ab}
Peels	<i>parasiticus</i>	57±1.34 ^{cd} _d	65±2.08 ^{cd} _c	78±2.45 ^e _b	89±2.01 ^{cd} _a
	<i>flavus</i>	63±1.09 ^{bc} _d	68±2.55 ^{bc} _{cc}	81±2.58 ^{cd} _{bc}	92±2.09 ^{bc} _{ab}
	<i>F. oryzae</i>	Nil	Nil	Nil	Nil
	<i>F. tritichum</i>	75±1.42 ^a _d	76±2.09 ^{ab} _c	90±1.55 ^{bc} _b	98±1.69 ^{ab} _a
	<i>oryzae</i>	69±1.95 ^{ab} _d	80±1.78 ^a _c	93±2.06 ^{ab} _{bc}	106±1.89 ^a _{ab}

Values are mean ± SD of three samples analyzed individually in triplicate at $p < 0.05$. Superscripts indicated significant difference ($p < 0.05$) between fungal strains while subscripts alphabets within the row depicted significant differences among different solvent.




Figure 5. Minimum inhibitory concentration (MIC) of *M. charantia* seeds, peels and leaves extracts against fungal strains.

3.9.2. Cytotoxicity Assay

By adopting the MTT assay Cytotoxicity of plant extracts was evaluated then by using baby hamster kidney cells (BHK-21) as described by Freshney and Frame (1982), though 10% DMSO was used as a positive control. Solutions of the tested materials were evaluated for cytotoxic potential.

Table 9. Cytotoxic potential of *M. Charantia* leaves.

Medicinal plant extract	Treatment (g/ml)	Cell survivalPercentage (%)
80%Methanolic leavesextract	24710	53.40±0.27

4. Conclusions

In this study, it was confirmed that different parts of *M. charantia* species exhibited excellent biological potential. The phytochemical analysis was carried out and showed considerable amount of TFC and TPC in different plant parts of *M. charantia*. Moreover leaves plant part exhibited effective antimicrobial and antioxidant activity. The major pharmacological components present in the leaves extracts of different parts of *M. charantia* showed higher antimicrobial effects as well as antioxidant potential. In this way this might accelerate the development of new drugs for numerous diseases caused by microorganisms. The present research work is economic, cost effective and can be applied in food and pharmaceutical industries by forming antimicrobial and antioxidative reagents.

References

- [1] Ahmad N, Hasan N, Ahmad Z, Zishan M, Zohrameena S (2016). *Momordica charantia*: For Traditional Uses And Pharmacological Actions. *Journal of Drug Delivery & Therapeutics* 6 (2): 40-44.
- [2] Ahmed D, Khan M M, and Saeed R (2015). Comparative Analysis of Phenolics, Flavonoids, and Antioxidant and Antibacterial Potential of Methanolic, Hexanic and Aqueous Extracts from *Adiantum audatum* Leaves. *Antioxidants*, 4, 394-409.
- [3] Alam M, Uddin R, Subhan N, Rahman MM, Jain P and Reza HM (2015). Beneficial Role of Bitter Melon Supplementation in Obesity and Related Complications in Metabolic Syndrome. *Journal of lipid volume* pages 18.
- [4] Aliyu AB, Musa AM, Ibrahim MA, Ibrahim H and Oyewale AO (2009). Preliminary phytochemical screening and antioxidant activity of leave extract of *Albizia Chevalieri* harms (Leguminoseae-Mimosoideae). *Bayero Journal of Pure and Applied Sciences*, 2 (1), 149 – 153.
- [5] Anilakumar KR, Kumar GP, Ilaiyaraja N (2015). Nutritional, pharmacological and medicinal properties of *Momordica charantia*. *International Journal of Nutrition and Food Sciences*. vol; 4 (1): 75-83.
- [6] Anjum F, Shahid M, Bukhari SA, Anwar S and Latif S (2013). Study of Quality Characteristics and Efficacy of Extraction Solvent/ Technique on the Antioxidant Activity of Bitter Gourd Seed. *J Food Process Technol*, vol4 (2).
- [7] Choo WS, Yap JY and Chan SY (2014). Antioxidant Properties of Two Varieties of Bitter Gourd (*Momordica charantia*) and the Effect of Blanching and Boiling on Them. *Pertanika J. Trop. Agric. Sci.* 37 (1): 121–131.
- [8] Chiampanichayakul S, Kataoka K, Arimochi H, Thumvijit S, Kuwahara T, Nakayama H, Vinitketkumnuen U and Ohnishi Y (2001). Inhibitory effects of bitter melon (*Momordica charantia* Linn.) on bacterial mutagenesis and aberrant crypt focus formation in the rat colon. *The Journal of Medical Investigation* Vol. 48, 88-96.

- [9] Chougule P, Pawar R, Limaye D, Joshi YM, and Kadam V (2012). In-Vitro Antioxidant Activity of Ethanolic Extract of *Centaurea* behen. *Journal of Applied and Pharmaceutical Sciences*, 2 (4), 106-110.
- [10] Daniel P, Supe U, Roymon MG (2014). A review on Phytochemical analysis of *Momordica charantia*. *International Journal Of Advances In Pharmacy, Biology And Chemistry* Vol. 3 (1).
- [11] Das DR, Sachan AK, Imtiyaz M, Shuaib M (2015). *Momordica charantia* as a Potential Medicinal Herb: An Overview. *Journal of Medicinal Plants Studies* 3 (5): 23-26.
- [12] De S, Chakraborty J, Das S (2000). Oral Consumption of Bitter Gourd and Tomato Prevents Lipid Peroxidation in Liver Associated with DMBA Induced Skin Carcinogenesis in Mice. *Asian Pacific Journal of Cancer Prevention*, Vol 1, 205- 208.
- [13] Dewanto V, Wu X, Adom KK, and Liu RH (2002). Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. *Journal of Agricultural and Food Chemistry*, 50 (10), 3010-3014.
- [14] Ebrahimzadeh MA, Nabavi SF, Nabavi SM, and Eslamib B (2010). Antihypoxic and antioxidant activity of *Hibiscus esculentus* seeds. *Grasas Y Aceites*, 61 (1), 30-36.
- [15] Fang EF, Zhang CZY, Wong JH, Shen JY, Li CH, Ng TB (2012). The MAP30 protein from bitter gourd (*Momordica charantia*) seeds promotes apoptosis in liver cancer cells in vitro and in vivo. *J. Cancer Letters* 324. 66–74.
- [16] Garcia EJ, Oldoni TLC, Alencar SMD, Reis A, Loguercio AD, and Grande RHM (2012). Antioxidant Activity by DPPH Assay of Potential Solutions to be Applied on Bleached Teeth. *Brazilian Dental Journal*, 23 (1), 22-27.
- [17] Giuliani C, Tani C, Bini LM (2015). Micromorphology and anatomy of fruits and seeds of bitter melon (*Momordica charantia* L., *Cucur bitaceae*). Published by Polish Botanical Society *Acta Soc Bot Pol* 85 (1): 3490.
- [18] Grover JK, Yadav SP (2004). Pharmacological actions and potential uses of *Momordica charantia*: a review. *Journal of Ethnopharmacology* 93 (2004) 123–132.
- [19] Gupta M, Shurma S, Gautam AK and Bhadauria R (2011). *Momordica charantia* Linn: nature silent healer. *International Journal of pharmaceutical Science Review and Research* Vol. 11 (1).
- [20] Hassan S M, Sultana B, Jahan N, and Iqbal T (2016). Evaluation of phenolic profile and antioxidant potential of medicinal plants. *Oxidation communication* 39, 3 (1), 2222-2236.
- [21] Hamissou M, Simth AC, Jr REC and Triplett II JK (2013). Antioxidative properties of bitter gourd (*Momordica charantia*) and zucchini (*cucurbita pepo*). *Emir. J. Food Agric.* 25 (9): 641-647.
- [22] Hasan SMR, Hossain MM, Akter R, Jamila M, Mazumder MEH, and Rahman S (2009). DPPH free radical scavenging activity of some Bangladeshi medicinal plants. *Journal of Medicinal Plant and Research*, 3 (11), 875-879.
- [23] Hossain MA, MostofaM, Debnath D, Alam AKMR, Yasmin Z and Moitry NF (2012). Antihyperglycemic and Antihyperlipidemic of Karala (*Momordica charantia*) Fruits in Streptozotocin Induced Diabetic Rats. *J. Environ. Sci. & Natural Resources*, 5 (1): 29–37.
- [24] Hsu B, and Coupar M (2006). Antioxidant activity of hot water extract from the fruit of the Doum palm, *Hyphaenethebaica*. *Food Chemistry*, 98, 317-328.
- [25] Jagessar RC, Mohameda A, Gomesb G (2008). An evaluation of the Antibacterial and Antifungal activity of leaf extracts of *Momordica charantia* against *Candida albicans*, *Staphylococcus aureus* and *Escherichia coli*. *Nature and Science* 6 (1), 1545-0740.
- [26] JayiniR (2012). Cytotoxic Activity Of *Momordica charantia* Seed Extract.
- [27] Joseph B, jinni D (2013). Antidiabetic effect of *momordica charantia* (bitter melon) and its medicinal potency. *the Asian Pacific Journal of Tropical Disease*
- [28] Krishnendu JR, Nandini PV (2016). Antioxidant Properties of Bitter Gourd (*Momordica charantia* L.). *International Journal of Advanced Engineering, Management and Science (IJAEMS)* Vol-2, Issue-10 ISSN: 2454-1311
- [29] Kai H, Akamatsu E, Torii E, Kodama H, Yukizaki C, Akagi I, Ino H, Sakakibara Y, Suiko M, Yamamoto I, Okayama A, Morishita K, Kataoka H and Matsuno K (2014). Identification of a Bioactive Compound against Adult T-cell Leukaemia from Bitter Gourd Seeds. *Plants*, 3, 18-26.
- [30] Kenny O, Smyth TJ, Hewage CM, Brunton NP (2014). Antioxidant properties and quantitative UPLC-MS/MS analysis of phenolic compounds in dandelion (*Taraxacum officinale*) root extracts. *Free Radicals and Antioxidants* Vo 4 / Issue 1
- [31] Kai M, Klaus HV, Sebastian L, Ralf H, Andreas R, Ulf PH (2007). Determination of DPPH Radical Oxidation Caused by Methanolic Extracts of Some Microalgal Species by Linear Regression Analysis of Spectrophotometric Measurements. *Sensors (basel)* 7 (10): 2080–2095.
- [32] Kim HY, Mok SY, Kwon SH, Lee DG, Cho EJ, and Lee S (2013). Phytochemical Constituents of Bitter Melon (*Momordica charantia*). *Natural Product Sciences* 19 (4): 286-289.
- [33] Kumar DS, Sharathnath KV, Yogeswaran P, Harani A, Sudhakar K, Sudha P, Banji D (2010). A Medicinal Potency of *Momordica charantia*. *International Journal of Pharmaceutical Sciences Review and Research* Vol 1 (2) 95-100.
- [34] Kumar KPS, Bhowmik D (2010). Traditional Medicinal Uses and Therapeutic Benefits of *Momordica charantia* Linn. *International Journal of Pharmaceutical Sciences Review and Research*. Vol 4 (3).
- [35] Kumar N and Khurana SMP (2016). Nutritive And Antidiabetic Benefits Of *Momordica charantia* L. (Bitter Gourd). *IAJPS*,
- [36] Kwatra D, Subramaniam D, Ramamoorthy P, Standing D, Moran E, Velayutham R, Mitra A, Umar S, and Anant S (2013). Methanolic Extracts of Bitter Melon Inhibit Colon Cancer Stem Cells by Affecting Energy Homeostasis and Autophagy. *Hindawi Publishing Corporation* 1-14.
- [37] Limtrakul P, Pitchakarn P and Suzuki S (2013). Kuguacin J, a Triterpenoid from *Momordica charantia* Linn: A Comprehensive Review of Anticarcinogenic Properties. *licensee InTech*.

- [38] Lu YL, Liu YH, Chyuan JH, Cheng KT, Liang WL, and Hou WC (2012). Antioxidant activities of different wild bitter gourd (*Momordica charantia* L. var. *abbreviata* Serin) cultivars. *Botanical Studies*. 53: 207-214.
- [39] Mada SB, Garba A, Mohammed HA, Muhammad A, Olagunju A and Muhammad AB (2013). Antimicrobial activity and phytochemical screening of aqueous and ethanol extracts of *Momordica charantia* L. leaves. *Journal of Medicinal Plants Research* Vol. 7 (10), pp. 579-586.
- [40] Mahmood A, Raja GK, Mahmood T, Muhammad Gulfranz and Khanum A (2012). Isolation and characterization of antimicrobial activity conferring component (s) from seeds of bitter gourd (*Momordica charantia*). *Journal of Medicinal Plants Research* Vol. 6 (4), pp. 566-573.
- [41] Mobaraki A, Moghaddam TN (2014). Karela, a Plant with Many Properties. *International Journal of Agriculture Innovations and Research* Volume 3, Issue 1, 2319-1473
- [42] Mohammady I, Elattar S, Mohammed S and Ewais M (2012). An Evaluation of Anti-Diabetic and Anti-Lipidemic Properties of *Momordica charantia* (Bitter Melon) Fruit Extract in Experimentally Induced Diabetes. *Life Science Journal*, 9 (2)
- [43] Naz S, Khan SN, Imani J, Mohy-Ud-Din G and Farooq S (2017). Geographical distribution of myrothecium leaf spot disease of *momordica charantia* l. Caused by myrothecium roridumtode in agro-ecological zones of punjab, Pakistan. *Pak. J.* 49 (4): 1599-1604,
- [44] Nishaa S, Vishnupriya M, Sasikumar JM, Christable HP, and Akkrishnan VNKG (2012). Antioxidant activity of ethanolic extract of *Maranta arundinacea*. L Tuberous Rhizomes. *Asian Journal of Pharmaceutical and Clinical Research*, 5 (4), 85-88.
- [45] Ozusaglam MA and Karakoca K (2013). Antimicrobial and antioxidant activities of *Momordica charantia* from Turkey. *African Journal of Biotechnology* Vol. 12 (13), pp. 1548-1558,
- [46] Panday, Rauniarb GP, Panday MDKR (2014). *Momordica charantia* (Karela); An Antidiabetic. *World Journal Of Pharmacy And Pharmaceutical Sciences*. Volume 4, Issue 1, 84-99.
- [47] Paul A, Raychaudhuri SS (2010). Medicinal Uses and Molecular Identification of Two *Momordica charantia* Varieties – a review. *Electronic Journal of Biology*, Vol. 6 (2): 43-51
- [48] Poonam T, Prakash GP and Kumar LV (2013). Interaction of *Momordica charantia* With Metformin In Diabetic Rats. *American Journal of Pharmacology and Toxicology* 8 (3): 102-106
- [49] Pornsuriya, Pramote, Pornsuriya, Pornthip and Numuen C (2011). Phenotypic diversity and classification of Thai bitter melon (*Momordica charantia* L.) landraces from three provinces in central region of Thailand. *Journal of Agricultural Technology* Vol. 7 (3): 849-856.
- [50] RajanS andPushpa A (2015). In vitro evaluation of enzymic antioxidants in the seed and leaf samples of *Syzygium cumini* and *Momordica charantia*. *International Journal of Scientific and Research Publications*, Volume 5, Issue 12.
- [51] Ramalingum N and Mahomoodally MF (2014). The Therapeutic Potential of Medicinal Foods. *Hindawi Publishing Corporation Advances in Pharmacological Sciences* Volume 2014, 18 pages.
- [52] Sathisha AD, Lingaraju HB, and Prasad KS (2011). Evaluation of Antioxidant Activity of Medicinal Plant Extracts Produced for Commercial Purpose. *European Journal of Chemistry*, 8 (2), 882-886.
- [53] Selvam K, Arunprakash S, Selvankumar T, Govarthanan M, and Sengottaiyan A (2017). Antioxidant prospective and secondary metabolites in *Abutilon indicum* at different environment. *International Journal of Pharmaceutical Sciences and Research*, 3 (7), 2011-2017.
- [54] Semiz A and Sen A (2007). Antioxidant and chemoprotective properties of *Momordica charantia* L. (bitter melon) fruit extract. *African Journal of Biotechnology* Vol. 6 (3), pp. 273-277.
- [55] Senguttuvan J, Paulsamy S, and Karthika K (2014). Phytochemical analysis and evaluation of leaf and root parts of the medicinal herb, *Hypochoeris radicata* L. for in vitro antioxidant activities. *Asian Pacific Journal of Tropical Biomedicine*, 4 (1), S359-S367.
- [56] Sharma S, Tandon S, SemwaB, Singh K (2011). *Momordica charantia* Linn.: A Comprehensive Review on Bitter Remedy. *Journal of Pharmaceutical Research And Opinion* 1:2 42–47.
- [57] Shobha CR, Vishwanath P, Suma MN, PrashantA, C. Rangaswamy, Gowdappa BH. 2015. In vitro anti-cancer activity of ethanolic extract of *Momordica charantia* on cervical and breast cancer cell lines. *International journal of health and allied science* Volume: 4, Issue: 4, Page: 210-217.
- [58] Sultana B, Anwar F, Ashraf M (2009). Effect of Extraction Solvent/Technique on the Antioxidant Activity of Selected Medicinal Plant Extracts. *Molecule*, 14, 2167-2180.
- [59] Suleman M, Nouren S, Hassan SM, Faiz AH, Sahr GA, Soomro, GA, Tahir MA, Iqbal M, and Nazir A (2018). Vitality and Implication of Natural Products from *Viburnum Grandiflorum*: an Eco-Friendly Approach. *Polish Journal of Environmental Studies*, 27 (3), 1-5.
- [60] Sherma B, and Kumar P (2009). Bioefficacy of *Lantana camara* L. against Some Human Pathogens. *Indian Journal of Pharmceutical Sciences*, 71 (5), 589-593.
- [61] Sumanth M and Chowdary GN (2010). Antimutagenic activity of aqueous extract of *Momordica charantia*. *International Journal for Biotechnology and Molecular Biology Research* Vol. 1 (4), 42-46.
- [62] Sylvie D D, Anatole PC, Cabral BP, and Veronique PB (2014). Comparison of in vitro antioxidant properties of extracts from three plants used for medical purpose in Cameroon: *Acalyphara cemos*a, *Garcini alucida* and *Hymeno cardialyrata*. *Asian Pacific Journal of Tropical Biomedicine*. 4 (2): S625-S632.
- [63] Tan S P, Stathopoulos C, Parks S, and Roach P (2014). An Optimised Aqueous Extract of Phenolic Compounds from Bitter Melon with High Antioxidant Capacity. *Antioxidants* vol, 3, 814-829.
- [64] Tokusoglu O, Unal M K and Yildirim Z (2003). HPLC–UV and GC–MS characterization of the flavonolaglycons quercetin, kaempferol and myricetin in tomato and tomato pastes and other tomato-based products. *Acta Chromato graphica*. 13: 196-207.

- [65] Tailor CS, Goyal A (2014). Antioxidant activity by DPPH Radical scavenging method of *Ageratum conyzoides* Linn Leaves. American Journal of Ethnomedicine1 (4): 244-249.
- [66] Vaghasiya Y, Chanda SV (2007). Screening of Methanol and Acetone Extracts of Fourteen Indian Medicinal Plants for Antimicrobial Activity. Turkish Journal of Biology, 31, 243-248.
- [67] Verma VN (2015). Chemical Composition and Quantitative Elemental Investigation of Bitter Melon. World Scientific News 84-94.
- [68] Zhang A, Fang Y, Wang H, Li H and Zhang Z (2011). Free-Radical Scavenging Properties and Reducing Power of Grape Cane Extracts from 11 Selected Grape Cultivars Widely Grown in China. Molecules, 16, 10104-10122.
- [69] Zhu Y, Dong Y, Qian X, Cui F, Guo Q, Zhou X, Wang Y, Zhang Y and Xiong Z (2012). Effect of Superfine Grinding on Antidiabetic Activity of Bitter Melon Powder. International Journal of Molecular Sciences 13 14204-14218.
- [70] Zafar F, Jahan N, Rahman KU, and Aslam S (2016). Synergistic free radical scavenging potential of polyphenolic phytotherapeutics in various plants combinations. Oxidation communication 39, 3 (1), 2213-2221.