

Performance of Selected Tef Genotype for High Potential Areas of Ethiopia

Yazachew Genet*, Tsion Fikre, Worku Kebede, Solomon Chanyalew, Kidist Tolosa, Kebebew Assefa

Ethiopian Institutes of Agricultural Research; Debrezeit Agricultural Research Centre, Debre Zeit, Ethiopia

Email address:

yazachewgenet@gmail.com (Y. Genet)

*Corresponding author

To cite this article:

Yazachew Genet, Tsion Fikre, Worku Kebede, Solomon Chanyalew, Kidist Tolosa, Kebebew Assefa. Performance of Selected Tef Genotype for High Potential Areas of Ethiopia. *Ecology and Evolutionary Biology*. Vol. 5, No. 3, 2020, pp. 35-42. doi: 10.11648/j.eeb.20200503.11

Received: July 17, 2020; **Accepted:** July 28, 2020; **Published:** August 17, 2020

Abstract: Genetic improvement of native crops is a promising strategy to combat hunger in the developing world. Tef is the major staple food crop for approximately 73 million people in Ethiopia. As an indigenous cereal, it is well adapted to diverse climatic and soil conditions; however, its productivity is extremely low mainly due to lack of high yielder genotypes, susceptibility to lodging, biotic and abiotic stresses. To circumvent these problem, an experiment comprising 20 tef genotypes including the standard and local checks were evaluated in a randomized complete block design with four replications at nine environment to develop high yielding, stable and farmers preferred variety (ies) for high potential areas. Combined analysis of variance revealed highly significant ($P \leq 0.01$) variations due to genotypes, environments for most of traits and significant ($p \leq 0.05$) genotype by environment interaction effects (GEI) for grain yield. AMMI analysis revealed 7.62%, 67.27%, 25.11% variation in grain yield due to genotypes, environments and GEI effects, respectively. The mean grain yield value of genotypes averaged over environments indicated that G12 (DZ-Cr-387 X Rosea (RIL-133) had the highest grain yield (2761 kg ha^{-1}) compared to the standard check variety Negus (2526 kg ha^{-1}). In addition this candidate variety proved stable across environments for grain yield during the variety evaluation experiment. Therefore, this genotype was evaluated by the national variety released committee for release as a new variety for the year of 2019/20 and the technical committee approved it for fully released as new variety in 2020. Thus, this variety should be used as a commercial variety for potential tef growing areas to increase tef productivity and production in the country.

Keywords: Tef, Genotypes, RIL, Multi Environment, GEI

1. Introduction

Background

Tef (*Eragrostis tef*) is being labeled as one of the latest super foods of the 21st century, like the ancient Andean grain quinoa, tef's international popularity is rapidly growing [3] mainly because the grains are free from gluten to which many people are allergic [24] a causal agent for celiac disease; hence, tef is becoming globally popular as a life-style crop [22]. Tef is a resilient crop that performs better than other cereals under local conditions including drought, waterlogging, and poor soil. Since it produces a reasonable yield when grown in areas that experience moisture scarcity, it is considered as a low risk crop [15, 16]. Tef is nutritious due to its high protein and mineral content [1, 9]. Tef is one of the most significant crops for farm income, food and nutrition security in Ethiopia. It

serves both as a staple and cash crop in the country.

Tef is versatility crop in adapting to adverse environmental conditions and staple food for ~73 million people in Ethiopia where it is annually cultivated by 7 million small-scale farmers on more than 30% of the total area allocated to cereal crops [4] In a country of more than 100 million people, tef accounts for about 15% of all calories consumed and, contribute well over 66% of the protein intake of the population consuming it as their staple food. The crop is preferred both by farmers and consumers because of its excellent nutritional quality (well balanced protein and minerals) and it makes good quality “injera”, pancake-like soft bread. The straw serves as an indispensable feed for cattle and has almost equal value as the grain.

Grain yield is a complex character which is dependent on a number of other characters and is highly influenced by many genetic factors as well as environmental fluctuations. On the other hand, the genotype x environment interaction (GEI) is an important aspect of both plant breeding program and the introduction of new crop cultivars [8, 12, 20]

Despite its importance, the productivity of tef is much lower than other cereals. The national average yield of tef is about 1.75 tha^{-1} , compared to maize (3.2 tha^{-1}) and wheat (2.4 tha^{-1}), respectively [4]. The major constraints limiting productivity and production of tef are; 1) limited availability of varieties suitable for different agro-ecologies; 2) limited use of improved varieties; 3) presence of biotic and abiotic stresses; and 4) inadequate seed and extension systems.

Tef research and development efforts in Ethiopia began in the late 1950s with the objectives of addressing the afore-mentioned constraints. Over the past 24 years, tef productivity increased by about 100%, from just 0.7 tha^{-1} in 1994 to 1.75 tha^{-1} in 2018. In tef improvement effort grain yield constituted the highest priority [13]. Therefore, tackling some of the high priority problems mentioned above is vital to increase tef productivity in the Country. Consequently, the objective of the study was to evaluate the performances of selected tef genotypes across multi-locations and identify candidate variety (ies) having broad and /or specific adaptation to different environments.

2. Materials and Methods

2.1. Plant Materials

Eighteen promising recombinant inbred lines (RILs) selected from preliminary yield trial plus two checks (local and standard check) were used. The 18 promising recombinant inbred lines were obtained through single seed descent (SSD) method from two different crosses. In both crosses Quncho (DZ-Cr-387/RIL 355) was used as the ovule parent. The cultivars Rosea and Alba described by [17] were used as pollen parent. The former cultivar is characterized by high number of florets per spikelet and hence used to pyramid yield traits into the popular variety Quncho released in 2006 [13]. Likewise, the cultivar Alba was the paternal

parent for six of the 18 RILs, and the cross of Quncho with cultivar Alba aimed at introgressing higher panicle length for yield as well as. thick and strong culm for increased lodging tolerance into the popular variety Quncho. The standard check variety was the variety Nigus released in 2017 [21] for agro-ecologies similar to the particular set of test locations and classified as high potential tef growing areas. On the other hand, the local check is a farmers' variety commonly grown around each of the respective test locations.

Table 1. Description code of the study tef genotypes.

Code	Genotypes
G1	DZ- Cr- 429 (RIL 125)/Negus (standard check)
G2	DZ-Cr-387 X Rosea (RIL-9)
G3	DZ-Cr-387X Roseau (RIL-22)
G4	DZ-Cr-387X Rosea (RIL-38)
G5	DZ-Cr-387X Rosea (RIL-24)
G6	DZ-Cr-387X Rosea (RIL-48)
G7	DZ-Cr-387X Rosea (RIL-52)
G8	DZ-Cr-387X Rosea (RIL-75)
G9	DZ-Cr-387X Rosea (RIL-92)
G10	DZ-Cr-387Xrosea (RIL-117)
G11	DZ-Cr-387 X Rosea (RIL-121)
G12	DZ-Cr-387 X Rosea (RIL-133)
G13	DZ-Cr-387 XAlba (RIL-347)
G14	DZ-Cr-387 XAlba (RIL-226)
G15	DZ-Cr-387 X Rosea (RIL-159)
G16	DZ-Cr-387 XAlba (RIL- 260)
G17	DZ-Cr-387 XAlba (RIL- 279)
G18	DZ-Cr-387 XAlba (RIL-249)
G19	DZ-Cr-387 XAlba (RIL- 216)
G20	Local

2.2. Experimental Locations and Seasons

Although the experiment was done for two seasons at each of the six locations and also additional other locations, the data for some of the years and locations were excluded because of the heterogeneity of variance in the combined analyses of grain yield data over all environments (locations and seasons). The test locations represent high potential tef growing areas with optimum rainfall and other climatic and edaphic conditions suitable for tef production (Table 2).

Table 2. The nine test environments used for the national variety trial for high potential areas.

Locations							
Code	Name latitude	latitude	longitude	Altitude (m.a.s.l)	Annual rainfall	mean Temperature	Soil type
E1	Akaki	8°54'N	38°45'E	2205	1025	18	vertisol
E2	Minjar-1	8°45'N	39°45'E	2000	1118	19.5	nitosol
E3	Holeta-1	09°03'N	38°30'E	2400	1102	14.5	nitosol
E4	Adadi-1	08°31'N	38°13'E	2383	1105	16.5	vertisol
E5	Minjar-2	8°45'N	39°45'E	2000	1118	19.5	nitosol
E6	Holeta-2	09°03'N	38°30'E	2400	1102	14.5	nitosols
E7	Adadi-2	08°31'N	38°13'E	2383	1105	16.5	vertisol
E8	Bichena	10°26'N	38°12'E	2543	1316	16.4	vertisol
E9	Adet	11°16'N	37°29'E	2174	1209	16.5	vertisol

*Climatic and edaphic information was obtained from their respective research and sub centers. Minjar-1=Minjar-2, Holeta-1=Holeta-2, Adadi-1=Adadi-2 are the same site, respectively.

2.3. Experimental Design and Management

The field experiment was conducted using a randomized complete block design with four replications of 2 m x 2 m (4m²) plot size during the two main seasons of 2017 and 2018. The field experiment was managed as per the research recommendation of agronomic practices of the respective test locations.

2.4. Data Collected

Data on agronomic yield and yield related traits were collected both on plot and individual plant base. Data on days to heading or panicle emergence using the sowing date as a reference, lodging index, grain and biomass yield were taken on plot bases. Days of heading and maturity were taken when each plot attained 50% heading (panicle emergency) and 90% physiological maturity respectively, and days were calculated beginning from the date of sowing. Lodging index was assessed using the method of [2] by considering assessments of both the lodging degree or the angle of leaning on 0 (completely upright) to 5 (completely flat on the ground) and the severity as the percentage of the plot stand manifesting each of the 0-5 degrees of lodging. Then, lodging index for each plot was taken as the product sum of the degree of leaning and the respective per cent severity divided by five. Grain yield (g) of each plot was measured on clean, sun dried seed and the measured grain yield value (g) has converted to kilogram per hectare for data analysis.

Plant height (cm), and panicle length (cm) were taken on the five individual samples of plants which were randomly taken from the central rows of each plot, and the averages of five sample plants were as used for analysis.

2.5. Data Analyses

For each trait analysis of variance was made first for individual location, and ultimately upon getting positive results from tests of homogeneity of variances using the method F-max of [11], a combined analysis of variance was made across the environments (locations and years) in order to determine the differences between genotypes across environments, among environments and their interaction. For the analysis of variance, Proc GLM (general linear model) suitable for the experimental design were employed [10] using SAS software version 9.00 [23] and the average performance for different traits presented below (Table 3). AMMI (additive main effects, multiplicative interactions analysis was used to adjust the main or additive genotype and

environmental effects by analysis of variance, in addition to the adjustment of the multiplicative effects for the G×E interaction by principal component analysis.

The sum of squares of the G×E interaction was divided into an singular axis or Interaction Principal Component Axis (IPCA), which reflects the standard portion in which each axis corresponded to a particular AMMI model. Mean comparison for traits showing significant differences in the analyses of variance were made using Least Significant Difference (LSD). GEA-R (2015) software version 2.0 was used for the stability analysis and GGE biplot analysis to visualize which genotypes performed best in which environment.

3. Result and Discussion

3.1. Components of Variation

ANOVA from additive main effect and multiplicative interaction (AMMI) for most of traits showed significant ($p \leq 0.01$) for genotypes and environments and significant ($P \leq 0.05$) effect for genotype by environment interaction (GEI). The effect of environment, genotypes and genotype by environment interaction accounted for 67.27%, 7.62% and 25.11% of the total sum squares (Table 3), respectively. A large sum of squares for environments indicated that the test environments were diverse with large differences among environmental means which causing most of the variation in grain yield. Therefore, this result designated the reliability of the multi environment experiments. The variation in temperature, rainfall, soil type, soil fertility, and moisture availability might be the main reasons for the presence of variation. The AMMI analysis also showed that the first interaction principal component (PC1) and second interaction principal component (PC2) explained 39.32% and 19.61% of the interaction sum squares, respectively. The mean squares for PC1 was highly significant ($p < 0.01$). Likewise, analysis of variance revealed highly significant ($p < 0.01$) effect GEI for aboveground biomass, days to heading, days to maturity, panicle length, plant height and lodging index. The significant interaction indicated that the genotypes respond differently across the different environments. The significant interaction indicated that the genotypes respond differently across different environments. The significant variability of genotypes traits showed in the present study for different traits of tef genotypes are in agreement with the previous report by different authors for genotype variability [12, 18].

Table 3. AMMI analysis of variance for grain yield grown at nine environments.

Source	DF	SS	M.S.	V.r.	F pr	Explained GEI SS%
Treatment	179	146663414	819349	4.14	<0.001	7.62
Genotypes (G)	19	11182431	588549	2.97	<0.001	67.27
Environments (E)	8	98656693	12332087	7.85	<0.001	
Block (E)	27	42427854	1571402	7.94	<0.001	
Interaction (GEI)	152	36824290	242265	1.22	0.05	25.11
PC1	26	14477805	556839	2.81	<0.001	39.32
PC2	24	7222800	300950	1.52	0.055	19.61
Error	513	101584390	198020			

DF = degree of freedom, S.S = Sum squares, V.r= F calculated value, Fpr = F probability Value

3.2. Mean performance of Selected Tef Genotypes

The mean grain yield performances of the 20 tef genotypes at each of the nine test environments are presented in Table 4. The overall mean grain yield of tef genotypes for the nine environments ranged from 1718 kg ha⁻¹ at E8 (Bichena-2018) to 3846 kg ha⁻¹ at E2 (Minjar-2017). Among nine environments, E2 (Minjar-2017), E1 (Akaki), E5 (Holeta-2), E4 (Adadimariam-2017) and E3 (Holeta-2017) were high yielding environments. While, E7 (Adadimariam-2018), E8 (Bichena-2017) and E9 (Adet-2017) were low yielding environments.

On the other hand, mean grain yield value of genotypes averaged over environments ranged from 2340 kg ha⁻¹ (G10) to 2761 kg ha⁻¹ (G12) (Table 4). The significant GEI in the present study indicates unstable performance of the tef genotypes across the testing environments (Figure 1). Thus, it implied that the genotypes respond differently across the different environments. The test genotype G12 (DZ-Cr-387 X Rosea RIL133) was the top yielder at E5 (Holeta-2018), and the second highest yielder at E1 (Akaki-2017), E4 (Adadimariam-2017), E7 (Adadimariam-2018) and E9 (Adet-2017) (Table 4). Overall, the genotype code G12 (candidate variety), although not at all of the environments, performed better than others at least at two low yielding environments (Adadimariam-2018, and Adet-2017) and three high yielding environment (Akaki -2017), Adadimariam-2017 and Minjar-2018). The huge variability in the grain yield among the 20 tef genotypes at the nine environments might be due to wide variability in climatic and soil conditions. This finding is in accordance with previous studies [7, 12, 19] that similarly reported which thereby

complicates the selection and recommendations stable genotype across environment.

In genotype x environment interaction (GEI) the result exhibited the genotypes gave statistically higher grain yield and above ground biomass than the standard check variety. In addition to this considering the current tef and straw price, 36 Birr kg⁻¹ and 5 birr kg ha⁻¹ (1 USD=27 birr), respectively, there was an economically meaningful difference among tested genotypes. Therefore, one promising candidate variety, DZ-Cr-387 X Rosea (RIL-133) gave grain yield (2761 kg ha⁻¹) and aboveground biomass 13802 kg ha⁻¹ compared to the standard check variety Negus depicting grain yield (2526.4 kg/ha) and aboveground biomass 11402 kg ha⁻¹, respectively. Therefore, DZ-Cr-387 X Rosea (RIL-133) has been evaluated in by the National Variety Release Technical Committee in the variety verification trial during 2019/2020 and released as a new variety in 2020. From the variety verification trial, the candidate variety showed promising performance than the newly released standard check variety Ebba.

The visualization of a ‘which-won-where’ pattern in multi-environment trials is essential to study adaptability of genotypes in the specific or across all test environments [25]. The vertex genotypes were the most responsive for being located at the greatest distance from the biplot origin. The genotypes with either the best or poorest performance in one or all environments were considered responsive [25] falling within the sectors. The GGE biplots of graph results was used to show the relative performance of all genotypes at a specific environment (Figure 1).

Table 4 Mean grain yield (kg ha⁻¹) performance of tef genotypes across nine environments.

Code	Environments								
	E1	E2	E3	E4	E5	E6	E7	E8	E9
G1	2980.0	3370	2544.3	2607.2	2718.8	2104.4	2116.7	2130.1	2166.3
G2	2828.8	3332.5	2479.0	2784.1	2796.9	2238.8	2529.5	1761.2	1804.8
G3	2583.8	3108.8	2605.9	2385.0	2298.1	2164.6	1753.3	2625.1	2101.9
G4	2925.0	3846.3	2436.6	2726.0	2310.6	2402.2	1895.0	2680.9	2192.0
G5	2782.6	3582.5	2901.2	2532.5	2732.5	2370.9	2080.4	2682.4	2093.6
G6	3133.8	3478.8	2830.9	2976.5	2384.4	2356.3	2158.5	2140.5	2224.9
G7	3021.3	3480.0	2488.8	2503.9	2553.8	1879.6	2059.5	2394.9	1943.9
G8	3091.3	3582.5	2367.9	2338.9	2513.1	2493.2	1926.3	2638.1	2320.3
G9	2567.5	3191.3	2591.4	2659.4	2376.3	1560.1	1718.2	2833.9	2037.4
G10	2898.8	3231.3	2157.8	2050.3	2393.1	2638.9	1792.4	1777.2	2116.1
G11	2602.5	3586.3	2549.9	2737.1	2510.0	2092.7	2435.5	2387.3	2003.4
G12	3087.5	3553.8	2640.5	3153.7	2963.8	2334.7	2512.2	2281.0	2315.9
G13	2842.5	3092.5	3113.7	3078.1	2145.0	2725.9	2326.5	2805.4	2287.3
G14	2803.8	3325.0	2923.6	2489.0	2865.6	2400.6	2244.3	1943.3	2614.7
G15	2465.0	3025.0	2678.5	2160.8	2383.1	2051.3	1735.7	2838.8	2308.3
G16	2906.3	3432.5	2726.7	3259.9	2647.5	2208.2	2076.0	2693.6	2287.3
G17	2853.8	3683.8	2741.4	2706.8	2423.1	2467.2	2343.3	2912.9	2007.6
G18	2637.8	2945.3	2628.0	2418.1	2613.8	2521.1	2105.3	2130.0	2204.1
G19	2825.0	3686.3	2654.9	2964.1	2915.0	2723.1	2190.7	1995.8	2419.8
G20	2808.8	3091.3	2801.8	2402.3	1987.5	2090.1	1881.8	2429.3	1920.0
CV	10	20	10	15	11	13	22.9	29	10
LSD	421.0	981.7	397.0	571.6	402.7	426.5	678.4	1020	320.6
SE	40.8	78.9	44.3	54.6	42.4	49.6	58.0	94.8	30.6

G1-G20 name of Genotypes, E1= Akaki (2017), E2= Minjar 2017, E3= Holeta (2017), E4= Adadi= 2017, E5= Minjar (2018), E6= Holeta (2018), E7=Adadi (2018), E8= Bichna (2018), E9= Adet (2018), CV= coefficient of variation, LSD=least Significant Difference and SE =Standard error.

Table 5. Mean grain yield and other agronomic characteristics of 20 tef genotypes averaged over nine environments.

Genotypes	Yldkg ha ⁻¹	AGBkg ha ⁻¹	PL (cm)	PH (cm)	DTH (days)	DTM (days)	LI (%)
G1	2526±116	11403	35	93	52	112	76
G2	2506±103	13045	40	106	55	117	74
G3	2403±84	12104	41	106	58	115	74
G4	2602±118	12778	38	100	58	114	76
G5	2640±110	12990	40	105	57	114	76
G6	2632±115	12514	38	101	55	113	76
G7	2481±99	12601	40	106	56	114	76
G8	2586±109	12295	38	105	55	115	75
G9	2393±109	13288	42	105	59	113	74
G10	2340±100	12618	41	109	54	114	72
G11	2545±103	11910	41	104	55	114	78
G12	2761±95	13802	40	106	55	115	75
G13	2713.0±106	13007	41	106	54	115	75
G14	2623±106	13250	40	105	57	116	73
G15	2405±93	12750	41	105	59	115	74
G16	2686±93	13347	42	112	56	114	74
G17	2682±109	13410	43	110	57	113	75
G18	2467±124	13073	43	111	55	115	74
G19	2708±112	11014	33	90	51	111	80
G20	2379±94	10837	36	95	52	114	80
Genotype (G)	**	**	**	**	**	**	**
Environment (E)	**	**	**	**	**	**	**
GEI	*	**	**	**	**	**	**
CV	20	18	11	5	4	2	10
LSD (0.05)	423	1936	4	5	2	2	7
R ²	53	88	73	84	94	98	68

Yldkg ha⁻¹ = yield kilogram per hectare, DTH= Days to heading, DTM= days to maturity, PH=plant height, GFP= grain filling period, PH=plant height, PL= Panicle length, LI= lodging index, R² (%)=the model explain the variability of the response data around its mean

3.3. Stability Analysis

Mean grain yield performance and its stability 20 tef genotypes over nine environments are shown in table 6 and Figure 2. From GGE biplot graph for stability analysis Average environmental axis (AEA) is a line passing through the origin and pointing to the positive direction with its distance equal to the longest vector. Besides, an ideal environment is a point on the AEA in the positive direction of the biplot origin and is equal to the longest vector of all environments [25]. This line was reported to be useful to evaluate mean grain yield and stability of genotypes [25]. According to such reports, genotypes considered to be stable are those appeared closer to the origin with the shortest vector from the AEC. Thus, Figure

2 in the present study shows the mean performance and stability of the genotypes. Based on this, G12 with the shortest vector from the AEC axis was identified as the most stable genotypes while G10 with the longest vector from AEC was the most unstable genotypes.

The mean grain yield value of genotypes averaged over environments indicated that G12 had the highest (2761kg ha⁻¹) and G10 the lowest grain yield (2340 kg ha⁻¹), respectively. Genotype superiority with the small measured value indicates the more stable genotypes (Table 6). Therefore, from the present study, G12 was the most stable and G10 most unstable genotypes, respectively.

Table 6. Stability coefficient analysis of mean grain yield of 20 tef genotypes tested across nine environments.

Genotypes	Grain yield Mean kgha ⁻¹	Standard Deviation	Genotype Superiority
G1	2526	446	135078 (12)
G2	2506	508	179646 (15)
G3	2401	388	198671 (16)
G4	2602	558	108536 (8)
G5	2640	459	81837 (5)
G6	2632	487	91877 (7)
G7	2481	516	162910 (13)
G8	2586	485	125513 (11)
G9	2393	530	226952 (19)
G10	2340	498	290293 (20)
G11	2545	456	122601 (10)
G12	2761	449	53531 (1)
G13	2713	3734	84241 (6)
G14	2623	410	119332 (9)
G15	2405	402	224489 (18)
G16	2686	468	63719 (2)

Genotypes	Grain yield Mean kg ha^{-1}	Standard Deviation	Genotype Superiority
G17	2682	471	72877 (3)
G18	2467	279	173804 (14)
G19	2708	492	78627 (4)
G20	2379	443	222293 (17)

N. B: Numbers in brackets give the position of each genotype, ranked according to the stability coefficient (running downwards from 1 = best).




Figure 1. The which-won-where view of the GGE biplot of 20 tef genotypes to show which genotypes performed best in which environments.

Beneficial advantage of new variety DZ-Cr.497 /DZ-Cr-387 X Rosea (RIL-133)

The New variety DZ-Cr.497/DZ-Cr-387 X Rosea (RIL-133)/ has got the following major advantages.

- 1) It showed advantage of 235kg ha^{-1} (9.3%) in grain yield and 2399 kg ha^{-1} (21%) in aboveground biomass yield over the standard check variety Negus as well as 16.1% in grain yield and 36.8% in aboveground biomass over the local check cultivar.
- 2) Moreover, the selected genotype showed highly stability (1st rank) among evaluated genotypes, indicating its suitability for multi environment in the high potential tef growing areas.
- 3) This genotype has also got immense farmers' preference

Table 7. Description of agronomic and morphological characteristics of the new variety DZ-Cr-387 x Rosea (RIL-133).

No.	Parameters	Description
I	Variety Name	
1	Breeders Name	DZ-.Cr-497
2	Pedigree	DZ-Cr-387 X Rosea (RIL-133)
3	Vernacular name given	Dinknesh
II	Adaptation conditions and agronomic practice	
4	Adaptation area	High & optimum tef growing areas
5	Altitude (m.a.s.l.)	1700-2500
6	Rain fall (mm)	700-1200
7	Soil type	Mainly vertisols and nitosols
8	Seed rate (Kg ha^{-1})	10-15
9	Planting method	both broad casting & row sowing
10	Row spacing (cm)	20
11	Planting date	July 10-30

and attention due to its overall performance and white caryopsis colour during the participatory variety evaluation.




Figure 2. Means versus stability of 20 tef genotypes tested at nine environments

Description of the new variety DZ-Cr-497/ DZ-Cr-387 X Rosea (RIL-133)

A summary of the description of the candidate variety including its pedigree, adaptation agro-ecological conditions, required cultural practices, and pheno-morphologic and agronomic traits is presented on Table 7.

No.	Parameters	Description
12	Fertilizer Use	recommended rate for tef
13	Pest reaction	Not significant
III	Qualitative traits	
14	Panicle form	Very loss
15	Lemma color	Variegated (yellow+red)
16	Anther color	Red
17	Caryopsis color	Very white
18	Growth habit	Erect
IV	Quantitative traits	Mean
19	Days to heading (days)	52
20	Days to maturity (days)	112
21	Plant height (cm)	93
22	Panicle length (cm)	35
23	1000 seed weight (g)	0.3
24	Grain yield on station (kg/ha-1)	2761
25	Grain yield on farm (kg/ha-1)	2140
26	Aboveground biomass (kg/ha-1)	13802

4. Conclusions and Recommendations

Crop yield is a complex trait that is influenced by a number of component characters along with the environment directly or indirectly. If high yielding stable recombinant inbred lines tef could be developed for diverse environments, it would be possible to provide diverse and stable varieties for the tef growing farmers. Stability analysis is a powerful approach to select the most stable high yielding recombinant inbreeds lines for specific as well as for diverse environments. In the present study, 20 tef genotypes including 18 promising RILs originating from two crosses and selected on the basis of previous preliminary variety trials as well as as standard check variety Negus and a local check (farmers' variety) from each location were field evaluated at nine environments (six location and two main seasons of 2017 and 2018). Combined analysis of variance revealed highly significant ($P \leq 0.01$) variations due to genotypes, environments for most of traits and significant ($p \leq 0.05$) genotype by environment interaction effects (GEI) for grain yield. AMMI analysis revealed 7.62%, 67.27%, 25.11% variation in grain yield due to genotypes, environments and GEI effects, respectively. Thus, the GEI mean squares showed tef genotypes exhibited differential performances across the different environments. Consequently, most of the genotypes showed environment specificity. The mean grain yield value of genotypes averaged over environments indicated that G12 had the highest (2761 kg/ha-1) and G10 the lowest yield (2340kg/ha⁻¹), respectively. It is noted that the variety G12 showed higher grain yield than all other varieties when averaged over all the environments.

One promising late set candidate variety, DZ-Cr-387 X Rosea (RIL-133) gave higher grain yield of (2761kg/ha⁻¹) compared to the standard check variety Negus (2526.4kg/ha⁻¹). Therefore, DZ-Cr-387 X Rosea (RIL-133) has been selected and evaluated by the National Variety Release Committee in 2019/2020 and released in 2020. Thus, it is recommended for high potential tef growing regions in the country. Multi environmental trial should be conducted continuously to get high yielding tef varieties for different tef

growing areas to increase production and productivity of tef.

Overall, the tef varieties released have shown steady and incremental genetic gain through tef breeding in Ethiopia of 0.90% year under lodging controlled (wire-mesh support) conditions from the earliest release in 1970 until 1995 [26, 27], and 0.58% per annum under lodging uncontrolled natural conditions from 1970 until 2013 [5, 6]. These figures are relatively good by the standards of most breeding programs for similar crops, except for the most important world crops like maize, wheat and rice. However, to bring breakthrough, instead of the steady increment, in in tef improvement further intensified crossing/hybridization in order to stack productivity traits/genes, break the apparent linkage between culm thickness and culm length for reduced lodging vulnerability, and use advanced breeding techniques including genomics are vital so as to get substantially high yielder and stable genotypes with the required qualities. Moreover, future research strategies on tef genetic engineering, high throughput mutant line development, and mining of the tef genetic resources including the wild relative species must be given due emphasis in the national tef breeding program.

Acknowledgements

The authors are indebted to the Agricultural Growth Program (AGP-II) for financial support and all collaborating centres and their respective tef research staff for the successful execution of the field experiment.

References

- [1] Abebe Y, Bogale A, Hambidgeb K, Michael S, Barbara J., Baileyd K & Gibson, R. S. (2007). Phytate, zinc, iron and calcium content of selected raw and prepared foods consumed in rural Sidama, Southern Ethiopia, and implications for bioavailability. *Journal of Food Composition and Analysis* 20, 3-4, 161-168.
- [2] Caldicott, J. J. B. and A. M. Nuttall. 1979. A method for the assessment of lodging in cereal crops. *Jour. of Nat. Inst. Agri. Botany*. 15: 88-91.

- [3] Collyns Dan., (2013). Quinoa brings riches to the Andes. Retrieved from <http://www.theguardian.com/world/2013/jan/14/quinoa-andes-bolivia-peru-crop>
- [4] CSA. 2018. Central Statistical Agency, Agricultural Sample Survey 2017/2018 (2010 E.C). Volume I. Report on Area and Production of Major Crops (Private Peasant Holdings, Meher Season). Statistical bulletin, 586, Addis Ababa, Ethiopia.
- [5] Fano Dargo (2013). Genetic Gain in Grain Yield Grain Yield Potential and Associated Traits of Tef [*Eragrostis tef* (Zucc.) Trotter] in Ethiopia, MSc. Thesis, Haramaya University, Haramaya, Ethiopia.
- [6] Fano Dargo, Firew Mekbib. and Kebebew Assefa (2016). Genetic Gain in Grain Yield Potential and Associated Traits of Tef [*Eragrostis tef* (Zucc.) Trotter] in Ethiopia. *Global Journals Inc.* (USA) 16: 2249-4626.
- [7] Fufa Hundera, Hailu Tefera, Kebebew Assefa, Tesfaye Tefera, Tiruneh Kefyalew Girma Taye (2000). Grain yield and stability analysis in late maturing genotypes of tef [*Eragrostis tef* (Zucc.) Trotter]. *J. Genet. Breed.* 54: 13-18.
- [8] Freeman GH (1985) the analysis and interpretation of interaction. *Journal of Applied Statistics.* 12: 3-10.
- [9] Geremew Bultosa, Hall A., & Taylor J. (2002). The World Factbook. Africa: Ethiopia. Retrieved from CIA website; <https://www.cia.gov/library/publications/the-world-factbook/geos/et.html>
- [10] Gomez, K. A., and A. A. Gomez. (1984). Statistical Procedures for Agricultural Research. 2nd ed., John Wiley and Sons, Inc., New York, USA.
- [11] Hartley, H. O. (1950). The maximum F-ratio as a short cut test for heterogeneity of variances. *Biometrika* 37: 308-312.
- [12] Habte Jifar, Kebebew Assefa, Kassahun Tesfaye, and Zerihun Tadele. (2019) genotype x environment interaction and stability analysis in grain yield of tef (*Eragrostis tef*) evaluated in Ethiopia. *JEAI*, 35 (5): 1-13; Article no. *JEAI*. 48459. DOI: 10.9734/JEAI/2019/v35i530214.
- [13] Kebebew Assefa, Yu J, K., Zeid, M., Getachew Belay, Hailu Tefera and Sorrells, M. E. (2011).
- [14] Breeding tef [*Eragrostis tef* (Zucc.) Trotter]: conventional and molecular approaches. *Plant Breed.* 130: 1-9.
- [15] Seyfu Ketema. (1997). Tef. *Eragrostis tef* (Zucc.) Trotter. Promoting the conservation and use of underutilized and neglected crops. 12. Institute of plant genetics and crop plant Research, Gatersleben/International Plant Genetic Resources Institute, Rome, Italy.
- [16] Seyfu Ketema. 1993. Tef (*Eragrostis tef*): Breeding, Genetic Resources, Agronomy, Utilization and Role in Ethiopian Agriculture. Institute of Agricultural Research, Addis Ababa, Ethiopia.
- [17] Tadesse Ebba (1975). Tef (*Eragrostis tef*) Cultivars, Morphology and Classification, Part II. Experiment Station Bulletin No. 66, Addis Ababa University, College of Agriculture. Dire Dawa, Ethiopia.
- [18] Tiruneh Kefyalew. Genotype x environment interaction in tef. (2001) Tefera H, Belay G, Sorrells M, (Eds). Narrowing the Rift: Tef Research and Development. Proceedings of the International Workshop on Tef Genetics and Improvement, Debre Zeit: EARO; 2001.
- [19] Tiruneh Kefyalew (1999). Assessment of Genotype x Environment Interaction for Yield and Yield Related Traits in Tef [*Eragrostis tef* (Zucc.) Trotter] Genotypes, MSc. Thesis, Alemaya University of Agriculture, Alemaya, Ethiopia.
- [20] McLaren CG, Chaudhary C, (1994) Use of additive main effects and multiplicative interaction models to analyse multiplication rice variety trials. Paper presented at the FCSSP Conference, Puerton Princesa, and Palawan, Philippines.
- [21] MoALR (2017). Ministry of Agriculture and Livestock Resource, Plant Variety Release, Protection and Seed Quality Control Directorate, Crop Variety Register Issue Mo. 20, Addis Ababa, Ethiopia.
- [22] Provost C. and Jobson E. 2014. Move over quinoa, Ethiopia's Tef poised to be next big super grain. The Guardian, January 23, 2014.
- [23] SAS Institute. (2002). SAS/STAT Guide for Personal Computers, Version 9.00 editions. Cary, N. C., SAS Institute Inc.
- [24] Spaenij-Dekking L., Kooy-Winkelaar Y., Koning F. 2005, 'The Ethiopian cereal Tef in celiac disease', New England *Journal of Medicine* 353 (16), 1748–1750.
- [25] Yan W, Tinker NA. Biplot analysis of multi- environment trial data: Principles and applications. *Can J Plant Sci.* 2006; 86: 623-645.
- [26] Yifru Teklu (1996). Genetic Gain in Grain Yield Potential and Associated Agronomic Traits of tef [*Eragrostis tef* (Zucc.) Trotter], MSc. Thesis, Alemaya University of Agriculture, Dire Dawa, Ethiopia.
- [27] Yifru Teklu and Hailu Tefera (2005). Genetic improvement in grain yield potential and associated agronomic traits of tef (*Eragrostis tef*). *Euphytica* 141: 247-254.