

# Assessment of Availability of Adequately Iodized Salt and Its Associated Factors at Household Level in Walmara Woreda, Finfine Special Zone, Ethiopia, 2020

Tesfaye Gudisa<sup>1,\*</sup>, Esubalew Tesfahun<sup>2</sup>, Robert Wondimu<sup>3</sup>, Zalalem Kaba<sup>4</sup>, Edosa Amente<sup>5</sup>, Haile Bikila<sup>6</sup>

<sup>1</sup>Walmara Worada Health Office, Holeta, Ethiopia

<sup>2</sup>Department of Public Health, Debre Berhan University, Debre Berhan, Ethiopia

<sup>3</sup>Maternal and Child Health Program, Sendafa Beke Town Health Office, Sendafa, Ethiopia

<sup>4</sup>East Wollega Zonal Health Office, Nekemte, Ethiopia

<sup>5</sup>Ethiopian Public Health Institute, Addis Ababa, Ethiopia

<sup>6</sup>Department of Public Health, Institute of Health Science, Wollega University, Nekemte, Ethiopia

## Email address:

tesfayegudisa2021@gmail.com (T. Gudisa), esubalew.tesfahun@gmail.com (E. Tesfahun), robertwondimu@gmail.com (R. Wondimu), kabazalalem@gmail.com (Z. Kaba), idoamante07@gmail.com (E. Amente), haile.bikila@gmail.com (H. Bikila)

\*Corresponding author

## To cite this article:

Tesfaye Gudisa, Esubalew Tesfahun, Robert Wondimu, Zalalem Kaba, Edosa Amente, Haile Bikila. Assessment of Availability of Adequately Iodized Salt and Its Associated Factors at Household Level in Walmara Woreda, Finfine Special Zone, Ethiopia, 2020. *European Journal of Preventive Medicine*. Vol. 9, No. 6, 2021, pp. 140-148. doi: 10.11648/j.ejpm.20210906.11

**Received:** November 8, 2021; **Accepted:** November 29, 2021; **Published:** December 11, 2021

---

**Abstract:** Background: Iodine Deficiency Disorders refer to a spectrum of health consequences resulting from inadequate intake of iodine. Around 2 billion people in 130 countries worldwide have insufficient intakes of iodine and Ethiopia also one of the countries more than half the population at risk of Iodine Deficiency Disorders (IDD). Recently monitoring of the iodine in Dessie and Kombolcha, Ethiopia indicated that 68.8% of households used adequately iodized salt which is below the international goal of 90% coverage. Universal salt iodization has been implemented to eliminate Iodine Deficiency Disorders. However, the adequacy of iodine in salts needs close monitoring to meet its intended goal. This study assessed availability of adequately iodized salt and its associated factors at household level in Walmara Woreda Finfine Special zone, Oromia, Ethiopia, 2020. A Community-based cross-sectional study was conducted in Walmara Woreda from February 01 to 30 in 2020. Multistage sampling technique Data collections were used to study participant. All independent variables with p-value<0.25 in the bivariate analysis were selected for multivariate logistic regression analysis. During multivariate logistic regression analysis the P-value less than 0.05 were considered as statistically significant and degree of association between dependent and independent variables were reported using adjusted Odds Ratio (AOR) and 95% CI. This study showed that the availability of adequately iodized salt at household level with parts-per-million (PPM) of >15 was 355 (65.9%). Households who live in urban areas were 8.74 times more likely to have adequately iodized salt than those rural areas (AOR=8.74, 95% CI: 2.92, 26.09). Storage of salt at dry place was 5.38 times more likely to have adequately iodized salt than that stored at moister place (AOR=5.38, 95% CI: 1.46, 19.88). Households who were using packed salt were 11.441 times more likely to have adequately iodized salt compared to those who were using non-packed salt (AOR=11.44 (95% CI: 3.97, 33.02)). The availability of adequately iodized salt at household level in Wolmara Woreda was low as compared to the Iodine Deficiency Disorders elimination strategy. Hence, Zonal and Woreda health office should give attention to monitoring and evaluation of iodine salt at production, transportation and consumption levels.

**Keywords:** Iodized Salt, Walmara, Household Level, Finfine Special Zone, Ethiopia

---

## 1. Introduction

Iodine is a trace element that is essential for the synthesis of thyroid hormones by the thyroid gland [1]. An iodine deficiency disorder is a spectrum of health consequences resulting from inadequate intake of iodine. Iodine deficiency disorders is manifested in multiple adverse health consequences like decreased child survival rates, goiter, abortion, stillbirth, malformation and overall impaired growth and development [2].

Globally more than 1.88 billion people are at risk of iodine deficiency and 246 million school-age children are estimated to have insufficient iodine [3, 4]. In Ethiopia, more than 35 million (62%) people are at risk of iodine deficiency and around 28 million people suffer from goiter, and 50,000 prenatal deaths are related to iodine deficiency each year in Ethiopia [5]. Salt is generally used as the vehicle for providing iodine and iodization of salt is an effective, feasible, economical, safe and broadly accepted by the community [4].

Around 2 billion people in 130 countries worldwide have insufficient intakes of iodine [6]. Nearly 38 million newborns in developing countries every year remain unprotected from the lifelong effect of brain damage due to Iodine Deficiency Disorders (IDD) [7]. In Africa about 260 million people have inadequate iodine intake resulting in iodine deficiency states, which are 10-15% lowering of average intellectual capacity [5]. The consequences of iodine deficiency disorders affect all stages of life from fetus to adulthood as well as old age [7].

Ethiopia is one of the countries more than half the population at risk of IDD (3). Recent monitoring of the iodine indicated that 68.8% of households used adequately iodized salt containing at least 15 ppm iodine Dessie and Kombolcha, considerably below the international goal of 90% coverage [2].

Monitoring Salt at the Household provides important information about the successful delivery of iodized salt to the consumer [8]. Hence the need to undertake this study was to provide current information regarding the availability of adequately iodized salt and its associated factor at household level in Wolmara Woreda, Finfine Special Zone, Oromia, Ethiopia. This study contributed information to monitor percentage of population remains at risk of inadequate intake of iodine. It gives baseline information for the next intervention plan based on the finding to promote Universal Salt Iodization (USI) for the Wolmara Woreda and stakeholders. The baseline information can also be used to forward recommendations for policy makers and implementers to enhance appropriate planning and implementation of intervention relevant to utilization of adequately iodized salt. This study finding can be used as starting point for other researchers who are interested to conduct further investigations.

## 2. Methods

### 2.1. Study Area and Period

This study was conducted in Wolmara Woreda, Finfine

Special Zone, Oromia Region, Ethiopia which is located at 30km from the Addis Ababa (the nation capital) to the west. According to the 2007 national population census, the Woreda has an estimated total population of 115,697 and 24,103 households. There are 4 public health centers, 24 health posts with 43 health extension workers to provide basic primary health care services and 7 private medium clinics. The public health service coverage of the Woreda is 87%. This study was conducted from February 01 to 30, 2020.

### 2.2. Study Design

Community based cross-sectional study was conducted.

### 2.3. Population

#### 2.3.1. Source of Population

All households who live in Wolmara Woreda were taken as the source population.

#### 2.3.2. Study Population

All sampled households in the selected kebeles were taken as the study population.

### 2.4. Inclusion and Exclusion Criteria

#### 2.4.1. Inclusion Criteria

All household selected and adult member of the household who were involved in the food preparation at household level were included. In the household where two or more household members were involved in food preparation, the representative of mother or wife was purposely selected for interview.

#### 2.4.2. Exclusion Criteria

- 1) Household who lived less than six months in the study area were excluded from the study.
- 2) Household member who were <18 years old and participated in food preparation was excluded from the study.

### 2.5. Sample Size Determination and Sampling Procedures

#### 2.5.1. Sample Size Determination

The study sample size was determined by using a single population proportion formula considering following assumption:

P-Currently monitoring of availability of adequately iodized salt at household level in Dessie and Kombolcha Towns=68.8% [9], the standard normal value at 95% confidence level (1.96), d=margin of error (0.05), Non-responder rate=10% and Design effect of the study=1.5, fitting in to the formula the final sample size was calculated as:-  

$$n = \frac{(Z \cdot \frac{\partial}{2})^2}{d^2} \cdot n_{\text{non-response}} \cdot \text{Design effect}$$

$$n = \frac{(1.96)^2}{0.05^2} \cdot 0.688 \cdot 1.5 = 330$$

Then adding 10% for non-response rate the sample size (n)=330+33=363. However, since the study utilized multi-stage sampling, this sample size was multiplied by 1.5 for the design effect. The final sample sizes for this study is 1.5\*363=545.

### 2.5.2. Sampling Procedures

Multi stage sampling technique with three stage sampling process was used. The Woreda has 26 Kebeles and all of them clustered to each health center. Then, the sample sizes of Kebeles were allocated proportionally to each health center. Kebeles were again selected randomly from each cluster by lottery method. Households were proportionally allocated to each Kebele based on the household size of them. Finally; systematic random sampling was used to select the household.

### 2.6. Data Collection Method

Data was collected using interviewer administered structured questionnaire and rapid iodine test kit. Four diploma and two BSc health professionals were participated in data collection and supervision respectively. During the data collection sample of salt used for cooking was obtained to test for the iodine content using MBI KITS (international salt test kits). The test kit contains a small white cup, 2 test solution ampoules of 10ml, a recheck solution ampoule of 10ml and color chart indicators for iodine content ranges of 0 ppm, less than 15 ppm and more than 15 ppm. The Procedure to test iodine content of salt are:

- 1) Fill small cup with salt, then spread the salt surface flat.
- 2) Add two drops of the test solution on the surface of the salt by piercing the white ampoule with a pin and gently squeezing the ampoule.
- 3) Compare the color on the salt with the color chart, within one minute and determine the iodine content.
- 4) If no color appears on the salt (after one minute) on a fresh sample add up to five drops of the recheck solution in red ampoule and then add two drops of test solution on the same spot. The color of the test sample is compared with the standard color chart for calculating the salt iodine content. Finally, salt with  $\geq 15$  PPM was categorized as adequately iodized salt, whereas  $<15$  PPM was considered non-iodized salt.

### 2.7. Data Quality Management

The questionnaire was developed in English and then translated in to Afaan Oromo (the regional language) and retranslated into English to check consistency. To assure data quality of the study; one day training was given for the data collectors and supervisors on the data collection tools and procedure. The questionnaires were pretested before the actual data collection on 5% of the sample size out of the study area to ensure its validity. The data collectors and supervisors checked the accuracy and completeness of filled questionnaires on daily basis. Any error in data collection was rechecked before proceeding with the next day's data collection activity.

### 2.8. Variables of the Study

#### 2.8.1. Dependent Variable

Availability of adequately iodized salt at household level.

#### 2.8.2. Independent Variable

- 1) Socio-demographic characteristics of study population (Age, Residence, Sex, Marital status, Ethnicity, family size, Education status, Occupation, Monthly income of the family, and role of respondent)
- 2) Knowledge about iodized salt And Iodine Deficiency disorder
- 3) Duration of storage salt
- 4) Place of storage salt.
- 5) Type of iodine salt (packed or not)
- 6) Types of container storage
- 7) Time salt added during food cooking.

### 2.9. Operational Definitions

- 1) Availability of adequately iodized salt: - Salt tested by rapid test kit had  $\geq 15$ ppm called adequately iodized salt [10].
- 2) Knowledge: - Study participants who scored  $\geq 50\%$  regarding iodized salt and IDD questions were considered as having good knowledge, whereas, participants who scored below 50% were considered as having poor knowledge [11].
- 3) Duration of storage iodized salt: - House hold those store the purchased salt more than two months were considered as having longer storage time, and those who store for less than two months were considered as having shorter storage time [12].
- 4) Storage of salt: - storage of iodized salt at household level without exposing to environmental factors such as heat, moisture and light [13].
- 5) Type of salt: - type of salt means based on whether the salt was packed or not. Packed salt means when the salt is packed by any plastic or with tin and not packed when the salt was not packed with anything.
- 6) Place of storage of salt: - place store iodized salt at household level without exposing to moisture or not. Storage at dry place when the storage place not exposed to moisture and if storage place of salt is exposed to moisture called storage at moisture.
- 7) Salt exposed to sun light/ heat: - storage of iodized salt at household level without exposing to environmental factors light/ Heat or not. Not Salt exposed to sun light/ heat when the storage place of salt not exposed to sun light/ heat and if storage place of salt is exposed to sun light/ heat called exposed to sun light/ heat.
- 8) Container of storage:- container used to store salt at house hold level, store with cover container when respondents use to store salt any type of container which has cover and without cover if not use with cover container.
- 9) Role of Respondent in food preparation:- preparation only when respondents participate only in food preparation, buy salt if respondents only participate in buy salt and participate both preparation and buy salt if the respondents participate both.

### 2.10. Data Processing and Data Analysis

After data collection, each questionnaire was checked for completeness and consistency manually. The data was entered into Epi data 3.1 and exported to SPSS statistical software Version 20.0 for analysis. The descriptive statistics was presented by using frequencies, percentages, tables and graphs. A binary logistic regression analysis was performed to determine the association. All the independent variables with  $p$ -value $<0.25$  in the bivariate analysis were the candidates for multivariate logistic regression analysis. The  $P$ -value less than 0.05 were considered as statistically significant. The degree of association between dependent and independent variables was reported using Odds Ratio (OR) and 95% CI.

#### Ethical Considerations:

The necessary explanation about the purpose, potential risk and procedure of study was explained and verbal consent was obtained from each respondent to ensure the right of participant. Participants were informed that participation was on voluntary basis and they can withdraw at any time if they are not comfortable about the questionnaire. In order to protect the confidentiality of information name and other identification was not be included in questioners and maximum effort was made to maintain privacy of the respondent during the interview. The collected data was stored and kept safe place where any the investigator is acceptable and the computer was protected by private pass word.

## 3. Result

### 3.1. Socio Demographic Characteristics of the Study Participants

Out of the 545 samples 539 of participant involved in the study, with a response rate of 98.8%. Out of study subject, 400 (74.2%) were rural residence area, 365 (67.7%) Oromo in ethnicity, 115 (21.3%) unable to read and write, 370 (68.4%) were married and 279 (51.8%) were farmers. Three hundred fifty seven (66.2%) study house hold had family size  $\leq$  five and 312 (57.9%) had an average monthly income of  $<5000$  Ethiopian Birr (ETB) [Table 1].

**Table 1.** Socio-demographic characteristics of participants at Walmara Woreda, Finfine Special Zone, Oromia, Ethiopia, 2020.

Variable	Frequency (n)	Percentage (%)
Sex		
Female	539	100.0
Residence		
Urban	139	25.8
Rural	400	74.2
Total	539	100.0
Age of respondents (in years)		
18-24	142	26.3
25-34	115	21.3
35-44	115	21.3
45-54	79	14.7
55-64	58	10.8
$>64$	30	5.6
Total	539	100.0

Variable	Frequency (n)	Percentage (%)
Ethnicity of the respondents		
Oromo	365	67.7
Amara	88	16.3
Tigre	40	7.4
Gurage	46	8.5
Total	539	100.0
Educational status		
Unable to read and write	115	21.3
Primary education	161	29.9
Secondary education	199	36.9
Diploma and above	64	11.9
Total	539	100.0
Marital status		
Currently married	370	68.6
Never married	84	15.6
Widowed	42	7.8
Separated	43	8.0
Total	539	100.0
Occupation status of house hold		
Government or private employee	47	8.7
Daily laborer	70	13.0
Merchant	143	26.5
Farmer	279	51.8
Total	539	100.0
Income status of house hold (in Ethiopian Birr)		
$<5,000$	312	57.9
5,000-9,999	181	33.6
$>10,000$	46	8.5
Total	539	100.0
Family size status		
$<5$ Family size	357	66.2
$>5$ Family size	182	33.8
Total	539	100.0
Role of respondents in food preparation		
Preparation only	102	18.9
Recruitment only	80	14.8
Preparation and recruitment	357	66.2
Total	539	100.0

### 3.2. Availability of Adequately Iodized Salt at the Household Level

Using rapid test kit adequately iodized salt ( $>15$ ) was found in 355 (65.9%) of the 539 tested salt sample at house hold level and rest 184 (34.1%) had inadequate ( $<15$ ppm) iodine salt. Regarding the type of salt tested 84.1% of packed salt had adequate iodine and only 24.4% of those non packed salt had adequately iodized at house hold level.

### 3.3. Respondents Information Regarding Iodine Salt and Iodine Deficiency Disorders

In this study, 100 (18.55%) of the participants had not heard about iodine. Among study subjects who heard about iodine 437 (51.9%) obtained information from friend and relatives only. From the total respondent 318 (59%) believed that all salt are iodized salt. One over two respondents those who do not use packed salt was believed that packed salt was not available and 52.1% respondents knew iodine important for health. Among study participant only 191 (35.4%) knew cause of goiter is iodine deficiency and one over two of respondents knew IDD preventable by universal salt iodization [Table 2].

**Table 2.** Respondents' information regarding iodine and IDD in Walmara Woreda, Finfine Special Zone, Oromia, Ethiopia, 2020.

Variable	Frequency (n)	Percentage (%)
Source of information respondent heard (n=437)		
From friend and relatives only	226	51.9
From health extension health worker only	93	21.1
From Television/Radio (media) only	21	4.8
From health worker and media television	92	21.1
Other	5	1.1
Total	437	100.0
Type of iodized salt respondents knew		
All are iodized	318	59.0
Packed salt iodized salt	140	26.0
Not packed salt	68	12.6
Don't know	13	2.4
Total	539	100.0
The reason respondent do not use packed iodized salt (n=169)		
Unknown where availability	89	52.7
Believe more expensive	71	42.0
Do not know	9	5.3
Total	169	100.0
Inadequate iodine intake iodine health problem		
Goiter	271	72.5
Cretinism	29	7.8
Reduce intelligence	39	10.4
Still birth	21	5.6
Sudden abortion	14	3.7
Total	374	100.0
Important of Iodine salt		
Important for health	203	52.1
Food for flyover	171	43.8
Advice from health professional	9	2.3
Other	7	1.8
Total	390	100.0
Courses of goiter		
Iodine deficiency	191	35.4
Hereditary	124	23.0
Environmental sanitation	113	21.0
Infection	45	8.3
Poisoning	66	12.2
Total	539	100.0
At risk group of population to goiter		
Children	97	18.0
Pregnant women	352	65.3
Men	35	6.5
Females	55	10.2
Total	539	100.0
Method of IDD preventable (n=373)		
use universal iodine salt	191	51.2
Eat iodine rich food	85	22.8
take iodine drug	97	26.0
Total	373	100.0
Knowledge of Respondent		
Good	402	74.6
Poor	137	25.4
Total	539	100.0

#### 3.4. Practice of Respondents Regarding the Use of Iodized Salt

Among study subject 370 (68.8%) of household use packed salt, nearly three forth of respondents buy the salt from market and only 232 (43%) check iodine content by reading the label on the pack. Regarding the practice of iodine salt, 491 (91.1%) households stored salt in a dry place,

434 (80.5%) store salt with cover, 152 (28.2%) of the respondents exposed salt for sunlight/heat at house hold level, all most all 99.1% household use salt without washing and 347 (64.4%) add salt to food at the end of food preparation[Table 3].

**Table 3.** Practice of respondents regarding the use of iodized salt in Walmara Woreda, Finfinne Special Zone, Oromia, Ethiopia, 2020.

Variable	Frequency (n)	Percentage (%)
Type salt respondent use		
Packed salt	370	68.6
Not packed salt	169	31.4
Total	539	100.0
Type of container salt was packed		
with plastic	327	88.6
with tin	42	11.4
Total	369	100.0
Method used to check iodine content		
Read the label	232	43.0
Asking the seller	134	24.9
Don't know	173	32.1
Total	539	100.0
Source of salts buy.		
From market	382	70.9
From supper market	96	17.8
From shop	61	11.3
Total	539	100.0
Storage place of salt in home		
Moisture place	48	8.9
dry place	491	91.1
Total	539	100.0
Container respondent used to store salt in home		
Container with cover	434	80.5
Container without cover	105	19.5
Total	539	100.0
Salt exposure to sun/heat		
Yes	152	28.2
No	387	71.8
Total	539	100.0
Duration store salt in home		
<2 month	461	85.5
>2 month	78	14.5
Total	539	100.0
Time respondent add to food during cooking		
At early of food cooking	47	8.7
At middle of food cooking	145	26.9
At end of food cooking	347	64.4
Total	539	100.0

#### 3.5. Factors Associated with the Availability of Adequately Iodized Salt at Household Level

The investigation on the presence of association between suspected factors and availability of adequately iodized salt revealed the following results. Residence of respondent, family size, respondents knew IDD preventable, method used to check salt iodine content, storage place, container used to store salt, salt exposure to sun or heat, duration salt storage and type of salt respondents use were significantly associated with availability of adequately iodized salt during multivariate analysis. Households who live in urban areas were 8.74 times more likely to have adequately iodized salt than those who were from rural areas (AOR=8.74, 95% CI: 2.92, 26.09).

Family size less than five 3.51 times more likely to have adequately iodized salt than those whose family sizes greater than five member (AOR=3.51, 95% CI: 1.45, 8.26). Storage salt in dry Place was 5.38times more likely to have adequately iodized salt than that stored at moisture Place in home (AOR=5.38, 95% CI: 1.46, 19.88). Use of container with a cover to store was 4.69 times more likely to have adequately iodized salt than that use a container without

cover to store (AOR=4.69, 95% CI: 1.64, 13.39). In relation to duration of salt storage those household who store salt for short time were 3.75 times more likely to have adequately iodized salt than those who store for longer period of time (AOR=3.75, 95% CI: 1.12, 11.32). House hold those who use packed salt was 11.44 times more likely to have adequately iodized salt compared to respondents those who use non packed salt (AOR=11.441 (95% CI: 3.97, 33.02) [Table 4].

**Table 4.** Factors associated with the availability of adequately iodized salt at household level in Walmara Woreda, Finfine Special Zone, Oromia, Ethiopia, 2020.

Variables	Availability of adequately Iodized salt at house hold level			
	Inadequate<15pp No (%)	Adequate>15ppm (%)	COR	AOR
Residence of respondents				
Urban	22 (4.08%)	117 (21.7%)	3.62 (2.20, 5.95)	8.74 (2.92, 26.09)
Rural	162 (30%)	238 (44.16%)	1	1
Family size status				
<5Familysize	111 (20.5%)	246 (45.64%)	1.48 (1.02, 2.15)	3.42 (1.48, 7.96)
>5Familysize	73 (13.54%)	109 (20.22%)	1	1
Respondents knew about IDD Preventable				
Yes	62 (11.50%)	311 (57.69%)	13.9 (8.96, 21.59)	8.20 (3.34, 20.16)
No	122 (22.63%)	44 (8.16%)	1	1
System of respondents use to check iodine salt content,				
Read the label	33 (6.12%)	199 (36.92%)	12.94 (7.94, 21.08)	3.71 (1.29, 10.68)
Asking the seller	33 (6.12%)	101 (18.74%)	6.57 (3.96, 10.90)	1.26 (0.437, 3.65)
Don t know	118 (21.89%)	55 (10.20%)	1	1
Storage place of salt at home during data collection				
At dry place	174 (32.28%)	338 (62.7%)	1.143 (0.51, 2.55)	5.38 (1.46, 19.88)
At moisture	10 (1.9%)	17 (3.15%)	1	1
Container used to store salt at home				
With cover	125 (23.19%)	309 (57.32%)	3.17 (2.046, 4.91)	4.48 (1.58, 12.02)
Without cover	59 (10.95%)	46 (8.53%)	1	1
Salt Exposure to sun/heat during storing in house				
No	120 (22.26)	267 (49.54)	1.62 (1.10, 2.38)	3.04 (1.38, 6.71)
Yes	64 (11.87%)	88 (16.33%)	1	1
Duration of storage salt at home				
Shorter time of storage	127 (23.5%)	334 (61.97)	7.14 (4.15, 12.26)	3.75 (1.13, 11.33)
Longer time storage	57 (10.57%)	21 (3.89%)	1	1
Type of salt respondents use				
Packed	56 (10.34%)	314 (58.25%)	17.51 (11.14, 27.82)	11.44 (3.97, 34.02)
Not packed	128 (23.75%)	41 (7.60%)	1	1

## 4. Discussion

The study finding indicates that the overall percentage of households who use adequately iodized salt (>15 ppm) was 355 (65.9%). This study result was nearly in line with the study conducted in Debra Tabor Town 66.4%, Assela town 62.9%, South Africa 62.4%, and Dessie and Kombolcha 68.8% [11, 9, 14, 15]. The finding was much higher than the study reports in Ethiopia from rural community of Maychew 33%, Gondar town 28.8%, and Lalo Assabi 8.7% [16, 13, 17]. In contrast to the above finding, adequacy of iodized salt in the study was lower than findings from Sri Lanka 88.7%, in Bia district Ghana 75.6%, the Iodized salt Coverage of EDHS 2016 89% [18, 19, 7]. This study also less than the WHO, UNICEF and ICCIDD recommendation for the elimination of IDD is possible when the proportion of households using iodized salt reaches 90%, and in Ethiopia USI target should be 90% at house hold level [2, 20]. The reason for these variations may be due to relatively

wide time elapsed between the current study and others. Additionally, controlling of iodine deficiency disorders has been given universal attention with strong monitoring and evaluation of iodine at every level (at production, transportation and consumption levels). The differences in agro-ecological situations, socio-economic status and practices among households may bring the above discrepancies in different areas [4].

Among study participant 402 (74.6%) of respondents have good knowledge about iodine salt and iodine deficiency disorder. This study result was higher than the study conducted in Jig Jiga town, Dessie and Kombolcha town [21, 9]. But, this study result was less than the study conducted in Addis Ababa which indicates the overall prevalence of knowledge among women in reproductive age was 78% [22]. The variation might be due to the nature of study settings, in Addis Ababa women in reproductive age group and who have opportunities to increase their knowledge through promotions of iodized salt on different media. It may also be due to variation on awareness creation and educational

activities so that most of the communities become aware of iodized salt and its importance to improve the health status of the people [22].

Among respondents who have good knowledge, three-fourth of household use adequately iodized salt. Similarly the study conducted in Ghana respondents those who have good knowledge 73.3% use adequately iodized salt and the study conducted in Gondar shows respondents those who have good knowledge were 1.94 times more likely have adequate iodized salt than those have poor knowledge [23, 19].

Residence of study participants was one of the associated factors with availability of adequately iodized salt at household level. The study participants who were from Urban areas were 8.74 (AOR=8.74, 95% CI: 2.92, 26.09) times more likely to have adequately iodized salt than those who were from rural areas. Similarly a study conducted in Lalo Assabi, Dessie and Kombolcha town indicates that in the urban areas respondents were 1.26 and 2.53 times more likely to have adequately iodized salt than those who were from rural areas respectively [20, 9]. Households whose family size less than five were 3.51 (AOR=3.51, 95% CI: 1.45, 8.26) times more likely to have adequately iodized salt than those households whose family size was greater than five.

Respondents who knew iodine deficiency disorder is preventable were 8.20 (AOR=8.20, 95% CI 3.33, 20.16) times more likely to have adequately iodized salt than those who didn't know.

Respondents who check the iodine content by reading the label on salt was 3.70 (AOR=3.70, 95% CI: (1.28, 10.68) times more likely to have adequately iodized salt than those did not use any method to check the iodine content during salt buy. This difference might be due to educational background of respondents, it also might be due to economic level, knowledge and practice of respondents.

Salt that stored in dry place was 5.38 (AOR=5.38, 95% CI: 1.46, 19.88) times more likely to have adequately iodized salt than that stored at moist place in home. This result was almost similar with a study conducted in Jig Jiga which indicates that salt stored at dry place at household level had 5.15 times adequately iodized salt than that stored at moisture place. But this study result was higher than study conducted in Lalo Assabi which shows that salt stored in dry place was 2.15 times more likely to have adequately iodized salt than the counterpart [21, 20]. The probable reason for this difference might be period of the investigation, geographical location of study areas and educational background of study participants.

On the other hand those household who use container with a cover to store salt was 4.69 (AOR=4.69, 95% CI: 1.64, 13.39) times more likely to have adequately iodized salt than those who store salt in a container with no cover. This study result was greater than a study conducted in Lalo Assabi which shows those who use container with a lid to store their salt at home were 1.45 times more likely to have adequately iodized salt than those who use container other than a lid [20] and it is also greater than a study conducted in Lay may chew

which shows those who use container with a cover to store their salt at home were 2.15 times more likely to have adequately iodized salt than those use container without Cover (26). Those households who were not exposing salt to sunlight or heat were 3.04 (AOR=3.04 95% CI: 1.38, 6.71) times more likely to have adequately iodized salt than those were exposing salt to sunlight or heat. This study finding was less than study conducted in Dire Dawa, Gonder and Dessie Aand Kombolcha which shows 7.26, 7.25 times more likely to have adequately iodized salt than those who expose salt to sun respectively [24, 9]. The reason behind this difference might be due to educational background of study participants.

Similarly duration storage of salt at home was significantly associated with availability of adequately iodized salt. Respondents who store salt for shorter time were 3.75 (AOR=3.75, 95% CI: 1.12, 11.32) times more likely to have adequately iodized salt than those who store for longer period of time. This study result was similar with a study conducted in Dire Dawa [24] which indicates that store salt for less than two month were 3.6 times more likely to have adequately iodized salt than that stored for more than two months; while it was more than a study result conducted in Maychaw [16] which shows 2.1 time more likely to have adequate iodine salt than those store more than two month respectively.

Using packed salt at was significantly associated with availability of adequately iodized salt. A respondents those use packed salt was 11.441 (AOR=11.441, 95% CI: 3.965, 33.014) times more likely to have adequately iodized salt compared to respondents those use non packed salt. A study conducted in Gonder those who use packed salt were 9.7 times more likely to have adequately iodized salt than those who use none packed salt [19]. A study conducted in Lalo Assabi those used packed salt were 1.38 times more likely to have adequately iodized salt than those who use non packed salt [20]. This might be due to good transportation system, storage, and keeping it in a suitable environmental condition.

## 5. Conclusions and Recommendations

### 5.1. Conclusions

Based on finding of this study, availability of adequately iodized salt at house hold was very low compared to IDD elimination strategy. Residence of respondent, family size, Knowledge of respondents about IDD preventable, method use to check salt iodine content, storage place of salt, container used to store salt, exposure of salt to sun or heat, duration salt store and type of salt respondents use were found to be significantly associated with adequacy iodine in edible salt. This showed that working at grass root level is vital to achieve the desired target of universal salt iodization. Educating the household's regarding proper practices of iodized salt, intervention of factors that affect the salt iodine content and proper use of salt in house hold level. It is important to develop regular and strict monitoring schedule on salt supply.

## 5.2. Recommendation

1. Health workers and community-based voluntary health workers should conduct regular community awareness creation and promotion activities to improve universal salt-iodization coverage and practice of iodized salt at house hold level.
2. Zonal health Department and Woreda health office should give attention to monitoring and evaluation of iodine salt at production, transportation and consumption levels.
3. The regulatory authority should appropriately implement legislation of national salt iodization program to ensure sufficient supply of adequately iodized salt throughout the country from the salt dietary production to consumption.
4. Further researches need to be done to identify factors that affect the adequacy of iodized salt at sale level (market, shop, super market).

## Lists of Abbreviations or Acronyms

- a) EDHS: - Ethiopian Demographic and Health Survey.
- b) EFMHACA: - Ethiopian Food, Medicine and Health care Administration and control Agency.
- c) HHIS: - House Hold Iodine Salt.
- d) ICCIDD:-International Center for the Control of Iodine Deficiency Disorders.
- e) IDD: - Iodine Deficiency Disorders.
- f) PPM: - Parts-Per-Million.
- g) RTK: - Rapid Test Kit.
- h) TGR: - Total Goiter Rate.
- i) UI:- Urinary Iodine
- j) UNAIDS:-United States Agency for International Development.
- k) UNICEF: - United Nations Children Emergency Fund.
- l) USI: - Universal Salt Iodization WHO: World Health Organization.

## Declarations

### *Ethics Approval and Consent to Participate*

The necessary explanation about the purpose, potential risk and procedure of study was explained and verbal consent was obtained from each respondent to ensure the right of participant. Participants were informed that participation was on voluntary basis and they can withdraw at any time if they are not comfortable about the questionnaire. In order to protect the confidentiality of information name and other identification was not be included in questioners and maximum effort was made to maintain privacy of the respondent during the interview. The collected data was stored and kept safe place where any the investigator is acceptable and the computer was protected by private pass word.

### *Consent for Publication*

Not Applicable.

## *Availability of Data and Materials*

The finding of this study is generated from the data collected and analyzed based on stated methods and materials. The original data supporting this finding are available from the corresponding author on reasonable request.

## *Competing Interests*

The authors declare that they have no competing interests.

## *Authors' Contributions*

TG participated in the design of the study, performed the data collection and the statistical analysis and served as the corresponding author of the manuscript. ET, RW, ZK, EA and HB supervised the study, ensured quality of the data, assisted in the analysis and interpretation of the data. All authors read and approved the manuscript.

## *Authors' Information*

TG is Head of Walmara Woreda Health Office, Oromia regional state, Ethiopia.

ET is Lecturer at Debre Berhan University, Department of Public Health, Ethiopia.

RW is Maternal and Child Health Program Coordinator, SendafaBeke Town Health Office, Ethiopia.

ZK is Deputy Head of East Wollega Zonal Health Office, Nekemte, Ethiopia.

EA is Ethiopian Public Health Institute, Addis Ababa, Ethiopia.

HB is Lecturer at Wollega University, Institute of Health Science, Department of Public Health, Nekemte, Ethiopia

## Acknowledgements

We are grateful to Walmara Woreda health office staff and administrate for their unfailing support and guidance during the whole activities of this thesis. We also indebted for the study participants for their co-operation without which this study could not have been possible.

## References

- [1] Nutrition Australia. Iodine Facts | Nutrition Australia [Internet]. Nutritian Australia. 2010. Available from: <http://www.nutritionaustralia.org/national/resource/iodine-facts>.
- [2] Paterson J, Crossland K. Micronutrient Malnutrition in the Pacific and Its Prevention. 2000; 43-65. Available from: <http://www.fao.org/tempref/docrep/fao/010/ai026e/AI026E03.pdf>.
- [3] Knowles JM, Garrett GS, Gorstein J, Kupka R, Situma R, Yadav K, et al. Household Coverage with Adequately Iodized Salt Varies Greatly between Countries and by Residence Type and Socioeconomic Status within Countries: Results from 10 National Coverage Surveys. J Nutr. 2017; 147 (5): 1004S-1014S.



- [4] WHO. GUIDELINE: Fortification of food-grade salt with iodine for the prevention and control of iodine deficiency disorders. Who Guidel [Internet]. 2014; (ISBN 978 92 4 150792 9): 1-43. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25473709>.
- [5] Yibrie A, Yifru S, Tigabu D. Module. Iodine Deficiency Disorders. For the Ethiopian Health Center Team. United States Agency for International Development. 2007. 53 p.
- [6] Zimmermann M. Key Barriers to Global Iodine Deficiency Disorder Control: A Summary. 2017; Available from: [https://www.spring-nutrition.org/sites/default/files/a2z\\_materials/508-gap-analysis-iodine-final.pdf](https://www.spring-nutrition.org/sites/default/files/a2z_materials/508-gap-analysis-iodine-final.pdf).
- [7] de Benoist B, Andersson M, Egli I, Takkouche B, Allen H. Iodine status worldwide. WHO Glob Database Iodine Defic Geneva World Heal Organ. 2004; 1-12.
- [8] Ababa A. GUIDELINES FOR THE PREVENTION AND CONTROL OF MICRONUTRIENT DEFICIENCIES in Ethiopia Federal Ministry of Health. Guidel Prev Control Micronutr Defic Ethiop Fed Minist Heal January. 2016; (January).
- [9] Mekonnen TC, Eshete S, Wasihun Y, Arefaynie M, Cherie N. Availability of adequately iodized salt at household level in Dessie and Combolcha Towns, South Wollo, Ethiopia. BMC Public Health. BMC Public Health; 2018; 18 (1): 1-9.
- [10] World Health Organization. Assessment of the iodine deficiency disorders and monitoring their elimination. A guide for programme managers. WHO, Geneva [Internet]. 2007; 107. Available from: [http://www.who.int/nutrition/publications/en/idd\\_assessment\\_monitoring\\_elimination.pdf](http://www.who.int/nutrition/publications/en/idd_assessment_monitoring_elimination.pdf).
- [11] Dessu S, Dawit Z, Alemu G. Assessment of Knowledge on Iodized Salt Utilization and Associated Factors among Households in Arba Minch Town, Southern Ethiopia. Vol. 3, International Journal of Research Studies in Medical and Health Sciences. 2018. 17-23 p.
- [12] WHO. Assessment of the iodine deficiency disorders and monitoring their elimination. WHO, Geneva. 2007; 2: 1-107.
- [13] Andersson M, Takkouche B, Egli I, De Benoist B. The WHO Global Database on iodine deficiency disorders: The importance of monitoring iodine nutrition. Scand J Nutr. 2003; 47 (4): 162-6.
- [14] EDHS. FEDERAL DEMOCRATIC REPUBLIC OF ETHIOPIA Demographic and Health Survey. 2016. 1-59 p.
- [15] Yaye AB, Baraki N, Endale BS. Availability of Adequately Iodized Salt at Household Level and Associated Factors in Dire Dawa, Eastern Ethiopia. Int J Public Heal Sci. 2016; 5 (4): 392.
- [16] BG, KA, AA, MK, YTY, HRS. Availability of Adequate Iodized Salt at Household Level and Associated Factors in Rural Communities in Laelay Maychew District, Northern Ethiopia: A Cross Sectional Study. J Nutr Heal Sci. 2015; 1 (4): 1-9.
- [17] WHO/NMH/NHD/EPG/14.5. Goitre as a determinant of the prevalence and severity of iodine deficiency disorders in populations. WHO/NMH/NHD/EPG/145 VMNIS [Internet]. 2014; 14-9. Available from: [http://apps.who.int/iris/bitstream/10665/133706/1/WHO\\_NM\\_H\\_NHD\\_EPG\\_14.5\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/133706/1/WHO_NM_H_NHD_EPG_14.5_eng.pdf).
- [18] Town T. Availability and Knowledge of Iodized Salt at Household Level and Associated Factors. 2019; 6 (1): 1-8.
- [19] Gebremariam HG, Yesuf ME, Koye DN. Availability of Adequately Iodized Salt at Household Level and Associated Factors in Gondar Town, Northwest Ethiopia. 2013 p. 1-6.
- [20] Regassa D M, H TW. Utilization of Adequately Iodized Salt on Prevention of Iodine Deficiency Disorders at Household Level and Associated Factors in Lalo Assabi District, West Ethiopia. J Nutr Food Sci. 2016; 06 (02).
- [21] Ahmed AT. Use of Iodized Salt at Household Level in Jig Jiga Town, Eastern Ethiopia. 2019; (April).
- [22] Bazezew MM, Yallew WW, Belew AK. Knowledge and practice of iodized salt utilization among reproductive women in Addis Ababa City. BMC Res Notes [Internet]. BioMed Central; 2018; 11 (1): 1-8. Available from: <https://doi.org/10.1186/s13104-018-3847-y>.
- [23] Sarah NA, Prince AK, Yao AS, Geoffrey AA, Wisdom TK, Margaret K. Knowledge on Iodized Salt Use and Iodine Content of Salt Among Households in the Hohoe Municipality, Volta Region - Ghana To cite this article : 2016; 2 (1): 1-10.
- [24] Ftwi G, Mengistie B, Abdo M, Roba KT. Household Salt Iodine Level and Associated Factors in Dire Dawa City Administration, Eastern Ethiopia. East African J Heal Biomed Sci. 2018; 2: 35-44.