



Impact of Nitrogen Fertilizer Use on Soil Ecosystem Integrity in the Greater Port Harcourt Region, Nigeria

Wanjala Muyoma Paul^{1, 5, *}, Odokuma Lucky¹, Etela Ibisime², Ramkat Rose³,
Blessing Adanta Odogwu⁴, Boadu Kwasi Opoku^{6, *}, Koranteng-Addo Epton Joseph⁶

¹Department of Microbiology, University of Port Harcourt, Port Harcourt, Nigeria

²Department of Animal Science, University of Port Harcourt, Port Harcourt, Nigeria

³Centre of Excellence in Phytochemicals, Textiles and Renewable Energy, Moi University, Eldoret, Kenya

⁴Department of Plant Science and Biochemistry, University of Port Harcourt, Port Harcourt, Nigeria

⁵Department of Physical and Biological Sciences, Bomet University College, Bomet, Kenya

⁶Department of Chemistry, University of Cape Coast, Cape Coast, Ghana

Email address:

piwanjala@gmail.com (Wanjala M. P.), koboadu@ucc.edu.gh (Boadu K. O.), Lucky.Odokuma@uniport.edu.ng (Odokuma L.),
ibisime.etela@uniport.edu.ng (Etela I.), chirirose@yahoo.com (Ramkat R.), blessing.odogwu@uniport.edu.ng (Blessing A. O.),
ekoranteng-addo@ucc.edu.gh (Koranteng-Addo E. J.)

*Corresponding author

To cite this article:

Wanjala Muyoma Paul, Odokuma Lucky, Etela Ibisime, Ramkat Rose, Blessing Adanta Odogwu, Boadu Kwasi Opoku, Koranteng-Addo Epton Joseph. Impact of Nitrogen Fertilizer Use on Soil Ecosystem Integrity in the Greater Port Harcourt Region, Nigeria. *International Journal of Biochemistry, Biophysics & Molecular Biology*. Vol. 6, No. 2, 2021, pp. 42-51. doi: 10.11648/j.ijbbmb.20210602.13

Received: June 17, 2021; Accepted: June 29, 2021; Published: October 5, 2021

Abstract: Fertilizers are used to increase crop yields in agriculture but can affect the integrity of soil ecosystems if misused. The present study was conducted to observe the impact of inorganic fertilizers in selected soils in Greater Port Harcourt Area on soil microbial population, in addition, the association between soil microbial population and Lethal Concentration 50 (LC₅₀) was observed. Microbial communities and LC₅₀ values were determined using normal analytical procedures. The study showed various fertilizer types that altered soil microbial population and LC₅₀ values. For agricultural areas the maximum NPKB mean value was 0.70±0.76 ppm which was higher than the mean value of the control location of 0.23±0.00 ppm. Urea tolerant fungi's (UREAF's) LC₅₀ mean value was 0.43±0.09 ppm in the agricultural area as compared to the control with a mean value of 0.46±0.00 ppm. Industrial and urban areas had significantly different mean values of 0.29 ± 0.19 ppm and 0.18±0.01 ppm and 0.23 ± 0.06 ppm and 0.23 ± 0.060.17 ± 0.010.17 ± 0.010.17 ± 0.01 ppm respectively (p = 0.001). This research concluded that the application of UREA and NPK fertilizers in study area soils lowered the pH of soils in A1, A2 and A3 agricultural sites. Application of UREA and NPK fertilizers in soils of the study areas increased the population of microorganisms in threshold concentrations but was detrimental for NPK and UREA fertilizers at concentrations of 1.50% or above. Further findings showed that microorganisms were more tolerant of NPK and UREA fertilizers in agricultural areas compared to urban and industrial areas which showed higher LC₅₀ values. This research concludes that fertilizers used in the study areas were related to soil microorganisms developing tolerance to NPK and UREA. Farmers should observe application of the recommended professional quantities of fertilizers to ensure the sustenance of the soil ecosystem integrity.

Keywords: Inorganic Fertilizers, Misuse, Impact, Soil Microbial Population, LC₅₀, Soil Ecosystem Integrity

1. Introduction

Fertilizers are used to increase crop yields in agriculture but may affect the integrity of soil ecosystems if misused [1]. The potential of various soil types to lose nutrients and

contaminants is a major factor impacting soil's ability to lose fertilizers [2]. In a situation of inorganic and organic phosphorus (P), this could be more closely linked to aluminium (Al), calcium (Ca) and iron (Fe) and cation exchange capacity for potassium and ammonium [3, 4].

Consequently, this may minimize nutrient loss to water and air, thereby reducing nutrient availability for crops [2]. In addition, soil fertilizer value depends on several reasons; the ratio and nature of soil organic and mineral components [5], the duration of the seasonal environment that determines crop uptake, weather conditions and application history [6]. Such nutrient variation caused by the introduction and erosion of inorganic fertilizers can impact the composition and population of soil microorganisms. A study at Ebony State, Nigeria, has shown that inorganic fertilizers contribute to the spatial distribution of soil microorganisms [7, 8]. Also, Xie *et al.*, [9], reported a decrease in the microbial population as soil contaminant levels. Cation levels in soil have also been attributed to influencing bacterial growth. A research by Hai-Hang, Zhang and Pan [10] showed that Al^{2+} , Ca^{2+} , Cu^{2+} , Zn^{2+} , Mn^{2+} , Mg^{2+} concentrations in the environment affect population and diversity of *Rhodotorula glutinins* and *Pseudomonas spp.* Organic matter levels, soil moisture, soil structure, and soil texture also affect hydrology and influence soil mineralization through influence on soil microorganism activity [11, 12]. The content of clay in soil plays a significant role in protecting organic matter from microbial degradation [13].

During crop production, application of fertilizers in soil affects soil microorganisms which are a cognitive indicator of soil health [14]. Long-term experiments show that mineral fertilizers increase the population of microorganisms relative to unfertilized soils, as opposed to grassland habitats where the population of microorganisms is decreased by an

increased N input [11]. With increased fertilisation using mineral fertilizers, the biomass of bacteria, actinomycetes and fungi increases. Fertilizers affect the bacteria to fungi ratio and even increase the Gram-positive to Gram-negative bacteria ratio [14]. Studies of the impact of fertilizers on changes in populations and communities of microorganisms remain inconsistent, as some studies report growing soil microorganisms with increased levels of fertilizers, while others report decreasing soil microorganism populations with increased concentrations of fertilizers in growth media [15, 16, 17], which are attributed to common site variables including soil and climatic properties [14].

Fertilization affects the composition of the microorganisms in the community through changes in soil properties that depend on the initial soil characteristics. The impact of application of fertilizer strongly depends on environmental factors that are considerably variable among different sites [14]. Therefore, research on the effect of fertilizer application on the population of microorganisms and diversity shifts in polluted environments is needed in order to have a clear understanding of the effects of fertilizers on the distribution of microorganisms in the soil. The scope of this research adds to the non-conclusive debate about the consequences of misuse of fertilizers in agricultural fields. Farmers must be aware of the impacts of fertilizer overuse or misuse on agricultural expanses. The results are essential for stakeholders and for decision taking to mitigate and restore deteriorated soils in the future.




Figure 1. Location of sampling sites in selected areas in Greater Port Harcourt area, Rivers State, Nigeria.

Table 1. GPS coordinates and economic activities for sampling sites in selected areas in Greater Port Harcourt area, Rivers State, Nigeria.

No	Selected study sites	Study area coding	Coordinates/ N latitude E longitude	Economic activities
Agricultural area				
1	Aluu	A1	4° 56' 11.160' 6° 57' 52.248'	Flow station
2	Eleme	A2	4° 44' 09.874' 7° 08' 58.494'	Village close to refinery
3	Emuoha	A3	5° 00' 00.018' 6° 49' 13.032'	Flow station
4	Control	CA	5° 00' 21.384' 6° 49' 00.000'	>1 km away from suspected areas
Industrial area				
1	Onne	I1	4° 46' 00.402' 7° 05' 43.092'	Hosts the NNPC Refinery
2	Agbada	I2	4° 56' 03.444' 6° 58' 42.060'	Hosts SPDC- flow station in a rural setting
3	Trans-Amadi	I3	4° 48' 20.455' 7° 02' 17.646'	Schlumberger/, Halliburton
4	Control	CI	4° 47' 13.788' 7° 07' 44.620'	>1 km away from suspected areas
Urban area				
1	GRA phase 2	U1	4° 49' 53.574' 6° 59' 45.552'	Inhabited areas Perecuma street
2	Diobu- Mile 1	U2	4° 47' 20.382' 7° 00' 13.164'	Petroleum refinery
3	Mguoba	U3	4° 50' 39.864' 6° 58' 20.232'	NTA
4	Control	CU	4° 49' 17.040' 6° 59' 24.168'	>1 km away from suspected areas

2. Material and Methods

2.1. Description of Study Site

This research was carried out at 9 selected test sites and 3 control sites in Port Harcourt, Nigeria's capital of Rivers State (Figure 1). The sites of the study were divided into three areas: urban (GRA phase 2, Diobu- Mile 1 and Mguoba), industrial (Eleme hosting the NNPC refinery, Agbada-SPDC- flow station) and agricultural (Aluu, Oquwi- Eleme, Emuoha- Eu). The sites of the study were distinguished by different economic activities, as shown in Table 1.

2.2. Sampling

In the wet season (April to October 2018), composite samples were collected through random sampling from each of the three areas; urban, industrial, and agricultural. Five (5) separate samples were collected randomly around each test field. The five individual samples were thoroughly mixed by coning and quartering in a sterile container to attain a homogenous composite mixture. A total of 12 composite samples; A1, A2, A3, I1, I2, I3 U1, U2 and U3 as test samples, and CA, CI and CU as control samples (Table 1) were collected from the top soil at depths of 0 to 15 cm using a standard auger 3 times during the rainy season. Homogenized composite samples (400 gm) were then packaged using a sterile wooden shovel into polyethylene bags. Samples were collected for microbial analysis using pre-sterilized materials to prevent contamination of the samples. Sampling locations were identified using a GPS and the recorded GPS readings. The samples were transported to the

laboratory for analysis.

2.3. Laboratory Analysis

2.3.1. Enumeration of Fertilizer Tolerant Bacteria

Pour plate method for enumeration of microbes was used to culture fertilizer-tolerant bacteria [18]. Under aseptic conditions (in a laminar flow cabinet) one gram of soil sample was weighed into a 9 ml sterile diluent (0.85% NaCl). The sample was then homogenized using a laboratory vortex mixer (Model 10101001, IP42) and serially diluted using sterile pipettes. Thereafter, 0.1 ml of inoculum aliquot was inoculated on mineral salt agar (MSA) that was then mixed with antifungal reagent (Fungin™) to inhibit fungal growth. Fixed dose procedure was used in which microorganisms were dosed in a step-by-step procedure using 0%, 1%, 5%, 10%, 15%, 20%, 25%, 30% and 60% NPK and ammonia fertilizer concentrations. Based on sighting study, the very first dose levels were selected as the dose projected to show toxicity effects, without causing mortality. Depending on the presence or absence of toxicity or mortality signs, microorganisms were then dosed in the maximum or least set dosages. The dose continued until the dose was established / seen to cause apparent toxicity or no more than one death, or if there were no symptoms at the maximum or minimum dose or death. Plates were then incubated in an inverted position at room temperature (28°C) for 5 to 7 days. Colonies were counted so that units forming colony could be obtained per gram of soil.

2.3.2. Enumeration of Fertilizer Tolerant Fungi

Pour plate method [18] was used to enumerate fertilizer-tolerant fungi. In this method, under aseptic conditions, 1 g of

soil sample was weighed into a 9 ml sterile diluent (0.85% NaCl). The sample was then homogenized using a vortex mixer (Model 10101001, IP42) and diluted in series using sterile pipettes. The Potato Dextrose Agar (PDA) was then inoculated in zero-point one (0.1 ml) aliquot of inoculum. A fixed dose procedure was used in which micro-organisms were dosed in a step-by-step procedure using fixed doses of [0%, 0.10%, 0.25%, 0.75%, 1%, 1.25%, 1.50%, and 1.75%] for NPK and UREA fertilizers. The initial dose levels were selected as the dose expected to produce toxicity effects, without causing mortality, based on sighting study. Microorganisms were then dosed at fixed doses higher or lower, depending on the presence or absence of toxicity or mortality signs. The dosage lasted until the dosage was confirmed to cause significant toxicity or no more than one death, or no signs were observed at the highest dose, or deaths occurred at the lowest dose. Plates were then incubated in an inverted position at room temperature (28 ° C) for 7 days. Colonies were counted using a colony counter to get units forming colony per gram of soil.

2.3.3. Determination of LC_{50} for Bacteria and Fungi

Percentages of dead organisms were determined at each concentration of the study and converted to probits [19] (Finney's Table). Regression analysis was conducted, where the probit analysis output was used to compare the amount of chemical required to create responses between micro-organisms from different areas of study to different concentrations of Ni / Cd in the culture medium [20]. Areas with lower LC_{50} values were found to be most toxic compared with areas with higher LC_{50} values [20]. Microorganisms were considered to be more tolerant to heavy metal exposure in areas with higher LC_{50} values. Lethal Concentration 50 (LC_{50}) was determined by calculating the corresponding x value for a 5.00 probit and then taking the inverse log of the associated concentration [20].

$$Y = ax + c \quad (1)$$

where y = From Finney's Table [19]

a = Calculated coefficients

x = Unknown value

c = Calculated coefficients

LC_{50} = Antilog of x

2.3.4. Data Analysis

Data analysis was conducted using the statistical software IBM SPSS Statistics for Windows, version 24.0. For multiple comparisons between the study areas and between seasons

(wet and dry seasons), data obtained from laboratory analysis was analyzed using ANOVA. Analysis of the correlation was used to check the relationship between chemical variables in the soils and the microbe population. Each of the data collected in the study were analyzed at value levels $p < 0.05$.

3. Results

3.1. UREA and NPK Tolerant Microorganisms

The values given in Tables 2 and 3 show the mean values (LC_{50}) of three independent tests. Lethal concentration 50 (LC_{50}) levels for each study site were determined and compared with LC_{50} levels for control sites. Findings indicate variability in bacterial tolerance to various concentrations of fertilizers UREA and NPK. The mean value of LC_{50} for NPKF in agricultural area was 0.20 ± 0.20 ppm which was lower than the control value of 0.47 ± 0.15 ppm. Lethal concentration 50 (LC_{50}) for NPKF was lower compared to the control samples in industrial and urban areas respectively (Figure 2). The LC_{50} for NPKB control was 0.66 ± 0.00 ppm higher than the 0.18 ± 0.22 ppm test sample. In agricultural areas the highest NPKB mean value was 0.70 ± 0.76 which was higher than the mean value of the control location (0.23 ± 0.00 ppm) (Table 2). The lowest lethal concentration (LC_{50}) in industrial area for NPKB showed a mean of 0.34 ± 0.15 ppm with the control sample being 0.24 ± 0.00 ppm. The mean LC_{50} for UREAF in the agricultural area was 0.43 ± 0.09 ppm, with a control mean of 0.46 ± 0.00 ppm. Similarly, Industrial had a mean of 0.29 ± 0.19 ppm with the control being 0.18 ± 0.01 ppm, respectively. The urban areas also had mean of 0.23 ± 0.06 ppm with the control having a mean of 0.17 ± 0.01 ppm respectively. All of which were significantly different ($p = 0.001$) {Table 2}. Test sites demonstrated greater tolerance for UREA compared with industrial and agricultural control samples. In the agricultural area, UREAB's lethal concentration 50 (LC_{50}) had a mean of 0.26 ± 0.21 ppm with a control having a mean of 0.64 ± 0.02 ppm. Urea tolerant bacteria (UREAB's) control mean was higher than that of the test-value (Table 2). The UREAB values for industrial and urban areas were 0.25 ± 0.14 ppm and 0.04 ± 0.01 ppm, and 0.49 ± 0.63 ppm and 0.12 ± 0.01 ppm, respectively (Table 2). No significant variations in LC_{50} levels for UREA ($p = 0.185$) and NPK ($p = 0.131$) for bacteria were observed in all study sites (Table 2).

Table 2. Variation of LC_{50} for fertilizer in fungi and bacteria in soil from agricultural, industrial and urban areas in Greater Port Harcourt area, Rivers State, Nigeria.

Study area	NPKF	NPKB	UREAF	UREAB
Agriculture	0.20 ± 0.20^a	0.70 ± 0.76^a	0.43 ± 0.09^{bc}	0.26 ± 0.21^{ab}
Control Agriculture	0.47 ± 0.15^b	0.23 ± 0.00^a	0.46 ± 0.00^c	0.64 ± 0.02^b
Industry	0.14 ± 0.14^a	0.34 ± 0.15^a	0.29 ± 0.19^{ab}	0.25 ± 0.14^{ab}
Control Industry	0.03 ± 0.03^a	0.24 ± 0.00^a	0.18 ± 0.01^a	0.04 ± 0.01^a
Urban	0.07 ± 0.09^a	0.18 ± 0.22^a	0.23 ± 0.06^a	0.49 ± 0.63^{ab}
Control Urban	0.06 ± 0.04^a	0.66 ± 0.00^a	0.17 ± 0.01^a	0.12 ± 0.01^{ab}
P Value	0.004	0.131	0.001	0.185

Data are Mean \pm standard deviations (Significant coefficients $P = 0.05$). Different letters show that there is significant difference between the mean values.

3.2. Comparison of NPK and UREA Tolerance Among Microorganisms in Study and Control Areas

Figures 2 (a) and (b) compares the mean variation of LC₅₀ for NPK tolerant fungi and bacteria to UREA. In the agricultural areas, NPK tolerant bacteria were higher compared with fungi (Figure 2a). Control samples in industrial areas showed lower LC₅₀ in both bacteria and fungi, respectively, compared to test areas. NPK recorded a

higher LC₅₀ value for bacteria as compared with fungi in urban areas (Figure 2a). The agriculture areas showed the highest mean value of LC₅₀ for NPK in fungi and bacteria. Figure 2b shows higher UREA tolerance values for fungi and bacteria in agricultural areas compared with industrial and urban areas, with the exception of urban areas where urea-tolerant bacteria are higher than those in agricultural and industrial areas (Figure 2).




Figure 2. Variation in LC₅₀ for NPK (a) and UREA (b) in fungi and bacteria in soil from agricultural, industrial and urban areas in Greater Port Harcourt Area, Rivers State, Nigeria.

3.3. Mean Concentrations of LC₅₀ for NPK and UREA Tolerant Fungi and Bacteria in the Study Sites

Table 3 shows the mean variability for NPK and UREA resistant fungi and bacteria at different test and control sites. The highest NPKF values were 0.30±0.129, 0.47±0.151 and 0.44±0.151 ppm respectively at I2, CA and A3. The highest UREA tolerant fungi LC₅₀ values were 0.54±0.01, 0.46±0.00, 0.54±0.05 and 0.40±0.00 ppm, respectively, recorded in I2, CA, A2 and A3. The highest mean values for NPK tolerant

bacteria were recorded with mean concentrations of 0.51±0.00, 1.69±0.05, 0.66±0.00 and 0.48±0.00 respectively in I1, A1, CU and U2. Highest LC₅₀ values were observed for UREA-tolerant bacteria in I3, CA, A1 and U2 which had values of 0.43±0.001, 0.64±0.02, 0.48±0.07 and 1.33±0.001 ppm. Agriculture sites showed the highest LC₅₀ frequencies (Table 3). The LC₅₀ values were significantly different in the study sites ($p \leq 0.000$). Mean pH values ranged from 5.22 through 7.96. Agricultural areas demonstrated the lowest pH of 5.22.

Table 3. Variation of LC₅₀ for fertilizer in fungi and bacteria in soil from agricultural, industrial and urban areas in Greater Port Harcourt Area, Rivers State, Nigeria.

	NPKF	UREAF	NPKB	UREAB	Soil pH -log ₁₀ [(ΣC _i)/(n)]
C1	0.031±0.034 ^{ab}	0.183±0.006 ^b	0.240±0.000 ^c	0.035±0.005 ^{ab}	7.96
I1	0.088±0.063 ^{ab}	0.150±0.000 ^a	0.510±0.000 ^b	0.144±0.022 ^{de}	7.28
I2	0.301±0.129 ^{cd}	0.543±0.012 ^h	0.157±0.029 ^c	0.188±0.053 ^c	6.49
I3	0.039±0.036 ^{ab}	0.180±0.000 ^b	0.360±0.000 ^d	0.430±0.001 ^e	7.47
CA	0.466±0.151 ^e	0.460±0.000 ^g	0.230±0.000 ^c	0.640±0.017 ⁱ	7.47
A1	0.127±0.081 ^{ab}	0.350±0.000 ^c	1.690±0.053 ^g	0.481±0.071 ^h	6.12
A2	0.027±0.028 ^{ab}	0.543±0.046 ^h	0.034±0.028 ^a	0.310±0.001 ^f	5.22
A3	0.436±0.151 ^{de}	0.400±0.000 ^f	0.367±0.012 ^d	0.001±0.000 ^a	5.89
CU	0.064±0.040 ^{ab}	0.173±0.006 ^{ab}	0.660±0.000 ^f	0.118±0.003 ^{cd}	7.32
U1	0.176±0.090 ^{bc}	0.233±0.006 ^c	0.020±0.000 ^a	0.062±0.003 ^b	7.12
U2	0.019±0.019 ^{ab}	0.290±0.000 ^d	0.480±0.000 ^c	1.330±0.001 ^j	7.52
U3	0.008±0.010 ^a	0.160±0.000 ^{ab}	0.050±0.000 ^a	0.076±0.005 ^{bc}	7.10
P Value	0.000	0.000	0.000	0.000	

Data are Mean ± standard deviations (Significant coefficients P=0.05). Different letters show that there was significant difference between the mean values.

3.4. Bacteria Population in Different Concentrations of UREA and NPK Fertilizers

The values given in Table 4 presents the means of three independent tests. The reductions in bacteria actively replicating and forming colonies as opposed to control indicated a decline in soil CFUs / g with an increase in UREA concentrations. There were variations in the study areas of maximum tolerable concentrations between bacteria

ranging from 0.1% to 1.75% UREA concentrations. Sites I3, CA, A3, U2 and U3 showed growth at a maximum concentration of 1.75%, ranging between 1.20×10^3 and 5.47×10^2 CFUs / g of soil, whereas CI, I1, I2, A1, CU and U1 showed growth at a maximum concentration of 1.50% of UREA, ranging between 1.03×10^3 and 6.60 CFUs / g of soil (Table 4). The table also represents mean values of LC_{50} calculated from values from three independent tests.

Table 4. Variation of bacterial populations in increasing concentrations of UREA fertilizer in soil from agricultural, industrial and urban areas in Greater Port Harcourt Area, Rivers State, Nigeria.

Sample	0%	0.25%	0.5%	0.75%	1%	1.25%	1.50%	1.75%	2.0%	LC_{50}
CI	7.17×10^7	5.73×10^6	4.17×10^5	3.27×10^4	3.13×10^3	2.43×10^3	1.13×10^3	0	0	0.04
I1	1.37×10^7	3.37×10^6	2.03×10^5	1.93×10^4	1.77×10^3	1.10×10^3	1.03×10^2	0	0	0.14
I2	7.43×10^6	2.26×10^6	3.12×10^5	4.03×10^4	3.67×10^3	2.37×10^3	1.27×10^3	0	0	0.19
I3	3.37×10^7	5.40×10^6	4.80×10^5	6.97×10^4	6.73×10^3	2.53×10^3	2.00×10^3	1.30×10^3	0	0.43
CA	6.17×10^7	1.15×10^7	9.53×10^5	2.26×10^4	1.77×10^3	1.50×10^3	6.87×10^2	5.47×10^2	0	0.64
A1	1.30×10^6	4.07×10^5	1.72×10^5	2.19×10^4	1.78×10^3	1.17×10^3	6.60×10^2	0	0	0.48
A2	7.07×10^7	1.77×10^7	3.63×10^6	6.90×10^4	6.67×10^3	6.40×10^3	5.90×10^3	3.73×10^3	0	0.31
A3	5.73×10^7	3.07×10^6	2.73×10^5	2.30×10^4	1.65×10^3	1.40×10^3	1.10×10^3	1.00×10^3	0	0
CU	3.33×10^6	1.34×10^6	2.73×10^5	2.42×10^4	2.07×10^3	1.60×10^3	1.03×10^3	0	0	0.12
U1	4.57×10^7	7.60×10^6	5.67×10^5	4.57×10^4	3.33×10^3	2.53×10^3	1.53×10^3	0	0	0.06
U2	1.87×10^7	5.07×10^6	1.03×10^6	3.70×10^4	2.67×10^3	1.90×10^3	1.73×10^3	1.47×10^3	0	1.33
U3	1.93×10^7	3.20×10^6	4.17×10^5	7.03×10^4	3.60×10^3	3.30×10^3	2.07×10^3	1.20×10^3	0	0.08

The values given in Table 5 shows the mean of three independent tests. The decreases in bacterial actively replicating and forming colonies as compared to control showed a reduction in soil CFUs / g with an increase in NPK fertilizer concentrations. There were differences in the study areas of maximum tolerable concentrations among bacteria ranging from 0.25% to 1.75% NPK concentrations. Sites CI, I2, I3, CA, A2, A3, CU, U1, U2 and U3 showed growth at a

maximum concentration of 1.50%, ranging between 1.73×10^3 and 4.57×10^3 CFUs / g of soil, whereas I1 and A1 reported growth at a maximum concentration of 1.25% of NPK fertilizers ranging between 1.23×10^3 and 2.10×10^3 CFUs / g of soil (Table 5). Furthermore, Table 5 represents mean LC_{50} values of 0% NPK fertilizers calculated from three independent test values.

Table 5. Variation of bacterial populations in increasing concentrations of NPK fertilizer in soil from agricultural, industrial and urban areas in Greater Port Harcourt Area, Rivers State, Nigeria.

Sample	0%	0.1%	0.25%	0.50%	0.75%	1.0%	1.25%	1.5%	1.75%	LC_{50}
CI	7.17×10^7	1.95×10^7	1.21×10^6	2.03×10^4	2.48×10^3	2.40×10^3	2.27×10^3	2.10×10^3	0	0.24
I1	1.37×10^7	5.20×10^6	3.37×10^5	2.87×10^4	2.37×10^3	1.77×10^3	1.67×10^3	0	0	0.51
I2	7.43×10^6	2.42×10^6	2.00×10^5	3.60×10^4	3.07×10^3	2.77×10^3	2.57×10^3	2.43×10^3	0	0.16
I3	3.37×10^7	8.47×10^6	6.17×10^5	5.17×10^4	4.50×10^3	4.23×10^3	4.00×10^3	3.50×10^3	0	0.36
CA	6.17×10^7	1.85×10^7	1.07×10^6	1.87×10^4	3.07×10^3	2.73×10^3	2.47×10^3	2.40×10^3	0	0.23
A1	1.30×10^6	7.03×10^5	3.57×10^5	2.20×10^4	2.03×10^3	1.57×10^3	1.23×10^3	0	0	1.69
A2	7.07×10^7	5.77×10^6	4.70×10^5	3.00×10^4	2.77×10^3	2.37×10^3	1.97×10^3	1.73×10^3	0	0.03
A3	5.73×10^7	1.03×10^7	2.80×10^5	2.20×10^4	3.70×10^3	3.50×10^3	3.47×10^3	3.27×10^3	0	0.37
CU	3.33×10^6	5.20×10^5	1.31×10^5	3.10×10^4	2.77×10^3	2.63×10^3	2.47×10^3	2.20×10^3	0	0.66
U1	4.57×10^7	5.80×10^6	5.03×10^5	4.10×10^4	3.60×10^3	3.37×10^3	3.17×10^3	2.90×10^3	0	0.02
U2	1.87×10^7	3.83×10^6	2.60×10^5	1.80×10^4	5.47×10^3	4.90×10^3	4.70×10^3	4.57×10^3	0	0.48
U3	1.93×10^7	2.19×10^6	3.03×10^5	2.77×10^4	2.77×10^3	2.43×10^3	2.03×10^3	2.00×10^3	0	0.05

3.5. Fungal Population in Increasing Concentrations of UREA and NPK Fertilizers

The values given in Table 6 show the means of three independent tests. The declines in fungal actively replicating and forming colonies relative to control showed a decline in soil CFUs / g with an increase in UREA fertilizer concentrations. There were variations in the study areas between 0.25% and 1.75% of UREA concentrations in the

maximum tolerable concentrations among the fungi. Sites CI, I1, I2, I3, CA, A1, A2, CU, U1, U2 and U3 showed growth at a maximum concentration of 1.75% with a count of 2.14×10^3 CFUs / g of soil. Additionally, Table 6. shows mean LC_{50} values of 0% NPK fertilizers calculated from three independent test values.

The values given in Table 7 represent the means of three independent tests. The decreases in fungal actively replicating and forming colonies as compared to the control showed a reduction in soil CFUs / g of soil with an increase in NPK

fertilizer concentrations. There were differences in the study areas between 0.10% and 1.75% for UREA concentrations in the maximum tolerable concentrations among fungi. Sites CI, I1, I2, I3, CA, A1, A2, A3, CU, U1, U2 and U3 showed growth at a maximum concentration of 1.75% with counts ranging

between 3.33×10^1 to 1.03×10^2 CFUs / g soil. There was no variation in the maximum NPK fertilizer concentration which showed zero fungi growth. Furthermore, Table 7. reflects mean LC_{50} values of percent NPK fertilizers determined from three independent test values.

Table 6. Variation in fungal populations in increasing concentrations of UREA fertilizer in soil from agricultural, industrial and urban areas in Greater Port Harcourt Area, Rivers State, Nigeria.

Sample	0.0%	0.25%	0.50%	0.75%	1.0%	1.25%	1.50%	1.75%	LC 50
CI	5.20×10^4	3.37×10^4	2.27×10^4	3.47×10^3	2.38×10^3	2.10×10^3	1.54×10^3	0	0.18
I1	6.17×10^5	4.50×10^4	2.68×10^4	3.43×10^3	2.67×10^3	1.81×10^3	1.23×10^3	0	0.15
I2	1.32×10^4	1.09×10^4	6.23×10^3	6.00×10^3	5.77×10^3	3.38×10^3	2.44×10^3	0	0.54
I3	7.50×10^4	3.57×10^4	5.93×10^3	1.71×10^3	1.34×10^3	1.20×10^3	1.09×10^3	0	0.18
CA	6.17×10^3	4.60×10^3	3.27×10^3	1.82×10^3	1.55×10^3	1.28×10^3	1.13×10^3	0	0.46
A1	1.45×10^5	1.13×10^5	6.33×10^4	3.10×10^4	2.01×10^3	1.36×10^3	1.18×10^3	0	0.35
A2	3.27×10^4	3.20×10^4	2.44×10^4	1.19×10^4	1.04×10^4	1.85×10^3	1.09×10^3	0	0.54
A3	8.23×10^3	5.53×10^3	4.57×10^3	2.49×10^3	2.14×10^3	0	0	0	0.4
CU	7.70×10^4	4.00×10^4	6.83×10^3	3.57×10^3	2.74×10^3	2.01×10^3	1.21×10^3	0	0.17
U1	1.44×10^4	5.67×10^3	3.57×10^3	2.61×10^3	6.43×10^2	4.30×10^2	2.37×10^2	0	0.23
U2	6.60×10^4	4.60×10^4	2.21×10^4	7.23×10^3	5.53×10^3	5.23×10^3	1.31×10^3	0	0.29
U3	6.23×10^4	3.60×10^4	5.03×10^3	2.90×10^3	1.57×10^3	1.16×10^3	2.97×10^2	0	0.16

Table 7. Variation in fungal populations in increasing concentrations of NPK fertilizer in soil from agricultural, industrial and urban areas in Greater Port Harcourt Area, Rivers State, Nigeria.

Sample	0%	0.10%	0.25%	0.50%	0.75%	1%	1.25%	1.50%	1.75%	LC ₅₀
CI	5.20×10^4	2.18×10^4	7.17×10^3	6.90×10^3	6.47×10^3	5.87×10^3	4.27×10^3	1.41×10^3	8.33×10^1	0.03
I1	6.17×10^5	9.90×10^4	2.54×10^4	6.83×10^3	6.23×10^3	6.00×10^3	5.17×10^3	1.48×10^3	6.67×10^1	0.09
I2	1.32×10^4	9.20×10^3	6.80×10^3	6.37×10^3	5.73×10^3	4.90×10^3	3.69×10^3	8.34×10^2	4.00×10^1	0.3
I3	7.50×10^4	5.87×10^4	8.37×10^3	5.01×10^3	3.87×10^3	2.99×10^3	2.15×10^3	8.08×10^2	7.67×10^1	0.04
CA	6.17×10^3	5.40×10^3	4.73×10^3	3.58×10^3	2.78×10^3	1.71×10^3	1.19×10^3	4.11×10^2	3.67×10^1	0.47
A1	1.43×10^5	9.23×10^4	3.08×10^4	4.88×10^3	2.91×10^3	2.28×10^3	1.63×10^3	4.93×10^2	4.67×10^1	0.13
A2	3.27×10^4	1.51×10^4	5.07×10^3	4.53×10^3	4.05×10^3	3.63×10^3	2.75×10^3	9.42×10^2	9.33×10^1	0.03
A3	8.23×10^3	6.67×10^3	5.48×10^3	4.86×10^3	3.14×10^3	2.28×10^3	1.49×10^3	4.04×10^2	3.33×10^1	0.44
CU	7.70×10^4	5.56×10^4	1.17×10^4	6.07×10^3	4.57×10^3	4.17×10^3	2.93×10^3	7.49×10^2	3.67×10^1	0.06
U1	1.44×10^4	8.77×10^3	5.50×10^3	4.12×10^3	2.95×10^3	1.95×10^3	1.34×10^3	5.39×10^2	7.00×10^1	0.18
U2	6.60×10^4	2.62×10^4	5.10×10^3	4.58×10^3	3.57×10^3	2.62×10^3	2.15×10^3	7.45×10^2	3.33×10^1	0.02
U3	6.23×10^4	2.55×10^4	5.40×10^3	4.70×10^3	4.20×10^3	3.63×10^3	2.68×10^3	1.05×10^3	1.03×10^2	0.01

4. Discussion

Inorganic fertilizers are important in the green revolution and growing farm yields, but misuse of fertilizers can result in a loss of the soil ecosystem integrity. Achieving food security is a key agenda that eludes Governments in sub-Saharan Africa [SSA] [21]. Low food crop productivity due to reduced organic fertilizers is one of the main contributors to SSA food scarcity in a region that has also experienced land degradation from oil spills for several decades [21], as well as post-harvest losses and unequal distribution of food. There is a rise in fertilizer use in countries that provide input subsidies, such as Kenya, Nigeria, Tanzania and Malawi [22], which is expected to increase further in the years ahead. During several years past, application rates for fertilizers were based on blanket guidelines [23]. Data that can help determine the correct levels of fertilizer and application for the specific plants and sites is crucial for improving efficiency of fertilizer use and minimizing negative environmental impacts.

Soil micro-organisms play a pivotal role in the ecosystem, including the bioremediation of contaminated sites, nutrient

cycling and the promotion of plant growth. Therefore, in terms of diversity and population, optimum levels of micro-organisms in soil need to be maintained for sustainable agriculture [24, 25, 26]. Changes in soil parameters are followed by differences in microorganism group composition and function [27]. Hence it is necessary to understand the population and diversity of microbial soil communities in order to enforce soil restoration. In this research, fungal and bacterial populations have been used to demonstrate the effect of consistent use of NPK and UREA fertilizers on tolerance of microorganisms with a view to ensuring sustainable use of fertilizers and conservation of the environment. The lethal concentration 50 (LC_{50}) was used to evaluate the effect of the fungi and bacteria exposure to fertilizer (NPK and UREA). Findings suggest that bacteria had a greater tolerance to NPK than fungi in agricultural areas. Test samples in urban areas showed lower LC_{50} in both bacteria and fungi, suggesting anthropogenic activity in the study sites may have had an effect on the soil. For NPK fertilizer, a higher value of LC_{50} was observed in the urban area for bacteria as compared with fungi. In agricultural areas the highest mean LC_{50} for NPK was observed in fungi, and the highest mean LC_{50} for NPK-tolerant bacteria was also

found in agricultural areas. The findings of this study also indicate higher UREA tolerance values for fungi and bacteria in agricultural areas relative to industrial and urban areas, except for some urban areas where UREA tolerant bacteria were higher than those in agricultural and industrial areas. The fungal and bacterial populations differentiated between research sites and were linked to fertilizer treatments in their natural habitats with pre-exposure. These findings are consistent with findings from Xue *et al.*, [28] that define soil microorganisms that will be affected by long-term field application of fertilizers. Tolerance found in urban bacteria can also be related to exposure to pollutants from domestic and industrial areas.

Nitrogen (N) fertilizer application significantly reduces soil pH, regardless of soil type [29, 30, 17]. This is well documented and is primarily the result of soil processes that produce protons, including oxidation of ammonium to nitrite, nitrate and nitrification [29, 30]. A study across China showed that soil pH decreases of 0.45–2.20 units resulting from 8 to 25 years of N fertilization [31]. Agricultural soils are subjects of the greatest N-induced pH decreases, suggesting that these soils were more exposed to N fertilizers than urban and industrial soils. This is consistent with other studies that show a strong link between acidification and application rate for N [31]. The explanation for this may be linked to the fact that the neutralizing effect of plant nitrate absorption had decreased when excess N was added [31]. Nitrogen fertilization in urban, industrial, and agricultural soils substantially decreased microbial populations. The LC₅₀ values, however, showed that this exposure to N fertilizers favoured the development of N tolerance among microbes. Thereafter, the distribution was affected by the activities in the above mentioned areas. This is consistent with several other studies that find repeated application of N fertilizers affecting the microbial soil population relative to control sites [17, 32, 33, 34]. Soil microorganisms may use UREA as a source of nitrogen, and may also be inhibited by the addition of high urea levels due to the detrimental characteristic of ammonia [35]. Nonetheless, the reduction in N-fertilization in the fungal and bacterial population was not synchronous, resulting in a significant reduction in the fungus-to-bacteria ratio in N-fertilized soils relative to non-fertilized soils. Fungi were more susceptible to changes in pH relative to bacteria [35, 36, 37]. As a result, fungal populations fell faster than bacterial populations with pH decreases induced by N fertilization, particularly given that the pH ranges in our study sites were much more beneficial to bacteria than fungi [35, 36].

Liang, [38] shows that fertilization with N is the most effective factor for increasing the relative abundance of soil bacteria. Several authors have noted changes in the abundance of certain bacterial classes, such as *Bacteroidetes*, some members of *Proteobacteria*, *Acidbacteria*, *Gemmatimonadetes* and *Verrucomicrobia*, when comparing N fertilized soils with controls [39, 40, 41]. Microorganisms were not described in our study, but fertilizer tolerance levels were measurements of vulnerability of the soil ecosystems. Remarkably observed high LC₅₀ values in agricultural areas

indicate that they react to the fertilizers used, as their apparent population growth in the presence of NPK and UREA fertilizers has been favoured. The existence of N in complex macromolecules, such as organic fertilizers, may have enriched various classes of species with prompt metabolizing adaptation in the presence of nutrients, as a potential reason for this result. Conversely, the presence of UREA, which is a simple source of N for micro-organisms, should have played a more neutral role rather than supporting a single micro-organism. For example, *bacteroidetes* are considered copiotrophic (r-selected), and their rise in abundance has already been documented in the presence of easily accessible N fertilization [39, 41]. On the other hand, *armatimonadetes* and *nitrospirae* appear to be negatively affected by N fertilizers, which were expected to be as readily available as N induces an out competition mainly of autotrophic *nitrospirae* microbes [39]. These are indications that different fertilizers affect different microorganisms in unique manner which is similar to findings of the current study.

The results found by Fierer *et al.*, [40] are consistent with our findings that after exposure to N fertilizers the tolerance of soil microorganisms to fertilizer toxicity could be related to microbial diversity and tolerance adaptations. Exposure to fertilizers may trigger changes in community makeup of soil microorganisms. By comparison other scientists [42] observed a decline in bacterial diversity with N additions. These two opposing versions indicate that the effects of N modifications on microbial populations are variable and dependent on the site as well as microbial tolerance production, or even other effects such as soil pH shifts, soil holding ability and water micronutrient status. For tuber and root crop growth, NPK fertilizers are more often used [14], while UREA is widely used by farmers as a synthetic fertilizer as it has a fairly high N content (45 %). Pieri, [43] observed that N fertilizers in the West African region have close links with acidification, with an average annual rise of 10% in Al saturation, it was proposed that after only a few years of cultivation they achieved a critical Al toxicity level of 30%. Ogbodo, [44], listed the most significant negative impacts associated with soil acidification as aluminium toxicity (Al) and manganese toxicity (Mg). Typical in these soils, complex aluminium, Al (OH), is converted to ionic form (Al³⁺) at low pH values.

As observed in soils with very high organic matter [42, 42, 44], the soil microbial population is very high. The LC₅₀ findings in the current research have led us to believe that the microbes can vary depending on the presence of organic material in the soils and are indigenous to the sampling sites. The suggestion could be that the microbes could maintain their populations in polluted environments, even under inorganic fertilizers and acidic conditions. The microbes were the primary beneficiaries of the applied fertilizer nutrients, readily available which they assimilated and achieved growth and multiplication at optimum concentrations but were impaired in concentrations beyond the threshold limits.

5. Conclusion and Recommendation

This research concludes that the application of UREA and NPK fertilizers in study area soils decreased the pH of soils in A1 (Aluu), A2 (Eleme) and A3 (Emohua) agricultural areas. Application of UREA and NPK fertilizers in soils of the study areas increased the population of microorganisms in threshold concentrations but is detrimental for NPK and UREA fertilizers at concentrations of 1.50% or above. Results showed that micro-organisms were more tolerant of NPK and UREA in agricultural areas compared with urban and industrial areas, which showed higher LC₅₀ values. This study concludes that fertilizers used in the study areas were linked to the development of NPK and UREA tolerance among soil microorganisms, and were characterized in the study areas by the community of NPK and UREA tolerant microorganisms. The study recommends isolation and characterization in agricultural areas of fungi and bacteria to create a consortium of fertilizer-tolerant microorganisms for the remediation of fertilizer-degraded land. It is crucial that the farmers should observe the application of recommended fertilizer quantities to ensure the soil ecosystem integrity is sustained.

Acknowledgements

This research has been carried out within the World Bank African Centre of Excellence for Oilfield Chemicals Research's PhD program, in line with the mandate of the World Bank to create the African Centre of Excellence at Port Harcourt University in Nigeria. The authors further acknowledge the Regional Universities Forum for Capacity Building in Agriculture (RUFORUM) through Professor Ikechukwu O. Agbagwa of University of Port Harcourt for their continual mentorship and financial support under the Carnegie Post-Doctoral funding.

References

- [1] Cassou, E., Jaffee, S. M., & Ru, J., (2017). The Challenge of Agricultural Pollution: Evidence from China, Vietnam, and the Philippines. *Reconstruction and Development / The World Bank group*. <https://doi.org/10.1596/978-1-4648-1201-9>
- [2] Crouse, D. A. (2018). Soils and Plant Nutrients, Chapter 1. In: K. A. Moore, & L. K. Bradley (eds). *North Carolina Extension Gardener Handbook*. NC State Extension, Raleigh, NC. <https://content.ces.ncsu.edu/extension-gardener-handbook/1-soils-and-plant-nutrients>
- [3] Omuto, C. T., & Vargas, R. R., (2018). Soil nutrient loss assessment in Malawi. Technical Report. FAO, UNEP and UNDP. 64 pp.
- [4] Weil, R. R., & Brady, N. C. (2017). *Soil Phosphorus and Potassium [M]. The Nature and Properties of Soils (15th Ed.)*, 14 (2002), 643–695.
- [5] Bhogal, A., Williams, J., Nicholson, F., Chadwick, D., Chambers, K., & Chambers, B. J., (2015). Mineralisation of organic nitrogen from farm manure applications. *Soil Use Manage.* 32 (Suppl 1), 32–43.
- [6] Food and Agriculture Organization [FAO], (2019). *Handbook on climate information for farming communities – What farmers need and what is available*. Rome. 184 pp. Licence: CC BY-NC-SA 3.0 IGO.
- [7] Alo M. N., Egbule U. C. C., Orji, J. O., & Aneke C. J., (2013). *Microbiological Analysis of Soil from Onu-Ebonyi Contaminated with Inorganic Fertilizer*. *American Journal of Infectious Diseases and Microbiology*. 1, no. 4: 70-74. <https://doi:10.12691/ajidm-1-4-3>.
- [8] Ouyang Y, & Norton J. M. (2020). Short-Term Nitrogen Fertilization Affects Microbial Community Composition and Nitrogen Mineralization Functions in an Agricultural Soil. *Appl Environ Microbiol.* 2020 Feb 18; 86 (5): e02278-19. <https://doi:10.1128/AEM.02278-19> PMID: 31836579; PMCID: PMC7028969.
- [9] Xie, Y., Fan, J., Zhu, W., Amombo, E., Lou, Y., Chen, L., & Fu, J., (2016). Effect of Heavy Metals Pollution on Soil Microbial Diversity and Bermudagrass Genetic Variation. *Front. Plant Sci.* 7: 755. <https://doi:10.3389/fpls.2016.00755>
- [10] Hai-Hang Li, Zhang, Z. Y., & Pan, L. P., (2010). Isolation, identification and characterization of soil microbes which degrade phenolic allelochemicals *Journal compilation © 2009. The Society for Applied Microbiology. Journal of Applied Microbiology.* 108 (2010) 1839-1849.
- [11] Huang, R., Wang, Y., Liu, J., Gao, J., Zhang, Y., Ni, J., Xie, D., Wang, Z., & Gao, M. (2020). Partial Substitution of Chemical Fertilizer by Organic Materials Changed the Abundance, Diversity, and Activity of NirS-Type Denitrifying Bacterial Communities in a Vegetable Soil. *Appl. Soil Ecol.* 152, 103589. <https://doi.org/10.1016/j.apsoil.2020.103589>
- [12] Li, Y., Liu, Y., Niu, Y. W. L., Xia, X., & Tian, Y., (2014). Interactive effects of soil temperature and moisture on soil N mineralization in a *Stipa krylovii* grassland in Inner Mongolia, China. *Journal of Arid Land*, 6, 571–580.
- [13] Sarkar, B., Singh, M., Mandal, S., Churchman, G. J., & Bolan, N. S. (2018). Clay Minerals—Organic Matter Interactions in Relation to Carbon Stabilization in Soils. *The Future of Soil Carbon*, 71–86. <https://doi:10.1016/b978-0-12-811687-6.00003-1>
- [14] Geisseler, D., Linquist, B. A., & Lazicki, P. A. (2017). Soil Biology & Biochemistry Effect of fertilization on soil microorganisms in paddy rice systems e A. *Soil Biology and Biochemistry*, 115, 452–460. <https://doi.org/10.1016/j.soilbio.2017.09.018>
- [15] Datta, A., Santra, S. C., & Adhya, T. K., (2013). Effect of inorganic fertilizers (N, P, K) on methane emission from tropical rice field of India. *Atmospheric Environment* 66, 123e130.
- [16] Das, S., & Adhya, T. K., (2014). Effect of combine application of organic manure and inorganic fertilizer on methane and nitrous oxide emissions from a tropical flooded soil planted to rice. *Geoderma* 213, 185e192.
- [17] Zhang, Y., Shen, H., He, X., Thomas, B. W., Lupwayi, N. Z., Hao, X., ... Shi, X. (2017). Fertilization shapes bacterial community structure by alteration of soil pH. *Frontiers in Microbiology*, 8 (JUL). <https://doi.org/10.3389/fmicb.2017.01325>
- [18] American Public Health Association [APHA]., (1998). *Standard Methods for Examination of Water and Waste Water*. American Public Health Association, American Water Works Association and Water Pollution Control Federation, 20th Edition, Washington DC, USA.

- [19] Finney D. J., (1952). Probit analysis: a statistical treatment of the sigmoid response curve. Cambridge University Press.
- [20] Vincent, B. K. (1980). Probit Analysis. <http://userwww.sfsu.edu/efc/classes/biol710/probit/ProbitAnalysis.pdf>
- [21] Food and Agriculture Organization (FAO) and Economic Commission for Africa (ECA), (2018). Regional Overview of Food Security and Nutrition. Addressing the threat from climate variability and extremes for food security and nutrition. Accra. 116 pp.
- [22] Druilhe, Z., & Barreiro-Hurlé, J., (2012). Fertilizer subsidies in sub-Saharan Africa, ESA Working Papers 288997, Food and Agriculture Organization of the United Nations, Agricultural Development Economics Division (ESA). <https://DOI:10.22004/ag.econ.288997>
- [23] Giller, K. E., Corbeels, M., Nyamangara, J., Triomphe, B., Affholder, F., Scopel, E., & Tittone, P. (2011). A research agenda to explore the role of conservation agriculture in African smallholder farming systems. *Field Crops Research*, 124 (3), 468–472. <https://doi:10.1016/j.fcr.2011.04.010>
- [24] Sahu, P., Singh, D., Prabha, R., Meena, K., & Abhilash, P., (2019). Connecting microbial capabilities with the soil and plant health: Options for agricultural sustainability. *Ecol. Indic.* 105, 601–612. <https://doi.org/10.1016/j.ecolind.2018.05.084>
- [25] Bhat, A. K., (2013). Preserving microbial diversity of soil ecosystem: a key to sustainable productivity. *International Journal of Current Microbiology and Applied Sciences*. ISSN: 2319 – 7706, Volume 2 Number 8 pp. 85-101.
- [26] Delgado-Baquerizo, M., Maestre, F. T., Reich, P. B., Jeffries, T. C., Gaitan, J. J., Encinar, D., ... Singh, B. K., (2016). Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nature communications*. 7, 10541. <http://doi:10.1038/ncomms10541>
- [27] Bach, E. M., & Wall, D. H. (2018). Trends in Global Biodiversity: Soil Biota and Processes. In *The Encyclopedia of the Anthropocene*; DellaSala Dominick, A., Goldstein Michael, I., Eds.; Elsevier: Oxford, UK, 2018; Volume 3, pp. 125–130. <https://doi.org/10.1016/B978-0-12-5409548-9.09822>
- [28] Xue, C., Zhang, X., Zhu, C., Zhao, J., Zhu, P., Peng, C., ... Shen, (2016). Quantitative and compositional responses of ammonia-oxidizing archaea and bacteria to long-term field fertilization. *Scientific Reports*, 6, 1–11. <https://doi.org/10.1038/srep28981>
- [29] Pasley, H. R., Cairns, J. E., & Camberato, J. J. et al., (2019). Nitrogen fertilizer rate increases plant uptake and soil availability of essential nutrients in continuous maize production in Kenya and Zimbabwe. *Nutr Cycl Agroecosyst* 115, 373–389. <https://doi.org/10.1007/s10705-019-10016-1>
- [30] Dong, W., Zhang, X., Wang, H., Dai, X., Sun, X., Qiu, W., & Yang, F. (2012). Effect of Different Fertilizer Application on the Soil Fertility of Paddy Soils in Red Soil Region of Southern China. *PLoS ONE*, 7 (9), 1–9. <https://doi.org/10.1371/journal.pone.0044504>
- [31] Yu, H., Gao, Q., Shao, Z., Ying, A., Sun, Y., Liu, J., ... Zhang, B. (2016). Decreasing nitrogen fertilizer input had little effect on microbial communities in three types of soils. *PLoS ONE*, 11 (3), 1–12. <https://doi.org/10.1371/journal.pone.0151622>
- [32] Geisseler, D., & Scow, K. M. (2014). Long-term effects of mineral fertilizers on soil microorganisms - A review. *Soil Biology and Biochemistry*, 75, 54–63. <https://doi.org/10.1016/j.soilbio.2014.03.023>
- [33] Wang, C., Liu, D., & Bai, E. (2018). Decreasing soil microbial diversity is associated with decreasing microbial biomass under nitrogen addition. *Soil Biology and Biochemistry*, 120 (72), 126–133. <https://doi.org/10.1016/j.soilbio.2018.02.003>
- [34] Zhang, Z., Zhang, X., Mahamood, M., Zhang, S., Huang, S., & Liang, W. (2016). Effect of long-term combined application of organic and inorganic fertilizers on soil nematode communities within aggregates. *Scientific Reports*, 6 (August), 1–12. <https://doi.org/10.1038/srep31118>
- [35] Fu, H., Lou, Z., & Hu, S. A (2020). Temporal-spatial analysis and future trends of ammonia emissions in China. *Sci. Total Environ.* 2020, 731, 138897. <https://DOI:10.1016/j.scitotenv.2020.138897>.
- [36] Veverka, K., Štolcová, J., & Růžek, P. (2007). Sensitivity of Fungi to Urea. *Plant Protect. Sci*, 43 (4), 157–164.
- [37] Bünemann, E. K., Schwenke, G. D., & Van Zwieten, L. (2006). Impact of agricultural inputs on soil organisms - A review. *Australian Journal of Soil Research*, 44 (4), 379–406. <https://doi.org/10.1071/SR05125>
- [38] Liang, R., Hou, R., Li, J., Lyu, Y., Hang, S., Gong, H., & Ouyang, Z. (2020). Effects of Different Fertilizers on Rhizosphere Bacterial Communities of Winter Wheat in the North China Plain. *Agronomy*, 10 (1), 93. <https://doi:10.3390/agronomy10010093>
- [39] Ramirez, K. S., Lauber, C. L., Knight, R., Bradford, M. A., & Fierer, N. (2010). Consistent effects of nitrogen fertilization on soil bacterial communities in contrasting systems. *Ecology*, 91, 3463–3470.
- [40] Fierer, N., Leff, J. W., Adams, B. J., Nielsen, U. N., Bates, S. T., Lauber, C. L., Owens, S., Gilbert, J. A., Wall, D. H. & Caporaso, J. G. (2012). Cross-biome metagenomic analyses of soil microbial communities and their functional attributes. *Proc. Natl. Acad. Sci. USA*, 109, 21390–21395.
- [41] Nemergut, D. R., Townsend, A. R., Sattin, S. R., Freeman, K. R., Fierer, N., Neff, J. C., Bowman, W. D., Schadt, C. W., Weintraub, M. N., & Schmidt, S. K. (2008). The effects of chronic nitrogen fertilization on alpine tundra soil microbial communities: Implications for carbon and nitrogen cycling. *Environ. Microbiol.* 10, 3093–3105.
- [42] Zano, M., Rosselli, R., Meneghesso, A., Sablok, G., Stevanato, P., Engel, M., ... Squartini, A. (2018). Response of Bacterial Communities upon Application of Different Innovative Organic Fertilizers in a Greenhouse Experiment Using Low-Nutrient Soil Cultivated with *Cynodon dactylon*, 1–11. <https://doi.org/10.3390/soilsystems2030052>
- [43] Pieri, Christian J. M. G. (1992). *Fertility of Soils: A Future for Farming in the West African Savannah*. Berlin: Springer-Verlag.
- [44] Ogbodo, E. N. (2013). Impact of the use of Inorganic Fertilizers to the Soils of the Ebonyi State Agro-Ecology, South-Eastern Nigeria., 3 (7), 33–39.