

Research Article

The Effects of Mother Tree Size and Flowering Period on Salt Tolerance of *Tamarix Austromongolica* Seeds in the Yellow River Delta

Qingzhi Lin^{1,2}, Yaling Yang³, Muzheng Hao⁴, Xin Han¹, Yuling Zhang^{5,*} ,
Jing Yang⁴, Mengmeng Zheng⁶, Qingshan Yang⁷, Xiaojun Lu⁶

¹Key Laboratory of National Forestry and Grassland Forestry Administration for Silviculture of the Lower Yellow River, Shandong Agricultural University, Taian, China

²Linyi County Industrial and Information Bureau, Dezhou, China

³Shandong Holy Land Planning and Design Institute Co., Ltd, Taian, China

⁴Dongying Promotion Center for Wetland City Construction, Dongying, China

⁵Shandong Qinghua Landscape Design Co., Ltd., Weifang, China

⁶Dongying Shengjing Forestry Co. Ltd., Dongying, China

⁷Shandong Academy of Forestry, Jinan, China

Abstract

Tamarix austromongolica is a two-season flowering plant with a spring and summer flowering period and both flowering periods produce viable seeds, with flowering extending from mid-May to October and seed maturation from mid-June to October. In order to reveal the germination pattern of *T. austromongolica* seeds in the Yellow River Delta region, this study investigated the salt tolerance of seeds of different age mother trees during the spring and summer flowering periods in the region. Spring and summer flowering seeds of different ground diameter mother trees in the *T. austromongolica* forest in the region were collected and subjected to seed germination experiments under different concentrations of NaCl solution. The results showed that the diameter grade of the mother tree had a significant impact on seed germination, but had no significant effect on early seedling growth (except for the root length of offspring seedlings during the summer flowering period). As the diameter level of the mother tree increases, the germination rate, germination index, and vitality index of seeds during the spring flowering period first increase and then decrease, with the highest values observed at a ground diameter of 10.0-19.9cm. However, the above indicators of summer flowering seeds gradually decreased with the diameter increasing. There were significant differences in salt tolerance of seeds during different flowering periods. The germination rate and germination index of spring flowering seeds significantly decreased between 11‰ and 15‰, but the effect of salt concentration on summer flowering seed germination was not significant. For the early growth of seedlings, as the salt concentration increased, the root length and seedling height of seed offspring seedlings during spring and summer flowering periods decreased after being higher than 5‰. Compared with spring flowering seeds, summer flowering seeds had higher germination ability and seedling weight under salt stress. Therefore, the salt tolerance of the seeds of *T. austromongolica* was determined by the size of the mother tree and the flowering period, which is an important strategy for its adaptation to the Yellow River Delta region.

*Corresponding authors: 497107211@qq.com (Yuling Zhang)

Received: 13 May 2024; **Accepted:** 28 May 2024; **Published:** 13 June 2024



Copyright: © The Author(s), 2024. Published by Science Publishing Group. This is an **Open Access** article, distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Keywords

Yellow River Delta, Mother Tree Size, Salt Stress, Florescence, Seed Germination, Seedling Size

1. Introduction

In the Yellow River Delta region, the water table is shallow and mineralization is high, and groundwater salts are easy to collect on the surface [1]. Soil salinization is a process in which salts from the soil subsoil or groundwater rise to the surface with capillary water, and water evaporation causes salts to accumulate in the surface soil [2]. The formation of soil salinization is influenced by a combination of natural and anthropogenic factors, such as climate, hydrology, geology, over-fertilization and indiscriminate cutting [3]. Excessive salinity in the soil can cause harm to plants, which is mainly divided into osmotic stress and ionic toxicity, the water potential in the soil is usually lower than that of the plant cells, and the water in the body of the plant in this environment is easy to seep out of the body, resulting in water shortage; salt stress breaks the dynamic balance between the ions in the soil, resulting in unbalanced ion uptake in the plant [4]. Salt stress inhibits plant seed germination and early seedling growth, but for some plants with strong salt tolerance, lower salt concentration has a certain promotion effect on their seed germination and seedling growth, such as Chinese *T. austromongolica* [3], salt pawpaw, salt spicebush [5].

Characteristics of the mother tree (age, diameter class size) are important factors affecting seed quality and seed germination [6]. Seed germination is key to population renewal and species continuation in the life history of seed plants [7-9]. The age of the mother tree is one of the main reasons for the differentiation of seed characteristics within populations, and the reproductive age of the mother tree affects the maturity of the mother tree itself and the efficiency of its utilization of resources; although the intraspecific habitats are relatively uniform, there are subtle differences in the spatial positions actually occupied by different plants, which affects the plants' access to resources, and consequently, their growth and reproduction [10, 11]. Seeds of mother tree individuals of different ages (young individuals, middle-aged and old individuals) showed differences in their adaptive capacity, and with the increase of mother tree diameter at breast height (DBH), the seed mass of *A. Tetracentron sinense* showed a tendency to increase firstly and then decrease later [12]. It has also been shown that the seed germination rate of *Emmenopterys henryi* is not related to the size of the mother tree [13]. Overall, the effect of mother tree size on seed quality is not consistent and is affected by a combination of tree species, growth environment and other aspects.

Tamarix austromongolica belongs to the genus *Tamarix* in the family *Tamarixaceae*, a deciduous shrub or small tree,

which is a dominant species in the Yellow River Delta region, with the ecological functions of windbreak and sand fixation, soil and water conservation and saline-alkaline land improvement [14]. Studies have found that salinity is a key factor affecting its growth and distribution in the region [15, 16]. Two-season flowering is a phenomenon in which a plant has two independent flowering periods within a single growing season, or a long-term sustained flowering period with two peak flowering periods, and can produce mature seeds throughout the flowering period [17]. *T. austromongolica* is a two-season flowering plant with a spring and summer flowering period and both flowering periods produce viable seeds, with flowering extending from mid-May to October and seed maturation from mid-June to October [18, 19]. In this paper, we investigated the germination process of spring and summer flowering seeds of *T. austromongolica* with different mother tree sizes under different salt concentrations, to clarify the effect of mother tree size on seed salt tolerance, to increase the knowledge of the germination ability of two-season flowering seeds, to reveal the mechanism of *T. austromongolica*'s adaptability, and to provide scientific basis for the protection and rational utilization of the forest stand.

2. Materials and Methods

2.1. Overview of the Study Area

The seeds used in the experiment were collected from the natural forest of *T. austromongolica* distributed around the Wetland City Construction Promotion Center in Dongying City, Shandong Province. The area has a continental monsoon climate, with rain and heat in the same season and four distinct seasons. The average annual temperature ranges from 11.7 to 12.6 °C, the annual precipitation is 550-600 mm, mostly concentrated in the summer, the precipitation varies greatly from year to year, which makes it easy to form droughts and floods, and the frost-free period lasts 206 d. The average annual temperature ranges from 11.7 to 12.6 °C.

2.2. Seed Collection

The seeds used in the experiment were randomly collected on July 23 and September 21, 2019 in the natural forest of *T. austromongolica* at three ground diameter classes, and 10

plants were collected at each diameter class. The ground diameter of diameter class I was 3.4-9.9 cm, that of diameter class II was 10-19.9 cm, and that of diameter class III was 20-35 cm. Seeds from sunny branches in the middle of the plant were taken and brought back to the laboratory to be dried in the shade for the germination test.

2.3. Experimental Design

Set the concentration of NaCl solution were control 0‰, 3‰, 5‰, 7‰, 9‰, 11‰, 13‰, 15‰ of eight gradients; according to the mother tree of *T. austromongolica* ground diameter is divided into I, II, III, the three grades of *T. austromongolica* have the seeds of spring and summer flowering period, a total of 48 treatments. Seeds were randomly selected (without going to seed hairs), 50 seeds in 1 group, 4 replications. The seeds were placed in Petri dishes with a diameter of 90 mm padded with 2 layers of filter paper and kept the filter paper moist, and the Petri dishes were placed in a light incubator sterilized with 75% alcohol, with the temperature set at 25 °C, 12 h of light and 12 h of darkness, and incubated for 8 d. During the process of seed germination, the number of seed germination was observed and recorded every 24 h. The number of seed germination was recorded by weighing and replenishment of seed loss per day, and the number of seed loss per day was recorded by weighing and replenishment of seed loss per day. Weighing method was used to replenish the water lost every day to keep the filter paper and seeds moist, and to maintain the relative stability of the concentration of each treatment during the experiment. At the end of the experiment, five representative seedlings with moderate growth were selected for each treatment to measure seedling height and root length; 10 seedlings were selected for each treatment and weighed separately for fresh weight using a 1-in-10,000 balance. Then germination rate (GR), mean germination time (MGT), germination index (GI) and vigor index (VI) were calculated.

GR (germination rate) = $G1/N \times 100\%$, where: G1 is the number of germination; N is the total number of seeds for test.

MGT (mean germination time) = $(\sum D \times n) / \sum n$, where: n is the number of normal germinated grains on each corresponding day; D is the number of days from seed placement.

GI (germination index) = $\sum (Gt/Dt)$, where: Gt is the number of net germination on each day; Dt is the corresponding germination day.

VI (vigor index) = $S \times GI$, where: S is the fresh weight of seedlings; GI is the germination index.

2.4. Data Processing and Analysis

Data were organized and statistically analyzed using Excel with SPSS 19.0 software. Two-factor analysis of variance (ANOVA) was performed on germination index, root length and seedling height of spring flowering seeds and summer

flowering seeds using SPSS 19.0. If the differences were significant, multiple comparisons were then performed using the least significant difference (LSD) method, and the level of significant difference test for analysis was $P < 0.05$.

3. Results and Analysis

3.1. Effects of Mother Tree Size and Salt Stress on Seed Germination of Flowering Seeds in Two Seasons of *T. Austromongolica*

For spring flowering seeds, mother tree size had a significant effect on germination rate, germination index and vigor index, and a non-significant effect on average germination time; salt stress had a highly significant effect on germination rate and germination index, and a significant effect on average germination time and vigor index (Table 1). The mother tree size was $II > III > I$ in germination rate, germination index and vigor index ($P < 0.05$), and the average germination time was shortest in diameter class I. Germination rate and germination index did not differ significantly between individual salt concentrations below 9‰ but were significantly higher than 11-15‰ treatments.

For summer-flowering seeds, mother tree size had highly significant ($P < 0.01$) effects on germination rate, mean germination time, germination index and vigor index, while salt stress had no significant ($P > 0.05$) effects on germination rate, mean germination time, germination index and vigor index (Table 1). The performance of mother tree size in germination rate, mean germination time, germination index and vigor index was $I > II > III$.

The differences in germination rate ($F=425.29$, $P < 0.01$), mean germination time ($F=8.34$, $P < 0.01$), germination index ($F=973.43$, $P < 0.01$), and vigor index ($F=79.59$, $P < 0.01$) between the two seasons of flowering seeds were highly significant. Summer flowering seeds had high germination percentage, mean germination time, germination index and vigor index compared to spring flowering seeds.

3.2. Effects of Mother Tree Size and Salt Stress on Early Seedling Growth of *T. Austromongolica*

For spring-flowering seeds, the effect of parent tree diameter class on root length, seedling height, and seedling biomass was not significant ($P > 0.05$), and the effect of salt stress on root length, seedling height, and seedling weight was highly significant ($P < 0.01$) (Table 1). Root length and seedling height decreased gradually with the increase of salt stress, and salt concentration below 9‰ was significantly higher than that of 11-15‰ treatment ($P < 0.05$). Seedling biomass CK was significantly greater than other salt concentrations ($P < 0.05$), 9‰ was greater than 13‰ and 15% ($P < 0.05$), and the differences among other salt concentrations

were not significant ($P > 0.05$).

For summer-flowering seeds, parent tree diameter class had a highly significant effect on root length and a non-significant effect on seedling height and seedling weight; salt stress had a highly significant effect on root length, a significant effect on seedling height, and a non-significant effect on seedling weight (Table 1). Between different diameter classes, the difference in root length between diameter class I and III was not significant ($P > 0.05$) and was highly significant lower than that of diameter class II ($P < 0.01$).

Root length and seedling height increased and then decreased with increasing salt concentration, and were highest ($P < 0.05$) at 3‰.

The differences in root length ($F=72.34$, $P<0.01$) and seedling weight ($F=55.15$, $P<0.01$) were highly significant between the two seasons of flowering seeds, but not in seedling height ($F=1.36$, $P>0.05$). Summer flowering seeds had high root length and seedling weight compared to spring flowering seeds.




Figure 1. Effect of parent tree size and salt stress on seed germination of *T. austromongolica* in two flowering seasons.




Figure 2. Effects of mother tree size and salt stress on flowering seedlings of *T. austromongolica* in two seasons.

Table 1. Two-factor ANOVA analysis of seed germination of *T. austromongolica* by mother tree size and salt stress.

Florescence	Indexes	Mother tree size		Salt stress		Mother tree size × salt stress	
		F-value	P-value	F-value	P-value	F-value	P-value
Spring-flowering seeds	Germination rate	4.91	<0.05	9.44	<0.01	1.92	<0.05
	Mean germination time	0.24	0.79	2.68	<0.05	0.67	0.80
	Germination index	5.00	<0.05	9.03	<0.01	2.22	<0.05
	Vogor index	4.61	<0.05	2.79	<0.05	1.61	<0.05
	Root length	0.33	0.72	3.80	<0.01	2.23	<0.05
	Seedling height	0.06	0.95	4.39	<0.01	1.67	0.09
	Seedling biomass	2.77	0.07	3.82	<0.01	2.24	<0.05
Summer-flowering seeds	Germination rate	79.35	<0.01	0.71	0.67	3.29	<0.01
	Mean germination time	12.02	<0.01	1.22	0.31	1.15	0.34

Florescence	Indexes	Mother tree size		Salt stress		Mother tree size × salt stress	
		F-value	P-value	F-value	P-value	F-value	P-value
	Germination index	65.87	<0.01	0.92	0.50	3.00	<0.01
	Vigor index	11.72	<0.01	1.47	0.20	0.85	0.61
	Root length	8.87	<0.01	10.777	<0.01	1.34	0.22
	Seedling height	1.46	0.24	3.04	<0.05	0.63	0.83
	Seedling biomass	1.65	0.20	2.10	0.06	0.80	0.67

4. Discussion

4.1. The Effect of Parent Tree Diameter Level on Seed Germination in Two Flowering Seasons of Salt-stressed *T. Austromongolica*

Seed germination is a key stage of plant natural regeneration and can directly reflect the seed quality. The difference in the diameter level of the mother tree affects seed quality and thus seed germination. In this paper, we found that the germination rate, germination index, and vigor index of the spring flowering seeds of *T. austromongolica* were II>III>I, and II was the largest in the diameter class. Studies on *Tetracentron sinense* [12], *Scaphium wallichii* [11], and *Camellia rosthorniana* [20] also found that seed germination was strongest in the middle of the radial class of the mother tree, which was thought to be related to the increased reproductive allocation at that stage. In a study of lateral cypress seeds of different parent tree sizes, It was found that the germination rate and germination index of *Platycladus orientalis* seeds increased with the increase of parent tree size [21]. It was shown similar results for *Pinus thunbergii* seeds, which were related to the increase of nutrient accumulation in the parent tree [22]. For the spring flowering seeds of *T. austromongolica*, the germination rate, germination index, and vigor index were I>II>III, and the highest quality of seeds was found in the diameter class I. For herbaceous plants *Ligularia virgaurea* [23], and *Gentianopsis paludosa* [24], it was found that the reproductive allocation decreased with the increase of individual size, but it was obviously affected by the habitat. Therefore, the variation of flowering seeds with parent tree size in both seasons of *T. austromongolica* may be the result of the joint influence of its own nutrient accumulation and habitat.

In this study, we found that the germination rate and germination index of *T. austromongolica* seeds in the spring flowering season at NaCl concentrations of 3%-9‰ were higher than those of the control and other salt concentration treatments, and the seed germination in the summer flowering

season was not affected by the salt concentration, which indicated that its seeds had a strong salt tolerance ability. In terms of salt tolerance, the summer flowering seeds of *T. austromongolica* were stronger than those of spring flowering, which was consistent with the results of a study on the spring and summer flowering seeds of *Tamarix arceuthoides* [25]. However, the study on the seeds of *Tamarix ramosissima* in two seasons found that the seeds in spring flowering stage were more salt-tolerant [26]. In this paper, we found that the average germination time of spring-flowering seeds of *T. austromongolica* was lower than that of summer-flowering seeds, and the germination rate was high, which was consistent with the previous findings [25]. However, it was found that spring flowering seeds germinated slower than summer flowering seeds in his study on the germination of *Tamarix arceuthoides* and *Tamarix ramosissima* under salt treatment [27]. *T. austromongolica* spring flowering seed formation in the Yellow River Delta region at the time of low rainfall and high soil salinity, while the summer flowering seed formation into the rainy season, the habitat improved significantly, so the seed quality is obviously affected by environmental factors.

4.2. Effect of Mother Tree Diameter Class on Early Growth of Seedlings of Offspring

In this paper, it was found that there was no significant effect of mother tree diameter class on root length, seedling height and seedling weight of spring flowering seeds, and there was also no significant effect on seedling height and seedling weight of summer flowering seeds, but the effect on root length was highly significant. In contrast, a study on *Pinus thunbergii* seedlings found that both seedling biomass and growth rate increased with increasing diameter class [22], and a similar finding was found for *Platycladus orientalis* seedlings [21]. This suggests a complex effect of parent tree size on offspring seedlings. With increasing salt concentration, seedling height and root length of both spring-flowering seed and summer-flowering seed progeny seedlings were significantly suppressed. Thus, salt stress plays a major role in seedling growth.

The level of root development determines the growth and development of plants [28]. In this paper, it was found that summer flowering seeds under salt stress had higher root length than spring flowering seeds progeny seedlings. Studies on *Morus alba* L. found that the degree of root development under salt stress was related to its salt resistance [29]. Therefore, summer flowering seedling progeny were more salt-adapted than spring ones. High seedling growth and seedling size favor seedlings to obtain more resources and improve competitiveness with surrounding plants [22, 30]. The biomass of seedling offspring of *T. austromongolica* was higher in the summer flowering season than in the spring, suggesting a greater adaptive capacity.

5. Conclusion

In this paper, it was found that *T. austromongolica* spring flowering seeds were the most salt-tolerant in the intermediate diameter class II of the mother tree; while the summer flowering seeds' salt tolerance decreased with the increase of the diameter class of the mother tree. For seedling growth, mother tree size had no significant effect. Therefore, mother tree size was an important factor affecting seed germination, but was influenced by flowering stage. Salt stress had a significant effect on seed germination during flowering in spring, but not in summer; it had a significant effect on seedling size, and growth was first promoted and then inhibited with increasing salt concentration. Therefore, the effect of soil salinity on seed germination was related to flowering stage, but was an important determinant of seedling growth. Compared with spring-flowering seeds, summer-flowering seeds had high salt tolerance and seedling size with greater salt tolerance. Therefore, the mother tree size and flowering period together determine the salt tolerance ability of *T. austromongolica* seeds, and the regulation of seed salt tolerance is an important strategy for the natural regeneration of its population adapted to the Yellow River Delta region, which is important for predicting the evolution and conservation of this population.

Abbreviations

DBH	Diameter at Breast Height
ANOVA	Analysis of Variance
GR	Germination Rat
MGT	Mean Germination Time
GI	Germination Index
VI	Vigor Index
LSD	Least Significant Difference

Author Contributions

The Manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Acknowledgments

The research was supported by funding from Shandong Provincial Natural Science Foundation (Project No: ZR2023QC133), Central Guiding Local Science and Technology Development Special Fund Project (Project No: YDZX2022111), Special project for comprehensive utilization of saline alkali land in Dongying City (Project No: 2023YJDZX03), Shandong Province Science and Technology Small and Medium sized Enterprise Innovation Ability Enhancement Project (Project No: 2023TSGC0044), and Germplasm Resources Project of Salt alkali tolerant Tree Species in the Yellow River Delta (Project No: 2019-370505-05-03-035206).

Conflicts of Interest

The authors declare that they have no competing interests.

References

- [1] Zhang Z, Song Y, Zhang H, Li X, Niu B. Spatiotemporal dynamics of soil salinity in the Yellow River Delta under the impacts of hydrology and climate. *Chinese Journal of Applied Ecology*, 2021, 32(4): 1393-1405.
- [2] Wang R, Kong S, Xu L, Li Y, Cheng W, Zhao E. Spatial distribution of soil salinity under different surface land cover types and micro-topography in the Yellow River Delta. *Transactions of the Chinese Society of Agricultural Engineering (Transactions of the CSAE)*, 2020, 36(19): 132-141.
- [3] Zhang X. Seed germination and seedling growth of *Tamarix chinensis* and *Nitraria tangutorum* under salt stress. Beijing Forestry University, 2016.
- [4] Mao L, Lu J, Jiang H. Mechanisms of plant responses to salt-alkali stress. *Molecular Plant Breeding*, 2020, 18(10): 3441-3448.
- [5] Zeng Y, Cai Z, Ma J, Zhang F, Wang B. Effects of salt and water stress on seed germination of halophytes *Kalidium foliatum* and *Halostachys caspica*. *Chinese Journal of Ecology*, 2006(09): 1014-1018.
<https://doi.org/10.1111/j.1365-2745.2005.01056.x>
- [6] Espahbodi K, Hosseini S M, MirzaeiI-nodoushan H, Tabari M, Dehghan-Shooraki Y. Tree age effects on seed germination in *Sorbus torminalis*. *General and Applied Plant Physiology*, 2007, 33(1-2): 107-119.
- [7] Flores J, Briones O. Plant life-form and germination in Mexican inter-tropical desert. Effects of soil water potential and temperature. *Journal of Arid Environments*. 2001;47: 485-497.
- [8] Guariguata M R. Seed and seedling ecology of tree species in neo tropical secondary forest: Management in plications *Ecological Applications*.2000, 10: 145-154.

- [9] Han G, Mao P, Liu S, Wang G, Zhang Z, Xue Q. Effects of sea water salinity and mother tree size on the seed germination and seedling early growth of *Pinus thunbergii* coastal protection forest. *Chinese Journal of Ecology*, 2009, 28(11): 2171-2176.
- [10] Zhou L, Li Q, Xie Z. Fruiting characteristics and seed germination capacity of *Abies fargesii* in Shennongjia Nature Reserve. *Biodiversity Science*, 2008, 16(5): 509-515.
- [11] Zhang Q, Huang P V, Chen C, Hien L T, Chen S. Effects of mother tree diameter classes and storage on seed germination rate of *Sterculia lychnophora* Hance. *Journal of Fujian Agriculture and Forestry University (Natural Science Edition)*, 2017, 46(2): 159-165.
- [12] Li H, Gan X, Zhang Z, Zhang C, Song L. Effects of altitudes and the DBH of seed trees on biological characteristics of *Tetracentron sinense* (Tetracentraceae) Seeds. *Plant Diversity and Resources*, 2015, 37(02): 177-183.
<https://doi.org/10.3724/SP.J.1259.2013.00295>
- [13] Guo L, Xu W, Fang H, Xiao Z, Tian Y, Huang X, Yin C, Hu G. Seed rain, seed bank and natural regeneration of *Emmenanthera henryi* in different age classes. *Acta Botanica Boreali-Occidentalia Sinica*, 2016, 36(11): 2273-2282.
- [14] Song X, Li S, Wei W, Guo J, Yu Y, Liu Z. Distribution characteristics of root system of *Tamarix chinensis* in Yellow River Delta and its influence factors. *Wetland Science*, 2017, 15(05): 716-723.
- [15] Xia J B, Zhang S Y, Zhang G C, Xu J W, Liu J T, Li C R. Growth dynamics and soil water ecological characteristics of *Tamarix Chinensis* Lour. forests with two site types in coastal wetland of Bohai golf. *Journal of Food, Agriculture & Environment*, 2013, 11: 1492-1498.
- [16] Zhao X. Effects of simulated groundwater on water and salt characteristics of soil-*Tamarix Chinensis* and *Tamarisk* growth in the Yellow River Delta. *Shandong Agricultural University*, 2018. <https://doi.org/10.3724/SP.J.1145.2012.00224>
- [17] Pico F X, Retana J. Seed ecology of a Mediterranean perennial herb with an exceptionally extended flowering and fruiting season [J]. *Botanical Journal of the Linnean Society*, 2003, 142(3): 273-280.
<https://doi.org/10.1046/j.1095-8339.2003.00172.x>
- [18] Jin Z, Wang X, Shen Q, Mou Z, Gao Q. Experimental Study on *Tamarix austromongolica*. *Soil and Water Conservation in China*, 1984(06): 36-39.
- [19] Liu M. Comprehensive research and large-scale promotion and application of *Tamarix* plants. Lanzhou: Lanzhou University Press, 1995.
- [20] Cao G, Zhong Z, Xie D, Liu Y. The relationship between reproductive allocation, fruit set and individual size of *Camellia rosthorniana* in different communities. *Acta Phytoecologica Sinica*, 2005, 29(3): 361-366.
- [21] Bai B, Zhang K, Liu W, Qin Y, Wang X, Gao Y, Zhao X, Zhang J, Cao B, Mao P. Effects of mother tree size on seed germination and seedling growth of *Platycladus orientalis*. *Seeds*, 2019, 38(06): 30-35.
- [22] Mao P, Mu H, Cao B, Liu Y, Fan Z, Wang S. Effects of sand burial and overstory tree age on seedling establishment in coastal *Pinus thunbergii* forests in the northern Shandong Peninsula, China. *The Forestry Chronicle*, 2016, 92(3): 357-365.
- [23] Liu Z, Du G, Chen J. Size-dependent reproductive allocation of *Ligularia virgaurea* in different habitats. *Acta Phytoecologica Sinica*, 2002, 26(1): 44-50.
- [24] Hou Q, Ye G, Ma X, Su X, Zhang S, Sun K. Size-dependent reproductive allocation of *Gentianopsis paludosa* in different habitats of the Qinghai-Tibetan Plateau. *Acta Ecologica Sinica*, 2016, 36(9): 2686-2694.
- [25] Yan C, Wei Y, Wang L. Study on germination of *Tamarix arceuthoides* seeds in Spring and Summer. *Arid Zone Research*, 2010, 27(05): 750-754.
- [26] Liu J. The comparative study of reproductive ecology in bi-seasonal flowering periods of *Tamarix ramosissima*. Shihezi University, 2018.
- [27] Wang L. Study on Bi-seasonal Flowering and Fruit Set Characteristics and Its Ecological Adaption of Two Species *Tamarix* in Xinjiang. Xinjiang Agricultural University, 2008.
- [28] Zhang Y, Liu B, Wang Y, Xu Z, Zhang X. Research progress of plant roots. *Tianjin Agricultural Sciences*, 2016, 22(11): 11-18.
- [29] Huang H, Fu J, Cao B, Zhao W, Hao M, Mao P, Wang T, Hao Y, Zhang X, Tan H. Effects of salt stress and temperature on seed germination and seedling growth of *Morus alba*. *Journal of Southwest Forestry University (Natural Science)*, 2021, 41(1): 1-9.
- [30] Toledo-Aceves T, Swaine M D. Above- and below-ground competition between the liana *Acacia kamerunensis* and tree seedlings in contrasting light environments [J]. *Plant Ecology* 2008.196: 233-244.
<https://doi.org/10.1007/s11258-007-9347-0>