

Research Article

Effect of Nitrogen Fertilizer and Seed Rates on Growth, Yield and Yield Components of Bread Wheat in Jimma Zone South Western Ethiopia

Sisay Gurmu^{*} , Muhidin Biya

Agronomy and Crop Physiology Department, Jimma Agricultural Research Center, Jimma, Ethiopia

Abstract

Wheat is one of the most important cereal crops in terms of production and consumption in Ethiopia. The country has the potential to produce a sufficient amount of wheat grain using irrigation, and government also gave attention to produce wheat and export it. But inappropriate agronomic practices, soil fertility depletion, and drought are among the constraints to the low yield of wheat crops in the country. In view of this, the field experiment was conducted during the irrigation season of 2020/2021 in two districts that are selected as irrigation potential and irrigation facilities namely, Sekacokorsa and Tiroafetadistricts of Jimma zone. The field experiment consisted of four N fertilizer rates (0, 23, 46, 69, and 92 kg N ha⁻¹) and three seed rates (125, 150, and 175 kg ha⁻¹). The experiment was laid out in a randomized complete block design with a factorial arrangement in three replications. From the field experiment results, the growth, yield and yield components of the crop were improved by optimizing plant population density and nitrogenfertilizerapplication. Numerically the highest grain yield and above ground biomass of 6.7 and 13.09 t ha⁻¹ at Tiroafeta and 3.67 and 9.61 t ha⁻¹ at Sekacokorsawas recorded in response to the seed rate of 175 kg ha⁻¹ respectively. On other hand in response to N fertilizer rate the maximum grain yield of 6.9 t ha⁻¹ and above ground biomass of 14.04 t ha⁻¹ at Tiroafetawas recorded from 92kg N ha⁻¹ and 69 kg N ha⁻¹ respectively. Whereas at Sekacokorsa the highest grain yield of 3.72 t ha⁻¹ and above ground biomass of 9.76 t ha⁻¹ was obtained from nitrogen fertilizer rate of 46kg N ha⁻¹. In conclusion, N fertilizer rate of 46kg N ha⁻¹ and seed rates of 125 kg ha⁻¹ in both locationsgavethe optimum yield. But due to the field experiment was done for one year only it should be repeated over the years to give sound recommendations under similar agro ecologies.

Keywords

Seed Rate, Nitrogen, Yield, Wheat and Fertilizer

1. Introduction

Bread wheat (*Triticumaestivum* L.) is one of the most important cereal crops of the world and is a staple food for about one third of the world's population [1]. It is one of the major cereal crops grown in the highlands up to mid lands and cur-

rently produced in clusters at different area of Ethiopia under irrigation. Wheat ranked 3rd in grain crop area and production after maize (*Zeamays*) and tef (*Eragrostistef*). The total grain crop area was 1.87 million hectare and production of 58.1

*Corresponding author: sis.sis1835@gmail.com (SisayGurmu)

Received: 7 February 2024; Accepted: 23 February 2024; Published: 11 April 2024



Copyright: © The Author(s), 2024. Published by Science Publishing Group. This is an **Open Access** article, distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

million quintal with average productivity of 31.1 quintal per hectare [2]. But its average yield is still very low and below the world's average [3]. The low yield may be due to the use of different or not optimum seed rate, varieties, diseases, low soil fertility and also improper water managements.

Plant density is a major factor determining the ability of the crop to capture resources and generate yield. Also it was the main responsible factors for low yield are less or more than optimum plant population and inadequate crop nutrition application especially nitrogen. It can be developed by using a suitable seeding rate. Growth and yield of wheat are affected by environmental conditions and can be regulated by seeding rate [4]. Maximum genetic potential of high yielding wheat varieties cannot be harvested without ensuring proper seeding rate. As the plant density increases, the competition for resources especially for nitrogen also increases that badly affect the ultimate yield. Provision of additional nitrogen can be hypothesized to further enhance the yield by increasing plant population but up to an optimum level. Further higher nitrogen can lead to the lodging of plants at higher seed rate [5]. Nitrogen occupies a conspicuous place in plant metabolism. Consequently to get more crop production, nitrogen availability is essential in the form of chemical fertilizers. All vital processes in plant are associated with protein, of which nitrogen is an essential constituent. The research work done indicates that appropriate level of nitrogen application has increased wheat yield significantly [6, 7]. Proper use of nitrogen is also considered for farm profitability and environment protection [8].

These all potentially limit crop productivity despite optimum use of nitrogen fertilizer and seed rate. Therefore, in order to set priorities among the different plant nutrients, it is important to identify the status of the limiting nitrogen nutrients in various soils and the seed rate. Thus, this research was carried out to investigate the optimum nitrogen fertilizer rate and seed rate applied for irrigated wheat for better yield and yield components at Jimma South western Ethiopia.

2. Materials and Methods

2.1. Description of the Study Site

The experiment was conducted in Tiroafeta and Sekacokorsa districts of Jimma zone in Oromia National Regional State of Ethiopia under irrigation condition from Mid of December to the end of December 2020. Tiroafeta district is situated at latitude 07°58' N and longitude 36°70' E with an elevation of 1675 meters above sea level. The experimental site reliably receives good rains up to 1955 mm per annum and minimum and maximum temperatures of 11.5 °C and 25 °C, respectively. Whereas Sekacokorsa district is located at latitude 07°30' N and longitude 36°05' E with an elevation of 1750 meters above sea level. The experimental site reliably receives good rains up to 1775 mm per annum and minimum and maximum temperatures of 12.7 °C and 25.7 °C, respectively. The soil of the experimental sites is nitosol type.




Figure 1. Monthly total rainfall and mean maximum and minimum temperatures in December 2020 up to April 2021 during irrigation wheat growing months at Sekacokorsa and Tiroafeta district.

2.2. Treatments and Experimental Design

The treatments were consisting of four rates of N fertilizer:

(0, 50, 100 and 150 and 200 kg urea ha⁻¹ or 0, 23, 46, 69 and 92 kg N ha⁻¹ respectively) and three seed rates (125, 150 and 175 kg ha⁻¹), arranged in a factorial randomized complete block design (RCBD) with three replications. The Gross area

of each plot size was 5 m x 3 m (15m²). There are 15 rows in each plot with 20 cm inter row spacing. The spacing between adjacent plots and replications were 0.5m and 1m respectively.

2.3. Experimental Materials

The bread wheat variety Kakaba was used as planting material. The variety was selected based on its adaptability to agro-ecological conditions and better performance in yield at the area. Kakabais the most common and widely distributed bread wheat variety in the country. The blended NPS fertilizer rates at 100 kg ha⁻¹ (19% of N, 38% of P₂O₅ and 7% of S) were uniformly supplemented for experimental plots except for without fertilizer (negative control plot) at sowing. Urea fertilizer as source of Nitrogen (46% N) was used.

2.4. Experimental Procedures and Field Management

The experimental field was ploughed with tractor and oxen to a fine tith four times and the plots were leveled manually. Then, a field layout was made and each treatment was assigned randomly to the experimental units within a block. Bread wheat seed was sown at the assigned treatment seed rate of rows of 20 cm spacing manually by drilling. The whole amount of blended (NPS) was applied at sowing and the 1/3 and 2/3of nitrogen was applied at the time of sowing and top-dressed at tillering (30 days after sowing) respectively. The spacing between ridge rows, plots and blocks were 50, 100, 200cm respectively. Weeding was done 3 times and harvesting and threshing was done manually. Other necessary cultural practices were applied to all plots uniformly as needed.

2.5. Data Collection

Plant height, number of total tillers, number of productive tillers, spike length (cm), number of kernels per spike, thousand seed weight (g), grain yield (t ha⁻¹), above ground bio-

mass (t ha⁻¹) and harvest index.

2.6. Statistical Analysis

Data were subjected to analysis of variance (ANOVA) by using the methods described by Gomez and Gomez [9] using SAS computer software 9.3. The comparisons among treatment means with significant difference for measured characters were done by LSD test at 5% level of significance.

3. Results and Discussions

The field trials were conducted at TiroAfeta and Sakacokorsa districts. The three seed rates and five Nitrogen rates were tested using bread wheat (kakaba variety) under irrigation on farmer's fields. The expected wheat stand per square meter was optimized. The effect of different rates of N fertilizer and seed rate on yield and yield related parameters of wheat variety were presented and discussed as follows.

The homogeneity test of the error variances for years indicated that the error variance was not homogenous and hence combined analysis of variance was not conducted over the locations. At both district (Sekacokorsa and Tiroafeta) N fertilizer rates and seed rate didn't show significant ($P > 0.05$) interaction effect on all collected growth, yield and yield related parameters. Grain yield and above ground biomass was highly significantly ($P < 0.01$) influenced by N fertilizer rate, but grain yield was not significantly ($P > 0.05$) affected by seed rate at both locations. Above ground biomass was highly significantly ($P < 0.01$) influenced by seed rate at Sekacokorsa district. At Tiroafeta district number of kernels per spike was highly significantly ($P < 0.01$) affected by N fertilizer rate and but not significantly ($P > 0.05$) affected by seed rate at Sekacokorsa. Whereas plant height, number of total tillers, number of productive tillers, spike length, thousand seed weight and harvest index was not significantly ($P > 0.05$) affected by seed rate and N fertilizer rate at both locations (Tables 1 and 2).

Table 1. Mean square of the effects of N fertilizer rates and seed rates on growth, yields attributes and yield of wheat at Sekacokorsaworeda, Jimma zone southwestern Ethiopia under irrigation condition.

Parameter	Seed rate (2)	N rate (4)	Rep (2)	Seed rate x N rate (8)	Error (28)
Plant height (cm)	3.569 ^{ns}	8.845 ^{ns}	40.991 ^{ns}	4.541 ^{ns}	14.1898
No. of total tillers	0.116 ^{ns}	0.304 ^{ns}	15.606 ^{**}	0.514 ^{ns}	0.4546
No. of productive tillers	0.170 ^{ns}	0.069 ^{ns}	11.991 ^{**}	0.427 ^{ns}	0.3795
Spike length (cm)	0.062 ^{ns}	1.0552 ^{ns}	1.154 ^{ns}	0.1901 ^{ns}	0.5571
No. of kernel per spike	7.0202 ^{ns}	18.798 ^{ns}	21.1209 ^{ns}	14.8022 ^{ns}	11.3380
Thousand seed weight (gm)	8.435 ^{ns}	3.497 ^{ns}	8.828 ^{ns}	6.094 ^{ns}	6.1456

Parameter	Seed rate (2)	N rate (4)	Rep (2)	Seed rate x N rate (8)	Error (28)
Grain yield (t ha ⁻¹)	0.1884 ^{ns}	0.3361 ^{**}	0.0341 ^{ns}	0.1126 ^{ns}	0.0925
AGB (t ha ⁻¹)	3.2135 ^{**}	2.8859 ^{**}	4.7894 ^{**}	0.4724 ^{ns}	0.6511
Harvest index	9.523 ^{ns}	1.873 ^{ns}	78.104 ^{**}	7.035 ^{ns}	6.5124

*Numbers in parenthesis = Degrees of freedom; Ns=not significant ($P > 0.05$); * = Significant ($P < 0.05$); ** = highly significant ($p < 0.01$) difference; AGB= Aboveground biomass; ha = Hectare

Table 2. Mean square of the effects of N fertilizer rates and seed rates on growth, yields attributes and yield of wheat at Tiroafetaworeda, Jimma zone southwestern Ethiopia under irrigation condition.

Parameter	Seed rate (2)	N rate (4)	Rep (2)	Seed rate x N rate (8)	Error (28)
Plant height (cm)	0.556 ^{ns}	41.522 ^{ns}	12.289 ^{ns}	51.472 ^{ns}	29.027
No. of total tillers	0.422 ^{ns}	0.557 ^{ns}	14.391 ^{**}	1.490 ^{ns}	1.1140
No. of productive tillers	0.296 ^{ns}	0.867 ^{ns}	9.689 ^{**}	1.124 ^{ns}	0.825
Spike length (cm)	0.800 ^{ns}	0.144 ^{ns}	1.267 [*]	0.244 ^{ns}	0.2905
No. of Kernel per spike	7.990 ^{ns}	108.107 ^{**}	16.042 ^{ns}	13.774 ^{ns}	13.8718
Thousand seed weight (gm)	5.243 ^{ns}	0.520 ^{ns}	0.860 ^{ns}	3.742 ^{ns}	3.3709
Grain yield (t ha ⁻¹)	0.464 ^{ns}	2.984 ^{**}	0.027 ^{ns}	0.246 ^{ns}	0.5780
AGB (t ha ⁻¹)	2.468 ^{ns}	19.669 ^{**}	2.064 ^{ns}	2.848 ^{ns}	4.3971
Harvest index	71.489 ^{ns}	29.649 ^{ns}	41.366 ^{ns}	32.317 ^{ns}	36.1422

*Numbers in parenthesis = Degrees of freedom; Ns=not significant ($P > 0.05$); * = Significant ($P < 0.05$); ** = highly significant ($p < 0.01$) difference; AGB= Above ground biomass; ha = Hectare

3.1. Plant Height

The interaction, seeding rate and nitrogen levels were found non significant (Tables 1 and 2) for both locations. However, comparatively taller plants were observed at seeding rate of 150 kg ha⁻¹ and 175 kg ha⁻¹ at Tiroafeta and Sekacokorsa respectively. Concerning N fertilizer rate taller plants were recorded from 92 kg N ha⁻¹ at both locations (Tables 3 and 5). The research findings are in coordinate with Lafondet *et al.* [10]; Geleta *et al.* [11] (2002). Ali *et al.* [12] studied that maximum plant height was obtained at a seeding rate of 125 kg ha⁻¹. Chandra *et al.* [13] concluded that plant height increased significantly by increasing nitrogen levels.

3.2. Number of Total Tillers

Crop yields are generally dependent upon many yield contributing agents. Among these numbers of tillers are the most important because of the contribution in final yield. The interaction, seeding rate and nitrogen levels were found non significant (Tables 1 and 2) for both locations. But, comparatively more number of tillers was observed at seeding rate of

175 kg ha⁻¹ and N fertilizer rate of 23 kg N ha⁻¹ at Tiroafeta district and at seeding rate of 150 kg ha⁻¹ and N fertilizer rate of 92 kg N ha⁻¹ at Sekacokorsa (Tables 3 and 5). Increase in number of tillers per unit area is due to increased seeding rate [14, 15]. N fertilizer rate also contributed in increasing tiller production up to an optimum level [16, 17]. Above optimum, the decrease in tillers might be due to the competition for space.

3.3. Number of Productive Tillers

The interaction, seeding rate and nitrogen levels were found non significant (Tables 1 and 2) for both locations. However, comparatively the maximum number of productive tillers per plant was observed at seed rate of 150 kg ha⁻¹ at both locations and N fertilizer rate of 69 kg N ha⁻¹ and 46 kg N ha⁻¹ at Tiroafeta and Sekacokorsa respectively (Tables 3 and 5). The current result is in agreement with Adinew [18] and Amare [19] who found that maximum productive tiller was recorded from minimum seed rate and vice versa under rain-fed conditions. On the other hand, Iqbalet *et al.* [20] reported that maximum productive tillers were obtained at 200 kg ha⁻¹ seed rates than at lower seeding rates under rain-fed conditions.

3.4. Spike Length

The interaction, seeding rate and nitrogen levels were found non significant (Tables 1 and 2) for both locations. However, numerically the maximum spike length was observed at seeding rate of 125 kg ha⁻¹ and N fertilizer rate of 69 kg N ha⁻¹ at Tiroafeta district whereas at Sekacokorsa the highest spike length recorded from the seed rate of 175 kg ha⁻¹ and N fertilizer rate of 92 kg N ha⁻¹ (Tables 3 and 5). These results are in agreement with those of Hussain *et al.* [21] and Ahmad *et al.* [14]. They concluded that spike length of wheat was increased significantly with increasing nitrogen levels.

3.5. Number of Kernels Per Spike

Seeding rate did not affect number of kernels per spike at both locations; however, nitrogen fertilization highly significantly affected kernels per spike at Tiroafeta but not significantly affected at Sekacokorsa (Tables 1 and 2). Highest number of kernels per spike (46.6) was observed at nitrogen fertilizer rate of 23 kg N ha⁻¹ followed by 69 kg N ha⁻¹ at Tiroafeta (Table 3). In agreement with this result, Dinkinesh [22] reported the highest number of kernels produced from the highest rate of NPSB fertilizers. In line with this, Seyoum [23] also reported that increasing the rates of N increased the number of kernels produced per spike.

Table 3. Main effect of seed rates on growth and yield components of bread wheat variety at TiroAfetaworeda.

Seed rates (kg ha ⁻¹)	Plant height (cm)	No. of total tillers	No. of productive tillers	Spike Length (cm)	Number of kernels per spike	Thousand seed weight (gm)
125	82.1	4.6	4.0	7.9	42.7	42.7
150	82.5	4.8	4.2	7.5	43.3	43.2
175	82.1	4.9	3.9	7.9	41.8	42.0
LSD (0.05)	4.03	0.79	0.679	0.403	2.786	1.373
Nitrogen Rates (kg ha ⁻¹)						
0	79.2	4.8	3.9	7.6	39.8 ^{cd}	42.4
23	81.4	5.0	4.1	7.8	46.6 ^a	42.8
46	82.7	4.3	3.8	7.8	38.5 ^d	42.9
69	82.8	4.9	4.6	7.9	45.1 ^{ab}	42.6
92	85.1	4.7	3.9	7.7	43.0 ^{bc}	42.4
Mean	82.2	4.74	4.04	7.73	42.6	42.6
LSD (0.05)	5.2	1.02	0.877	0.52	3.6	1.77
CV (%)	6.55	22.3	22.5	6.97	8.74	4.31

3.6. Thousand Seed Weight

The interaction, seeding rate and nitrogen levels were found non significant (Tables 1 and 2) for both locations. However, comparatively the highest thousand seed weight was recorded from 150 kg ha⁻¹ seed rate and N fertilizer rate of 46 kg N ha⁻¹ at Tiroafeta district (Table 3). At Sekacokorsa district the highest thousand seed weight was obtained from seed rate of 175 kg ha⁻¹ and N fertilizer rate of 23 kg N ha⁻¹ (Table 5). The insignificance of N rate might be due to good grain filling period and favorable environmental condition. In line with this Bereket *et al.* [24] also reported that increasing N rate from 92 kg N ha⁻¹ to 138 kg N ha⁻¹ decreased thousand kernels weight.

3.7. Grain Yield

Grain yield of wheat is the function of its unique yield component in response to nitrogen level and seeding rate for the yield of the crop. Grain yield was highly influenced by N fertilizer rate at both locations but not significantly influenced by seed rate and interaction of seed rate and N fertilizer rate (Tables 1 and 2). The higher grain yield (7.10 t ha⁻¹) was obtained from N fertilizer rate of 69 kg N ha⁻¹ which was statically at par with N rate of 92 and 46 kg N ha⁻¹ at Tiroafeta (Table 4). N application of 46 kg N ha⁻¹ increased the grain yield by 19.8% over the control at Tiroafeta district. At Sekacokorsa the highest grain yield of 3.72 t ha⁻¹ was recorded from N fertilizer rate of 46 and 92 kg N ha⁻¹ which

was statically at par with N rate of 69 kg N ha⁻¹ (Table 6). Application of 46 kg N ha⁻¹ increased the grain yield by 11.1% over the control at Sekacokorsa. At both locations the seed rate of 175 kg ha⁻¹ gave highest grain yield even if non significant difference was not seen between different seed rates (Tables 4 and 6). The maximum grain yield might be attributed to the improvement in number of tillers, spike length, and 1000 grain weight. Similar results were obtained by

Pandyet al. [25]. He reported that increasing nitrogen rates increased grain yield. Maqsoodet al. [26] found that application of 150 kg N ha⁻¹ gave the highest grain yield. Nitrogen fertilizer up to 150 kg ha⁻¹ increased the grain yield, whereas higher levels of N fertilizer decreased both yields [27]. N application increased the grain yield by 9.6% in wheat and deficiency depressed the yield in cereals [28].

Table 4. Main effect of seed rates on yield and harvest index of bread wheat variety at TiroAfetaworeda.

Seed rates (kg ha ⁻¹)	Grain Yield (t ha ⁻¹)	Above ground BiomassYield (t ha ⁻¹)	Harvest index
125	6.61	12.28	55.0
150	6.36	12.66	50.8
175	6.70	13.09	51.8
LSD (0.05)	0.569	1.568	4.53
Nitrogen Rates (kg ha ⁻¹)			
0	5.76 ^b	10.71 ^c	53.9
23	6.15 ^b	11.59 ^{bc}	54.5
46	6.90 ^a	13.13 ^{ab}	52.9
69	7.10 ^a	14.04 ^a	51.1
92	6.90 ^a	13.95 ^a	50.2
Mean	6.56	12.68	52.53
LSD (0.05)	0.734	2.025	5.85
CV (%)	11.59	16.5	11.44

3.8. Above Ground Biomass

Above ground biomass represent overall growth performance of the plant as well as the crop and is considered to be the essential yield parameter to get useful information about overall growth of wheat. It is highly inclined by crop nutrition and seed rate. Above ground biomass was highly influenced by N fertilizer rate at both locations but not significantly influenced by seed rate at Tiroafeta and highly influenced by seed rate at Sekacokorsa district (Table 1 and 2). At Tiroafeta the highest above ground biomass (14.04 t ha⁻¹) was obtained at N fertilizer rate of 69 kg ha⁻¹ which was statistically at par with N fertilizer rate of 46 and 92 kg ha⁻¹ while lowest above ground biomass (10.71 t ha⁻¹) was recorded

fromcontrol (Table 4). At Sekacokorsa district the highest above ground biomass (9.76 t ha⁻¹) was obtained at N fertilizer rate of 46 kg ha⁻¹ which was statistically at par with N fertilizer rate of 69 and 92 kg ha⁻¹ while lowest above ground biomass (8.57 t ha⁻¹) was recorded from control (Table 6). Regarding seed rate the highest above ground biomass (9.61t ha⁻¹) was recorded from 175 kg ha⁻¹ which was statistically at par with 150 kg ha⁻¹ seed rate at Sekacokorsa district (Table 6). Increased in biomass production might be attributed to the increased plant population due to higher seeding rate. These results were in agreement with Islam et al. [17], Mojiri and Arzani [27] and Soyluet al. [28]. Ottesonet al. [29] found that biological yield was increased by increasing nitrogen up to optimum levels. Nayyaret al. [30] obtained the highest biological yield at higher seeding rate of 100 kg ha⁻¹.

Table 5. Main effect of seed rates on yield and yield components of bread wheat variety at Sakacokorsaworeda.

Seed rates (kg ha ⁻¹)	Plant height (cm)	Number of total tillers	Number of productive tillers	Spike length (cm)	Number of kernels per spike	Thousand seed weight (gm)
125	82.6	3.4	2.8	9.1	43.5	34.6
150	82.2	3.5	3.0	9.1	42.2	35.9
175	83.2	3.3	2.9	9.2	43.3	36.0
LSD (0.05)	2.82	0.50	0.461	0.56	2.52	1.85
Nitrogen rates (kg ha ⁻¹)						
0	81.8	3.3	2.8	8.8	42.9	35.8
23	82.3	3.5	2.9	8.9	41.6	36.0
46	81.9	3.4	3.0	9.1	42.3	34.5
69	83.3	3.1	2.8	9.1	42.8	35.3
92	84.0	3.6	3.0	9.7	45.4	35.8
Mean	82.7	3.4	2.91	9.1	43.0	35.5
LSD (0.05)	3.64	0.65	0.59	0.72	3.25	2.39
CV (%)	4.6	19.89	21.2	8.2	7.8	6.99

3.9. Harvest Index

The ability of a cultivar to convert the dry matter into economic yield is indicated by its harvest index. The higher the harvest index value, the greater the physiological potential of the crop for the converting dry matter to grain yield. The interaction, seeding rate and nitrogen levels were found non significant (Tables 1 and 2) for both locations. However, comparatively the highest harvest index was recorded from 125 kg ha⁻¹ seed rate and N fertilizer rate of 23 kg ha⁻¹ at Tiroafeta district (Table 4). At Sekacokorsa district the highest harvest index was obtained from seed rate of 125 kg ha⁻¹ and N fertilizer rate of 92 kg ha⁻¹ (Table 6). Increase in nitrogen level increased harvest index [16].

Table 6. Main effect of seed rates on yield and yield components of bread wheat variety at Sakacokorsaworeda.

Seed rates (kg ha ⁻¹)	GrainYield (t ha ⁻¹)	Above ground Biomass (t ha ⁻¹)	Harvest index
125	3.44	8.72 ^b	39.6
150	3.57	9.41 ^a	38.2
175	3.67	9.61 ^a	38.3
LSD (0.05)	0.23	0.60	1.92
Nitrogen rates (kg ha ⁻¹)			
0	3.35 ^b	8.57 ^b	39.1
23	3.36 ^b	8.71 ^b	38.8
46	3.72 ^a	9.76 ^a	38.1
69	3.68 ^a	9.68 ^a	38.3
92	3.72 ^a	9.52 ^a	39.2
Mean	3.56	9.25	38.7
LSD (0.05)	0.29	0.78	2.48
CV (%)	8.5	8.7	6.59

4. Conclusions

From the present study, we observed that the grain yield and yield components of bread wheat that were planted under irrigation were highly improved through optimizing nitrogen application and plant population. At both district (SekaCokorsa and Tiroafeta) N fertilizer rates and seed rate didn't show significant interaction effect on all collected growth, yield and yield related parameters. Grain yield and above ground biomass was highly significantly influenced by N fertilizer rate, but grain yield was not significantly affected by seed rate at both locations. Above ground biomass was highly significantly influenced by seed rate at Sekacokorsa district. At Tiroafeta district number of kernels per spike was highly significantly affected by N fertilizer rate and but not significantly affected by seed rate at Sekacokorsa. The higher grain yield (7.10 t ha^{-1}) was obtained from N fertilizer rate of 69 kg N ha^{-1} which was statically at par with N rate of 92 and 46 kg N ha^{-1} at Tiroafeta whereas at Sekacokorsa the highest grain yield of 3.72 t ha^{-1} was recorded from N fertilizer rate of 46 and 92 kg N ha^{-1} which was statically at par with N rate of 69 kg N ha^{-1} . In conclusion, N fertilizer rate of 46 kg N ha^{-1} and seed rates of 125 kg ha^{-1} in both locations gave the optimum grain yield. As a conclusion, since this study was conducted in one season, it should be repeated in more seasons for further recommendation under similar agro ecologies.

Abbreviations

AGB: Above Ground Biomass
 ANOVA: Analysis of Variance
 CV: Coefficient of Variation
 Ha: Hectare
 LSD: Least Significant difference
 NS: Not Significant
 RCBD: Randomized Complete Block Design

Acknowledgments

Of all, the authors greatly acknowledge irrigation wheat project, Ethiopia Institute of Agricultural Research, Jimma Agricultural Research Center for its financial support and immense role in administrative support incredibly appreciated. The authors would also like to extend their gratitude to all staff members of the cereal agronomy research team for their assistance in setting up, planting, collecting the data, and maintaining the field experiments.

Conflicts of Interest

The authors declare no conflicts of interest.

References

- [1] Hussain, M. I., Shah, S. H., Hussain, S. A. J. J. A. D. and Iqbal, K. H. A. L. I. D., 2002. Growth, yield and quality response of three wheat (*Triticumaestivum* L.) varieties to different levels of N, P and K. *International Journal of Agriculture and Biology*, 4(3), pp. 362-364.
- [2] CSA (Central Statistic Agency), 2021. Agricultural sample survey report on area and production of major crops for the period 2020/2021 cropping season. Vol 5. Statistical bulletin 578, Addis Ababa; 2021.
- [3] FAOSTAT, 2022. Online <http://faostat.fao.org/site/291/default.aspx>. Published 2020/1
- [4] Ozturk, A., O. Caglar and S. Bulut, 2005. Growth and yield response of facultative wheat to winter sowing, freezing sowing and spring sowing at different seeding rates. *J. Agron. Crop Sci.*, 192: 10-16.
- [5] Nazir, M. S., A. Jabbar, Z. Waheed, A. Gaffar and M. Aslam, 2000. Response of late sown wheat to seeding density and nitrogen management. *Pak. J. Biol. Sci.*, 3: 998-1001.
- [6] Gwal, H. B., R. J. Tiwari, R. C. Jain and F. S. Prajapati, 1999. Effect of different levels of fertilizer on growth, yield and quality of late sown wheat. *RACHIS Newsletter*, 18: 42-44.
- [7] Ali, A., M. A. Choudhry, M. A. Malik, R. Ahmad and Saifullah, 2000. Effect of various doses of nitrogen on the growth and yield of two wheat cultivars. *Pak. J. Biol. Sci.*, 3: 1004-1005.
- [8] Makowski, D., D. Wallach and J. M. Meynard, 1999. Model of yield, grain protein and residual mineral nitrogen responses to applied nitrogen for winter wheat. *Agron. J.*, 91: 337-385.
- [9] Gomez, K. A. and Gomez, A. A., 1984. *Statistical Procedures for Agricultural Research*. 2nd ed. John Wiley and Sons. New York.
- [10] Lafond, G. P., 1994. Effects of row spacing, seeding rate and nitrogen on yield of barley and wheat under zero-till management. *Can. J. Plant Sci.*, 74: 703- 711. [Wheat, Barley and Triticale Absts., 1(1): 45; 1995].
- [11] Geleta, B., M. Atak, P. S. Baenziger, L. A. Nelson, D. D. Baltenesperger, K. M. Eskridge, M. J. Shipman and D. R. Shelton, 2002. Seeding rate and genotype effect on agronomic performance and end-use quality of winter wheat. *Crop Sci.*, 42: 827-832.
- [12] Ali, M. A., R. Rehman, M. Siddique and M. Rashid, 1996. Effect of seeding rate and row spacing on wheat yield. *J. Agric. Res.*, 34: 351-357.
- [13] Chandra, S., M. L. Varshncy, J. Singh and S. K. Singh, 1992. Response of wheat varieties to different levels of nitrogen. Narendra Deva. *J. Agric. Res.*, 7(1): 169-171. [Wheat, Barley and Triticale Absts., 11: 4321; 1994].

- [14] Ahmad, M. M., N. Yousaf and M. S. Zamir, 2000. Response of wheat growth, yield and quality to varying application of nitrogen and phosphorous. *J. Agric. Res.*, 38: 289-29.
- [15] Khan, H., M. A. Khan, I. Hussain, M. Z. Khan and M. K. Khattak, 2000. Effect of sowing method and seeding rates on grain yield and yield components of wheat variety Pak-81. *Pak. J. Biol. Sci.*, 3: 1177-1179.
- [16] Singh, C. B., J. Kumar, A. A. Khan, R. A. Katiyar and A. K. Katiyar, 2002. Effect of nitrogen and dates of sowing on yield and quality of wheat (*Triticumaestivum*L.) seeds. *Progressive Agric.*, 2: 92-93.
- [17] Islam, Z. U., S. Khan, J. Bakhat and W. A. Shah, 2002. Frequency of various nitrogen levels, lodging and seed quality in wheat. *Asian J. Plant Sci.*, 1: 510-512.
- [18] Adinew A., 2019. "Effect of seed source and rates on productivity of bread wheat varieties at Kersa, Eastern Ethiopia," *Journal of Biology, Agriculture and Healthcare*, vol. 9, no. 3, pp. 1-9.
- [19] Amare A., 2017. "Response of bread wheat varieties to different seeding rates for growth, yield and yield components in Kombolcha district, North-Eastern Ethiopia," *Journal of Biology, Agriculture and Healthcare*, vol. 7, no. 23, pp. 79-91.
- [20] Iqbal N., Akbar N., Ali M., Satter M., and Ali L., 2010, "Effect of seed rate and row spacing on yield and yield components of wheat," *Pakistan Journal of Agricultural Research*, vol. 48, no. 2, pp. 151-156.
- [21] Hussain, S., A. Sajjad, M. I. Hussain and M. Saleem, 2001. Growth and yield response of three wheat varieties to different seeding densities. *Int J. Agric. Biol.*, 3: 228-229.
- [22] DinkineshAbera, 2018. Effects of Blended Fertilizer (NPSB) Rates on Yield, Yield Components and Grain Quality of Durum Wheat (*Triticumturgidum* L. Var. Durum) Varieties in MinijarShenkora District, Central Ethiopia. A Thesis Submitted to the School of Plant Sciences Post Graduate Program Directorate Haramaya University.
- [23] SeyoumAlemu, 2017. Effect of blended NPS and N fertilizer rates on yield components, yield and grain protein content of bread wheat (*Triticumaestivum* L.) in Bore district, Guji Zone, Southern Ethiopia. MSc Thesis, Alemaya University, Alemaya, Ethiopia.
- [24] Bereket Haileselassie, Dawit Habte, Mehretab Haileselassie, and Gebremedhin Gebremeskel, 2014. Effects of Mineral Nitrogen and Phosphorus Fertilizers on Yield and Nutrient Utilization of Bread Wheat (*Triticumaestivum*) on the Sandy Soils of Hawzen District, Northern Ethiopia. *Agriculture, Forestry and Fisheries*, 3(3): 189-98.
- [25] Pandey, R. K., J. W. Maranvilla and A. Admou, 2001. Response of durum wheat genotypes to nitrogenous fertilizers. *Rastenievdni-Nauki*, 38: 203-207.
- [26] Maqsood, M., A. Ali, Z. Aslam, M. Saeed and S. Aslam, 2002. Effect of irrigation and nitrogen levels on grain yield and quality of wheat. *Int. J. Agric. Biol.*, 4: 164-165.
- [27] Mojiri, A. and Arzani, A., 2003. Effects of nitrogen rate and plant density on yield and yield components of sunflower. *J. Sci. Technol. Agric. Nat. Resour.*, 7: 115-125.
- [28] Soylu, S., Sade B., Topal A., Akgun N., and Gezgin, S., 2005. Response of irrigated durum and bread wheat cultivar to boron application in a low boron calcareous soil. *Turk. J. Agri.*, 29: 275-286.
- [29] Otteson, B. N., M. Mergoum and J. K. Ransom, 2007. Seeding rate and nitrogen management effects on spring wheat yield and yield components. *Agron. J.*, 99: 1615-1621.
- [30] Nayyar, M. M., Asghar M. and Tahir, M. J., 1992. Effect of sowing date and seeding rate on wheat grain yield under irrigated conditions. *J. Agric. Res.*, 30: 329- 333.