

A Nutrient Combination That Can Affect Yield of *Olitorius Jute*

Md Saheb Ali¹, Md Marfudul Hoque^{1,2}, Md Abdul Alim¹, Md Mahbubul Islam^{1,3,*}

¹Agronomy Division, Bangladesh Jute Research Institute, Dhaka, Bangladesh

²Department of Agriculture Extension Kaptai, Rangamati, Bangladesh

³PTC Division, Bangladesh Jute Research Institute, Dhaka, Bangladesh

Email address:

mahbub_agronomy@yahoo.com (Md M. Islam)

*Corresponding author

To cite this article:

Md Saheb Ali, Md Marfudul Hoque, Md Abdul Alim, Md Mahbubul Islam. A Nutrient Combination That Can Affect Yield of *Olitorius Jute*. *Plant*. Vol. 7, No. 3, 2019, pp. 42-46. doi: 10.11648/j.plant.20190703.13

Received: July 26, 2019; Accepted: August 19, 2019; Published: August 29, 2019

Abstract: Mineral fertilizer use is expanding in Jute growing countries to satisfy fiber demands. Fertilizer consumers are being asked to improve fertilizer use efficiency through better management in their fields to provide a healthy economy. Following the thought a field experiment was established to determine the influence of mineral fertilizer (NPKS) on growth and yield of the advanced *C. olitorius* breeding line of O-043-7-9. We selected three different locations under Bangladesh Jute Research Institute to examine jute crop response to optimum fertilizer combination and the yield variability by 10 different nutrient combinations. Because of the essentiality of N, P, K, and S fertilizer combination on Jute fibre production and their effects on its growth were studied. The lower levels of N, P, K and S were zero applications in specific treatments. N, P, K and S were positively correlated with yield level. Over time, the simple effect of N exhibited an increased positive trend only when applied limited to 100kg/ha, while simple effects of P and K increased sharply at the rate 10 and 30 kg/ha respectively. The zero fertilizer treatment yielded decline for each mineral in specific treatment in the trial. Fertilizer N, P, K and S were utilized more efficiently in fully balanced combination. After the findings from three different locations for growth, yield and agronomic characteristics, we can draw the conclusion that the combination dose of N₁₀₀ P₁₀ K₃₀ S₁₅ Kg/ha may need to be applied for breeding line O-043-7-9 cultivation in Bangladesh.

Keywords: Yield, Jute, Nitrogen, Potassium, Phosphorus, Sulfur

1. Introduction

Among the total jute fibre production in the world, ninety two percent produced in Bangladesh and India. Raw jute signifies fibre produced from, *Corchorus capsularis*, *Corchorus olitorius*, *Hibiscus sabdariffa* and *Hibiscus cannabinus*. Renowned jute growing areas in Bangladesh are Faridpur, Jessore, Manikganj, Kishoregan, Mymensingh and Rangpur regions. *Corchorus olitorius* is one of the most popular fibre crops in every districts of Bangladesh. *Corchorus olitorius* is mainly known for its fibre product, jute and for its leafy vegetables [1]. Several species of *Corchorus* are used as vegetable, of which *Corchorus olitorius* is most frequently cultivated. *C. olitorius* is consumed as a healthy vegetable in Japan, because it contains

abundant carotenoids, vitamin B1, B2, C and E, and minerals. On the other hand, accidental death of cattle has occurred when the cattle were feed with vegetation containing the seeds, because the seeds contain cardiac glycoside [2]. The dark-green leaves of *C. olitorius* have varying proportion of Ca, Fe, -carotene, vitamin C, fibre and protein required for health [1, 3]. Nutrition is an important aspect of Jute production system and this includes adequate supply of essential nutrients like nitrogen, phosphorus, potassium, sulfur etc to the plant.

The availability of these nutrients to plant contributes a lot to its growth and yield. Therefore adequate amount of nutrients need to be supplied to plant at the right quantity and also at the right time to favour both growth and yield. NPKS have influence on the growth and yield of fibre crops [4-40]. Fertilizer studies showed positive responses of *Corchorus*

olitorius to nitrogen [41]. Phosphorus is important in root development and helps hasten maturity of the fruit. Soils require extraneous inputs of phosphorus for optimum *Corchorus olitorius* growth [42]. The role of nitrogen and phosphorus in crop fertilization, leading to increased absorption of elements can be attributed to increase top growth particularly as a result of nitrogen absorption [43]. Similarly, application of any essential element should have a marked effect on yield if the soils were deficient in the element. Therefore, it is necessary to determine the growth and yield performance of the advance olitorius line O-043-7-9 by nutrient combination.

2. Methods

The experiment was conducted at three locations of Bangladesh Jute Research Institute to determine the nutritional requirement for optimum growth and yield of the advanced *C. olitorius* breeding line of O-043-7-9. The experiment was laid out in randomized complete block design with three replications. A total of 10 treatment combinations along with a control were distributed randomly in each equal plot as one replication. Each replication was divided into 10 unit plots. At the time of final land preparation, the land was well prepared and fertilizers were applied as per treatment (Table 1). NPKS fertilizers were applied in the form of urea, TSP, MAP and gypsum respectively. Half of urea was applied at sowing and the rest half was top dressed at 45 days after sowing while all other fertilizers were applied at the time of sowing. At harvesting time, ten plants were selected at random from each plot and tagged in the field to make a note of yield and yield contributing parameters.

Table 1. Treatment combinations (NPKS in Kg/ha).

T1: N0P10K30S20	T6: N100P15K30S20
T2: N75P10K30S20	T7: N100P10K0S20
T3: N100P10K30S20	T8: N100P10K45S20
T4: N125P10K30S20	T9: N100P10K30S0
T5: N100P0K30S20	T10: N100P10K30S15

3. Results and Discussions

Our research findings showed that among the ten nutrient combinations of chemical fertilizer, nutrient combination for treatment T₁₀ performed best for the advanced breeding line O-043-7-9 yield and yield influencing factors. The plant growth and yield were affected by N levels over the control. Nitrogen @ 100 kg/ha treatment (T₁₀) produced significant highest fibre yield. But highest dose of N 125 kg/ha showed significantly lower yield than the dose N 100 kg/ha. The growth indicator i.e. plant height and base diameter were found significantly highest by N 100 kg/ha (T₁₀) that was identical with the treatments T₃, T₄, T₆, T₇ and T₈. From the results, it was observed that 100 kg N/ha may be an adequate amount for fibre production for the advance breeding line O-043-7-9. The different results noticed that application of higher dose of fertilizer consequenced on growth and yield of

fibre production of the advance olitorius line O-043-7-9 (Figures 1-4).




Figure 1. Plant height of olitorius breeding line O-043-7-9 using different chemical fertilizer treatments. Each datum was calculated from three independent experiments. The results are expressed as the mean \pm S.E.M. * $p < 0.05$ significance by the Student's *t*-test.




Figure 2. Base diameter of olitorius breeding line O-043-7-9 using different chemical fertilizer treatments. Each datum was calculated from three independent experiments. The results are expressed as the mean \pm S.E.M. * $p < 0.05$ significance by the Student's *t*-test.




Figure 3. Fibre yield of olitorius breeding line O-043-7-9 using different chemical fertilizer treatments. Each datum was calculated from three independent experiments. The results are expressed as the mean \pm S.E.M. * $p < 0.05$ significance by the Student's *t*-test.




Figure 4. Stick yield of olitorius breeding line O-043-7-9 using different chemical fertilizer treatments. Each datum was calculated from three independent experiments. The results are expressed as the mean \pm S.E.M. * $p < 0.05$ significance by the Student's *t*-test.

The yield and yield influencing factors of O-043-7-9 were affected due to addition of different phosphorus levels. The dose of P 10 kg/ha contributed maximum yield of fibre (2.78t/ha) and stick (6.67 t/ha) at Manikganj (Figure 3 & Figure 4). Results also explored that the longest plant (3.14m) and base diameter (18.39mm) were found significantly higher with 10 kg P/ha at Kishoreganj (Figure 1 & Figure 2).

To observe potassium (K) requirement, there was applied three rates of K, such as 0, 30, and 45 kg/ha. The rate 30 kg K/ha gave the statistically highest plant height (3.08m), base diameter (16.56mm) fibre yield (2.69 t/ha) and stick yield (6.72 t/ha) at Dinajpur. But the supreme rate of K 45 kg /ha yielded lower fibre yield (2.52 t/ha) compare to K 30 kg/ha (Figure 3). Study noticed that the advance olitorus breeding line O-043-7-9, needs K 30 kg/ha to produce maximum yield.



Figure 5. Intensive supervision signifying in experimental field for good agronomic practices by Dr. Md. Mahbubul Islam, Chief Scientific Officer & Head of Agronomy Division, Bangladesh Jute Research Institute, Manik Mia Avenue, Dhaka-1207, Bangladesh.

Highest plant height (3.14m), base diameter (18.39mm) fibre yield (2.93t/ha) and stick yield (6.31t/ha) were observed with 15 kg S/ha (Figures 1-4). Results showed that the combined dose of NPK and S 100-10-30-15 kg/ha may be a suitable dose for the cultivation of advance breeding line O-

043-7-9. The nutrient combination and their ratio is one of the key factors for olitorus breeding line O-043-7-9 production that is supported by previous report [4, 32, 33]. This outcome is owing to competent direction and time to time suggestion about experimental events (Figure 5)

4. Conclusion

The overall treatments had significant positive impact over nitrogen control (T_1) on growth and yield and effect of combined fertilizers on them were explained. The most important parameter, fibre yield (2.93 t/ha) and stick yield (6.72 t/ha), were recorded highest with $T_{10}:N_{100}P_{10}K_{30}S_{15}$ treatment. Considering all these aspects, specially yield, $T_{10}:N_{100}P_{10}K_{30}S_{15}$ kg/ha treatment seems to be the best combination for the yield potential of the advance breeding line O-043-7-9.

Author Contributions

All authors contributed equally.

Conflict of Interest Statement

The authors declare that they have no competing interests.

Acknowledgements

Thanks to Jute Research Stations of BJRI and Ministry of Agriculture, Bangladesh is acknowledged.

References

- [1] Schippers RR (2000). African indigenous vegetables: An overview of the cultivated species. University of Greenwich. England 193-205.
- [2] Matsufuji H, Sakai S, Chino M, Goda Y, Toyoda M, et al. (2001). Relationship between cardiac glycoside contents and colour of *Corchorus olitorius* seeds. *J Health Sci* 47: 89-93.
- [3] Adebajo A, Shopeju E (1993). Sources of mycoflora associated with some sundried vegetables in storage. *Int Biodeterior Biodegrad* 31: 255-263.
- [4] Ali MS, Gani MN, Islam MM (2017). Efficiency of BJRI Kenaf-4 Yield Under Different Fertilizer Levels. *A J Agric For* 5: 145-149.
- [5] Ali MS, Hossen M, Ahmed B, Gani MN, Islam MM (2017). Jute Seed Yield Response to Irrigation and Nitrogen Fertilization in Field-Grown Environment. *Int J Biol Environ Eng* 2: 9-13.
- [6] Akanbi WB, Olaniyan AB, Togun AO, Ilupeju AEO, Olaniran OA (2009). The effect of organic and inorganic fertilizer on growth, calyx yield and quality of Roselle (*Hibiscus sabdariffa* L.). *Am Eurasian J Sustain Agric* 3: 652-657.
- [7] Sakimin SZ, Patrie SS, Juraimi AS, Alam MA, Aslani F (2017). Growth performance of roselle (*hibiscus sabdariffa* L.) in response to oil palm by-product media and controlled release fertilizer. *Bangladesh J Bot* 46: 427-431.

- [8] Sajid M, Rab A, Jan I, Iqbal A, Shah F, et al. (2017). Response of various growth parameters of roselle (*Hibiscus sabdariffa* L.) to organic and inorganic phosphorus. *Fresenius environmental bulletin* 26: 878-884.
- [9] Atta A, Diallo AB, Bakasso Y, Sarr B, Saadou M, et al. (2010). Microelement contents of roselle (*Hibiscus sabdariffa* L.) at different growth stages. *Afr J Food Agric Nutr Dev* 10.
- [10] Haruna IM, Maunde SM, Yahuza S (2011). Growth and calyx yield of Roselle (*HIBISCUS SABDARIFFA* L.) as affected by poultry manure and nitrogen fertilizer rates in the southern guinea savanna of Nigeria. *Can J Pure Appl Sci* 5: 1345-1348.
- [11] Javadzadeh SM, Pour MS (2017). Morpho-agronomic characteristics of two Roselle varieties (*Hibiscus sabdariffa* L.) in tropical Iran. *Int J Adv Res Biol Sci* 4: 99-104.
- [12] Pushpa K, Murthy NK, Murthy RK, Sharanappa, Vishwanath AP (2016). Nutrient uptake in mesta (*Hibiscus cannabinus*) as influenced by varieties, spacing and nutrient sources. *Environ Ecol* 34: 489-493.
- [13] Khalil SE, Yousef RM (2014). Study the effect of irrigation water regime and fertilizers on growth, yield and some fruit quality of *Hibiscus sabdariffa* L. *Int J Adv Res* 2: 738-750.
- [14] Ibrahim EB, Abdalla AWH, Ibrahim EA, Naim AME (2013). Interrelationships between yield and its components in some Roselle (*Hibiscus sabdariffa* L.) Genotypes. *World J Agric Res* 1: 114-118.
- [15] Dahmardeh M (2012). Effect of mineral and organic fertilizers on the growth and calyx yield of roselle (*Hibiscus sabdariffa* L.). *Afr J Biotechnol* 11: 10899-10902.
- [16] Musa A, Ogbadoyi EO (2011). Effect of Nitrogen Fertilizer on the Level of Some Nutrients, Anti-Nutrients and Toxic Substances in *Hibiscus sabdariffa*. *Asian J Crop Sci* 4: 103-112.
- [17] Abbas MK, Ali AS (2011). Effect of Foliar Application of NPK on some Growth Characters of Two Cultivars of Roselle (*Hibiscus sabdariffa* L.). *Am J Plant Physiol* 6: 220-227.
- [18] Atta S, Seyni HH, Bakasso Y, Sarr B, Lona I, et al. (2011). Yield character variability in Roselle (*Hibiscus sabdariffa* L.). *Afr J Agric Res* 6: 1371-1377.
- [19] Gad N (2011). Productivity of Roselle (*Hibiscus sabdariffa* L.) Plant as affected by Cobalt and Organic Fertilizers. *J Appl Sci Res* 7: 1785-1792.
- [20] Haruna IM, Ibrahim HY, Rahman SA (2011). The yield and profitability of roselle (*Hibiscus sabdariffa* L.) at varying poultry manure and nitrogen fertilizer rates in the southern guinea savanna of Nigeria. *J Agric Technol* 7: 605-609.
- [21] Ahmed YM, Shalaby EA, Shanan NT (2011). The use of organic and inorganic cultures in improving vegetative growth, yield characters and antioxidant activity of roselle plants (*Hibiscus sabdariffa* L.). *Afr J Biotechnol* 10: 1988-1996.
- [22] Oyewole CI, Mera M (2010). Response of roselle (*Hibiscus sabdariffa* L.) to rates of inorganic and farmyard fertilizers in the Sudan savanna ecological zone of Nigeria. *Afr J Agric Res* 5: 2305-2309.
- [23] Atta S, Sarr B, Bakasso Y, Diallo AB, Lona I, et al. (2010). Roselle (*Hibiscus Sabdariffa* L.) yield and yield components in response to nitrogen fertilization in Niger. *Indian J Agric Res* 44: 96-103.
- [24] Haruna IM, Ibrahim H, Rahman SA (2009). The yield and profitability of Roselle (*Hibiscus sabdariffa* L.) at varying poultry manure and nitrogen fertilizer rates in the Southern Guinea Savanna of Nigeria. *Electronic J Environ Agric Food Chem* 8: 1136-1139.
- [25] Hassan FAS (2009). Response of *Hibiscus sabdariffa* L. plant to some biofertilization treatments. *Ann Agric Sci* 54: 437-446.
- [26] Alam AKMM, Khandker S, Gani MN, Ahmed SA (2000). Uptake addition and balance of nutrients under integrated fertilizer management in jute based cropping patterns. *B J Sci Tech* 2: 147-153.
- [27] Sarker AK, Bandopadhyay PK (2000). Effect of potassium, boron and crop age on the yield and quality of white jute (*Corchorus capsularis*). *Indian Agric* 26: 212-216.
- [28] Das K, Guha B, Pathak D (1996). Response of *Capsularis* jute to potassium fertilization. *Ann Agric Res* 17: 188-189.
- [29] Sarkar SK, Ghosh RK, Sounda G, Maitra S, Rux DK, et al. (1997). Effect of levels of nitrogen, potassium and soil moisture tension on growth, nutrient uptake and water use efficiency of jute. *J Interacademia* 1: 183-188.
- [30] Zheng ZH, Huang YX, Peng XJ (1984). Effects of additional K application to jute. *China's Fiber Crops* 2: 14-15.
- [31] Chew WY, Malek MAA, Ramli K (1982). Nitrogen and potassium fertilization of Congo jute (*Urena lobata*) and kenaf (*Hibiscus cannabinus*) on Malaysian peat. *MARDI-Res. Bulletin* 10: 317-322.
- [32] Islam MM, Alam ATMM (2012). Agronomic Research and Advancement in Jute Crop in Bangladesh, (Sub-Theme Paper). *Souvenir, 11th Conference on Advances in Agronomic Research under Changing Environment in Bangladesh*. At Bangladesh Agricultural Research Institute, Joydebpur, Gazipur-1701, Bangladesh. 101.
- [33] Islam MM, Rahman MM (2008). Hand book on agricultural Technologies of Jute, Kenaf and Mesta crops. Bangladesh Jute Research Institute, Manikmia Avenue, Dhaka-1207, Bangladesh.
- [34] Islam MM (2013). Jute of Bangladesh: Its contribution to environment and challenges. The Reflector, Reflecting the issues and development in societies and states. *Md. Eanyet Ullah, Bangladesh* 2: 8-10.
- [35] Islam M M (2019). Varietal Advances of Jute, Kenaf and Mesta crops in Bangladesh: A review. *Int J Bioorganic Chem*, 4 (1), 24-41.
- [36] Gani MN, Ali MS (2018). Md. Mahbubul Islam. Nutritional Requirements of Newly Released Tossa Jute MG-1. *American Journal of Food, Nutrition and Health*, 3 (2), 31-34.
- [37] Islam MM, Ali MS (2017). Economic Importance of Jute in Bangladesh: Production, Research Achievements and Diversification. *International Journal of Economic Theory and Application*, 4 (6), 45-57.
- [38] Ali MS, Gani MN, Islam MM (2017). Nutrient Management on Growth and Yield of BJRI Tossa Pat 6. *Nutr Food Technol Open Access*, 3 (3).
- [39] Ali MS, Hoque MM, Gani MN, Islam MM (2017). Variation in Inorganic Fertilizer Is an Important Regulator of Yield Potential in BJRI Mesta-3. *American Journal of Environmental Engineering and Science*, 4 (6), 78-84.

- [40] Gani MN, Ali MS, Islam MM (2017). Nutrient Requirement of NPK&S on Advance Capsularis Breeding Line BJC-5105. *Biosystems Engineering*, 2 (5), 44-47.
- [41] NIHORT (1986). *Advances in fruit and vegetable research at NIHORT*. A commemorative publication, Ibadan, Nigeria.
- [42] Zapata F, Axman H (1995). ³²P Isotopic techniques for evaluating the agronomy effectiveness of rock phosphate materials. *Fertilizer Research* 41: 189-195.
- [43] Olaniyi JO (2000). Growth, seed yield and nutrient composition of egusimelon cultivars as affected by NPK 15-15-15 fertilization. *Journal of Agricultural Biotechnology and Environment* 2: 11-16.